1 Correspondence: Dr. Alan M. Jones

2 Address: Department of Biology

3 The University of North Carolina at Chapel Hill

4 Coker Hall, CB#3280

5

6 Phone : (919) 962-6932

7 Fax: (919) 962-1625

8 E-mail: alan_jones@unc.edu

9

10 The author responsible for distribution of materials integral to the findings presented in this article is Dr.

11 Alan Jones alan_jones@unc.edu

12 **Running Title: Atypical Heterotrimeric G Proteins**

13 **Keywords:** Arabidopsis, AGB1, G $\beta\gamma$, AtGPA1, AtRGS1, Extra-large G protein, XLG2, dominant
14 negative G alpha

15

A decoy heterotrimeric Gα protein has substantially reduced nucleotide binding but retains nucleotide-independent interactions with its cognate RGS protein and Gβγ dimer

Fei Lou^a, Tigran M. Abramyan^b, Haiyan Jia^a, Alexander Tropsha^b, and Alan M. Jones^{a,c*}

Departments of ^aBiology and ^cPharmacology, University of North Carolina at Chapel Hill, NC, U.S.A.

^bDivision of Chemical Biology and Medicinal Chemistry, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, NC, USA.

*Correspondence: Dr. Alan M. Jones email, alan_jones@unc.edu

ABSTRACT

Plants uniquely have a family of proteins called extra-large G proteins (XLG) that share homology in their C-terminal half with the canonical Gα subunits; we carefully detail here that Arabidopsis XLG2 lacks critical residues requisite for nucleotide binding and hydrolysis which is consistent with our quantitative analyses. Based on microscale thermophoresis, Arabidopsis XLG2 binds GTPγS with an affinity 100-1000 times lower than that to canonical Gα subunits. This means that given the concentration range of guanine nucleotide in plant cells, XLG2 is not likely bound by GTP *in vivo*. Homology modeling and molecular dynamics simulations provide a plausible mechanism for the poor nucleotide binding affinity of XLG2. Simulations indicate substantially stronger salt bridge networks formed by several key amino-acid residues of AtGPA1 which are either misplaced or missing in XLG2. These residues in AtGPA1 not only maintain the overall shape and integrity of the apoprotein cavity but also increase the frequency of favorable nucleotide-protein interactions in the nucleotide-bound state. Despite this loss of nucleotide dependency, XLG2 binds the RGS domain of AtRGS1 with an affinity similar to the Arabidopsis AtGPA1 in its apo-state and about 2 times lower than AtGPA1 in its transition state. In addition, XLG2 binds the Gβγ dimer with an affinity similar to that of AtGPA1. XLG2 likely acts as a dominant negative Gα protein to block G protein signaling. We propose that XLG2, independent of guanine nucleotide binding, regulates the active state of the canonical G protein pathway directly by sequestering Gβγ and indirectly by promoting heterodimer formation.

INTRODUCTION

The canonical heterotrimeric guanosine nucleotide-binding protein complex, consisting of $G\alpha$, $G\beta$ and $G\gamma$ subunits, serves as a molecular on-off switch in the cell. The inactive or “off-state” form consists of the guanosine diphosphate (GDP) bound to the $G\alpha$ subunit in complex with the $G\beta\gamma$ dimer. For the active or “on-state”, exchange of GDP for GTP in $G\alpha$, either spontaneously or catalyzed by a guanine nucleotide exchange factor, changes the $G\alpha$ conformation leading to dissociation, partly or entirely (1), from the $G\beta\gamma$ dimer and thus enabling both $G\alpha$ and $G\beta\gamma$ to propagate signaling to downstream components (2-4). Signaling is terminated when the $G\alpha$ subunit hydrolyzes GTP thus returning to the inactive GDP-bound state. The rate of GTP hydrolysis is an intrinsic property of each $G\alpha$ subunit but it can be accelerated by Regulator of G protein Signaling (RGS) proteins (5, 6). The $G\alpha$ structure required for nucleotide binding and hydrolysis and for interaction with $G\alpha$ - $G\beta\gamma$ and $G\alpha$ -RGS interactions are well understood (7, 8).

In humans, there are multiple genes encoding G protein subunits resulting in 23 $G\alpha$, 5 $G\beta$ and 12 $G\gamma$ subunits. The $G\alpha$ subunits are divided into four subclasses (G_s , G_i , G_q and $G_{12/13}$) based on function and sequence similarity. However, in Arabidopsis, there is only one canonical $G\alpha$ (AtGPA1) which approximates the sequence of the ancestral $G\alpha$ subunit that evolved into these four animal $G\alpha$ subclasses (9). AtGPA1 has a near identical structure to that of human $G_{i\alpha 1}$ (10). In addition to the canonical $G\alpha$ subunit AtGPA1, the Arabidopsis genome encodes three atypical Extra-large G proteins (XLG1, XLG2, and XLG3) (11). The other components of the Arabidopsis G protein core are a $G\beta$ subunit (AGB1) (12), one of three $G\gamma$ subunits (AGG1, AGG2, and AGG3) (13), and one receptor-like RGS protein (AtRGS1) (14).

The presence of these atypical G proteins makes G protein signaling in plants unique and paradoxical (11, 15, 16). Specifically, the N-terminal half of XLG proteins lacks homology to any characterized domain but contains a putative nuclear localization signal and a cysteine-rich region while

the C-terminal half of XLG proteins shares homology (i.e. evolutionary history, (16)) with canonical Gα subunits (~30% identity). However, there is controversy to what extent that these atypical Gα homologs bind and hydrolyze nucleotides and interact with AtRGS1 and AGB1 (17, 18).

For canonical Gα subunits, there are three major conformational changes between the GDP and GTP-bound states of the protein located in what are called Switch I, II and III. Switch I and Switch II directly contact the bound guanine nucleotide and include residues critical for catalyzing GTP hydrolysis, while Switch III contacts Switch II when in the activated conformation (19). These switches are linked between the nucleotide-binding domain and the RGS binding domain and are represented by five conserved sequence motifs named G1 to G5 (20). The G1–G3 boxes provide critical contacts for the β and γ phosphates of the guanine nucleotide and are essential for the coordination of Mg²⁺. The G4 and G5 loops are involved primarily in binding the guanine ring. The G2 and G3 boxes overlap with Switches I and II that are also the key Gβγ binding sites. The RGS domain directly binds to the three switch regions and stabilizes them in a transition state conformation.

It is paramount to resolve unequivocally if XLG proteins bind guanine nucleotide and relevant signaling elements such as RGS and Gβγ to elucidate its atypical mechanism. Here, we combine structure-based and physicochemical experimental methods along with molecular simulations to analyze the binding of XLG2 with both the nucleotide and with a candidate binding partner, AtRGS1/Gβγ dimer. We describe for the first time in great detail the structural issues that should raise concern among those who claim that XLG proteins are nucleotide-dependent switches. In fact, we show that XLG2 binds nucleotide so poorly that it is essentially nucleotide free in the cell, yet despite its nucleotide-free, “empty” state, XLG2 interacts with its partners AtRGS1 and AGB1 with an affinity similar to AtGPA1 in its transition state. We used molecular dynamic simulations to explain how this binding is disrupted and how these protein-protein interactions are maintained.

RESULTS AND DISCUSSION

XLG proteins lack critical residues for coordination of the γ and β phosphates on the guanine nucleotide

A multiple sequence alignment of XLGs G α domain, AtGPA1, and human G α 1 is shown in Fig. 1 with the G1-G5 motifs and switches I-III regions highlighted (noted as SwI-III). To compare the protein structures between XLGs and canonical G α subunits, we created high-quality models of the G α homology domains of XLG2 using MODELLER and the aligned sequences shown in Fig. 1. The human RGS4 and G α 1 transition state (Ligand: AlF₄ and GDP) complex (PDB [1AGR]) was used as template to generate the models of XLG2. Models were created using the *Automodel* script based on the template of human G α 1 in complex with AlF₄ and GDP (PDB [1AGR]). For evaluation and selection of the "best" model, we calculated the objective function (molpdf) DOPE score, GA341 assessment score between the model and the template (Fig. S1). The final model (XLG2-1) was selected given the lowest average value of the molpdf and the DOPE assessment scores.

As shown in Fig. S2, human G α 1 and Arabidopsis GPA1 have two domains: a Ras-like domain and an all-helical domain. Animal G α subunits and AtGPA1 are extremely similar in structure (RMSD= 1.8 Å (10)). The Ras-like domain is essential for the nucleotide and RGS proteins binding which contains the five guanine nucleotide binding motifs (G1-G5) and three flexible switch regions (SwI-III) (Fig. S2A). The all-helical domain is important for the intrinsic nucleotide exchange rate (21, 22). XLG2-1 shares a similar overall 3D structure with human G α 1 and AtGPA1 even though the sequence identity is ~ 30%. XLG2-1 contains a globally similar Ras-like domain and α helix domain. The three switch regions and the G1-G5 boxes are highlighted (Fig. S2B). The RMSD between G α 1 and XLG2-1 is 0.67 Å. However, despite similar global structure between XLG2-1 and human G α 1, many of the conserved motifs which are essential for nucleotide binding and hydrolysis are missing, including key residues within the G1, G3 and G5 motifs for nucleotide binding and some dominant residues in the P loop,

Switch I and Switch II for coordinating water and Mg^{2+} to catalyze GTP hydrolysis (Fig. 1 and Fig. 2B). These critical differences between the canonical $G\alpha$ and the XLG $G\alpha$ domain are described in detail next.

Figure 1. Alignment between human *G12* Arabidopsis *AtGPA1* and the C terminal $G\alpha$ domain of the three XLGs The G1-G5 motifs are shown in black boxes. The switches I-III regions (SwI-III) are highlighted (SwI in red, SwII in yellow and SwIII in blue). A percentage of equivalent residues is calculated per columns, considering physico-chemical properties. Blue boxes highlight residues with the same physico-chemical properties and red solid highlighting means the same residues. The contact residues to RGS protein are labeled with white boxes $\square$ and the contact residues with $G\beta$ are labeled with red boxes $\blacksquare$. The residues which are conserved in human *G12* and *AtGPA1* for GTP/GDP binding and hydrolysis but are missing in XLGs are highlighted with yellow and blue respectively. The residues essential for the catalysis of the nucleotide are highlighted as Rcat and Qcat. (The C domain of XLGs start with first C residues in the paper, C436 in XLG1, C435 in XLG2 and C396 in XLG3).

The highly conserved G1 motif is a phosphate-binding region containing a flexible structure designated “P-loop” (23). The G1 motif has a consensus sequence of GXXXXGKS/T for the heterotrimeric $G\alpha$ subunits (7). The P-loop envelopes the phosphates allowing the main chain and side-chain nitrogen atoms to interact tightly with the negatively-charged phosphates (Fig 2 A, B). In animal $G\alpha$ subunits as well as in AtGPA1, the sequence in the P loop and G1 motif are invariantly set to “GAGESGKS” (Fig 1, see G1 box). However, in XLG2-1, the G42 residue of $G\alpha 1$ in the P loop is replaced by E471 and the K46 residue of $G\alpha 1$ is replaced by A475, respectively (Fig 1 and Fig 2B). The G42 residue of $G\alpha 1$ or G47 residue of AtGPA1 in the P loop play a dominant role in binding the substrate with the main chain forms hydrogen bond with the γ phosphate oxygen atom (23). This G residue is shown in Fig 2A and B. More importantly, only a G residue side chain is small enough to avoid steric clash with the nucleotide and mutation of the corresponding P-loop residue in $G\alpha 1$, G42 to V, also drastically reduces its GTP hydrolysis activity (24-26). Structural studies of G42V mutant in $G\alpha 1$ suggest that the introduced valine side chain sterically prevents appropriate positioning of Q204 which coordinates a nucleophilic water molecule during GTP hydrolysis and steric pressure will induce the reconfiguration of switch II (6, 25, 26). Thus, we assume that the substitution of the large side chain of E471 in XLG2-1 reduces GTPase activity (Fig 2B).

Figure 2. Comparison of the G motifs of AtGPA1 PDB [2XTZ] and the XLG2 G alpha domain model with human G α 1 (PDB [1AGR]) Grey: AtGPA1, Magenta: XLG2-1 model, Light orange: G α 1. The substrate GDP and AlF₄ are shown as sticks and spheres. Mg²⁺ is shown as green sphere. Wn: nucleophilic water. The main different residues in G1 motif of XLG2 compared with G α 1 and AtGPA1 are shown as sticks **(A, B)**. Both AtGPA1 and G α 1 have the same G and K residues in G1 motif **(A)**. the G42 and K46 in G α 1 were replaced by E471 and A475 in the counterpart position of XLG2 **(B)**. The main different residues in G2 motif of XLG2 compared with G α 1 and AtGPA1 are shown as sticks **(C, D)**. Both AtGPA1 and G α 1 have the same R residues (known as arginine finger) and similar charged K180 and R192 in G2 motif **(C)**. But in XLG2 no arginine finger exists rather a Glu is at this position. Also, the charged K or R was replaced by a L **(D)**. The main different residues in G3 motif of XLG2 compared with G α 1 and AtGPA1 are shown as sticks **(E, F)**. Both AtGPA1 and G α 1 have the same DVGG residues in G3 motif **(E)**. But in XLG2 the DVGG was replaced by R669/N671/P672 relatively **(F)**. The main different residues in G5 motif of XLG2 compared with G α 1 and AtGPA1 are shown as sticks in G and H. Both AtGPA1 and G α 1 have the same A residues in G5 motif **(G)**. While in XLG2 the conserved A was replaced by Q818 **(H)**.

Additional differences were found with the P loop of the XLG proteins. The lysine (K46 of G α 1 and K51 of AtGPA1) residue in the G1 motif directly interacts with the β - and γ -phosphate oxygens of the GTP and thus is crucial for the required free energy change (6) (Fig. 2A). Given that there are two dominant residues mutations in the nucleotide pocket of XLG2 (G42 in G α 1 to E471 and K46 to A475) (Fig. 3B), we hypothesize that XLG2 binds the nucleotide with a reduced affinity *in vitro* and that XLG2 is nucleotide free *in vivo*.

The G3 box contains the signature sequence DXGG conserved throughout the heterotrimeric G-protein superfamily. Similar to the P loop, residues with the G3 motif interact with the γ -phosphate of GTP but also orients the Mg²⁺ ion that is critical for coordination of the guanine nucleotide. In AtGPA1 and G α 1, the G3 box is invariant “DVGG” (Fig. 2E), however in the XLG2 G α domain, the residues are replaced by “RLNP” (Fig. 2F). The conserved Asp residue of canonical G α subunits provides the water-molecule-mediated coordination of Mg²⁺ and therefore, the substitution of Asp for this critical Arg disrupts the ability to bind Mg²⁺ (6, 7, 27). Moreover, the main chain amide of the signature Gly residue is essential for nucleotide binding through hydrogen bonding to the γ -phosphate oxygen of the GTP (27). The main chain amide of this Gly is hydrogen bonded to the γ -phosphate and mutation of the two Gly residues in the G3 box confer dominant negative phenotypes (7, 27, 28). Gilman’s group showed that GDP-bound G α s G226A mutant (the second Gly in the G3 DVGG motif) has a higher affinity for

G β y than the wild-type subunit and is incapable of undergoing a GTP-induced conformational change (29). Taken together, stark differences in the three dominant residues in the G3 motif in the XLG2 protein nucleotide binding pocket which is conserved among all XLG proteins (Fig. 1) suggest that XLGs exist in the empty nucleotide state.

The G5 motif consensus is S/C/T-A-K/L/T. In G α 1, the G5 motif sequence is C-A-T (A = residue 326) and in AtGPA1, it is T-A-L (A = residue 355, Fig. 2G and H). In either form, the main chain of A326 in G α 1 is essential for the binding of GTP/GDP specifically forming a hydrogen bond with the oxygen of the guanidine nucleotide and S substitution at this site weakens the affinity for GTP γ S through steric crowding (30). Also, the equivalent A366S mutation in the G5 motif of Gas decreases Gas's affinity for GDP and GTP γ S by steric crowding and shifting G α towards the empty nucleotide pocket state (30, 31). However, in XLG2, the equivalent residues are C-Q-V (Q = residue 818, Fig. 2H). Thus, this substitution of A326 with Q818 in XLG2-1 is predicted to create a steric clash for nucleotide binding providing further inference that XLG is nucleotide-free.

XLG proteins lack key residues to catalyze GTP hydrolysis

Two amino acids, one from the G α subunit (the conserved catalytic glutamine residue in Switch II region which is named "Qcat") and one from the RGS protein (the so-called "Asn thumb"), together with nucleophilic water and a Mg²⁺ in the catalytic center are essential elements for the catalytic reaction (6) (Fig. 3). In G α 1, the Qcat in Switch II is Q204 is essential for catalytic activity in the G α subunit. A conserved Arg residue in Switch I region designated "Rcat" here, is also a major determinant of the catalytic activity. A water molecule designated "Wn" occupies the position for the nucleophile engaged in an in-line attack on the phosphate. The Asn thumb (N128 in RGS4) in the RGS domain reorients the Qcat allowing the carboximido moiety to form hydrogen bonds with AlF₄ mimicking a γ phosphate oxygen atom and Wn. Rcat forms electrostatic interactions with the β phosphate oxygen and with one

of the fluoride substituents of AlF_4 (Fig.3). Mutations in these residues of Switch I and Switch II are known to drastically alter GTPase activity (6).

Figure 3. Interactions between the catalysis center of the Ras domain and critical residues of RGS proteins. (A) The critical contact residues between Gai1 and RGS4 (PDB 1AGR) are shown in light orange. Arg 178 (Rcat) is within hydrogen bonding distance of the leaving group β - γ bridge oxygen and Q204 (Qcat) is a hydrogen bond donor to a fluorine (or O-) Al substituent and accepts a hydrogen bond from the presumptive water nucleophile (Wn). The hydrogen bond network (yellow dashed lines) involving N128 (Asn thumb of RGS4), Qcat, G42 and the γ phosphate (modeled by AlF_4) orient Wn for nucleophilic attack and stabilize developing charge at the β - γ bridge leaving group oxygen. RGS4 residues Asn 128 constrain the conformation of Gai1 Q204 (Qcat) to the pre-transition state conformation. AtGPA1 contains the same catalysis network (A) however the catalysis network was disrupted in XLG2 with the loss of the Gln cat and Arg finger and replaced by R673 and E629 respectively **(B)**. Grey: AtGPA1, Magenta: XLG2, Light orange: Gai1. The substrate GDP and AlF_4 are shown as sticks and spheres. Main catalysis residues between Gai1, AtGPA1, XLG2 and RGS4 are highlighted as sticks. Wn: nucleophilic water.

All XLG proteins lack both essential Rcat and Qcat for the catalysis (Fig. 3B). In XLG proteins, the Rcat residue in Switch I is E, creating a charge reversal that disrupts electrostatic interactions with the β and γ phosphates of the guanine nucleotide. The equivalent mutation in Gai1 exist as a stable protein in a nucleotide-free state and lacks the capacity to form the active conformation (19). In all XLG proteins, the Qcat residue of Switch II is R/K which is unable to coordinate with either the Asn thumb of the RGS protein or the nucleophilic water to hydrolyze GTP. Both Q204R and R178C mutations abrogate nucleotide hydrolysis (19). The structural characteristic of the XLG proteins catalysis center suggests

that XLGs lack the ability both to coordinate with RGS to hydrolyze GTP and the intrinsic GTPase activity of G γ subunits.

XLG2 has a much lower binding affinity towards nucleotide than canonical G subunits yet interacts with similar affinities towards G $\beta\gamma$ and AtRGS1

Assessments of nucleotide binding to XLG proteins to date lack quantitation for binding constants (32). Similarly, XLG protein interaction with AtRGS1 and G $\beta\gamma$ have been indirect measurements (11, 16, 17). To correct this deficit, we used microscale thermophoresis (MST) to measure the binding affinity of XLG2 and AtGPA1 with guanine nucleotides (GDP and GTP γ S) and with binding partners AtRGS1 and G $\beta\gamma$. The advantages of this new technique are the capability of obtaining accurate affinities in the low affinity (μ M-mM Kd) range with small amounts of protein. Note that, unlike MST, traditional radioisotope binding assays are not accurate for low affinity interactions. Raw data with the quality control parameters provided are in Fig 4 and S3 and are summarized in Table 1. The observed Kd of AtGPA1 binding GTP γ S was ~ 21 nM. This is within the range of Kds reported for animal G subunits (10-100 nM, (33)). XLG2 bound GTP γ S with a Kd of ~ 2 μ M which is 100 times lower affinity than GTP γ S binds AtGPA1 when tested under the same conditions and nearly 1000 times lower when measured using radioactive ligand (34). Moreover, the affinity of XLG2 to GDP is ~100 fold lower (177 μ M) than for GTP γ S. Quantitative analyses clearly show that XLG2 is severely impaired in guanine nucleotide binding (Table 1).

	GTP γ S	GDP	RGS1-C domain G α in apo state	RGS1-C domain G α in transition state	G $\beta\gamma$ G α in GDP state
AtGPA1	21 $\pm$ 18 nM	28 $\pm$ 12 μ M	125 $\pm$ 81 nM	67 $\pm$ 18 nM	2 $\pm$ 1.22 μ M
XLG2	2350 $\pm$ 460 nM	177 $\pm$ 33 μ M	198 $\pm$ 45 nM	NA	0.7 $\pm$ 0.09 μ M

Table 1. Summary of the binding affinity (Kd) among AtGPA1 and XLG2 with nucleotide, RGS1-C domain and Gβγ. Top row in the inset indicates the tested interactors. The values were determined from the binding isotherms shown in Figure 4. The values are averages and StdDev for all the experimental replicates. Each experiment was replicated at least once.

Figure 4. Binding isotherms for nucleotide, RGS1-C domain and Gβγ to AtGPA1 and XLG2. Microscale Thermophoresis was used. (A) Binding isotherm and Kd value of AtGPA1 binding GTPγS and (B) GDP. (C) Binding isotherm and Kd value of XLG2 binding GTPγS and (D) GDP. (E) Binding isotherm and Kd value of RGS1 C terminal domain to AtGPA1 apo state and (F) XLG2 apo state. (G) Gβγ binding to AtGPA1 and (H) XLG2 Gα domain. S/N: signal to noise ratio. Each experiment was repeated at least once. Binding curve and Kd were fitted as described in Methods. Error bars represent StdDEV. Each experiment was repeated at least once.

256

257 With this poor affinity toward guanine nucleotides, the concentration of GTP in plant cells would
 258 need to be 100 times greater than in animal cells for XLG2 to be GTP bound, however, for several
 259 reasons, this explanation of a mechanism to compensate the weak GTP affinity by XLG proteins is not
 260 reasonable. First, such a saturating concentration of GTP would eliminate the switch-like behavior of
 261 the canonical plant Gα subunit. Second, protein translation uses the same machinery in both plant and
 262 animal cells and the GTP hydrolyzed for its proof reading and is sensitive to its cytoplasmic
 263 concentration. Similarly, plant and animal microtubules requires GTP binding and hydrolysis. Both
 264 translation and cytoskeletal dynamics would cease at this high concentration of GTP. Third, nucleotide
 265 synthesis uses product inhibition to control the levels accordingly. A 100-fold higher concentration of
 266 GTP would be incompatible with enzymes involved in nucleotide synthesis. Fourth, the highest known
 267 concentration of GTP in plant cells is equivalent to only one Kd for GTP binding to XLG2 (35-37). As
 268 such, the concentration of GTP in plant cells, especially non-dividing cells, may be rate-limiting for full
 269 occupancy of XLG2 by GTP. For these reasons, we conclude that XLG2 is not likely bound by GTP *in*
 270 *vivo*.

271 Assmann's group reported the unusual finding that the three XLG proteins bind and hydrolyze
 272 GTP using Ca²⁺ instead of Mg²⁺ as a coordinating factor (32). To test this, we performed MST
 273 experiments to measure the binding affinity of XLG2 with nucleotide in the presence of Ca²⁺. The results
 274 showed lower binding affinity towards GTPγS (~186 μM) with Ca²⁺ vs. Mg²⁺ (Fig S3). This indicates that
 275 Ca²⁺ may not act as the cofactor for XLGs binding GTPγS. Ca²⁺ induced relatively higher binding affinity
 276 for GDP (~28 μM), albeit still poor, compared to Mg²⁺ as a cofactor (Fig S3).

277 Interestingly, despite XLG2 having much lower binding affinity towards GTPγS and GDP
 278 compared with AtGPA1, it had a similar binding affinity to the C-terminal RGS domain of AtRGS1 and
 279 to the Arabidopsis Gβγ dimer (AGB1/AGG1). AtGPA1 bound AtRGS1 with a Kd ~125 nM with in
 280 AtGPA1 in its apo state and ~ 67 nM in its transition state (Fig.S3). XLG2 has a similar Kd of ~198 nM

towards AtRGS1 when in its apo state (Table 1 and Fig. 4). The K_d for a transition state XLG2 was not determined because this state is not relevant due to its nucleotide independence. Moreover, XLG2 showed a ~0.7 μM binding affinity towards Gβγ similar to that of AtGPA1 which is ~2 μM (Table 1 and Fig. 4). This suggests that XLG2 exists as a nucleotide-independent inhibitor of G signaling through its ability to sequester Gβγ directly or indirectly by binding to AtRGS1 thus enabling freed AtGPA1 to sequester Gβγ.

A mechanistic explanation: Relative instability of XLG2 confers the reduced nucleotide interaction

We applied several computational modeling and simulation approaches to understand the underlying molecular mechanisms differentiating AtGPA1 and XLG2 proteins. We sought to provide structural and molecular dynamics rationales for the experimentally observed differences in nucleotide binding preferences by the two proteins. To this point, we performed microseconds of molecular dynamics (MD) simulations of four molecular complexes, involving GDP and GTP nucleotides, each in complex with both AtGPA1 and the homology-modeled XLG2-1 Gα domain, followed by comparative analyses of the respective MD trajectories. The main finding of our simulations is that the molecular dynamic behaviors of XLG2-1 differs from that of AtGPA1. We observed that overall XLG2-1 was more mobile in comparison with AtGPA1, which generally retained its original crystallographic structure over the course of simulations. Furthermore, to distinguish the two proteins with respect to their nucleotide binding capabilities, we focused on analyzing the behavior of the ligand binding site both in the context of the intra-protein and ligand-protein interactions in order to more clearly understand the key factors contributing to the experimental findings of the lower nucleotide binding affinity in XLG2.

In preparation for MD simulations, the structure of XLG2 obtained by homology modeling, was subjected to molecular mechanics minimization following several protocols as described in the Methods section in order to avoid unnatural clashes between atoms resulting from homology modeling. To

understand the overall dynamics of the proteins, we analyzed RMS fluctuations per residue and calculated RMSD using all C-alpha atoms of the proteins (**Fig. S4, S5**), which showed that the general fold of AtGPA1 was more stable and the amino-acid residues displayed lower mobility compared to XLG2. We then sought to understand the dynamics of the nucleotide binding site and explored the key differences in the interactions formed within the binding site. First, we visualized the binding sites of the two proteins to explore the main differences in terms of the amino-acid residue composition (**Fig. 5A**). The following differences in the similarly-positioned, binding-site residues were determined between AtGPA1 and XLG2: E48 to K472, D162 to R601, R190 to E629, F253 to E705, R260 to K714, K288 to K742 in guanine and ribose binding sites, and K51 to A475, S52 to T476, T193 to S632, D218 to R669, Q222 to R673 in Mg²⁺ and phosphates binding sites (**Fig. 1, 5**).

Figure 5. Difference in dynamics between the nucleotide-bound AtGPA1 and XLG2: Insights from MD simulations. Homology modeling and MD simulations reveal the main differences in amino-acid residue composition and the nucleotide binding site dynamics of AtGPA1 (grey) and XLG2 (magenta) **(A)** Aligned minimized AtGPA1 crystal structure (PDB ID 2xtz) and the homology modeled XLG2. The zoomed in plots separately display phosphate and Mg^{2+} binding site (top) and guanine and ribose binding site (bottom) on the example of GTP-bound complexes, highlighting the most prominent differences in the amino-acid residues. **(B)** The relationship between the nucleotide-protein interaction energies (designated as $E_{\text{interaction}}$ on the scatter plots, and calculated as the sum of the Coulomb and LJ terms) and mobility of the nucleotide ($\text{RMSD}_{\text{ligand}}$) and binding site ($\text{RMSD}_{\text{binding site}}$) display substantial separation among AtGPA1-GTP (black points and solid line), XLG2-GTP (purple points and dashed line), AtGPA1-GDP (grey points and solid line), and XLG2-GDP (pink points and dashed line) complexes (Fig. S6-8).

To understand the differences in the binding site dynamics, we first calculated RMSD of the heavy atoms of residues located in the binding sites (**Fig. 5B and Fig. S6**), which we defined as the protein residues within 4 Å from GTP (see Methods). We observed that AtGPA1 and XLG2 nucleotide binding

sites differed in conformational dynamics and had distinctly different configurations as elaborated in the following paragraph. Next, we aimed to understand the impact of the difference in dynamics of the binding site residues on the nucleotide mobility and nucleotide binding preferences (**Fig. 5B & Figs. S7, S8**). Through exploring the relationship between the nucleotide-protein interaction energy (calculated as the sum of intermolecular Coulomb and LJ terms of the molecular mechanics energy of the nucleotide-protein complexes) and mobilities of the binding site and ligand, we observed substantial differences across the four complexes formed when AtGPA1 and XLG2 bound to both GDP and GTP. The molecular systems occupied distinct regions on each of these two landscapes. Importantly the ranking order of the means of two parameters, (i) the nucleotide mobility in the pocket as characterized by the RMSD of the nucleotide (from smallest to largest), and subsequently (ii) the nucleotide-protein interaction energies characterized as the sum of all LJ and Coulomb terms of the nucleotide-protein interactions (from more negative to less negative), agreed with the ranking order in terms of our experimental binding affinities ($K_d \pm \text{StDev}$) as follows: **1st**) GPA1-GTP (0.021 $\pm$ 0.018 μM), **2nd**) XLG2-GTP (2.4 $\pm$ 0.5 μM), **3rd**) GPA1-GDP (28 $\pm$ 12 μM), **4th**) XLG2-GDP (177 $\pm$ 33 μM) (**Table 1**).

This result added confidence to our structural and simulations-derived interpretations of the molecular complex formations. We would like to emphasize, however, that such calculations of the intermolecular interaction energies are merely estimates of the relative strengths of ligand-protein interactions in the bound state, which by no means is equivalent to the assessment of the change in Gibbs free energy of binding (38-41).

Figure 6. Apo and nucleotide-bound proteins obtain distinct configurations defining the nucleotide binding preferences of AtGPA1 and XLG2. (A) The top five most populated metastable states of the nucleotide binding site obtained in cluster analysis of MD trajectories indicate that nucleotide-bound AtGPA1 obtains a stable frequently visited conformational state, whereas XLG2 complexes tend to transition between conformationally diverse states with lower probabilities. Interestingly, both apo proteins obtain multiple states with equivalently low probabilities (Fig. S6,S9). Aligned centroids of the largest metastable states are presented in panels B-D. (B) The most populous apo states show stable E48-R190-R260 and D162-K288 salt bridge networks in the guanine binding site (left image) of AtGPA1 (grey), and a more destabilized salt bridge network between similarly positioned residues in XLG2 (magenta) primarily contributed by R601-K742 electrostatic repulsion. K51-D218 salt bridge in the phosphate and Mg^{2+} binding sites (right image) enables a more structures AtGPA1, while the neutral A475 and a repulsion between R669 and R673 cause a more disintegrated XLG2. (C) GDP- and (D) GTP-bound complexes retain the strong salt bridge network in AtGPA1 and less stable electrostatic interactions in XLG2. K51 reorients and forms an additional bond with phosphates in AtGPA1, which is prevented by the equivalently positioned neutral A475 in XLG2. K472 breaks its bonds with E629 and re-arranges to interact with closer located phosphates. The absence of γ -phosphate in GDP makes the nucleotide more mobile, losing the frequency of its contacts. R673 in GTP-bound XLG2, however, forms a relatively stable bond with the γ -phosphate seemingly increasing the nucleotide binding affinity. The residues shown in darker shades in panels B-D (D162 and R190 of GPA1; R601 and E629 of XLG2) make the key intra-protein interactions defining the binding site shape. The differences in the frequency of the aforementioned interactions, on the example of GTP-bound complexes, are clearly seen through heatmaps of Δ contacts (minimum distances) (E) between the non-hydrogen atoms of the nucleotide and binding site residues, and (F) within the binding site

residues. 100 states of the most populated clusters were used to generate the heatmaps. The red squares highlight the most prominent changes in the interactions stipulating the importance of the residues to the stability of the active sites and to the interactions with the nucleotide.

Cluster analysis of the generated MD trajectories (42-44) (see Methods for details) revealed metastable states with distinct configurations of the binding sites linked to the experimental nucleotide binding preferences (**Fig. 6**). For the most populated metastable state of each molecular complex, we explored the specific intra-protein chemical interactions which directly impact the dynamics of the active sites, and their impact on the nucleotide-binding interactions. The results show that nucleotide-bound AtGPA1 achieves stable dominant (i.e., frequently visited) conformations, whereas XLG2 complexes tend to transition between conformationally-diverse states with lower probabilities. Such low frequency populations of the top clusters are associated with a more dynamic binding pocket in XLG2. Interestingly, both apo-proteins assume multiple states with low probabilities across the top five clusters (**Fig. S6A, S9, Supplemental Movies 1-6**).

The most populous states of the apo-protein show a stable salt bridge network in AtGPA1, but no network and fewer coherent salt bridges are found in XLG2-1. In the guanine binding site of AtGPA1, salt bridges formed between E48 of the P-loop, R190 of Switch I, and R260 of Switch III, as well as between D162 and K288, while a more destabilized salt bridge network appeared between similarly positioned residues in XLG2-1, namely: E629 of Switch I, K472 of P-loop, K714 of Switch III and E705 (**Fig. 6B, S9-12**). In XLG2-1, a positive charge at K714 (position equivalent to R260 in AtGPA1) is not capable of forming a salt bridge with K472 (position equivalent to E48 in AtGPA1). In the phosphate and Mg^{2+} binding sites, the K51-D218 salt bridge enables a more structured apo-AtGPA1, while for XLG2, the neutral sidechain of A475 (position equivalent to K51 in AtGPA1) together with a repulsion between R669 and R673 precludes a stabilizing salt bridge. In general, AtGPA1 salt bridges formed by the two loop residues D162 and R190 (and equivalently placed E629 in XLG2) drawing the two domains closer together to subsequently increase the number of interactions between the all-helical domain and the nucleotide.

The nucleotide-bound complexes retain the aforementioned strong salt bridge network in AtGPA1 and weak electrostatic interactions in XLG2 (**Fig. 6C,D, Fig. S9-12**). In the bound state, the K51 sidechain of AtGPA1 was re-arranged to form an additional bond with phosphates, which is an additional salt bridge lacking by the neutral A475 in XLG2-1. In the GTP-bound XLG2-1 model, E705 lost its K472 interaction to γ -phosphate which also attracted R673 for further stabilization. Moreover, E48 in AtGPA1 avoided the negatively-charged phosphates which further promoted electrostatic interactions with both R190 and R260 to stabilize this cavity. In the GDP-occupied state, due to the lack of the γ -phosphate, GDP is more mobile and loses a number of its contacts with the active site residues in both G α proteins, but more so in XLG2-1 due its structurally-unstable binding site.

The heatmaps of Δ contacts from experiments determining the difference in the minimum distances between the two G α proteins and **(i)** the atoms of the nucleotide and binding site residues, and **(ii)** intra-protein interactions within the binding site residues, clearly show the contrast between the interaction frequencies within these two proteins. As highlighted in Fig 6E and F, the most prominent changes in the interactions important to the stability of the active sites and to the interactions with the nucleotide, in addition to the previously indicated interactions, reside with F253 in AtGPA1 which was overall closer to both the nucleotide and the binding site residues compared with similarly positioned E705 in XLG2-1. This aromatic residue makes frequent pi-cation interactions with R190 in both apo- and ligand-bound protein and occasional pi-cation interactions with the Mg²⁺ in the ligand-bound state.

Our analyses shows that the D162-K288 salt bridge (**Fig. 7A**) is one of the key interactions maintaining the shape of the apo-AtGPA1 nucleotide-binding site. In contrast in XLG2-1, two positively charged residues, R601 and K742, situated in positions equivalent to D162 and K288 of AtGPA1 caused electrostatic repulsion, pushing away the all-helical domain of the protein from the Ras-like domain, resulting in an increased mobility and a less structured nucleotide binding site in XLG2-1. The R601-K742 distance in XLG2-1 was highly correlated with the fluctuations of the binding site in XLG2-GDP and to a lesser extent in XLG2-GTP (**Fig. 7B and C**). The reason for the former is the lack of an

extra anchor in terms of γ -phosphate in GDP to enable the electrostatic repulsion between R601 and K742 to be the main contributor to the instability of XLG2-GDP binding pocket. This agrees with the experimentally observed poor binding affinity of GDP to XLG2. Although this correlation still exists in XLG2-GTP, it is less pronounced due to the presence of the extra phosphate in GTP. In XLG2, K714 makes ionic bonds with phosphates, however, this does not seem to be sufficient to retain the binding site integrity distorted by the aforementioned repulsion.

Figure 7. Key intra-protein distances responsible for the experimental nucleotide binding affinities as determined from MD simulations. Distribution of the minimum distances between the residues emphasized in Fig. 6 and text. In general, a stronger salt bridge network in AtGPA1 maintains the shape of its nucleotide-bound binding site (see text for more details). **(A)** Distance between D162 and K288 and the equivalently placed R601 and K742 in XLG2. **(B)** The repulsion between R601 and K742 is highly correlated ($R=0.95$) with the binding site RMSD in XLG2-GDP and **(C)** to a lesser extent in XLG2-GTP ($R=0.46$). The 2D correlation plots were constructed using kernel-density estimation with Gaussian kernels. **(D)** Distance between E48 and R190 in AtGPA1 and similarly positioned K472 and E629. **(E)** Distance between E48 and R260 in AtGPA1 and K472 and E705 (aligned with F253 of AtGPA1) in XLG2. The distributions show that both of the salt bridges (panels E and D) are dominant in the apo proteins, and are less persistent in nucleotide-bound XLG2. **(F)** The distributions of the minimum distance between Mg^{2+} and γ -phosphate binding site residues (D218, S52, T193, and D218 in GPA1; T476, S632, R669 in XLG2) and Mg^{2+} counterion. For clarity in panels A, D, E, and F the

values of probabilities on the y-axis are hidden. Apo AtGPA1 is plotted with solid light grey lines, AtGPA1-GDP—solid grey, AtGPA1-GTP—solid black, apo XLG2—dotted pink, XLG2-GDP—dotted magenta, XLG2-GTP—dotted purple; the probability densities for XLG2 are shaded for contrast.

Another distinction between the two proteins is in the Mg^{2+} binding site (**Fig. 7F & Fig. S13**) discussed above. In AtGPA1, D218 formed an H-bond with S52 (which interacts with Mg^{2+}) and a Coulomb interaction with Mg^{2+} , whereas the ‘bulkier’ and positively charged sidechain of R669 in XLG2 (position equivalent to D218 in AtGPA1) did not form stable interactions with either T476 or S632 (which interact with Mg^{2+}) and caused an electrostatic repulsion with Mg^{2+} . The distribution of the minimum distance between the Mg^{2+} binding site residues (S52, T193, D218, and Q222 in AtGPA1; T476, S632, R669, and R673 in XLG2) and the Mg^{2+} counterion clearly explain this effect.

To interpret the observed equivalent AtRGS1 binding capability of the two Gα proteins (**Fig. 3, 4, Table 1**), we estimated the structural stability of the specific regions that are involved in AtRGS1 binding (**Fig. S14**). We showed that the three equivalently placed AtRGS1 binding site residues of apo-AtGPA1 and apo-XLG2-1 similarly maintained their structural integrity over the course of our simulations. Such conformationally-preserved regions in the apo-proteins position them to bind AtRGS1 when it is tethered close to either protein.

Conclusion

XLG2 binds GTP *in vitro* poorly such that at the estimated concentration of GTP in plant cells, XLG2 is not expected to be nucleotide bound. However, XLG2 binds regulatory partners, AtRGS1 and Gβγ. Therefore, XLG2 is a decoy that negatively regulates by sequestering the Gβγ dimer directly and also indirectly by promoting AtGPA1 interacting with Gβγ through freeing AtGPA1 from the AtRGS1::AtGPA1 complex. While this concept shares similarities for control of G signaling by dominant negative mutations of canonical G protein in animals (45), it is unique in that the negative control is provided *in trans* by a genetically-encoded, atypical G protein.

Taken together, our modeling data provide credible interpretations for the experimentally observed strengths of guanine nucleotide binding to AtGPA1 and XLG2. Several key intra-protein and nucleotide-protein interactions in AtGPA1 were shown to be attributed to the higher structural stability of the binding site of the protein and to more persistent contacts of the protein with the nucleotide and magnesium. We show mechanistically that among the chief intra-protein interactions preserving the stability of the binding site in both apo- and nucleotide-bound-states of AtGPA1 include the following ionic bonds: D162-K288, R190-E48-R260, and K51-D218. Because XLG2 is important for disease resistance and development (17, 46, 47), engineering these equivalent residues may lead to improvements in crop performance.

Acknowledgments

This work was supported by NIGMS (R01GM065989) and NSF (MCB-0718202) awarded to Alan. M. Jones. The deep computational analyses were supported by the resources of the UNC Longleaf cluster (<https://its.unc.edu/research-computing/longleaf-cluster/>)

References

1. Lambert NA (2008) Dissociation of heterotrimeric G proteins in cells. *Sci Signal* 1(25):re5.
2. Sprang SR (1997) G proteins, effectors and GAPs: structure and mechanism. *Current Opinion in Structural Biology* 7(6):849-856.
3. Urano D, Chen J-G, Botella JR, & Jones AM (2013) Heterotrimeric G protein signalling in the plant kingdom. *Open Biology* 3(3).
4. Urano D & Jones AM (2014) Heterotrimeric G protein-coupled signaling in plants. *Annual Review of Plant Biology* 65(1):365-384.
5. Kleuss C, Raw AS, Lee E, Sprang SR, & Gilman AG (1994) Mechanism of GTP hydrolysis by G-protein alpha subunits. *Proc Natl Acad Sci U S A* 91(21):9828-9831.
6. Sprang SR (2016) Invited review: Activation of G proteins by GTP and the mechanism of G α -catalyzed GTP hydrolysis. *Biopolymers* 105(8):449-462.
7. Sprang SR (1997) G protein mechanisms: insights from structural analysis. *Annu Rev Biochem* 66:639-678.
8. Gilman AG (1987) G proteins: transducers of receptor-generated signals. *Annu Rev Biochem* 56:615-649.
9. Temple BR, Jones CD, & Jones AM (2010) Evolution of a signaling nexus constrained by protein interfaces and conformational states. (Translated from eng) *PLoS computational biology* 6(10):e1000962 (in eng).
10. Jones JC, et al. (2011) The crystal structure of a self-activating G protein alpha subunit reveals its distinct mechanism of signal initiation. *Sci Signal* 4(159):ra8.
11. Chakravorty D, Gookin TE, Milner MJ, Yu Y, & Assmann SM (2015) Extra-Large G proteins expand the repertoire of subunits in Arabidopsis heterotrimeric G protein signaling. *Plant Physiology* 169(1):512-529.
12. Ullah H, Chen J-G, Temple B, Boyes D, & Alonso J (2003) The β subunit of the Arabidopsis G protein negatively regulates auxin-induced cell division and affects multiple developmental processes. *Plant Cell* 15:393.
13. Thung L, Trusov Y, Chakravorty D, & Botella J (2012) G γ 1 + G γ 2 + G γ 3 = G β : The search for heterotrimeric G-protein γ subunits in Arabidopsis is over. *J. Plant Physiol.* 169:542.
14. Johnston CA, et al. (2007) GTPase acceleration as the rate-limiting step in Arabidopsis G protein-coupled sugar signaling. *Proceedings of the National Academy of Sciences* 104(44):17317-17322.
15. Maruta N, Trusov Y, Brenya E, Parekh U, & Botella JR (2015) Membrane-localized extra-large G proteins and G $\beta\gamma$ of the heterotrimeric G proteins form functional complexes engaged in plant immunity in Arabidopsis. *Plant Physiology* 167(3):1004-1016.
16. Urano D, et al. (2016) Saltatory evolution of the heterotrimeric G protein signaling mechanisms in the plant kingdom. *Science Signaling* 9(446): ra93.
17. Liang X, et al. (2018) Ligand-triggered de-repression of Arabidopsis heterotrimeric G proteins coupled to immune receptor kinases. *Cell Research* 28(5):529-543.
18. Urano D, et al. (2016) Plant morphology of heterotrimeric G protein mutants. *Plant and Cell Physiology* 57(3):437-445.
19. Coleman DE, et al. (1994) Structures of active conformations of G α 1 and the mechanism of GTP hydrolysis. *Science* 265(5177):1405-1412.
20. Berman DM, Wilkie TM, & Gilman AG (1996) GAIP and RGS4 are GTPase-activating proteins for the G α subfamily of G protein alpha subunits. *Cell* 86(3):445-452.
21. Jones JC, Jones AM, Temple BR, & Dohlman HG (2012) Differences in intradomain and interdomain motion confer distinct activation properties to structurally similar G α proteins. *Proc Natl Acad Sci U S A* 109(19):7275-7279.
22. Jones JC, Temple BR, Jones AM, & Dohlman HG (2011) Functional reconstitution of an atypical G protein heterotrimer and regulator of G protein signaling protein (RGS1) from Arabidopsis thaliana. *J Biol Chem* 286(15):13143-13150.
23. Saraste M, Sibbald PR, & Wittinghofer A (1990) The P-loop--a common motif in ATP- and GTP-binding proteins. *Trends Biochem Sci* 15(11):430-434.

24. Seeburg PH, Colby WW, Capon DJ, Goeddel DV, & Levinson AD (1984) Biological properties of human c-Ha-ras1 genes mutated at codon 12. *Nature* 312(5989):71-75.
25. Bosch DE, *et al.* (2012) A P-loop mutation in Galpha subunits prevents transition to the active state: implications for G-protein signaling in fungal pathogenesis. *PLoS Pathog* 8(2):e1002553.
26. Raw AS, Coleman DE, Gilman AG, & Sprang SR (1997) Structural and biochemical characterization of the GTPgammaS-, GDP.Pi-, and GDP-bound forms of a GTPase-deficient Gly42 --> Val mutant of Gialpha1. *Biochemistry* 36(50):15660-15669.
27. Noel JP, Hamm HE, & Sigler PB (1993) The 2.2 Å crystal structure of transducin-alpha complexed with GTP gamma S. *Nature* 366(6456):654-663.
28. Barren B & Artemyev NO (2007) Mechanisms of dominant negative G-protein alpha subunits. *J Neurosci Res* 85(16):3505-3514.
29. Lee E, Taussig R, & Gilman AG (1992) The G226A mutant of Gs alpha highlights the requirement for dissociation of G protein subunits. *J Biol Chem* 267(2):1212-1218.
30. Posner BA, Mixon MB, Wall MA, Sprang SR, & Gilman AG (1998) The A326S mutant of Gialpha1 as an approximation of the receptor-bound state. *J Biol Chem* 273(34):21752-21758.
31. Iiri T, Herzmark P, Nakamoto JM, van Dop C, & Bourne HR (1994) Rapid GDP release from Gs alpha in patients with gain and loss of endocrine function. *Nature* 371(6493):164-168.
32. Heo JB, Sung S, & Assmann SM (2012) Ca²⁺-dependent GTPase, Extra-large G Protein 2 (XLG2), promotes activation of DNA-binding protein related to Vernalization 1 (RTV1), leading to activation of floral integrator genes and early flowering in Arabidopsis. *Journal of Biological Chemistry* 287(11):8242-8253.
33. Malinski JA, Zera EM, Angleson JK, & Wensel TG (1996) High affinity interactions of GTPγS with the heterotrimeric G Protein, transducin: Evidence at high and low protein concentrations *J Biol Chem* 271(22):12919-12924.
34. Johnston CA, Willard MD, Kimple AJ, Siderovski DP, & Willard FS (2008) A sweet cycle for Arabidopsis G-proteins: Recent discoveries and controversies in plant G-protein signal transduction. *Plant Signal Behav* 3(12):1067-1076.
35. Ashihara H & Nygaard P (1989) Purine nucleotide and RNA synthesis in suspension cultured cells of carrot. *Physiologia Plantarum* 75:31-36.
36. Meyer R & Wagner KG (1985) Analysis of the nucleotide pool during growth of suspension cultured cells of *Nicotiana tabacum* by high performance liquid chromatography. *Physiologia Plantarum* 65(4):439-445.
37. Yin Y, Katahira R, & Ashihara H (2014) Metabolism of purine nucleosides and bases in suspension-cultured *Arabidopsis thaliana* cells. *Eur. Chem. Bull.* 3(9):925-934.
38. Ganotra GK & Wade RC (2018) Prediction of Drug-Target Binding Kinetics by Comparative Binding Energy Analysis. *ACS Med Chem Lett* 9(11):1134-1139.
39. Henriksen NM, Fenley AT, & Gilson MK (2015) Computational Calorimetry: High-Precision Calculation of Host-Guest Binding Thermodynamics. *J Chem Theory Comput* 11(9):4377-4394.
40. Mobley DL & Gilson MK (2017) Predicting Binding Free Energies: Frontiers and Benchmarks. *Annu Rev Biophys* 46:531-558.
41. Fujitani H, *et al.* (2005) Direct calculation of the binding free energies of FKBP ligands. *J Chem Phys* 123(8):084108.
42. Abramyan TM, Snyder JA, Thyparambil AA, Stuart SJ, & Latour RA (2016) Cluster analysis of molecular simulation trajectories for systems where both conformation and orientation of the sampled states are important. *Journal of Computational Chemistry* 37(21):1973-1982.
43. Daura X, *et al.* (1999) Peptide folding: When simulation meets experiment. *Angewandte Chemie International Edition* 38(1-2):236-240.
44. Shao J, Tanner S, Thompson N, & Cheatham T (2007) Clustering molecular dynamics trajectories: 1. Characterizing the performance of different clustering algorithms. *J Chem Theory Comput.* 3(6):2312-2334.
45. Barren B & Artemyev NO (2007) Mechanisms of dominant negative G-protein α subunits. *Journal of Neuroscience Research* 85(16):3505-3514.
46. Liang X, *et al.* (2016) Arabidopsis heterotrimeric G proteins regulate immunity by directly coupling to the FLS2 receptor. *eLife* 5:e13568.

47. Liang Y, Gao Y, & Jones AM (2017) Extra large G-Protein interactome reveals multiple stress response function and partner-dependent XLG subcellular localization. *Frontiers in Plant Science* 8(1015).