Early vegetation recovery after the 2008-2009 explosive eruption of the Chaitén Volcano.

Ricardo Moreno-González^{1*}, Iván A. Díaz², Duncan A. Christie^{2,3}, Rafael E. Coopman^{2,4}, Antonio Lara^{2,3}

1. Department of Palynology and Climate Dynamic, University of Göttingen, Germany.

2. Instituto de Conservación, Biodiversidad y Territorio, Universidad Austral de Chile, Valdivia, Chile.

3. Center for Climate and Resilience Research (CR)²

4. Marilyn Ball's Lab, Plant Science Division, Research school of Biology, The Australian National University

* Corresponding author: Ricardo Moreno González

E-mail: rmoreno@uni-goettingen.de

Abstract

In May 2008, Chaitén volcano entered in eruptive process, one of the world largest eruptions in the last decades. The catastrophic event left different type of disturbance and caused diverse environmental damage distributed heterogeneously in the surrounding areas of the volcano. We went to the field to assess the early vegetation responses a year after the eruption, in September 2009. Particularly, we evaluated the lateral-blast disturbance zone. We distributed a set of plots in three disturbed sites, and one in an undisturbed site. In each of these sites, in a plot of 1000m² we marked all stand tree, recording whether they were alive, resprouting or dead. Additionally, in each site 80 small-plots (~4m²) we tallied the plants regeneration, its coverage, and the log-volume. We described whether the plant regeneration was growing on mineral or organic substrate. In the blast-zone the eruption created a gradient of disturbance. Close to the crater we found high devastation marked by no surviving species, scarce standing-dead trees and logs, as well as no tree regeneration. On the other extreme of the disturbance gradient, the trees with damaged crown were resprouting, small-plants were regrowing and seedlings were more dispersed. The main regeneration strategy was the resprouting of trunks or buried roots, while few seedlings were observed in the small plots and elsewhere in disturbed areas. However, the assessment was too soon after the eruption and updated monitoring is required to confirm observed patterns. The main findings of this study are: i) a mosaic of pioneering-wind dispersed species, scattered survivors regrowing and spreading from biological legacies, and plant species dispersed by frugivorous birds, likely favored by the biological legacies; (ii) the early succession is influenced by the interaction of the species-specific life history, altitudinal gradient and the different intensity of disturbance.

Keywords: 2008-Chaiten Volcano, Volcanic disturbance, vegetation dynamics, *Nothofagus* dynamics, South American temperate rainforest, early succession.

1. Introduction

Volcanic eruptions are large-scale disturbance that modify the composition, the structure and ecosystems processes (Swanson et al., 2013; Turner and Dale, 1998; Turner et al., 1997). As volcanoes enter in eruptive phase complex mechanism involve the events in a gradient of multiple type of disturbances such as explosive blasts, thermal and toxic chemical waves, landslides, glowing avalanche deposits, debris flows, lava flows, and air falls of volcanic tephra (Dale et al., 2005; Swanson et al., 2013). A volcanic eruption could have effects at very large spatial and temporal scales (Peet, 1992; Robock and Oppenheimer, 2003), yet directly on surrounded forest throughout burning, burying and/or blowing down trees or other plants (Dale et al., 2005). However, since this kind of large disturbance are infrequent (Turner et al. 1998) or have been seldom monitored (Chrisafulli and Dale, 2018), our understanding of vegetation responses is still limited. Despite the importance of volcanic eruptions, few studies have analyzed the effect of eruptions short after the event (Dale et al., 2005; Chrisafulli and Dale 2018). On the other hand, the effect of volcanism has been studied by comparing different stands in chronosequences assuming that different forest patches would reflect the successional stage after the disturbance (e.g. Drury and Nisbet 1973; Franklin and Swanson, 2010; Pickett 1988; Walker et al 2010).

Forest recovery is considerably slow following disturbances that heavily impact soils as well as aboveground vegetation (Chazdon, 2003). Despite volcanoes are considered as catastrophic disturbance event, it is possible that diverse biological legacies can remain after the event. The

persistence and heterogeneous distribution of these biological legacies might modulate and help the recolonization of the plants (Foster et al., 1998; Franklin et al., 1985; Turner et al., 1998). For example, the life-history traits of the pre-disturbed community will result in particular survivors perhaps able to re-colonize and expand over the new conditions on the disturbed environment (Foster et al., 1998; Turner et al., 1998). Therefore, the arrangement and the amount of the biological legacies would drive the rate and the pathways of the new colonizers.

One of the most active tectonic and volcanic areas in the world correspond to the subduction area of the Nazca and the South American plate (e.g., Stern, 2004), which also originated the longest and the second highest mountain range in the world, the Andes. The volcanoes in the Chilean Andes are quite active. The Southern and Austral Volcanic Zones comprise 74 active volcanoes since the Post-Glacial up to the present and at least 21 have had one large explosive eruption (Fontijn et al., 2014). In the southern portion of the Andes, it is located the South American Temperate rainforest (SATR) (Armesto et al., 1998). This cordillera plays a critical role in SATR dynamics, where current successional models after volcanic disturbance suggest the *Nothofagus* are dominant pioneer trees that benefit from disturbances to persist (Veblen et al., 2016). The SATR represent a biogeographical island, rich in endemic species and dominated by evergreen, broad-leaved species (Armesto et al., 1996; Villagran and Hinojosa, 1997, Arroyo et al., 2004). South of 40°S, extensive areas of old-growth forest dominate the mountainous landscape. These forests are not directly disturbed by modern human activity, and represent one of the last pristine areas with pre-industrial biogeochemical conditions (Perakis and Hedin 2002)

On May 1st 2008 started the last eruption of the Chaiten Volcano (42° 50'S), near the small town of Chaiten, in southern Chile. This eruption was the largest rhyolite eruption since the

great eruption of Katmai Volcano in 1912, being the first rhyolite eruption in which some attributes of its dynamics and impacts have been monitored (Major and Lara, 2013). The eruption consisted of an approximately 2-week-long explosive phase that generated as much as 1 km³ bulk volume tephra (~0.3 km³ dense rock equivalent) followed by an approximately 20-month-long effusive phase that erupted about 0.8 km³ of high-silica rhyolite lava that formed a new dome within the volcano's caldera (Major and Lara, 2013). Chaitén volcano was surrounded by extensive old-growth undisturbed forest by humans, providing a unique opportunity to evaluate the effects of volcanism over undisturbed landscape, then providing an opportunity to understand how volcanism may have affected the forest, under pre-industrial conditions.

In this study, we characterized the early establishment of the vegetation after the eruption of the Chaitén volcano along a disturbance gradient one year after the eruption started, from nearby the crater to the closest old-growth forest with vegetation completely alive. We predict that plant species diversity and the number of living trees decrease close to the crater, and pioneering *Nothofagus* species are colonizing the areas affected by the eruption. Our objectives included the creation of a baseline for further long-term monitoring of vegetation recovering, and to identify and analyze the importance of biological legacies for forest recovery.

2. Material and methods

2.1 Study site

The Chaitén volcano is located in the west side of the Andean Range in southern Chile, at 42° 50' S and 72° 39' W (Fig 1) and is part of the so-called Southern Volcanic Zones. The Chaitén

Volcano is a dome of 1100 m height, located at 10 km NW of the town of Chaitén, populated by approximately 5000 inhabitants before the eruption. The climate of the area corresponds to a humid temperate with a strong oceanic influence (Luebert and Plischoff, 2006). The landscape is characterized by sharp mountains shaped by glaciers and valleys originated by fluvial and glacio-fluvial outwash (Swanson et al., 2013). Vegetation within the influence area of the Chaitén volcano is dominated by a dense old-growth temperate rainforest of northern Patagonian and Valdivian types (Luebert and Plischoff, 2006; Swanson et al., 2013; Veblen et al., 1983; Veblen et al., 1996). These forests composition is dominated by broad-leaved evergreen trees such as *Nothofagus dombeyi* (Mirb.) Oerst. (Nothofagaceae), *Laureliopsis philippiana* (Looser) Schodde, *Gevuina avellana* (Molina), *Amomyrtus luma* (Molina) D. Legrand and Kausel, *Luma apiculata* (DC.) Burret, *Drymis winteri* J.R. Forst. and G. Forst., *Eucryphia cordifolia* Cav. and *Weinmannia trichosperma* Cav. (both Cunoniaceae). These species are normally arranged in multi-layered canopy strata and old, large-emergent trees (up to c. 40 m height) and abundant snags and logs. The understory is densely covered by ferns such as *Lophosoria quadripinnata* (J.F.Gmel.) C.Chr. and *Blechnum magellanicum* (Desv.) Mett., seedlings, saplings and bamboo tickets (*Chusquea* spp.).

129

Figure 1. Study area.

130

131

132 **2.2 Study design**

133 We carried out the field work in September 2009. We first conducted a general inspection of
 134 the study area (del Moral, 1981). Later, we selected four sites with different degrees of volcanic
 135 impact following similar classifications of previous studies (del Moral, 1981). The first site
 136 corresponds to a forest with low volcanic impact that only received few millimeters of tephra
 137 deposition. This site was called “Old-growth forest”. It was located 5 km from the center of the

new dome at an elevation of 600 m a.s.l. (Figure 1b). The second site was characterized by the presence of abundant dead standing trees, and higher tephra deposition. This site was called “Standing-dead” located at 3.5 km from the center of the new dome, at 160 m. The third site was more affected by the eruption, and was characterized by high tephra deposition, where most trees were broken and throw down by the eruption. This site was called “Blow down” and was located at 2.3 km from the center of the new dome at an elevation of 300 m a.s.l. Finally, the fourth and most affected site was in the border of the crater, at 1.5 km of the center of the new dome and at 750 m a.s.l. This site presented the higher tephra deposition and all trees were blow down by the eruption and heavily buried under volcanic ash. This site was called “Total destruction”.

In each one of these four areas was installed a 50 x 20 m permanent plot where we measured the diameter at the breast height (dbh) of all trees ≥ 5 cm. Every tree was identified at the species level when possible and marked. We registered if the stems were alive, resprouting after the eruption, or snag (standing dead). To estimate the presence and abundance of tree regeneration, we traced four transects of 120 m long. Transects were distributed parallel, separated 20 m from each other. In each transect, we mounted five circular plots of 1.5m radius (4.71m^2) every 30 m. On these plots we counted all seedlings and saplings present, and recording the substrate where they rooted (forest soil, logs, or volcanic tephra). We registered other non-tree plant species in the circular plots, estimated its abundance as the relative cover, and we also recorded the substrate where they rooted were organic or mineral (e.g., forest soil, logs, or volcanic tephra). We assessed the volume of all exposed logs in the plot, until each border within the plot. Assuming a cylindrical form of each log, we calculated the volume of

logs present inside the plot and standardizing its measurements to m³ of logs per m² of plot surface.

2.3 Data Analysis

Trees seedlings, herbs and shrubs species richness was compared among sites using rarefaction analysis (Colwell 2006). Rarefaction curves allow for comparing species richness among environments avoiding the bias in the number of species detected due to the increased probability of species detection in areas where individuals were more abundant, by plotting the number of species detected in function of the number of individuals observed (Colwell and Coddington, 1994, Gotelli & Colwell 2011, Colwell et al., 2012). This procedure includes Monte Carlo simulations delivering an average value of species richness with a confidence interval of 95% (Colwell 2006). Rarefaction analyses were conducted using the software EstimateS Win 8 (Colwell 2006). Non-vascular species were excluded from the rarefaction analyses since not all of them were classified up to the species level, and one genus could include several species. Plant composition among sites was compared using Sørensen's similarity index based on the presence/absence of plant species. Rarefaction and similitude analyses were both carried out with the software Estimates 8.2 (Colwell 2006). In attempting to define if the plant composition of one site was a subset of the composition of other site, we conducted a Nested Analysis in NestCalc free access software (Atmar and Patterson, 1995).

3. Results

3.1 General description of the effects of the eruption

The eruption of the Chaitén volcano has an impressive effect on the forest. The eruption in the study area produced a gradient of destruction. Around the crater all the trees were blown down

and heavily buried by tephra. Other trees were partially buried, broken in half, with the trunks covered with impacts of many little stones looking similar to the fire of gun shots (Fig x). Moving away and down from the crater, trees were blown down and densely covered by ash(Fig x), while further away trees were standing dead presenting progressively finer branches(Fig x). Many small patches of the organic litter were exposed and sparsely distributed and protected from the direct effect of the eruption by root discs and large trunks, or were exposed by small ravines that flush away the volcanic ash. These small patches looked like clumps of organic soil, with roots of ferns and other small plants that were not buried by the tephra. These clumps were covered by plants that survived in the area affected by the eruption, and their frequency increases when moving away from the crater towards the undisturbed forests.

Figure 2. Upper panel showing impressive impacts close to the crater. Comparison of the volcanic dome and vegetation cover before (bottom-left panel) and during the 2008-2009 eruption (bottom-right panel).

In the surrounding, we did not observed evidence of fire caused by the eruption, but the trunk and bark of several trees looked charred within the influence area. Trees were dead and dry by the eruptive explosions, but some trees in Blow-down and Standing-dead areas maintained the bark in the opposite side of the crater, perhaps with possibilities to resprouting hereafter. Tephra buried the fallen trees in the whole study area, but principally in Blow-down and Total-destruction areas. Tephra was very compacted, seeming as pavement that covered the fallen forests. Heterogeneous tephra deposition exposed some trunks, clumps and remnants with organic-forest soil likely from the original forest. Shrubs and epiphytes were spread everywhere in the Standing-dead area. Ferns, bamboo and tree seedlings were growing in remnants of exposed organic soil, usually below or at one side of trunks or unearth roots. Surviving trees also increases when moving away and down from the crater.

3.2 Species composition

We found 10 species of trees in the area, and 50 species of smaller plants, including seedlings, herbs and ferns (Table 1). Rarefaction analysis for tree species showed that the Standing-dead plot holds one more species than the Old-growth forest plot (Fig. 3). This same analysis showed the Blow-down plot holds very few individuals, but the initial slope and shape of the curve was similar to the other two curves. It indicates that at similar number of individuals, all curves showed a similar number of tree species. Rarefaction analysis of small vascular plants showed the Old-growth forest site had the richest species composition, while decreasing in the Standing dead and Blow-down plots (Fig. 3). The Total-destruction plot showed practically no species.

Table 1. List of plant species present in the Chaitén volcano and their frequency in the big plot and in 20 plots of 3.14m² in the different studied areas.

Families	Species	Old-growth	Standing dead	Blow-down	Near crater
SEEDLINGS					
Cunoneaceae	<i>Weinmannia trichosperma</i>		2		
	<i>Caldcluvia paniculata</i>	4	4		
Eucryphiaceae	<i>Eucryphia cordifolia</i>	5			
Flacourtiaceae	<i>Azara lanceolata</i>			1	
Monimiaceae	<i>Laureliopsis philippiana</i>	6			
Podocarpaceae	<i>Podocarpus saligna</i>	1			
Winteraceae	<i>Drimys winteri</i>	3			
Proteaceae	<i>Embothrium coccineum</i>		1	4	
	<i>Lomatia ferruginea</i>	5			
	<i>Gevuina avellana</i>	1			
Myrtaceae	<i>Myrceugenia parviflora</i>	3			
	<i>Myrceugenia planipes</i>	12			
	<i>Amomyrtus luma</i>	16	4		
Araliaceae	<i>Raukaua laetevirens</i>	3			
HERBS AND SMALL SHRUBS					
Gesneraceae	<i>Mitraria coccinea</i>	8			
	<i>Asteranthera ovata</i>	1	1		
	<i>Campsidium valdivianum</i>	5			
Poaceae	<i>Chusquea uliginosa</i>	7	1	1	
Apocinaceae	<i>Elytropus chilensis</i>	6	1		
Fitolaceae	<i>Ercilla syncarpellata</i>	3			
Saxifragaceae	<i>Ribes magellanicum</i>		1		
Bromeliaceae	<i>Greigia landbeckii</i>	4			
Griselineaceae	<i>Griselinia racemosa</i>	1			
	<i>Griselinia ruscifolia</i>	4	1	1	
Gunneraceae	<i>Gunnera tinctoria</i>			1	

Rubiaceae	<i>Nertera granadensis</i>	1		
Hydrangeaceae	<i>Hydrangea serratifolia</i>	12		
Luzuriagaceae	<i>Luzuriaga poliphylla</i>	8	1	1
	<i>Luzuriaga radicans</i>	14		
FERNS				
Dicksoniaceae	<i>Lophosoria quadripinnata</i>	6	4	3
Hymenophyllaceae	<i>Hymenophyllum tortuosum</i>			2
	<i>Hymenophyllum magellanicum</i>	2		
	<i>Hymenoglossum cruentum</i>	6		
	<i>Hymenophyllum caudiculatum</i>	3		
	<i>Hymenophyllum cuneatum</i>	2		
	<i>Hymenophyllum dentatum</i>	5		
	<i>Hymenophyllum dicranotrichium</i>	12		
	<i>Hymenophyllum krauseanum</i>	5		
	<i>Hymenophyllum pectinatum</i>	3		
	<i>Hymenophyllum plicatum</i>	3		
	<i>Serpilopsis caespitosa</i>	5		
Aspleniaceae	<i>Asplenium daeroides</i>	4		
	<i>Pleurosorus papaverifolius</i>	4		3
Dryopteridaceae	<i>Megalastrum spectabile</i>	7		
Blechnaceae	<i>Blechnum mochaenum</i>	1		
	<i>Blechnum arcuatum</i>	1		
	<i>Blechnum chilense</i>	2	1	1
	<i>Blechnum magellanicum</i>	1		
Dennstaedtiaceae	<i>Hypolepis rugosula</i>			1
Total general		201	22	20
				0

218

219

220

221

Figure 3. Rarefaction analysis showing the relationship between number of tree seedlings species (a) and small vascular plant species (b) as function of the number of individuals. Both panel the shaded area is indicating one standard error). All sites affected by the eruption of Chaitén Old-growth, Standing-dead and Blow-down are compared, except Total-destruction was not included because of the absence of plants.

222

223 Plant composition among Old-growth and Standing-dead plots showed some similarities (34%),

224 but less related with Blow-down (20%) and the Total-destruction plot (0%). Standing-dead and

225 Blow-down plots showed the highest similarities in species composition (Table 2). In the

disturbed plots, the plant species recorded were a subgroup of the species present in the Old-growth forest plot (Nestedness calculator $T = 13.85^\circ$). The most frequent plant was the fern *L. quadripinnata* that survived the direct impact of the eruption in small clumps of organic matter protected by trunks or root disc of trees, and were not completely covered by tephra. Bamboo and ferns were sprouting from roots in the remnants of exposed forest soil.

Table 2. Sørensen similarity index for the plant composition among the different sites in the Chaitén influence area.

Site	Old-growth	Standing dead	Blow down	Total destruction
Old-growth	1			
Standing dead	0.342	1		
Blow down	0.2	0.526	1	
Total destruction	0	0	0.1	1

3.3. Forest structure

The forest structure of the Old-growth site showed individuals of all sizes, with several large emergent trees (Fig. 4). Most trees in the Old-growth plot were alive and the snags most probably represent trees that died before the eruption, while in the other plots most trees were killed by the eruption. The Standing-dead plot showed similar dbh distribution than the Old-growth plot, with higher frequency of small dbh. The Blow-down plot showed a very small number of standing trees, concentrated in intermediate size class, while the Total-destruction plot practically had no individual standing trees (Fig. 4).

Figure 4. Diameter at the breast height (dbh) distribution for living and standing dead trees in the study sites influenced by the eruption of the Chaitén volcano.

standing dead trees in the Total destruction plot. The volume of fallen trees increased in the Blow-down and in Standing-dead plots, but it was much lower in the Old-growth and the Total-destruction plots (Fig. 5). In the Total-destruction plot most logs were heavily buried by a dense tephra layer.

Table 3. Total basal area in 1000 m² of tree species of the study plots in the Chaitén volcano area, classified as alive, resprouting after the eruption, or snag

	Total-destruction	Blow-down		Standing-dead		Old-growth	
	Snag	Snag	Resprouting	Snag	Resprouting	Snag	Alive
<i>Amomyrtus luma</i>				0,02	0,34		0,47
<i>Amomyrtus meli</i>					0,06		
<i>Caldcluvia paniculata</i>							2,54
<i>Drimys winterii</i>				0,14			
<i>Lomatia ferruginea</i>				0,,04	0,01		0,02
<i>Myrceugenia planipes</i>							0,84
<i>Eucryphia cordifolia</i>		0,55		0,59	0,61	0,26	6,37
<i>Laureliopsis philippiana</i>		0,25		0,87			9,64
<i>Nothofagus dombeyi</i>				3,46			
<i>Weinmannia trichosperma</i>		0,95	0,41		2,38		0,71
<i>not identified</i>	1,41	0,94		1,45			
Total	1,41	2,69	0,41	6,57	3,39	0,26	20,59

Figure 5. Volume of fallen dead trees (logs) in the study sites influenced by the eruption of the Chaitén volcano in the study sites.

3.4. Seedlings and small plants

We found a total of 90 seedlings from 5 trees and shrubs species in the disturbed plots, while we registered 174 seedlings of 12 species in Old-growth plots. The Standing-dead plots had 83 seedlings from five species, while the Blow-down plots presented 7 seedlings from two species. We did not find any seedling in the total destruction plot. In the Old-growth plots, seedlings were largely dominated by the shade tolerant *A. luma*, *M. parviflora* and *M. planipes*. In the Standing-dead plots, seedlings were dominated by *C. paniculata* (48% of the total), but also present the shade-tolerant tree *A. luma* (24%), the pioneer tree species *W. trichosperma* and *E. coccineum* (18% and 8% respectively). In the Blow-down plot, *E. coccineum*, a pioneer species, and *A. lanceolata* were the only recorded species.

We found different mechanism of forest regeneration in the plots. In the Old-growth plot most regeneration was from seeds, while in the Standing-dead plots most regeneration become from

resprout and few tree seedlings from seeds (Table 4). In the Old-growth plots, most of the tree seedlings corresponded to species with fleshy fruits dispersed by birds, while in the Standing-dead plots most seeds were wind-dispersed. In the Blow-down plots, all seedlings came from seeds (Table 4), but a half of the seedlings corresponded to species dispersed by birds (e.g. *Ribes magellanicum*) and the other half were species dispersed by wind (e.g. *E. coccineum*). Second, in all plots most seedlings were growing on organic substrata (Table 4), such as organic soil on the ground, root discs and logs, while few seedlings were growing directly either on the tephra or on the rocks. In the Blow-down plot we found 7 seedlings of *E. coccineum*, growing just in remnants of organic litter; in Standing-dead plot 11 seedlings growing directly in the tephra, while the other 72 seedlings were resprouting directly from the standing tree's trunks. In the Old-growth plots we found just one species growing in an exposed rock while the rest of the seedlings were growing in organic substrates (Table 4).

Table 4. Number of seedlings per regeneration type and substratum type in the plots.

	Regeneration type		Substratum type	
	Resprouting	Seed	Mineral	Organic
Total-destruction	-	-	-	-
Blow-down	-	7	-	7
Standing-dead	74	9	11	72
Old-growth	31	127	1	157

Other small plant species in the studied plots (Table 1) showed differences along the disturbance gradient (Table 2). For example, the Old-growth plot has species in all categories of plant cover, with few species covering over 75% of the ground, and many species covering between 1-5% (Fig. 6). In contrast, Standing-dead and Blow-down plots were covered by fewer species, and

287 concentrated in the lower ranges of coverage (Fig. 6). In the Total destruction plot we only
 288 found a non-vascular plant (*Marchantia* sp.) with a cover between 5-25 % (Fig. 6) growing on
 289 a log exposed to the surface. The dominant understory species in the Old-growth forest were
 290 the bamboo *Chusquea* spp. and the fern *L. quadripinnata*. These species were also present in
 291 the Standing-dead and Blow-down plots, although scattered and with a low percent of coverage
 292 in the disturbed plots.

Figure 6. Plant cover (including vascular and non-vascular species) at the ground level of the study areas affected by the Chaitén volcano.

4. Discussion

4.1 Early vegetation establishment: the relevance of biological legacies

Few studies have been conducted short after an eruption in the southern Andes, however successional model predict that disturbed areas in low and mid altitude (<1000 m a.s.l.) forests would be dominated by *Nothofagus* species due to the shade-intolerance (e.g. Veblen and Ashton, 1978). However, in this study after the first year we did not register *Nothofagus* species despite were abundant in the surroundings non disturbed areas. It might be due to cyclic seed production with not constant seed rain along the years (Donoso, 1993; Veblen, 1985), and the short-term and limited spatial extension we were able to monitor the vegetation. However, our results pointed that regeneration occurs mainly due to an *in situ* development of plants associated to biological legacies such as logs, soil clumps and individuals that survive to the eruption. Other areas disturbed by Chaiten volcano presented similar patterns of types and abundance of biological structures (Swanson et al., 2013).

The initial establishment of the tree species, beside of the disturbance gradient, may be influenced by the interaction between life history of the species and the altitudinal gradient. Life history attributes such as seed dispersal, flowering phenology, growth form, resprouting ability or light-temperature stress resistance are a mechanism that influence the recovery response of the post-disturbance vegetation (Eriksson 2000; McIntyre et al., 1994; McLachlan and Bazely, 2001; Pausas and Lavorel, 2003; Tilman, 1997). In the disturbed plots, the principal mechanism of regeneration was the resprouting capability of some species (Tables 1 and 4). This is a frequent strategy in many plant species in response to disturbances of different types and intensities (Bellingham and Sparrow, 2000; Bond and Midgley, 2001). The seed dispersion syndrome could be also an important mechanism that influence the colonization after disturbance. Chilean forests have one of the highest proportions of plant dispersed by

frugivorous birds compared to other temperate rainforests (Armesto et al. 1987, Willson et al., 1996). Birds consume a high proportion of fruits and may disperse the seed to specific places, such as to a perch (Armesto et al., 2001; Lindenmayer and Franklin, 2002). The seed availability might increase in sites where remnant snags, shrubs or perching trees were available (e.g., Bustamante-Sánchez, 2011). Finally, among the wind dispersed seeds, we only found seedlings of *E. coccineum* in disturbed plots. While this tree is adapted to stressful condition, the seed is able to disperse for long-distances (Donoso, 2006).

For non-tree species, Old-growth plot had more species than Standing-dead plot. On the contrary, for tree species Standing-dead plot is richest than Old-growth, revealing the effect of the eruption on the small plant species. On the other hand, in Blow-down and in Total-destruction plot, i.e. from mid to high disturbance intensity at mid to high altitude, and without or sparse biological legacies, we found low species richness and coverage, and sparse tree regeneration. This mosaic of legacies in the landscape offers opportunities for the recolonization of many different species in the affected area. The different species composition would promote different pathways to early succession with different communities, but also different recovery rates. We propose that in low to mid elevation areas, the vegetation recovery will be faster, dominated by species shade-tolerant to semi-tolerant and dispersed by birds, mostly due to biological legacies. On the other hand, at higher altitude, the vegetation recovery would be slower, recolonization made by shade-intolerant and seed dispersed by winds because the most stressful conditions of temperature at higher altitudes.

In synthesis, the results point to an early succession with (i) a mosaic of pioneering-wind dispersed species, scattered survivors regrowing and spreading from biological legacies, and

plant species dispersed by frugivorous birds, likely favored by the biological legacies; (ii) the early succession is influenced by the interaction of the species-specific life history (including invasion by exotic species), altitudinal gradient and the different intensity of disturbance in a broad area.

Acknowledgements

Field work was financed by CONICYT PDA-24. A.L. and D.C. acknowledge support from grant CONICYT/FONDAP/15110009. RM acknowledge the scholarship DAAD/Becas Chile during late stages of this manuscript.

REFERENCES

- Armesto, J. J., Rozzi, R., Miranda, P., & Sabag, C. (1987). Plant/frugivore interactions in South American temperate forests. *Revista Chilena de Historia Natural*, 60(2), 321-336.
- Aravena, J.C., Carmona, M., Perez, C.A., Armesto, J.J., 2002. Changes in tree species richness, stand structure and soil properties in a successional chronosequences in northern Chiloé Island, Chile. *Revista Chilena de Historia Natural*. 75, 339-360.
- Armesto, J.J., Arroyo, M.T.K., Villagran, C., 1996. *Ecología de los Bosques Nativos de Chile*. Editorial Universitaria, Chile. 470pp.
- Armesto, J.J., Díaz, I.A., Papic, C., Willson, M.F., 2001. Seed rains of fleshy and dry propagules in different habitats in the temperate rainforest of Chiloé Island, Chile. *Austral Ecology* 26, 311-320.
- Armesto, J.J., Pickett, S.T.A., 1991. Foreword. *Revista Chilena de Historia Natural* 644, 389.
- Arroyo, M.T.K., P. Marquet, C. Marticorena, J. Simonetii, L. Cavieres, F.A. Squeo & R. Rozzi, 2004. Chilean winter rainfall-valdivian forests. In (Mittermeier, R.A., P. Robles Gil, M. Hoffmann, J. Pilgrim, T. Brooks, C. Goettsch, J. Lamoreux & G.A.B. da Fonseca, eds), "Hotspots Revisited", pp. 99-103. CEMEX, México.
- Atmar, W., Patterson, B.D., 1993. The measure of order and disorder in the distribution of species in fragmented habitat. *Oecologia* 96, 373-382.
- Atmar, W., Patterson. B.D., 1995. The nestedness temperature calculator: a visual basic program, including 294 presence-absence matrices. AICS Research, Inc., University Park, NM, and The Field Museum, Chicago.
- Bellingham, P., Sparrow. A.D., 2000. Resprouting as a life-history strategy in woody plant communities. *Oikos* 89(2), 409-416.
- Bond, W.J., Midgley, J.J., 2001. Ecology of sprouting in woody plants: the persistence niche. *Trends in Ecology & Evolution* 16(1), 45-51.
- Bustamante-Sánchez, M.A., Armesto, J.J., Halpern, C.B., 2011. Biotic and abiotic controls on tree colonization in three early successional communities of Chiloé Island, Chile. *Journal of Ecology* 99(1), 288-299.
- Chazdon, R.L., 2003. Tropical forest recovery: legacies of human impact and natural disturbances. *Perspective in Plant Ecology, Evolution and Systematics* 6, 51-71.
- Colwell, R.K., Chao, A., Gotelli, N.J., Lin, S., Mao, C., Chazdon, R.L., Longino, J.T., 2012. Models and estimators linking individual-based and sample-based rarefaction, extrapolation and comparison of assemblages. *Journal of Plant Ecology* 5, 3-21.

- 400 Colwell, R.K., and Coddington, J.A., 1994. Estimating terrestrial biodiversity through
401 extrapolation. *Phil. Trans. R. Soc. Lond. B* 345: 101-118.
- 402 Crews, T., Kitayana, K., Fownes, J., Riley, R., Herber, D., Mueller-Dombois, D., Vitousek,
403 P., 1995. Changes in soil phosphorus fractions and ecosystem dynamics across a long
404 chronosequence in Hawaii. *Ecology* 76: 1407-1424
- 405 Dale V.H., Swanson, F.J., Crisafulli, C.M., 2005. Ecological responses to the 1980 eruption
406 of Mount St. Helens. Springer, 342pp.
- 407 del Moral, R. 1981. Initial life returns to Mount St. Helens. *Natural History* 90, 36-49.
- 408 del Moral, R., Grishim, S., 1999. Volcanic Disturbance and Ecosystem recovery, in: L.R.
409 Walker (Ed.) *Ecosystem of disturbed Ground*. Elsevier, pp. 137-160.
- 410 Dinerstein, E., Olson, D., Graham, D., Webster, A., Primm, S., Bookbinder, M., Ledec, G.,
411 1995. A conservation assessment of the terrestrial ecoregions of Latin America and the
412 Caribbean. The International Bank for Reconstruccion and Development. Waschington, DC:
413 The World Bank.
- 414 Donoso, C., 1993. Bosques templados de Chile y Argentina, variación, estructura y dinámica.
415 Tercera edición. Editorial Universitaria, Santiago-Chile, 484p.
- 416 Donoso, C., 2006. Las Especies arbóreas de los Bosques Templados de Chile y Argentina.
417 Autoecología. Marisa Cúneo Ediciones, Valdivia-Chile, 678p.
- 418 Donoso, C., Lara, A., 1999. Silvicultura de los Bosques Nativos de Chile. Editorial
419 Universitaria, Santiago-Chile, 421p.
420
- 421 Donoso PJ, Promis A (2013) Silvicultura en Bosques Nativos. Avances en la investigación en
422 Chile, Argentina y Nueva Zelandia. Estudios en Silvicultura de Bosques Nativos, Vol. 1. Ed.
423 Marisa Cuneo, Valdivia, Chile. p 253
- 424 Eriksson, O., 2000. Funtional roles of remnants plant population in communities and
425 ecosystem. *Global Ecology and Biogeography* 9, 443-449.
- 426 Foster, D., Knight, D., Franklin, J.F., 1998. Landscape patterns and legacies resulting from
427 Large, Infrequent forest disturbance. *Ecosystems* 1, 497-510.
- 428 Franklin, J.F., 1990. Biological Legacies: a critical Management concept from Mount St.
429 Helens. *Transactions 55th North American Wildlife and Natural Resources Conference*, 216-219.
- 430 Franklin, J.F., Armesto, J.J., 1996. La retención de elementos estructurales del bosque durante
431 la cosecha: Una alternativa de manejo para los bosques nativos chilenos. *Ambiente y*
432 *Desarrollo* 12 (2), 69-79.
- 433 Franklin, J. F., MacMahon, J. A., Swanson, F. J., Sedell, J. R., 1985. Ecosystem responses to
434 the eruption of Mount St. Helens. *National Geographic Research* 1:198-216.

- 435 Franklin, J.F., Mitchell, R.J., Palik, B.J., 2007. Natural disturbance and Stand development
436 principles for ecological forestry. United States Department of agriculture, Forest Service,
437 Northern Research Station. General Report NRS-19. 48p.
- 438 Franklin, J.F., Lindenmayer, D., MacMahon, J.A., McKee, A., Magnuson, J., Perry, D.A.,
439 Waide, R., Foster, D., 2000. Threads of continuity: ecosystem disturbance, recovery, and the
440 theory of biological legacies. *Conservation Biology in Practice* 1, 8-16.
- 441 Franklin, J.F., Spies, T.A., van Pelt, R., Carey, A., Thornburgh, D.A., Rae Berg, D.,
442 Lindenmayer, D.B., Harmon, M.E., Keeton, W.S., Shaw, D.B., Bible, K., Chen, J., 2002.
443 Disturbance and structural development of natural forest ecosystem with silvicultural
444 implications, using Douglas-fir forest as an example. *Forest Ecology and Management* 155,
445 399-423.
- 446 Franklin, J.R., Swanson, M.E., 2010. Long-term ecological research in the forest of the
447 United States: key lessons for its application in Chile and around the world. *Revista Chilena*
448 *de Historia Natural* 83, 185-194.
- 449 Gallardo, M.B., Pérez, C., Nuñez-Avila, M., Armesto, J.J. 2012. Desacoplamiento del
450 desarrollo del suelo y la sucesión vegetal a lo largo de una cronosecuencia de 60 mil años en
451 el volcán Llaima, Chile. *Revista Chilena de Historia Natural* 85: 291-306
- 452 Gotelli, N.J., Entsminger G.L., 2005. EcoSim: Null models software for ecology. Version 5.0.
453 Acquired Intelligence Inc. & Kelsey-Bear.
454 <http://homepages.together.net/~gentsmin/ecosim.htm>.
455
- 456 Grosse, H. y Quiroz, I. 1999. Silvicultura de los bosques de segundo crecimiento de roble, raulí
457 y coigüe en la región centro-sur de Chile. p 95-125. *Silvicultura de los bosques nativos de Chile*.
458 Editorial Universitaria. 421p
- 459 Haberle, S.G., Szeicz, J.M., Bennett, K.D., 2000. Late Holocene vegetation dynamics and
460 lake geochemistry at Laguna Miranda, XI Region, Chile. *Revista Chilena de Historia Natural*
461 73, 655-669.
- 462 Johnson, E.A., Miyanishi, K., 2008. Testing the assumptions of chronosequences in
463 succession. *Ecology Letters* 11: 419-431.
- 464 Kitzberger, T., Veblen, T.T, Villalba, R., 1995. Tectonic influences on tree growth in northern
465 Patagonia, Argentina: the roles of substrate stability and climatic variation. *Canadian Journal*
466 *of Forest Research* 25, 1684-1696
- 467 MacArthur, R.H., Wilson, E.O., 1967. The theory of island biogeography. Princeton
468 University Press, Princeton
- 469 Major, J., Lara, L.E., 2013. Overview of Chaiten Volcano, Chile, and its 2008-2009 eruption.
470 *Andean Geology* 40(2), 196-215.

- 471 McIntyre, S., Lavorel, S., Tremont, M., 1994. Plant Life-History Attributes: Their
472 relationship to disturbance response in herbaceous vegetation. *Journal of Ecology* 83(1), 31-
473 44.
- 474 McLachlan, S.M., Bazely, D.R., 2001. Recovery patterns of understory herbs and their use
475 as indicators of deciduous forest regeneration. *Conservation Biology* 15, 98-110.
- 476 Müller-Dombois, D., Ellenberg, H., 1974. Aims and methods in vegetation ecology. The
477 Blackburn Press. 580p.
- 478 Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, A.B., Kent, J., 2000.
479 Biodiversity hotspot for conservation priorities. *Nature*, 402, 853-58.
- 480 Lara, L.E., 2009. The 2008 eruption of the Chaitén Volcano, Chile: a preliminary report.
481 *Andean Geology* 36(1), 125-129.
- 482 Lindenmayer, D.B., Franklin, J.F., 2002. Conserving Forest Biodiversity: A Comprehensive
483 Multiscaled Approach. Island Press, Washington, 351p.
- 484 Luebert, F., Pliscoff, P., 2006. Sinopsis bioclimática y vegetacional de Chile. Editorial
485 Universitaria, Santiago-Chile, 316p.
- 486 Pausas, J.G., Lavorel, S., 2003. A hierarchical deductive approach for functional types in
487 disturbed ecosystem. *Journal of Vegetation Sciences* 14(3): 409-416.
- 488 Peet, R., 1992. Community structure and ecosystem function, in: Glenn-Lewin D., Peet R.,
489 Veblen, T. (Eds.), *Plant succession theory and prediction*. Chapman & Hall, London, pp. 103-
490 151.
- 491 Perakis, S.S., Hedin L., 2002. Nitrogen loss from unpolluted South American forest mainly
492 via dissolved organic compounds. *Nature* 415, 416-419.
- 493 Pickett, S.T.A., Collins, S.L., Armesto, J.J., 1987. Models, mechanism and pathways of
494 succession. *The Botanical Review* 53(3), 335-371.
- 495 Pickett, S.T.A., 1988. Space-for-time substitution as an alternative to long term studies, in:
496 Likens, G.E. (Ed.), *Long-term Studies in Ecology*. Springer, New York, NY, pp. 110–135.
- 497 Robock, A., Oppenheimer, C., 2003. Volcanoes and Earth's Atmosphere. Geophysical
498 monograph 139.
- 499 Runkle, J.R., 1985. Disturbance regimes in temperate forests, in: Pickett, S.T.A., White, P.S.
500 (Eds.), *The Ecology of Natural Disturbance and Patch Dynamics*. Academic Press, Orlando,
501 FL, pp. 17-33
- 502 Stern, C.R., 2004. Active Andean volcanism: its geologic and tectonic setting. *Andean*
503 *Geology*. 31(2), 161-206

- 504 Swanson, F., Franklin, J., 1992. New Forestry principles from ecosystem analysis of Pacif
505 Northwest Forest. *Ecological applications* 2(3), 262-274
- 506 Swanson, F., Jones, J., Crisafulli, Ch., Lara A., 2013. Effects of volcanic and hydrology
507 processes on forest vegetation: Chaiten Volcano, Chile. *Andean Geology* 40(2), 359-391
- 508 Swanson, M., Franklin, J., Beschta, R., Crisafulli, C., DellaSala, D., Hutto, R., Lindenmayer,
509 D., Swanson, F., 2011. The Forgotten stage of of forest succession: Early-successional
510 ecosystems on forest site. *Biological Sciences Faculty Publications*. Paper 278.
511 http://scholarworks.s.umt.edu/biosci_pubs/278
- 512 Tilman D., 1997. Community invasibility, recruitment limitation, and grassland biodiversity.
513 *Ecology* 78, 81–92
- 514 Turner, M., Dale, V., 1998. Comparing large, infrequent disturbance: what have we learned?.
515 *Ecosystem* 1, 493-496.
- 516 Turner, M., Dale, M., Everham, E., 1997. Fires, hurricanes, and volcanoes: Comparing large
517 disturbances. *BioSciences* 47(11), 758-768.
- 518 Veblen, T.T., 1985. Stand dynamics in Chilean *Nothofagus* forest, in: Pickett, S.T.A., White,
519 P.S. (Eds.), *The ecology of natural disturbance and patch dynamics*. Academic Press,
520 Orlando, FL, pp 35-51.
- 521 Veblen, T.T., Ashton, D.H., 1978. Catastrophic influences on the vegetation of the Valdivian
522 Andes, Chile. *Vegetatio* 36(3): 149-167.
- 523 Veblen, T.T., Ashton, D.H., Schlegel, F., Veblen, A., 1977. Plant succession in a timberline
524 depressed by vulcanism in South-Central Chile. *Journal of Biogeography* 4: 275-294.
- 525 Veblen, T.T., Donoso, C., 1987. Alteración natural y dinámica regenerativa de las especies
526 chilenas de *Nothofagus* de la Región de Los Lagos. *Bosque* 8(2), 132-142.
- 527 Veblen, T.T., Donoso, C., Kitzberger, T., Rebertus, A.J., 1996. Ecology of southern Chilean
528 and Argentinean *Nothofagus* forests, in: Veblen T.T., Hill, R.S., Read, J., (Eds.) *The ecology
529 and biogeography of Nothofagus forests*. Yale University Press, New Haven, Connecticut,
530 USA, pp. 293-353
- 531 Veblen, T.T., Donoso, C., Schlegel, F., Escobar, B., 1981. Forest dynamics in South-Central
532 Chile. *Journal of Biogeography* 8, 211-247.
- 533 Veblen, T.T., Schlegel, F., Oltremari, J., 1983. Temperate broad-leaved evergreen forest of
534 South America, in: Ovington, J.D., (Ed.) *Temperate broad-leaved evergreen forest*. Elsevier,
535 Amsterdam, pp 5-31.
- 536 Villagran, C., Hinojosa, F., 1997. Historia de los Bosques Templados de Sudamerica, II:
537 Análisis fitogeográfico. *Revista Chilena de Historia Natural*, 70, 241-267.

538 Willson, M.F., De Santo, T. L., Sabag, C. E., Armesto, J.J., 1996. Avian communities in
539 temperate rainforest of North and South America. In: Lawford, R., Alaback, P. B., Fuentes,
540 E., (Eds.) High-latitude rainforests and associated ecosystems of the west coast of the
541 Americas. Springer, New York. pp. 228–247