1 **Title:** Mosaic evolution, preadaptation, and the evolution of evolvability in apes

2 **Author:** Caroline Parins-Fukuchi*¹

3 ¹Department of Geophysical Sciences, University of Chicago, Chicago, IL, USA

4 *Author for correspondence: parinsfukuchi@uchicago.edu

5 **Data archived online:** https://figshare.com/articles/parins-fukuchi_supp_zip/8063726

6

7 **Acknowledgements**

8 I wish to thank D Alvarado-Serrano, M Cosman, CW Dick, DC Fisher, M Foote, E

9 Greiner, D Jablonski, LM MacLatchy, H Marx, J Saulsbury, SA Smith, G Slater, N

10 Walker-Hale, M Webster for conversations that have greatly benefitted this work. I also

11 thank S Worthington for help accessing his skeletal measurements that comprised a large

12 portion of the character matrix analyzed here. The manuscript benefitted immensely from

13 careful reviews provided by P Wagner and PD Polly.

14

15

16

17

18

19

20

21

22

Abstract: A major goal in post-synthesis evolutionary biology has been to better understand how complex interactions between traits drive movement along and facilitate the formation of distinct evolutionary pathways. I present analyses of a character matrix sampled across the haplorrhine skeleton that revealed several modules of characters displaying distinct patterns in macroevolutionary disparity. Comparison of these patterns to those in neurological development showed that early ape evolution was characterized by an intense regime of evolutionary and developmental flexibility. Shifting and reduced constraint in apes was met with episodic bursts in phenotypic innovation that built a wide array of functional diversity over a foundation of shared developmental and anatomical structure. Shifts in modularity drove dramatic evolutionary changes across the ape body plan in two distinct ways: 1) an episode of relaxed integration early in hominoid evolution coincided with bursts in evolutionary rate across multiple character suites; 2) the formation of two new trait modules along the branch leading to chimps and humans preceded rapid and dramatic evolutionary shifts in the carpus and pelvis. Changes to the structure of evolutionary mosaicism may correspond to enhanced evolvability that has a ‘preadaptive’ effect by catalyzing later episodes of dramatic morphological remodeling.

Keywords: integration, evolvability, preadaptation, modularity, hominoidea

Introduction

The formation of new species and phenotypes is often hypothesized to be limited by the availability of unoccupied ‘adaptive zones’, classically represented as peaks on a fitness landscape (Wright 1932, Simpson 1944). Frequently referred to as ‘ecological opportunity’ (Losos 2010), the opening of new adaptive zones through shifts in biotic interactions and the environment is often invoked to explain large evolutionary radiations (Rainey and Travisano 1998, Yoder et al., 2010, Wagner and Harmon et al 2012, but see Slater 2015). The role of ecological processes in generating morphological novelty can be contrasted with an increasing focus on ‘constructional’ factors that include functional interactions and architectural constraints (Seilacher 1970, 1991). While character evolution may sometimes reflect simple optimization processes when natural selection operates independently on separate traits, the evolutionary pathways realized in nature are also constrained by the complex interactions between functionally dependent traits and the limitations of an organism’s developmental architecture. When placed into a broader evolutionary literature, these limiting factors might be coarsely partitioned into developmental (Gould and Lewontin 1979, Alberch 1980, Olson 2012) and functional (Charlesworth et al., 1982, Maynard-Smith et al., 1985, Arnold 1992) constraints.

Developmental and functional constraints limit the capacity for a clade to evolve novel phenotypes. Together, these factors might be said to determine the level of ‘constructional opportunity’ available at a given time. Contrasted to ecological opportunity, constructional opportunity reflects how the structure of developmental and functional interactions between characters limits the ‘evolvability’ of a clade—its capacity to evolve novel phenotypes. While ecological opportunity is often evoked as a

limiting factor on a clade's phenotypic diversification, constructional opportunity, as measured by the overall level of constraint and internal integration between phenotypes, may also limit the range of evolutionary modifications available to a population faced with ecological pressure.

Evolutionary constraints are shaped by shared developmental pathways, multivariate selection, and underlying genetic processes such as epistasis and pleiotropy. Such interactions between traits and genes often create a common tendency to drive the formation of modules of characters that are internally integrated in their evolutionary trajectories. Such integration is typically identified by calculating numerical correlations in their variance (Olson and Miller 1958). Integration between traits constrains the combination of phenotypes accessible to each. For example, for two traits that are strongly positively integrated, it is unlikely to observe an individual that possesses phenotypes for each trait that lie at opposing ends of the spectrum of variation. As long as this correlation exists, the evolution of the organism is constrained in the sense that only certain combinations of phenotypes are accessible. At the population level, integration can be caused by multivariate selection or physical/constructional linkages such as pleiotropy or allometry (Cheverud 1984). However, it is generally not possible to distinguish between these in macroevolutionary study systems.

Phenotypic integration generally occurs in a modular fashion (Wagner et al. 2007). Instead of covarying uniformly across the entire organism, integration often manifests as a patchwork of modules of internally integrated traits that are free to evolve largely independently (Goswami et al. 2014, 2015). Such modular structures lead to the

pattern of mosaic evolution when separate modules evolve at different rates as lineages diverge (Felice and Goswami 2017, Parins-Fukuchi, in press). When analyzed phylogenetically, these patterns result in ‘mosaic disparity’—a scenario where individual modules of traits each display a unique pattern in phylogenetic disparity across lineages (Fig. 1).

The related concepts of mosaicism, constraint, and integration can help to explain the emergence of phenotypic novelty across clades. For example, when ecological opportunity is abundant, evolutionary radiations can be driven primarily by the release of constraints (Wagner et al. 2003). In addition, bursts in evolutionary rate can also coincide with the break-up of existing modules and the subsequent formation of a new modular structure (Wagner 2018). Characterizing the role of integration and mosaic evolution in shaping diversity across clades will ultimately contribute to a deeper understanding of phenotypic evolution at the constructional level. This may improve our understanding of evolutionary processes more generally by revealing the extent to which fluctuations in constructional opportunity contribute to macroevolutionary patterns alongside ecological opportunity.

In this study, I sought to determine the extent to which the evolutionary trajectories blazed by ape species were shaped by early shifts in the structure of integration and the timing of development. To address these questions, I reconstructed the mosaic macroevolutionary patterns in morphological, neurological, and developmental phenotypes across living and fossil great apes throughout the Miocene. To help place these empirical results in a theoretical context, I also present a set of simple simulations

to generate a theoretical expectation for the patterns in disparity expected when Markovian diffusion is constrained by trait integration imposed by both functional and structural/developmental interactions. Taken together, these analyses suggest the capability for shifting patterns in constraint imposed by integration to create constructional opportunities that generate repeated and dramatic episodes in morphological innovation.

Apes have been extensively studied in the context of both developmental constraint and environmental adaptation. Young et al. (2010), found that apes display weak integration between the fore- and hindlimb in comparison to cercopithecoids. Reduced integration can confer greater functional flexibility, and so may have facilitated the proliferation of the diverse locomotor function across living and fossil apes. Changes in developmental timing have also been important in ape evolution. Developmental heterochrony appears to have driven the emergence of many key aspects of hominin morphology in the cranium and post-cranium (Gould 1977, Berge 1998). This long-standing interest in the developmental basis of ape evolution has not excluded environmental hypotheses. Environmental fluctuations throughout the Miocene and Pliocene have also frequently formed the basis for adaptive hypotheses across apes (Andrews 1992, Ruff 1994, Michel et al., 2014). The body plans of hominins, and undoubtedly apes in general, have been shaped through modifications to separate suites of traits occurring at different times (Holloway 1973, McHenry 1975). These diverse threads suggest that a multiplicity of causes, perhaps including both ecological opportunity and constructional factors, has shaped the evolution of hominoid body plans.

As a result, apes are an excellent exemplar taxon in which to pluralistically examine the mosaic patterns in constraint and innovation that shape body plans in living and fossil vertebrates.

Methods

Morphological data

I gathered a dataset of 149 quantitative morphological traits from the literature (Lewton 2010, Worthington 2012) spanning the cranium, forelimb, and pelvis. These traits were sampled across 10 extant taxa. I also collected fossil cranial and forelimb traits from Worthington (2012), retaining all taxa with at least 25% matrix occupancy. Taxa were all studied at the generic level, with each extant genus represented by a single exemplar species. Each morphological trait represented either a dimensionless ratio or geometric mean, depending on how the author of each dataset handled intraspecific variation and body mass correction. I scaled the traits to display an empirical variance of 1 across all taxa. This simplified the analyses by reducing the complexity of the dataset, while retaining the same comparative information. Since the original traits were dimensionless, both the unscaled and the scaled datasets would facilitate examination of *relative*, rather than absolute, evolutionary rates, but the scaled traits improve Markov-chain Monte Carlo (MCMC) mixing and simplify the identification of mosaic suites.

Neurological data

To directly study the evolutionary patterns in related developmental and phenotypic traits, I also gathered neurological data from the literature (Capellini et al., 2010, Boddy et al., 2012). For neurological phenotypes, I collected encephalization

quotients (EQ) estimated in 76 primate species by Boddy et al. (2012). EQ is calculated by fitting a nonlinear model to brain size and body size and taking the deviations (residuals) from the best fit curve and so measures the enlargement of brain size while controlling for allometry with body size. I compared EQ patterns to those in a postpartum encephalization development (ED) metric calculated as:

$$ED = EQ_{adult} - EQ_{neonatal} \quad (1)$$

ED is therefore intended to reflect the amount of brain tissue grown by members of each species following birth relative to their body size. The ED metric was scaled to unit variance across all sampled taxa. Neonatal brain and body mass data representing 24 primate species were acquired from Capellini et al. (2010).

Fossil placement and divergence time estimation

Since the evolutionary relationships between the extant taxa represented in the morphological dataset have all been extensively studied and are well-resolved, I used these as a fixed scaffolding to place the fossil taxa. Fossil placements were inferred from the continuous trait dataset using the cophymaru program (Parins-Fukuchi 2018). I used the ‘binary weights’ procedure to filter out reliable traits from ones likely to mislead by using the extant tree as a fixed point of reference. Markov-chain Monte Carlo (MCMC) simulations were run for 1,000,000 generations and checked manually for convergence. The posterior tree sample was summarized as the maximum clade credibility tree using sumtrees.py (Sukumaran and Holder 2010).

For downstream comparative analyses, I also estimated divergence times on the full 19-taxon tree. I downloaded cytochrome B sequences from Genbank representing

each of the 10 extant taxa. Molecular dating was performed using Beast version 2.4.8 (Bouckaert et al. 2014) using the fossilized birth-death (FBD) prior. The topology was fixed to the summary tree generated during the fossil placement step. Temporal occurrence ranges for the fossil taxa were assembled from the literature. These fossil and extant occurrence ranges were used in the dating analysis to infer the diversification and sampling rate parameters used in the FBD prior.

Measuring integration throughout the hominoid radiation

As an evaluation of the overall strength of integration in the skeletal traits across the haplorrhine phylogeny, I estimated ancestral states for each character under Brownian motion along a dated tree of nine extant haplorrhine taxa (*Pan paniscus* was removed because it was missing pelvic measurements). I then estimated the strength with which the reconstructed phenotypes at each internal node were correlated with the values at each descendant node or tip. The correlations indicated how strongly the reconstructed evolutionary changes in skeletal traits were integrated along each branch in the tree. Stronger correlations corresponded to tighter overall integration. The Python script that I developed for the reconstruction of ancestral states and calculation of character correlations is available in the data supplement.

Evolutionary rates across mosaic morphological suites

I used the *greedo* program to recover mosaic patterns in disparity from the continuous trait dataset (Parins-Fukuchi 2019). *Greedo* is a phylogenetic clustering/mixture model estimation approach that uses the Akaike information criterion (AIC) to iteratively merge and split clusters of traits to find a best-supported set of

character suites based on shared patterns in disparity (i.e., where integrated traits within the same module are similarly disparate across taxa), where each is represented by a tree with branch lengths (Fig. 1). Traits that display the highest improvement in log-likelihood when assigned to a separate cluster are prioritized during the splitting steps to first identify likely sources of heterogeneity in the dataset. During the merging step, only clusters that show an improved AIC score when placed under the same model are joined. This process recovers an estimate of the number of suites, the membership of each trait, and a tree with branch lengths scaled to units of disparity for each suite. Fossil taxa were not included in the mosaic analyses, because they were too fragmentary in their sampling to inform disparity patterns when the traits were split into suites. I performed many runs of the *greedo* procedure to avoid presenting results obtained from a suboptimal peak on the likelihood surface. Since several of the top clusterings yielded close AIC support, I performed an additional model-averaging step using Akaike weights to summarize the results as a graph where each trait occupies a node where each link is weighted to reflect the total weighted AIC support across all the models. Comparative analyses were performed using a single summary clustering that resulted from application of the Markov clustering (MCL) algorithm to the support network (Dongen 2000).

To examine the anatomical composition of the mosaic suites of characters, I performed a statistical enrichment procedure to identify skeletal regions that were over and under-represented in each suite relative to the proportions for each present in the entire dataset. The proportion of the 149-trait dataset represented by each skeletal region was used to generate a set of ‘expected’ values. I then compared these to the observed

proportions within each suite to examine trends in the deviations of the proportions occupied by each region. I did not assess the statistical significance of these deviations, because 1) my aim was to examine the general patterns in the composition of each character suite rather than to determine absolute significance, and 2) the small sample size of the dataset generated expected chi-square test statistic values that were nearly entirely <5 , making the test inappropriate. To examine the relative contributions of separate anatomical regions in shaping the macroevolutionary patterns reconstructed in each suite, I also performed a principal component analysis (PCA) on the traits contained within each suite. I then transformed the loadings to calculate the total variance contributed by each trait across all axes of the PCA, summing them to produce the contribution of each anatomical region.

I transformed the branch lengths for each mosaic suite, which were scaled to reflect disparity (total variance, v , accumulated over time t), to rates (v per unit t) using the results from the divergence time analysis. I also calculated evolutionary rates averaged over the entire morphological dataset while including the fossil taxa to coarsely reconstruct evolutionary tempo throughout the Miocene. Although this step missed valuable information recovered by the mosaic analyses, including the fossil taxa in this manner enabled finer resolution into the coarse patterns in evolutionary tempo throughout the Miocene.

Evolutionary rates in neurological data

I estimated macroevolutionary rates across primates in ED and EQ using BAMM version 2.5.0 (Rabosky 2014). Since the trees that I constructed for the morphological

analysis contained only a fraction of the taxa present in the neurological datasets, I used the dated primate supertree packaged in BAMM that was originally sourced from Mos and Mooers (2006) and pruned the tips to match the taxa present in each dataset. MCMC simulations were run until the estimated sample size (ESS) well exceeded 200 for each parameter. Results were presented by plotting the mean rate estimated along each branch and the maximum *a posteriori* configuration of rate shifts using BAMMtools version 2.5.0 in R (Rabosky et al., 2014).

Lineage diversification rate analyses

I performed analyses of origination and extinction rates in hominoid and cercopithecoid fossil records to examine the correspondence between lineage diversification and the patterns in disparification emphasized in the analyses of morphological rates and mosaicism. Fossil occurrence data spanning primates were acquired from the Paleobiology Database (paleobiodb.org) on July 31, 2019. This dataset was partitioned into two subsets: one encompassing Hominoidea and the other representing Cercopithecoidea. Rates of origination and extinction were inferred from each subset separately using PyRate (Silvestero et al. 2014). The inferential model was constrained to contain uniform origination and extinction rates across time, thus ignoring the possibility of diversification shifts within each clade. Preservation was constrained to being time-homogeneous and was assumed to be uniform within each clade. While ignoring potential information regarding diversification rate heterogeneity through time, this simplified model matches classic paleobiological work that assumes simple birth-death-sampling models and was therefore appropriate for the comparison of average

diversification rates in sister clades while avoiding possible model overfitting that may obfuscate the straightforward test sought here.

Simulated Markovian diffusion

To examine the expected effect of constructional opportunity stemming from both developmental and functional constraint on phenotypic evolution, I designed a set of evolutionary simulations based on a simple Markovian diffusion. In this system, quantitative traits belonging to a single population of organisms evolve stochastically along a fitness landscape. Each generation, values for each trait are proposed. Values that increase the overall fitness of the population are accepted. To mimic the effect of drift, values that decrease fitness are also accepted with some (user-specified) probability. Constraint stemming from developmental integration caused by pleiotropy can be mimicked by proposing values for traits that covary, while functional constraints are modeled by evolving traits along a shared, multivariate adaptive landscape, modelled here using a mixture of multivariate Gaussian distributions. In a completely unconstrained system, each trait evolves according to its own univariate landscape, with its value drawn independently of all others in each generation. Performing simulations in this way facilitated an illustration of the potential for functional and developmental integration to constrain evolutionary rates in an adaptive landscape. Although distinguishing between these modes of integration is generally difficult or impossible in paleontological study systems, the simulations performed here provide insight by testing 1) the potential for varying systems of integration to catalyze or constrain adaptive change and 2) the extent to which functional and developmental integration generate

overlapping and distinct patterns in evolutionary rate and disparity. When placed in a macroevolutionary context, the simulations here should be thought of as a demonstration of the possible microevolutionary processes that may have driven the higher-level patterns in evolutionary rate occurring along a single phylogenetic branch in the mosaic analyses. The script used to generate these simulations is available in the data supplement.

Results and Discussion

Strength of integration throughout the haplorrhine radiation

Calculating character correlations from the skeletal data along each branch of the extant haplorrhine phylogeny revealed a single episode of reduced integration at the root of Hominoidea, where the correlation decreased from 0.12 at the ancestor of catarrhines, to -0.04 at the root of hominoids (Fig. 2). This suggests that the reduction in integration across apes between the fore- and hindlimbs revealed by Young and colleagues (2009) extended across the skeleton along the branch leading to the stem hominoids. Such episodic relaxed integration was not detected in the other haplorrhine lineages, with cercopithecoids (0.2-0.33 correlation) and platyrrhines (0.16-0.31 correlation) displaying higher integration than hominoids (-0.04-0.25 correlation) overall. However, it is possible that more complete sampling of taxa and characters across both of these clades may reveal changes in modularity and integration not detected here.

Composition of mosaic suites

The analysis of mosaic disparity recovered five suites of traits (Fig 3b). The cranium was the largest contributor of both traits and variance across all suites except for

C4, which was represented by the pelvis alone (Fig. 3b and 3c). This reflects the higher sampling of cranial traits in the dataset. While the cranium has previously been shown to display substantial modularity (Felice and Goswami 2017), the results here suggest that individual cranial modules may result from the formation of broader evolutionary complexes shared with the post-cranium. The suites were distinct in their composition across postcranial anatomy: C0 was represented strongly by the wrist, ulna, and humerus; C1 was represented postcranially by the scapula; C2 by carpal and pelvic traits. The cranium was represented even more strongly than expected in C3 and contributed nearly half of the variance (Fig. 3b).

Rates of morphological evolution

The phylogenetic rate calculations for each suite revealed a strong pattern of evolutionary mosaicism (Fig. 3a). Three of the suites, C0, C1, and C3, displayed the highest rates after the divergence of the great apes from hylobatids, but before the divergence of gorillas from humans and chimpanzees. The two suites that did not experience the shared great ape rate increase, C2 and C4, experienced large bursts in *Homo*. This finding is consistent with general knowledge of human evolution, as both suites are dominated by cranial traits and pelvic traits related to birthing and locomotor function (Table S1).

The rate analyses (Fig. 3a) show that humans have experienced major macroevolutionary bursts. However, the unique characteristics of the human body plan are built over an anatomical structure shared by all great apes that was shaped in the Miocene during similarly dramatic episodes. For example, although suite C0 increased in

evolutionary rate along the human branch, its evolution was shaped by a sustained period of elevated rate between 20 and 10 million years ago. This suggests that much of the ‘groundwork’ underlying the derivation of humans’ unique features may predate the divergence between our lineage and chimpanzees’. The mosaic analysis also demonstrates that substantial visible phenotypic novelty can result from the evolutionary remodeling of a small subset of the anatomy. The most elevated evolutionary rates in humans were observed in suites C2, C3, and C4, which cumulatively comprise fewer than one-third of the traits sampled in the matrix (Table S1). C4, the smallest suite (10 traits) displayed by far the largest increase in rate along the *Homo* lineage. Perhaps notably, C4 also was the most static character suite throughout the earlier stages in the hominoid radiation. The mosaic pattern recovered here shows that the uniqueness of the human body plan was shaped by dramatic evolutionary shifts in only a small number of traits.

While the large amount of missing data among the fossil taxa made an additional mosaic analysis infeasible, their inclusion revealed an otherwise hidden shift in evolutionary rate that occurred at the root of Hominoidea (approximately 30 million years ago). The fossil data also recapitulated the burst in evolutionary rate during the mid-Miocene that was displayed by C0 and C1 in the mosaic analysis (Fig. 3 and 4). When averaging over all the traits and including fossil taxa, these two episodes are the most dramatic macroevolutionary events when tracing the evolutionary lineage leading to *Homo*, suggesting the importance of early shifts in the ape body plan in shaping the functional morphologies of living taxa.

Several of the major rate shifts that occurred throughout the great ape clade corresponded to increased evolvability conferred by relaxed integration among early apes. However, comparison of the human-specific evolutionary rates in cranial, pelvic, and neurological traits to the phylogenetic character correlations in Figure 2 shows that the elevated morphological rates in humans were immediately preceded by an *increase* in the strength of integration. This illustrates an alternative means through which shifts in systems of modularity promote rapid and dramatic evolutionary remodeling of morphology. The suites that showed the most dramatic rate increases in humans, C2 and C4, were evolutionarily inactive prior to the most recent common ancestor (MRCA) of chimps and humans. This suggests that, while early ape evolution was characterized by relaxed integration leading to rapid evolutionary change in existing modules, human morphological evolution was shaped by the formation of two small modules of characters from the pelvis and carpus that underwent an initial rate increase following their formation along the branch leading to chimps and humans and then a more dramatic rate increase in hominins. Although the overall structure of integration was well-defined along the chimp-human MRCA branch, the formation of *new*, smaller modules contributed to dramatic changes across the body plan in humans. The occurrence of these initial shifts in modularity and rate along the chimp-human MRCA shows that chimpanzees share the modified structure of integration that facilitated the evolution of key human traits. This result is surprising given the typical assumption that chimpanzees are morphologically more similar to other great apes than are humans.

Rates of neurological evolution

The great apes differentiated rapidly from other catarrhines in both EQ and ED. *Homo* displays the highest rate of both EQ and ED evolution. However, the human rate shift in both traits occurred as part of an older trend of rapid neurological evolution in African apes. For EQ, humans experienced a substantial increase in evolutionary rate relative to the rest of the African apes, despite the statistical evidence for a shared rate shift at the root of the clade (Fig. 5b). The shared macroevolutionary regime shared by all African apes is most apparent in encephalization development (Fig. 5a), where the estimated evolutionary rate in *Homo* increased only slightly after splitting from *Pan* (Fig. 5c). It appears that the increased encephalization developed throughout the post-natal period in humans reflects a general trend among great apes in the evolutionary plasticity of neurological development. As a result, the ability to develop a relatively large mass of brain tissue does not itself appear to be a strong limiting factor in the evolution of large encephalization in humans. Instead, the morphological analyses suggest that pelvic traits (C4 in Fig. 1) demanded a more dramatic alteration in macroevolutionary regime, requiring both the formation of a new suite of characters and massive burst in rate.

Structural and ecological opportunity in ape evolution

The elevated evolutionary rates experienced by apes early in their divergence correspond to a general relaxation of constraint in early Miocene stem and ancestral hominoids. The reduction in integration across the skeleton and burst in the evolutionary rate of neurological development that occur at the base of the hominoid clade correspond in timing to the elevated rates in morphological evolution observed in most of the skeletal characters examined here (Fig. 3 and 4). Freed from previous functional and

developmental limitations, ape body plans would have rapidly diversified when placed in the context of the repeated environmental fluctuations that occurred throughout the mid and late Miocene (Michel et al. 2014, Hunt 2016) before becoming functionally canalized through development and stabilizing selection. This pattern is consistent with a scenario in which morphological features evolve by stochastic diffusion on a Simpsonian adaptive landscape that is alternately dampened and released by relaxed and shifting patterns in character integration.

Rapid and frequent environmental fluctuations throughout the Miocene suggest a possible abundance of available adaptive zones during the evolution of both cercopithecoids and hominoids. Paleoeological information extracted from deposits containing key hominoid taxa are so variable that environmental turnover often outpaces the effects of time-averaging (Michel et al. 2014). Early Miocene ape and monkey species evolved in overlapping environmental conditions and shared similar ecomorphological and dietary habits. However, as their lineages diverged, apes evolved a diverse set of locomotor suites, while evolution in locomotor features has remained comparatively static across old world monkeys (Hunt 2016). The patterns in developmental and mosaic morphological evolution revealed here suggest that the substantial differences between the two taxa in phenotypic disparity were shaped more by differences in patterns in integration and constraint rather than from the ecological opportunities available to either. While hominoids and cercopithecoids inhabited a similar range of habitats throughout their evolution, and so likely experienced a similar abundance of ecological opportunity, greater constructional opportunity is a

distinguishing feature that correlates with hominoids' vast disparity in skeletal morphology.

The scenario entertained here for apes is consistent with recent work that suggests that the initial burst in phenotypic disparity that often accompanies the origin of a new lineage can result from an early relaxation of constraints through the dissolution of integrated modules that is followed by reformation of character suites that lead to rapid, correlated evolutionary changes between constituent characters (Wagner 2018). In the cited work, the author distinguishes this 'breakup-relinkage' model with one where evolutionary rates are elevated by a relaxation of constraint alone. The 'breakup-relinkage' model is directly supported by the phylogenetic integration pattern, which show that an initial decrease in integration at the root of hominoids was followed by an increase at the following node (Fig. 2). Therefore, the recanalization of functional variation into a set of suites with distinct functional and developmental properties early in hominoid variation is likely equally important to the initial relaxation in constraint in having driven the remarkable morphological disparity observed across hominoid species.

"Preadaptation" and developmental enablers in apes

Bursts in phenotypic change and diversification are often preceded by evolutionary enablers that facilitate the construction of more diverse morphological forms or fulfill a necessary condition for the later emergence of a more derived trait. Such patterns have been suggested to stem from higher-level processes variously referred to as "preadaptation" (Bock 1959), "exaptation" (Gould and Vrba 1982), and "developmental enablement" (Donoghue 2005). The evolution of developmental, morphological, and

behavioral traits is often hierarchical through functional inter-dependencies and temporally autocorrelated, with changes in traits at relatively lower levels ‘setting the stage’ for higher-level traits by providing the developmental or structural conditions necessary for their emergence. The stepwise macroevolutionary bursts observed here show that hominoid evolution has been defined by sequential, episodic releases in constraint that likely set the stage for later innovations that compounded upon earlier rearrangements in the structure of developmental and functional integration.

The related concepts of preadaptation, exaptation, and developmental enablement all focus on identifying the origin of single characters that facilitate the later emergence of more derived states. However, the patterns presented here suggest that such processes may often be better characterized as reflecting larger changes in developmental and functional character-linkage systems rather than by the sequential derivation of individual characters. The dramatic bursts in evolutionary rate that occur in the largest character suites coincide with or immediately follow the reduction in integration that occurred at the root of hominoids. This suggests that a general shift in functional and/or developmental integration occurred near this time. This shift precedes and, although speculative, may have resulted in a general predisposition across apes that facilitated the later functional divergence of later species into an unusually broad range of habits. Therefore, rather than stemming from a single “preadaptive” character, the morphological disparity of apes and the emergence of features derived in single lineages (such as bipedalism in humans) may be better explained by a shift in developmental and functional integration patterns earlier in ape evolution.

Enhanced evolvability conferred by shifts in the structure of integration contributed to the emergence of key human traits. The apparent formation of suites C2 and C4, represented disproportionately by the cranium and pelvis, along the branch leading to humans and chimpanzees coincided with an initial burst in evolutionary rate. Both of these events immediately preceded the dramatic bursts in *Homo* (Fig. 3). Likewise, EQ and ED have been evolving rapidly throughout the ape radiation, with ED experiencing an initial increase in evolutionary rate before the splitting of gorillas, chimps, and humans followed by more dramatic upticks of rate in humans in both traits (Fig. 5c). While distinct in its details, this pattern in chimps and humans provides further evidence for the preadaptive potency of shifts in systems of modularity and integration in driving dramatic anatomical rearrangements. It also illustrates that the broad range of possible evolutionary pathways opened by either an overall reduction in integration (at the hominoid root) or the formation of new modules (in chimps and humans), rather than movement toward a specific set of phenotypes, is the means through which such predispositions facilitate evolutionary remodeling.

It is possible that the initial bursts in ED and suite C2 and C3 evolution, coupled with the reduced integration in the MRCA of chimp and humans, may reflect a set of structural rearrangements that facilitated, at least in part, the later emergence of human-specific neurological and cognitive features. This scenario echoes the results of Rice (1998), who demonstrated that humans and chimpanzees, to the exclusion of gorillas, both inherited a shared ‘additional’ stage of brain development from our most recent common ancestor. It is therefore possible that, in addition to sharing a similar derived

structure of modularity, chimpanzees possess more of the neurological and cranial
groundwork that contributed to the evolution of humans' particular mode of cognition
than has previously been appreciated.

The patterns uncovered here suggest that the concepts of preadaptation and
exaptation may sometimes be more compellingly applied by shifting focus from
individual features to patterns in integration and modularity across broad suites of
characters. Although the concept is essentially unchanged as applied to integration
patterns, it may be more accurate in the example here to refer instead to an 'evolutionary
predisposition' given the lack of evidence for a specific sequence of selective regimes
(Smith et al. 2018). This semantic argument is likely also valid in more traditional
examples of single trait exaptations or preadaptations. And so, a generalization from
preadaptation or exaptation to evolutionary predisposition that 1) incorporates integration
patterns and 2) shifts emphasis from adaptation, which makes assumptions about the
(often unknowable) set of historical microevolutionary processes, to the sequence of
observable states in both characters and integration patterns can provide a stronger
conceptual basis for future work that considers contingency and sequential order in the
evolutionary pathways of complex traits. Although this reconceptualization might
superficially appear to de-emphasize evolutionary process in favor of pattern, I would
instead argue that constraining the scale of inquiry and incorporating a broader range of
biological complexity can facilitate the reconstruction of elusive higher-level
macroevolutionary processes that shape organismal diversification and disparification
over deep timescales. This formulation also avoids the conceptual and practical

frustrations that occur when attempting to invoke the concepts of preadaptation and exaptation to explain evolutionary history over deep timescales by reducing dependency on knowledge of lower-level population processes.

Diversification and disparification in haplorrhine evolution

Cercopithecoids have diversified in species number at over twice the rate as hominoids over the same time scale (Table 1). One limitation of these analyses stems from cercopithecoids' somewhat poor fossil record as compared to hominoids. Nevertheless, the difference in preservation rates estimated in each clade (~3 in apes compared to ~2.6 in cercopithecoids) should ameliorate some of the potential effect of this bias on parameter estimates. The general pattern of higher net diversification in cercopithecoids is also consistent with previous neontological results (Purvis et al. 1995), suggesting that the paleontological estimates presented here are adequately robust to these effects. This difference in lineage diversification rates between both taxa demonstrates that increased morphological disparification is, overall, not correlated with faster lineage diversification in haplorrhine primates. While the evolutionary predisposition conferred upon apes by their relaxed integration corresponds to an increase in morphological disparity, this effect does not extend to lineage diversity.

On the surface, the overall similarity in the environments, dietary niches, and geographical ranges of hominoids and cercopithecoids suggests that cercopithecoids have a selective advantage in the species-sorting dynamics between the two clades. However, the long-term persistence of hominoids and their remarkable innovation in locomotor function relative to cercopithecoids throughout this timespan suggests an equilibrium in

their relative abundances (Van Valen 1975, Chesson 2000). Although cercopithecoids experienced less dramatic innovation in postcranial morphology, they display a broader range of derived digestive physiologies, including both intestinal and dental features associated with food processing (Hunt 2016). Given these observations, it is possible that the differential in diversification rate between cercopithecoids and hominoids stems from differences in the timescales of their life histories and other factors not directly related to interspecific competitive ability. If this is the case, hominoids may achieve steady persistence through higher evolvability conferred by reduced integration and increased developmental flexibility when faced with changing environments, while cercopithecoids may do so through a combination of their higher net diversification rates and capability to exploit a range of dietary resources through their digestive physiologies. Such differences would be consistent with an equilibrium model of coexistence rather than a simple species selection model that predicts the ultimate extinction of hominoids through competitive exclusion by cercopithecoids. Nevertheless, more rigorous comparison of each of these attributes in both lineages is needed to further constrain the range of possible explanatory factors.

Constrained Markovian diffusion and entropy

The patterns in character disparity shown here evoke a Markovian diffusion that is dampened by constraint. The tightly integrated nature of vertebrate body plans suggests major constraints in their ability to fill morphospace. However, the pattern displayed by apes suggests that the structure of integration may sometimes shift, allowing diffusion into new areas. Morphological evolution might then be conceived as a multivariate

diffusion in morphospace where movement is directionally biased due to fitness differences modelled by a traditional Simpsonian adaptive landscape and certain regions are rendered inaccessible by structural interactions between characters. Although this diffusive model, including the feedback links between phenotype, development, and environment, has been considered previously (Fisher 1986), its behavior has not been well explored in this context.

The Monte Carlo simulations presented here provide a theoretical illustration of the population-level dynamics stemming from one possible set of integration scenarios and their potential effects on the evolutionary disparity/rate along a single branch in Figures 3 or 4. The results shed light on the empirical analyses by demonstrating the overlapping and distinct empirical patterns that can result from varying levels of developmental and functional integration in a simple closed system in the absence of environmental fluctuation (Fig. 6). As is expected from quantitative genetic theory (Cheverud 1984, Maynard-Smith et al. 1985), developmental and functional constraints can generate similar patterns (Fig. 6a and 6b). In all these cases, the overall entropy observed in the system over time remains low, with the system tending to become stuck or moving away very slowly from a suboptimal location (Fig 4c and 4e) or crawling slowly in concert along a gradient (Fig 6a and 6b). However, when both types of constraint are released, the system displays high entropy, with each trait able to independently jump between peaks (Fig. 6d). Such unconstrained movement would be expected to result in higher phenotypic disparity over macroevolutionary timescales by

freeing individual lineages to generate novel character combinations when exposed to distinct and changing environments.

The patterns in increased evolutionary rate and entropy across traits displayed by the less constrained simulated diffusions provide mechanistic explanations that are consistent with the bursts in evolutionary rate encountered throughout hominoid evolution. Although it is not possible in the empirical example to distinguish between functional and developmental integration as is done in the simulations, the general pattern of reduced integration that gives way to increased evolutionary rate is consistent between the two. Alternation between the varying strengths of integration as explored in the simulations would likely generate substantial novelty during periods of relaxed integration. Episodes of reduced developmental constraint (Figs. 2 and 5a) may have generated an initial burst of constructional opportunity that led to rapid morphological diversification as ape populations were exposed to highly variable environments throughout the mid-Miocene. If the processes modelled in the simulations drove the patterns revealed in the empirical analyses, this sequence of events would represent true preadaptation in the sense that relaxed integration leads to increased mean population fitness. Under the evolutionary models explored here and elsewhere (Wagner 2018), stabilizing effects such as developmental canalization (Waddington 1959) or functional covariance (Cheverud 1984) would then be expected to re-constrain the newly divergent phenotypes after an initial burst in diversification. This dampened diffusive model was explored in early work in theoretical morphology that focused on the constrained filling of morphospace (Raup 1968).

593 The simulated diffusions and empirical pattern in evolutionary predisposition
594 revealed here hint at the source of teleological concepts in human evolution. If the
595 morphological and neurological traits can be assumed to have followed a multivariate
596 Markovian diffusion that is at least qualitatively similar to the simulations in Figure 6
597 (especially Fig. 6a), the pattern in evolutionary rates along the path leading to humans
598 may paint a misleading view that such changes have followed a progressive trend leading
599 to a human-defined apex. However, the mosaic analysis here suggests that humans’
600 biological uniqueness can be attributed to a relatively small number of anatomical
601 rearrangements built over a longer diffusion into an area of developmental-morphospace
602 occupied by all apes that is characterized by increased structural opportunity. Instead of
603 treating humans as exceptional, this view suggests that the structural opportunities that
604 emerged early in ape evolution may have freed the ancestors of currently extant taxa to
605 blaze unique trajectories along a complex, multivariate adaptive landscape that would
606 have been otherwise inaccessible. Structurally, all ape species might therefore be viewed
607 as equally progressive in their evolution, with species-level differences in form and
608 function shaped by stochastic differences in separate realized evolutionary paths along a
609 shared adaptive landscape. Alternatively, separate taxa may have forged new and unique
610 adaptive landscapes following speciation events as structural differences between newly
611 isolated species drove the creation of new functional niches early in hominoid evolution.
612 The derived structure of integration and neurological development shared by chimps and
613 humans highlights the importance of contingency in human evolution—chimps’ sharing
614 of much of the same underlying constructional architecture shows how this derived

predisposition allowed, but did not determine, the emergence of humans' unique suite of phenotypes.

The empirical and simulated analyses presented here evoke a picture of morphological evolution that involves episodic diffusion across adaptive zones. However, the empirical and theoretical analyses allow for alternating and interacting roles for both ecological and constructional opportunity. This view can supplement existing conceptions of adaptive radiation and key innovation by providing an alternative to ecological opportunity as a lone driving factor in the disparification of form. Instead, phenotypic disparity produced through diffusion into a set of abundant adaptive zones can be directed and limited by the availability of structural opportunity. Nature surely merges these simplified extremes when shaping patterns in biodiversity (Seilacher 1991), and so further integrative study of the interplay between ecological and constructional factors will be critical in developing a pluralistic understanding of the complex patterns and processes that have shaped the vast diversity across the tree of life.

References

Alberch P. 1980 Ontogenesis and morphological diversification. *American Zoologist* 20, 653-667.

- 637 Andrews P. 1992 Evolution and environment in the Hominoidea. *Nature* 360, 641–646.
- 638 (doi:10.1038/360641a0)
- 639 Arnold SJ. 1992 Constraints on phenotypic evolution. *American Naturalist*, 140(Supp):
- 640 S85-S107.
- 641 Berge C. 1998 Heterochronic processes in human evolution: An ontogenetic analysis of
- 642 the hominid pelvis. *American Journal of Physical Anthropology* 105, 441–459.
- 643 (doi:10.1002/(sici)1096-8644(199804)105:4<441::aid-ajpa4>3.0.co;2-r)
- 644 Bock J. 1959 Preadaptation and multiple evolutionary pathways. *Evolution* 13, 194-211.
- 645 Boddy AM, McGowen MR, Sherwood CC, Grossman LI, Goodman M, Wildman DE.
- 646 2012 Comparative analysis of encephalization in mammals reveals relaxed
- 647 constraints on anthropoid primate and cetacean brain scaling. *Journal of*
- 648 *Evolutionary Biology* 25, 981–994. (doi:10.1111/j.1420-9101.2012.02491.x)
- 649 Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut
- 650 A, Drummond AJ. 2014 BEAST 2: A Software Platform for Bayesian
- 651 Evolutionary Analysis. *PLoS Computational Biology* 10, e1003537.
- 652 (doi:10.1371/journal.pcbi.1003537)
- 653 Capellini I, Venditti C, Barton, RA. 2010 Phylogeny and metabolic scaling in mammals.
- 654 *Ecology*, 91, 2783-2793.
- 655 Charlesworth B, Lande R, Slatkin M. 1982 A Neo-Darwinian Commentary on
- 656 Macroevolution. *Evolution* 36, 474–498. (doi:10.1111/j.1558-
- 657 5646.1982.tb05068.x)

- 658 Chesson P. 2000 Mechanisms of maintenance of species diversity. *Annual Review of*
659 *Ecology and Systematics*, 31, 343-366.
- 660 Cheverud JM. 1984 Quantitative genetics and developmental constraints on evolution by
661 selection. *Journal of Theoretical Biology* 110, 155–171. (doi:10.1016/s0022-
662 5193(84)80050-8)
- 663 Crouch NMA, Ricklefs RE. 2019 Speciation Rate Is Independent of the Rate of
664 Evolution of Morphological Size, Shape, and Absolute Morphological
665 Specialization in a Large Clade of Birds. *The American Naturalist* 193, E78–E91.
666 (doi:10.1086/701630)
- 667 Donoghue MJ, Ree RH. 2000 Homoplasy and Developmental Constraint: A Model and
668 an Example from Plants. *American Zoologist* 40, 759–769.
669 (doi:10.1093/icb/40.5.759)
- 670 Donoghue MJ. 2005 Key innovations, convergence, and success: macroevolutionary
671 lessons from plant phylogeny. *Paleobiology* 31, 77-93. (doi: 10.1666/0094-
672 8373(2005)031[0077:KICASM]2.0.CO;2)
- 673 Felice RN, Goswami A. 2017 Developmental origins of mosaic evolution in the avian
674 cranium. *Proceedings of the National Academy of Sciences* 115, 555–560.
675 (doi:10.1073/pnas.1716437115)
- 676 Felsenstein J. 1985 Phylogenies and the comparative method. *The American Naturalist*,
677 125, 1-15.
- 678 Fisher, DC. 1986 Progress in organismal design. Pp. 99-117 in: *Patterns and Processes in*
679 *the History of Life*. Raup DM and Jablonski D, eds. Springer, Berlin, Heidelberg.

- 680 Goswami A, Binder WJ, Meachen J, O’Keefe FR. 2015 The fossil record of phenotypic
681 integration and modularity: A deep-time perspective on developmental and
682 evolutionary dynamics. *Proceedings of the National Academy of Sciences* 112,
683 4891–4896. (doi:10.1073/pnas.1403667112)
- 684 Goswami A, Smaers JB, Soligo C, Polly PD. 2014 The macroevolutionary consequences
685 of phenotypic integration: from development to deep time. *Philosophical*
686 *Transactions of the Royal Society B: Biological Sciences* 369, 20130254–
687 20130254. (doi:10.1098/rstb.2013.0254)
- 688 Gould SJ, Vrba ES. 1982 Exaptation—a Missing Term in the Science of Form.
689 *Paleobiology* 8, 4–15. (doi:10.1017/s0094837300004310)
- 690 Gould SJ and Lewontin RC. 1979 The spandrels of San Marco and the Panglossian
691 paradigm: a critique of the adaptationist programme. *Proceedings of the Royal*
692 *Society of London. Series B. Biological Sciences* 205, 581–598.
693 (doi:10.1098/rspb.1979.0086)
- 694 Gould, SJ. 1977 *Ontogeny and Phylogeny*. Harvard University Press, Cambridge, USA.
- 695 Harmon LJ. 2003 Tempo and Mode of Evolutionary Radiation in Iguanian Lizards.
696 *Science* 301, 961–964. (doi:10.1126/science.1084786)
- 697 Holloway RL. 1973 Endocranial volumes of early African hominids, and the role of the
698 brain in human mosaic evolution. *Journal of Human Evolution* 2, 449–459.
699 (doi:10.1016/0047-2484(73)90123-1)
- 700 Hunt KD. 2016 Why are there apes? Evidence for the co-evolution of ape and monkey
701 ecomorphology. *Journal of Anatomy* 228, 630–685. (doi:10.1111/joa.12454)

- 702 Lewton KL. 2010 Locomotor function and the evolution of the primate pelvis. (Doctoral
703 dissertation, Arizona State University).
- 704 Losos JB. 2010 Adaptive Radiation, Ecological Opportunity, and Evolutionary
705 Determinism. *The American Naturalist* 175, 623–639. (doi:10.1086/652433)
- 706 Mahler DL, Ingram T, Revell LJ, Losos JB. 2013 Exceptional Convergence on the
707 Macroevolutionary Landscape in Island Lizard Radiations. *Science* 341, 292–295.
708 (doi:10.1126/science.1232392)
- 709 McHenry H. 1975 Fossils and the mosaic nature of human evolution. *Science* 190, 425–
710 431. (doi:10.1126/science.809842)
- 711 Michel LA et al. 2014 Remnants of an ancient forest provide ecological context for Early
712 Miocene fossil apes. *Nature Communications* 5. (doi:10.1038/ncomms4236)
- 713 Rice, SH. 2002. The role of heterochrony in primate brain evolution. In: Human
714 evolution through developmental change. N. Minugh-Purvis & K.J. McNamara,
715 eds. Johns Hopkins Univ. Press, Baltimore.
- 716 Seilacher A. 1991 Self-Organizing Mechanisms in Morphogenesis and Evolution. In
717 Constructional Morphology and Evolution, pp. 251–271. Springer Berlin
718 Heidelberg. (doi:10.1007/978-3-642-76156-0_17)
- 719 Silvestro D, Salamin N, Schnitzler J. 2014 PyRate: a new program to estimate speciation
720 and extinction rates from incomplete fossil data. *Methods in Ecology and*
721 *Evolution* 5, 1126–1131.
- 722 Smith JM, Burian R, Kauffman S, Alberch P, Campbell J, Goodwin B, Lande R, Raup D,
723 Wolpert L. 1985 Developmental Constraints and Evolution: A Perspective from

724 the Mountain Lake Conference on Development and Evolution. *The Quarterly*
725 *Review of Biology* 60, 265–287. (doi:10.1086/414425)

726 Smith SA, Brown JW, Yang Y, Bruenn R, Drummond CP, Brockington SF, Walker JF,
727 Last N, Douglas N, Moore MJ. 2018 Disparity, Diversity, and Duplications in the
728 Caryophyllales. *New Phytologist* 217, 836-854.

729 Olson ME. 2012 The developmental renaissance in adaptationism. *Trends in Ecology and*
730 *Evolution* 27, 278–287. (doi:10.1016/j.tree.2011.12.005)

731 Olson EC and Miller RL. 1958. *Morphological Integration*. University of Chicago Press.

732 Parins-Fukuchi C. 2018 Bayesian placement of fossils on phylogenies using quantitative
733 morphometric data. *Evolution* 72, 1801–1814. (doi:10.1111/evo.13516)

734 Parins-Fukuchi C. In press. Detecting mosaic patterns in macroevolutionary disparity.
735 *American Naturalist*.

736 Purvis A, Nee S, Harvey PH. 1995 Macroevolutionary inferences from primate
737 phylogeny. *Proceedings of the Royal Society of London, Series B: Biological*
738 *Sciences* 260, 329-333.

739 Rabosky DL, Santini F, Eastman J, Smith SA, Sidlauskas B, Chang J, Alfaro ME. 2013
740 Rates of speciation and morphological evolution are correlated across the largest
741 vertebrate radiation. *Nature Communications* 4. (doi:10.1038/ncomms2958)

742 Rabosky DL. 2014 Automatic Detection of Key Innovations, Rate Shifts, and Diversity-
743 Dependence on Phylogenetic Trees. *PLoS One* 9, e89543.
744 (doi:10.1371/journal.pone.0089543)

- 745 Rabosky DL, Grundler M, Anderson C, Title P, Shi JJ, Brown JW, Huang H, Larson JG.
746 2014 BAMMtools: an R package for the analysis of evolutionary dynamics on
747 phylogenetic trees. *Methods in Ecology and Evolution* 5, 701–707.
748 (doi:10.1111/2041-210x.12199)
- 749 Rainey PB, Travisano M. 1998 Adaptive radiation in a heterogeneous environment.
750 *Nature* 394, 69–72. (doi:10.1038/27900)
- 751 Raup DM. 1968 Theoretical morphology of echinoid growth. *Journal of Paleontology* 42,
752 50-63.
- 753 Ruff CB. 1994 Morphological adaptation to climate in modern and fossil hominids.
754 *American Journal of Physical Anthropology* 37, 65–107.
755 (doi:10.1002/ajpa.1330370605)
- 756 Simpson GG. 1944 *Tempo and Mode in Evolution*. Columbia University Press. New
757 York.
- 758 Slater GJ, Friscia AR. 2019 Hierarchy in adaptive radiation: A case study using the
759 *Carnivora* (Mammalia). *Evolution* 73, 524–539. (doi:10.1111/evo.13689)
- 760 Slater GJ. 2015 Iterative adaptive radiations of fossil canids show no evidence for
761 diversity-dependent trait evolution. *Proceedings of the National Academy of*
762 *Sciences* 112, 4897–4902. (doi:10.1073/pnas.1403666111)
- 763 Sukumaran J, Holder MT. 2010 DendroPy: a Python library for phylogenetic computing.
764 *Bioinformatics* 26, 1569–1571. (doi:10.1093/bioinformatics/btq228)
- 765 Stebbins GL. 1983 Mosaic evolution: An integrating principle for the modern synthesis.
766 *Experientia* 39, 823–834. (doi:10.1007/bf01990398)

- 767 Stebbins GL. 1984 Mosaic evolution, mosaic selection and angiosperm phylogeny.
768 Botanical Journal of the Linnean Society 88, 149–164. (doi:10.1111/j.1095-
769 8339.1984.tb01568.x)
- 770 Van Valen L. 1975 Group selection, sex, and fossils. *Evolution*, 29, 87-94.
- 771 Vos RA. 2006 Inferring large phylogenies: the big tree problem (Doctoral dissertation,
772 Simon Fraser University)
- 773 Waddington CH. 1959 Canalization of Development and Genetic Assimilation of
774 Acquired Characters. *Nature* 183, 1654–1655. (doi:10.1038/1831654a0)
- 775 Wagner CE, Harmon LJ, Seehausen O. 2012 Ecological opportunity and sexual selection
776 together predict adaptive radiation. *Nature* 487, 366–369.
777 (doi:10.1038/nature11144)
- 778 Wagner GP, Pavlicev M, Cheverud JM. 2007 The road to modularity. *Nature Reviews*
779 *Genetics* 8, 921-931.
- 780 Wagner GP, Amemiya C, Ruddle F. 2003 Hox cluster duplications and the opportunity
781 for evolutionary novelties. *Proceedings of the National Academy of Sciences* 100,
782 14603–14606. (doi:10.1073/pnas.2536656100)
- 783 Wagner GP. 1988 The influence of variation and of developmental constraints on the rate
784 of multivariate phenotypic evolution. *Journal of Evolutionary Biology* 1, 45–66.
785 (doi:10.1046/j.1420-9101.1988.1010045.x)
- 786 Wagner PJ. 2018 Early bursts of disparity and the reorganization of character integration.
787 *Proceedings of the Royal Society B: Biological Sciences* 285, 20181604.
788 (doi:10.1098/rspb.2018.1604)

- 789 Worthington S. 2012 New approaches to late Miocene hominoid systematics: Ranking
790 morphological characters by phylogenetic signal (Doctoral dissertation, New
791 York University).
- 792 Wright, S. 1932 The roles of mutation, inbreeding, crossbreeding and selection in
793 evolution. Proc. 6th Int. Cong. Genet. 1: 356–366.
- 794 Yoder JB, et al. 2010 Ecological opportunity and the origin of adaptive radiations.
795 Journal of Evolutionary Biology 23, 1581–1596. (doi:10.1111/j.1420-
796 9101.2010.02029.x)
- 797 Young NM, Wagner GP, Hallgrímsson B. 2010 Development and the evolvability of
798 human limbs. Proceedings of the National Academy of Sciences 107, 3400–3405.
799 (doi:10.1073/pnas.0911856107)

800

801

802

803

804

805

806

807

808

809

810 **Tables**

Clade	Origination	Extinction	Net diversification	Preservation
Hominoidea	0.395	0.338	0.057	3.121
Cercopithecoidea	0.251	0.128	0.123	2.551

Table 1. Lineage diversification and preservation rate parameters estimated in apes (Hominoidea) and Old World monkeys (Cercopithecoidea). Parameters were summarized as the mean values from four million Markov-Chain Monte Carlo generations after discarding 20% as burn-in.

Figures

merged models cease to improve. The best clustering achieved from this procedure represents the best-supported heterogeneous model of character integration.

Figure 2. Strength of integration across haplorrhine evolution. Branch labels correspond to the strength and polarity of correlation in the reconstructed changes along each branch. Higher correlations in either direction (closer to either 1 or -1) indicate stronger evolutionary integration, while lower correlations (closer to 0) indicate weaker integration.

Figure 3. A) Mosaic evolutionary rates calculated while walking back from *Homo* to the most recent common ancestor of New World monkeys, Old World Monkeys, and apes. B) Variance contributed to each suite by each skeletal region, estimated from a principal component analysis. C) Skeletal regions over-represented in each mosaic evolutionary suite. Values above zero correspond to regions that occupy a higher proportion of their clusters than expected.

852

853 *Figure 4.* Branch colors correspond to rates of morphological evolution across all
854 characters along individual branches. Phylogenetic positions of fossil genera reflect
855 placements on extant reference tree.

856

857

Figure 5. Rate shifts in A) encephalization development, and B) encephalization quotient.
C) Branch-specific evolutionary rates when tracing the lineage from *Homo* to the root.

Figure 6. Monte Carlo simulations of integrated and atomized systems of three continuous traits. Each box is a simulated diffusion between three traits under A) moderate developmental linkage and shared selection, B) no developmental linkage and shared selection, C) no developmental linkage and selective independence, D) high developmental linkage and selective independence, and E) high developmental linkage and shared selection.

871

872

suite	skeletal region	trait
0	Ulna	Olecranon length
0	Ulna	Middle shaft shape
0	Ulna	Olecranon shape
0	Ulna	Distal shape
0	Ulna	Styloid process length
0	Ulna	Proximal shaft shape
0	Ulna	Coronoid
0	Ulna	Trochlea notch shape
0	Humerus	Head P D shape
0	Humerus	Trochlea
0	Humerus	Intertubercular angle
0	Humerus	Lat. trochlea keel dev.
0	Humerus	Bicipital groove shape
0	Humerus	Humeral Head Size
0	Humerus	Head torsion
0	Humerus	Trochlea sup. notch angle
0	Humerus	Zona conoidea dist. depth
0	Humerus	Grt. tuberosity proj.
0	Carpus	Capitate head M L width
0	Carpus	Capitate M C 3 shape
0	Carpus	CAPCM4
0	Carpus	Hamate DP depth COR
0	Carpus	Capitate head A P / P D
0	Carpus	Capitate dorsal nonarticular

0 Carpus	Capitate M C 3 depth
0 Carpus	Lunate dist.
0 Carpus	Hamate triquetral facet angle
0 Carpus	Capitate M C 2 facets
0 Cranium	Grt. palatine foramen position
0 Cranium	Zygomatic root origin elevation
0 Cranium	Basicranial length
0 Cranium	Piriform aperture shape
0 Cranium	Postglenoid projection
0 Cranium	Basicranium vs. palate
0 Cranium	Bregma position
0 Cranium	Lacrima fossa & orbital floor
0 Cranium	Facial projection
0 Cranium	Zygomax. tubercle A P width
0 Cranium	Basicranium vs. nuchal
0 Cranium	Nasoalveolar clivus length
0 Cranium	Head M L shape
0 Cranium	Zygomatic
0 Cranium	Face height vs. depth
0 Cranium	Piriform max. M L elevation
0 Cranium	Neurocranial shape
0 Cranium	Asterion angle
0 Radius	Lunate facet shape
0 Radius	Lunate facet angle
0 Radius	Neck shape
0 Radius	Scapholunate facet areas
0 Radius	Lunate facet dorsal proj.

0 Radius	Distal facets area
0 Radius	Total length
0 Radius	Radial neck length
0 Radius	Shaft curvature M L
0 Radius	Head articular surface P D
0 Scapula	Trapezius insert angle
0 Scapula	Coraco
0 Scapula	Inf. glenoid cor
0 Scapula	Acromian size
0 Scapula	Inferior angle
0 Scapula	Sup. notch shape
0 Scapula	Spinatus ratio
0 Scapula	Glenoid vent. bar angle
0 Pelvis	pubic symphysis length
1 Ulna	Ulna head height
1 Ulna	Semilunar keel angle
1 Ulna	Radial notch shape
1 Ulna	Distal shaft shape
1 Humerus	Dist. articular shape
1 Humerus	Trochlea dist. waisting
1 Humerus	Med. trochlea keel dev.
1 Humerus	Olecranon fossa depth
1 Humerus	Olecranon fossa shape
1 Carpus	Capitate P D length
1 Carpus	Hamate M C 4
1 Carpus	Lunate radial facet M L
1 Carpus	Lunate triquetral facet shape

1 Cranium	Piriform base M L width
1 Cranium	M1 shape
1 Cranium	Lsr. palatine foramen
1 Cranium	Molar crown areas
1 Cranium	Entoglenoid width
1 Cranium	Entoglenoid projection
1 Cranium	Nasal bone M L width
1 Cranium	Orbital margin M L width
1 Cranium	Orbit area
1 Cranium	Interorbital width
1 Radius	Distal surface shape
1 Radius	Styloid process length
1 Radius	Radial Head Shape
1 Manubrium	Manubrium shape
1 Scapula	Scapula Body shape
1 Scapula	Trapezius med. insert
1 Scapula	Superior angle
1 Scapula	Subscapular depth
1 Scapula	Vertebral border length
1 Scapula	Glenoid fossa shape
1 Scapula	Coracoid length
1 Pelvis	illium length
1 Pelvis	bi
1 Pelvis	anteroposterior pelvic diameter
1 Pelvis	pelvic length
2 Humerus	Epicondyles projection
2 Humerus	Olecranon perforation

2 Ulna	Shaft curvature A P
2 Carpus	Capitate M C 3 waisting
2 Carpus	Capitate waisting
2 Carpus	Lunate PD Length
2 Carpus	Hamate hamulus P D proj.
2 Carpus	Capitate M C 2 margin
2 Carpus	Capitate M C 2 orientation
2 Carpus	Lunate dist. facet shape
2 Cranium	Orbit shape
2 Cranium	Pterion post. config
2 Cranium	Palate shape
2 Cranium	Lacrima fossa visibility
2 Cranium	Palatine width
2 Cranium	Postorbital constriction
2 Cranium	Palate depth
2 Radius	Head depth P D
2 Scapula	Glenoid fossa depth
2 Pelvis	ilium width
2 Pelvis	ischium length
2 Pelvis	AIIS to hip joint
2 Pelvis	inferior pubic ramus length
2 Pelvis	acetabulum diameter
2 Pelvis	superior pubic ramus length
3 Scapula	Spinoglenoid notch depth
3 Scapula	Infraglenoid tuberosity M L
3 Pelvis	lower iliac height
3 Pelvis	inferior pubic ramus cross

3 Ulna	Radial notch depth
3 Cranium	Temporal fossa width
3 Cranium	Face vs. nasal height
3 Cranium	Face height vs. width
3 Cranium	Palate width
3 Cranium	Entoglenoid lat. pterygoid
3 Humerus	Medial trochlea keel & medial epicon
4 Pelvis	ischium cross
4 Pelvis	superior pubic ramus cross
4 Pelvis	upper illiac height
4 Pelvis	lower ilium cross
4 Pelvis	ischium dorsal projection
4 Pelvis	pubic symphysis
4 Pelvis	auricular surface area
4 Pelvis	biacetabular diameter
4 Pelvis	maximum transverse diameter
4 Pelvis	sacrum width

873

874 **Table S1:** List of traits and their skeletal regions that comprise each of the mosaic suites detected
875 from the skeletal data.

876