Radiologic Evaluation of the Influence of Cleft Type on Nasal Dorsum Growth

Lingling Pu, M.D^{a,b}, Renkai Liu M.D^{a,b}, Bing Shi, M.D^{a,b}, David W Low, M.D^c,
Chenghao Li, M.D, PhD^{a,b,*}

^a State Key Laboratory of Oral Disease, West China College of Stomatology, Sichuan University, Chengdu 610041, the People's Republic of China.

^b Department of Cleft Lip and Palate Surgery, West China Stomatological Hospital, Sichuan University, Chengdu 610041, the People's Republic of China.

^c Division of Plastic Surgery, The Children's Hospital of Philadelphia

***Correspondence to:**

1. Cheng-hao Li, M.D., Ph.D., Department of Cleft Lip and Palate Surgery, West China Stomatological Hospital, Sichuan University

E-mail: leechenghao_cn@yahoo.com

Fax: 86-28-85502570

Telephone: 86-28-85501462

Address: No. 14, Section 3, Ren Min Nan Road, Chengdu 610041, People's Republic of China.

18
19
20
21
22
23
24
25
26
27
28

29

30

31 **Abstract**

32 *Purpose:* The study was designed to evaluate whether intrinsic morphological
33 characteristics of the nasal dorsum are affected by cleft type, specifically cleft lip only
34 (CL) and cleft lip with cleft palate(CL/P).

35 *Methods:* 576 cleft patients (278 CL only, 298 CL/P), and 333 individuals without
36 orofacial clefts were retrospectively enrolled. Lateral cephalometric radiographs of all
37 individuals were taken to evaluate the nasal length and nasal dorsum height. Dunn's
38 test was used to analyze the difference ($p < 0.001$).

39 *Results:*In CL and control, the angulation of the nasal bone and nasal dorsum increase
40 by age similarly (5y-18y, $p>0.05$). In CL, the total dorsal length is significantly shorter
41 (5y-18y, $p<0.001$). Although the upper nasal dorsum is similar (except in 5y-6y), the
42 lower nasal dorsum is shorter (5y-18y, $p<0.001$).

43 In CLP, there is no significant difference in the nasal bone angle compared with
44 controls between 5y-7y. However, it develops insufficiently as children grow (8y-18y,
45 $p<0.001$). The nasal dorsum angle is notably smaller (5y-18y, $p<0.001$). Nasal bone
46 length is not significantly different from control at all stages except during ages 11y-
47 13y ($p<0.05$). Total nasal dorsal length is similar to the control at skeletal maturity
48 (17y-18y, $p>0.05$), although it is shorter during 8y to 16y ($p<0.05$). The upper nasal

dorsum is overdeveloped (14y-18y, $p<0.05$), whereas the lower nasal dorsum is underdeveloped (5y-18y, $p<0.001$).

Conclusion: CL inhibits the growth of nasal dorsum length, leading to short nose deformity. CL/P patients are prone to saddle-nose deformity because of the diminished nasal height (decreased nasal angle).

Keywords: Cleft nose; nasal deformity; cephalometric; cleft lip and palate

Introduction

Cleft lip is frequently accompanied by nasal deformities. The congenital anatomic deficiency or aberrancy, potential changes related to growth, the cleft itself, and even scarring from previous procedures are the main factors which lead to a wide variability in secondary cleft nasal deformities and the complexity of surgical techniques over the past few years¹. Subsequently, secondary surgery for the cleft nasal deformity undeniably presents a formidable challenge to the plastic surgeon, and the results are not as ideal as expected due to lack the comprehensive inward characters that hidden under complex deformed manifestations.

Due to its central location, the nose plays a prominent role in facial aesthetics ².

How does one distinguish the different factors that contribute to the cleft nasal

deformity, including cleft type, intrinsic potential changes, or surgical damage? Nowadays, surgeons mainly define the cleft nasal malformation with regard to the alar base, columella, nostril, nasal tip, nasal floor, and nasal septum , attaching more importance to the dysmorphia of the nasal tip ^{3,4}. However, because of the complexity of this anatomic structure, it is so difficult to define the key factors. As an important part of nose, the nasal dorsum plays a major role in nasal and facial harmony ⁵. Analysis of rhinoplasty results has shown that even slight differences in nasal shape can transform the look of an individual's face ⁶, the key point being that one might be able to distinguish the effect of different cleft types on nasal dorsum deformity, because of its simpler anatomic structure.

Based on the above reasons, this study focused on evaluating morphologic characteristics of the nasal dorsum, to analyze the role of cleft type on nasal dorsum growth in cleft lip patients with and without cleft palate. The study population included patients with cleft lip only (CL), cleft lip and cleft palate (CL/P), and healthy individuals. The soft and hard tissue of the nasal dorsum was analyzed through lateral cephalometric radiographs to obtain objective data of the hard and soft tissue morphology of the three groups in different ages, then compared. Lateral cephalometric radiographs of all individuals were taken to evaluate the nasal length, including the length of the nasal bone, the nasal dorsum, upper nasal dorsum and lower nasal dorsum. The angulation of the nasal bone and the nasal dorsum were evaluated as the indexes of nasal dorsum height. The results indicate that CL inhibits the growth of nasal dorsum

length, leading to a short nose deformity, while CL/P tends to result in a saddle nose because of decreased nasal height. These findings help characterize nasal dorsum development, provide comprehensive characteristics of the secondary nasal deformity in cleft patients, and potentially improve the outcome of secondary reconstructive surgery.

Methods

Ethics statement

Samples were collected in accordance with the guidelines of The West China Hospital of Stomatology Institutional Board(WCSHIRB). The experimental protocol was approved by local ethics committee (WCSHIRB, Sichuan University,China). Informed consent was obtained from all subjects or, if subjects are under 18, from a parent and/or legal guardian.

Sample

The study sample comprised a total of 909 Chinese children aged between 5 and 18 years at the West China Hospital of Stomatology, Sichuan University, Chengdu, China, between 2011 and 2016, who were divided into CL only, CL/P, and a control group. The CL group was composed of 278 children with cleft lip, and the CL/P group was comprised of 298 children with combined cleft lip and palate. They were non-syndromic and had no other congenital anomalies. Following our cleft center protocol, the CL group was treated with a modified Millard technique at 3-6 months ⁷. The CL/P group underwent the same lip repair technique, and then underwent a Sommerlad

palatoplasty at 9-12 months. None received any other secondary surgery such as lip revision, fistula repair, rhinoplasty, or orthopedic treatment except for bone grafting at 9-12 years of age. The control group was composed of 333 healthy children without cleft or any other congenital anomalies of the same age range as the CL and CL/P groups randomly chosen from Department of Orthodontics in West China Stomatology. These children underwent simple orthodontic treatment and had normal skeletal relationships, symmetric faces, and no history of craniofacial surgery. All groups were divided by age from 5 to 7 years, 8 to 10 years, 11 to 13 years, 14 to 16 years, and 17 to 18 years (Table 1).

Cephalometric analysis

Because our study was a retrospective case-control study using the archive, all the patients took lateral cephalometric radiographs just for clinical needs. Lateral cephalometric radiographs were taken for each subject under standardized conditions with the head oriented along the Frankfort horizontal plane (FH) parallel to the floor. Subjects were asked to relax their lips in a resting position, and to place their teeth in centric occlusion. An EASYMTIC 3298-125 Cephalometry X-ray machine (Chemetron Co., Chicago, IL, USA) was used for all subjects. In order to reduce the influence of maxillary hypoplasia, a reliable craniofacial reference plane “Sella–nasion S-N” was selected, and maxillary and nasal parts were separated by a vertical line through point nasion. Three hard and three soft tissue landmarks were digitized by one observer. Anthropometric landmarks on the nose were defined ^{8,9}. Nasal Dorsum was

measured by its length and angular of the hard and soft tissue. Fig.1 shows the landmarks that were used in the cephalometric analysis directly and indirectly, including four linear measurements and two angular measurements. The angulation of the nasal bone and the nasal dorsum were evaluated as the indexes of nasal dorsum height. The parameter measurements are shown in Fig.2. Each parameter was measured three times repeatedly and the mean was recorded, P25 (First Quartile), and P75 (Third Quartile).

Statistical analysis

All statistical analyses were performed with Statistical Package for Social Sciences (SPSS) software version 22.0. ANOVA analysis was used to determine the differences of age distribution in the three groups. Differences in the cephalometric results among the three groups were based on Dunn's test. The significant difference was defined at 95% level.

Reliability

To calculate the method error, 100 cephalograms were selected randomly and measured twice, to examine the intra-class correlation coefficient (ICC) ¹⁰. The ICC results for test-retest reliability ranged between 0.90 and 0.98, suggesting dependable reliability and reproducibility of the adopted measuring strategy. (Table 2)

Results

There was no significant difference in the age composition of CL, CL/P and Control groups. Nasal morphology in three groups was comparable (Table 3). Fig 2 shows the

growth tendency of each index in the CL, CL/P and Control group. Fig 3 is the nasal profile map of three groups in 17y-18y.

1. CL patients show shorter nose and normal nasal angulation.

In CL, compared with Control, the angulation of the nasal bone and nasal dorsum increase similarly by age (5y-18y, $p>0.05$), (Fig2a, 2b), while the total dorsum length is significantly shorter (5y-18y, $p<0.001$), (Fig2c,2d). In CL, the upper nasal dorsum is similar to Control (except in 5y-6y), (Fig2e), but the lower nasal dorsum is shorter (5y-18y, $p<0.001$), (Fig2f).

2. CL/P patients have flatter angulation, but normal-length nose.

In CL/P, there is no significant difference in the angulation of the nasal bone compared with Control in the 5y-7y age range. However, it develops insufficiently as children grow (8y-18y, $p<0.001$), (Fig2a). The angulation of the nasal dorsum is notably smaller than that in non-cleft children (5y-18y, $p<0.001$), (Fig2b). Nasal bone length is not significantly different from the control peers at all stages except the peers between 11y and 13y ($p<0.05$), (Fig2c). At skeletal maturity, the nasal dorsum grows as long as the Control group (17y-18y, $p>0.05$), although it is shorter prior to that (8y-16y, $p<0.05$), (Fig2c).

The upper nasal dorsum is overdeveloped (14y-18y, $p<0.05$) while the lower nasal dorsum is underdeveloped (5y-18y, $p<0.001$), (Fig2d,2e), the net effect being a total dorsum length similar to controls at skeletal maturity.

Discussion

Our previous works have analyzed the craniofacial and soft tissue morphology of patients with CL/P and CP^{11,12}. As mentioned earlier, this study was designed to distinguish cleft type factors associated with nasal dorsum deformity. With secondary rhinoplasty mainly aimed at patients with CL regardless of CP, and to eliminate the influence of maxillary retrusion after primary repair on nasal dorsum shape, our study encompassed patients with CL, CL/P, and healthy peers.

In this study, in CL only patients, we found the nasal bone and nasal dorsum were significantly shorter (Fig.2c,2d). This is consistent with previous findings: prior studies have demonstrated that in fetuses, newborns, children, and male adults, compared with normal peers, patients with isolated cleft lip had a significantly shorter nasal bone^{13,14}. Nasal tip position is one of the important indicators for the measurement of the length of nasal dorsum. Compared with healthy ones, CL (with or without CP) showed significant upward deviation in the nasal tip, suggesting that CL patients have a congenital tendency toward a short nose¹⁵, but previous researchers have not discussed the nasal features of CL and CL/P separately. Meanwhile, few studies have presented the data from our research, specifically that the nasal dorsum height in CL does not differ from controls (Fig2a, 2b).

The nasal bone length in CL/P was not significantly different compared with control (Fig.2c), which was consistent with the conclusion of other authors who have demonstrated that patients with CP +/- CL showed normal length of the nasal

bone^{13,14,16}. The length of the nasal dorsum of CL/P showed growth retardation, which was significantly shorter than that of controls in 8y-16y, but there was no significant difference in the 5y-6y and 17y-18y groups (Fig.2d). Moreira, et al.¹⁷ analyzed the lateral cephalometries of 70 white children with CL/P who had undergone primary operation and found that they had similar nasal dorsum length. But Ferrario, Chiarella, Claudia, Laura, Armando¹⁸ found that CL/P patients have a shorter nasal dorsum. Considering the limit of the above sample sizes, we tried to resolve this issue by enrolling a larger sample size, and analyzed the development of hard and soft tissues of the nose in different groups in detail. Additionally, in order to reduce the influence of maxillary hypoplasia, we presented a new evaluation method by utilizing Bookstein, FL and Nadia, H's design. The results showed nasal dorsum height in CL/P to be lower than controls (Fig2a,2b). Therefore, in addition to maxillary hypoplasia, CL/P also demonstrates a flatter nasal dorsum.

CL and CL/P both had a shorter lower nasal dorsum than control (Fig.2f). Shape changes of the nasal dorsum are most closely related to angulation changes of the lower dorsum¹⁹, which may emphasize the malformation of the lower nasal dorsum leading to the whole nasal deformity. The upper nasal dorsum of CL/P was longer than controls, and the difference in 14y-18y was statistically significant (Fig.2e). However, there was no significant difference in the length of the entire nasal dorsum in 17y-18y (Fig.2d). We conclude that overdevelopment of upper nasal dorsal length in CL/P compensates for hypodevelopment of lower nasal dorsal length, and the net result leads to a similar

length of the entire nasal dorsum compared with controls when growth is completed.

However, the developmental mechanism of the upper nasal dorsum deserved further

elucidation.

The deformity of the nasal dorsum in CL/P is mainly due to underdevelopment of

the height of the hard and soft tissues of the nose, and patients with CL present a shorter

nose instead of a flatter one. The characteristic features of the nose for CL, CL/P, and

control groups in 17y-18y are shown in Fig.3, providing a basis for a specific approach

to secondary rhinoplasty. Most of all, our study confirms that different types of clefts

indeed influence the features of nasal dorsum deformity. A flatter nasal dorsum

contributes to a flatter profile in patients with CL/P. Hence according to our results, for

CL, secondary rhinoplasty should lengthen the nasal dorsum, and for CL/P, the aim of

surgery should make the nose more prominent.

Conclusion

In this study, we evaluated whether morphologic characteristics of the nasal dorsum

were affected by cleft types in different ages after primary operation. The results

indicate that isolated CL inhibits the growth of nasal dorsum length which leads to a

short nose deformity, while the C/LP patients tends to develop a saddle nose because

of reduced dorsal angulation which leads to a decreased nasal height.

237 **References**

- 238 1. Pawar, S. S., and Wang, T. D. Secondary cleft rhinoplasty. *Jama Facial Plastic*
239 *Surgery*. 2014; 16(1):58.
- 240 2. Krane, N. A., J. D. Markey, L. B. Moneta, and M. M. Kim. Aesthetics of the Nasal
241 Dorsum: Proportions, Light, and Shadow. *Facial Plastic Surgery*. 2017
242 33(2):120-4 .
- 243 3. Henry C, Samson T, Mackay D. Evidence-based medicine: The cleft lip nasal
244 deformity. *Plastic & Reconstructive Surgery*. 2014; 133(5):1276-88.
- 245 4. Sykes JM, Tasman AJ, Suarez GA. Cleft Lip Nose. *Clinics in plastic surgery*.
246 2016;43(1):223-35.
- 247 5. Pereira ND, Tinoco C, Oliveira ECD, Paço J. Intermediate osteotomies in
248 rhinoplasty: a new perspective. *European Archives of Oto-Rhino-Laryngology*.
249 2017;274(7):2953-8.
- 250 6. Aston SJ, Guy CL. The nasal spine. *Clinics in plastic surgery*. 1977; 4(1):153-62.
- 251 7. He, X., Shi, B., Li, S., Zheng, Q.: A geometrically justified rotation advancement
252 technique for the repair of complete unilateral cleft lip. *J Plast Reconstr Aesthet*
253 *Surg*. 2009;62(9):1154-60.
- 254 8. Bookstein FL. *Morphometric Tools for Landmark Data*. 1991.
- 255 9. Hasanzadeh, N., Majidi, M. R., Kianifar, H., and Eslami, N. Facial soft-tissue
256 morphology of adolescent patients with nonsyndromic bilateral cleft lip and
257 palate. *Journal of Craniofacial Surgery*. 2014; 25(1):314-7.

- 258 10. Hopkins, Dr Will G. Measures of Reliability in Sports Medicine and Science.
259 Sports Medicine. 2000; 30(1):1.
- 260 11. Liu R., Wamalwa P, Lu DW, Li CH, Hu HK, Zou S.. Soft-Tissue Characteristics of
261 Operated Unilateral Complete Cleft Lip and Palate Patients in Mixed Dentition.
262 Journal of Craniofacial Surgery. 2011; 22(4):1275-9.
- 263 12. Liu, R. , Lu, D. , Wamalwa, P. , Li, C. , Hu, H. , and Zou, S.. Craniofacial morphology
264 characteristics of operated unilateral complete cleft lip and palate patients in
265 mixed dentition. Journal of Craniofacial Surgery. 2011;112(6):e16-25.
- 266 13. Hansen L, Skovgaard LT, Nolting D, Hansen BF, Kjaer I. Human prenatal nasal
267 bone lengths: normal standards and length values in fetuses with cleft lip and
268 cleft palate. The Cleft palate-craniofacial journal : official publication of the
269 American Cleft Palate-Craniofacial Association. 2005;42(2):165-70.
- 270 14. Nielsen BW, Mølsted K, Skovgaard LT, Kjaer I. Cross-sectional study of the
271 length of the nasal bone in cleft lip and palate subjects. The Cleft palate-
272 craniofacial journal : official publication of the American Cleft Palate-
273 Craniofacial Association. 2005;42(4):417.
- 274 15. Miyamoto J, Nakajima T. Anthropometric evaluation of complete unilateral
275 cleft lip nose with cone beam CT in early childhood. Journal of Plastic
276 Reconstructive & Aesthetic Surgery. 2009;63(1):9-14.
- 277 16. Goyenc, Y. B., H. G. Gurel, and B Memili. Craniofacial morphology in children
278 with operated complete unilateral cleft lip and palate. Journal of Craniofacial

- 279 Surgery. 2008;19(5):1396-401.
- 280 17. Moreira I, Suri S, Ross B, Tompson B, Fisher D, Lou W. Soft-tissue profile
281 growth in patients with repaired complete unilateral cleft lip and palate: A
282 cephalometric comparison with normal controls at ages 7, 11, and 18 years.
283 American Journal of Orthodontics and Dentofacial Orthopedics.
284 2014;145(3):341-58.
- 285 18. Ferrario VF, Chiarella S, Claudia D, Laura V, Armando C. Three-dimensional
286 nasal morphology in cleft lip and palate operated adult patients. Ann Plast Surg.
287 2003;51(4):390-7.
- 288 19. Buschang, P. H., La Cruz R De, A. D. Viazis, and A Demirjian. Longitudinal shape
289 changes of the nasal dorsum. American journal of orthodontics and dentofacial
290 orthopedics : official publication of the American Association of Orthodontists,
291 its constituent societies, and the American Board of Orthodontics. 1993;
292 104(6):539-543.

294 **Author information**

295 **Affiliations**

296 State Key Laboratory of Oral Disease, West China College of Stomatology, Sichuan
297 University
298 Department of Cleft Lip and Palate Surgery, West China Stomatological Hospital,
299 Sichuan University

300 Lingling Pu Email: plldangna@gmail.com

301 Renkai Liu Email:liurenkai_cn@hotmail.com

302 Bing Shi Email: shibingcn@vip.sina.com

303 Chenghao Li Email: leechenghao_cn@yahoo.com

304

305 Division of Plastic Surgery, The Children's Hospital of Philadelphia

306 David W Low Email: LOW@email.chop.edu

307

308 **Contributions**

309 All authors contributed extensively to this work. Conceived and designed the

310 experiments: L.L.P. and C.H.L. Performed the experiments: L.L.P. and R.K.L.

311 Analyzed the data: L.L.P. and B.S. Interpreted the data: L.L.P. and C.H.L. Wrote the

312 paper: L.L.P., D.W.L, and C.H.L. All authors reviewed the manuscript.

313 **Competing Interests**

314 The authors declare no competing interests.

315 **Corresponding author**

316 Correspondence to Chenghao Li

317

318

319

320

321

Fig1. The profile cephalometric radiographs and the cephalometric radiographs. (A)Reference points on the profile cephalometric radiographs. S =sella, the center of sella turcica; n =nasion, junction of frontal, maxillary, and nasal bones; N=soft nasion, closest point on soft tissue outline from hard tissue nasion; na=nasale, point at the most anterior inferior part of the nasal bone; NA= soft nasale, closest point on soft tissue outline from hard tissue nasale; Prn=pronasale, most anterior point on the contour of nose. (B) Measurements of angles and lines on the profile cephalometric radiographs. L1=vertical line of S to n, n is the foot point; L2=parallel to L1; na-n-L1(degrees):angulation of nasal bone; the angle between na-n-L1 ; prn-N-L2(degrees): angulation of nasal dorsum; the angle between prn-N-L2 ; n-na(mm):length of nasal bone, from the nasion to nasale; N-prn(mm):length of nasal dorsum,from soft-tissue nasion to pronasale; N-NA(mm):upper nasal dorsum; NA-prn(mm):lower nasal dorsum. (C) Measurements of angles and lines on the lateral cephalometric radiograph.

Fig.2 (A) Angulation of the nasal bone changes by years in different groups;(B) Angulation of the nasal dorsum changes by years in different groups;(C) Length of the nasal bone changes by years in different groups;(D) Length of the nasal dorsum changes by years in different groups;(E) Length of the upper nasal dorsum changes by years different groups;(F) Length of the lower nasal dorsum changes by years in different groups.

Fig.3 The characteristic features of 17y-18y group determined from our results. The CL patients have a shorter nasal dorsum. The CL/P patients have a flatter nasal dorsum.

Table1. Sample distribution by age

	5-7y				8y-10y				11y-13y				14y-16y				17y-18y			
	N	mean	SD	p	N	mean	SD	p	N	mean	SD	p	N	mean	SD	p	N	mean	SD	p
CL	44	5.9	0.80		35	8.9	0.73		67	12.0	0.80		78	15.3	0.75		54	17.5	0.50	
CLP	68	6.1	0.84		63	8.9	0.84		63	11.9	0.74		58	15.1	0.83		46	17.5	0.50	
				0.17				0.79				0.90				0.17				0.91
Control	64	6.1	0.77		74	9.0	0.83		75	12.0	0.82		73	15.0	0.82		47	17.5	0.51	
Total	176	6.1	0.81		172	9.0	0.81		205	12.0	0.79		209	15.1	0.80		147	17.5	0.50	

Table1 shows the sample distribution by age. There is no difference in sample distribution by age in each Group

Table2. Reliability analysis

intraobserver concordance			
Measurement	ICC	95%CI	
na-n-L1(degrees)	0.98	0.97	0.99
prn-N-L2(degrees)	0.9	0.86	0.93
n-na(mm)	0.97	0.96	0.98
N-prn(mm)	0.97	0.96	0.98
N-NA(mm)	0.97	0.96	0.98
NA-prn(mm)	0.96	0.94	0.97

Table2 shows the reliability analysis. The ICC results for test-retest reliability ranged between 0.90 and 0.98, suggesting dependable reliability and reproducibility of the adopted measuring strategy.

Table3. Statistical descriptions of the angle of nasal bone, the angle of nasal dorsum, the length of nasal bone, the length of nasal dorsum, the length of upper nasal dorsum and the length of lower nasal dorsum by ages and results of Dunn's-test between CLP, CL and Control separately

		CLP			CL			Control			CLP vs Control	CL vs Control
		Median	P25	P75	Median	P25	P75	Median	P25	P75		
na-n-L1(degrees)	5y - 7y	8.98	5.04	12.54	10.01	8.66	13.96	11.08	7.67	14.10	NS	NS
	8y - 10y	7.43	4.63	11.13	10.48	7.59	13.39	12.08	8.43	14.72	<0.001***	NS
	11y - 13y	11.63	7.50	15.60	14.34	10.04	18.10	15.69	12.51	19.41	<0.001***	NS
	14y - 16y	10.37	7.75	12.68	18.21	13.40	21.30	17.67	14.59	20.90	<0.001***	NS
	17y - 18y	10.38	7.51	15.02	17.19	13.62	20.98	20.33	15.89	22.72	<0.001***	NS
prn-N-L2(degrees)	5y - 7y	15.19	12.42	17.50	19.64	16.80	21.43	19.62	16.84	22.72	<0.001***	NS
	8y - 10y	15.04	11.44	17.65	18.38	16.69	20.37	20.41	18.21	22.71	<0.001***	NS
	11y - 13y	17.26	14.34	21.10	21.82	19.27	24.73	22.08	19.20	24.99	<0.001***	NS
	14y - 16y	15.74	12.26	19.27	23.08	20.24	26.87	23.75	20.04	25.84	<0.001***	NS
	17y - 18y	16.49	12.41	20.16	22.00	18.27	26.50	24.14	21.61	25.97	<0.001***	NS
n-na(mm)	5y - 7y	20.98	19.55	22.48	19.50	17.47	20.54	21.46	20.19	23.04	NS	<0.001***
	8y - 10y	22.76	20.58	24.85	21.05	19.71	22.95	23.17	21.63	23.99	NS	0.003**
	11y - 13y	24.00	22.17	25.60	22.85	21.44	24.56	24.79	23.09	27.22	0.045*	<0.001***
	14y - 16y	24.72	22.02	26.53	23.09	21.16	25.73	25.73	23.18	27.70	NS	0.005**
	17y - 18y	25.61	24.34	27.75	22.94	20.86	25.89	25.51	23.26	27.28	NS	0.008**
N-prn(mm)	5y - 7y	32.78	30.70	35.79	28.83	27.20	30.65	34.08	31.32	36.59	NS	<0.001***
	8y - 10y	34.00	30.44	35.64	32.91	30.77	35.63	36.66	34.46	38.73	<0.001***	<0.001***
	11y - 13y	35.58	32.54	38.04	35.42	33.66	37.67	39.49	37.09	41.40	<0.001***	<0.001***
	14y - 16y	38.05	36.69	40.79	36.78	34.94	39.63	40.58	37.96	42.95	0.012*	<0.001***
	17y - 18y	40.46	38.10	45.20	38.23	35.71	41.07	41.29	39.55	44.14	NS	<0.001***
N-NA(mm)	5y - 7y	14.75	12.69	17.03	10.81	9.73	12.80	14.34	13.07	15.89	NS	<0.001***
	8y - 10y	15.38	12.70	17.39	13.65	12.20	15.81	14.89	13.69	16.76	NS	NS
	11y - 13y	15.92	13.59	18.37	15.43	13.86	16.84	16.01	13.77	18.12	NS	NS
	14y - 16y	17.00	15.16	19.37	15.55	13.63	17.82	15.21	14.02	16.95	0.012*	NS
	17y - 18y	19.20	16.02	20.62	15.21	13.00	16.44	15.76	13.98	17.07	0.01**	NS
NA-prn(mm)	5y - 7y	18.18	16.82	19.30	17.59	16.23	18.93	18.97	18.02	21.52	0.007**	<0.001***
	8y - 10y	18.29	16.79	19.22	19.64	17.07	20.71	21.61	20.40	22.68	<0.001***	<0.001***
	11y - 13y	20.01	18.10	21.31	20.27	19.04	22.18	23.49	22.01	24.81	<0.001***	<0.001***
	14y - 16y	21.02	19.23	22.81	21.68	20.02	22.96	24.42	23.16	26.21	<0.001***	<0.001***
	17y - 18y	22.84	20.61	24.72	23.30	21.19	25.28	25.40	24.00	27.31	<0.001***	<0.001***

Abbreviations: na-n-L1(degrees):nasal bone angle;prn-N-L2(degrees):nasal dorsum angle; n-na(mm):the length of nasal bone; N-prn(mm):the length of nasal dorsum; N-NA(mm):the length of upper nasal dorsum; NA-prn(mm):the length of lower nasal dorsum.*p<.05, **p<.01, ***p<.001. NS =no significance

453 **References**

- 454 1. Pawar, S. S., and Wang, T. D. Secondary cleft rhinoplasty. *Jama Facial Plastic*.
455 *Surgery*. 2014; 16(1):58.
- 456 2. Krane, N. A., J. D. Markey, L. B. Moneta, and M. M. Kim. *Aesthetics of the Nasal*
457 *Dorsum: Proportions, Light, and Shadow*. *Facial Plastic Surgery*. 2017
458 33(2):120-4 .
- 459 3. Henry C, Samson T, Mackay D. Evidence-based medicine: The cleft lip nasal
460 deformity. *Plastic & Reconstructive Surgery*. 2014; 133(5):1276-88.
- 461 4. Sykes JM, Tasman AJ, Suarez GA. Cleft Lip Nose. *Clinics in plastic surgery*.
462 2016;43(1):223-35.
- 463 5. Pereira ND, Tinoco C, Oliveira ECD, Paço J. Intermediate osteotomies in
464 rhinoplasty: a new perspective. *European Archives of Oto-Rhino-Laryngology*.
465 2017;274(7):2953-8.
- 466 6. Aston SJ, Guy CL. The nasal spine. *Clinics in plastic surgery*. 1977; 4(1):153-62.
- 467 7. He, X., Shi, B., Li, S., Zheng, Q.: A geometrically justified rotation advancement
468 technique for the repair of complete unilateral cleft lip. *J Plast Reconstr Aesthet*
469 *Surg*. 2009;62(9):1154-60.
- 470 8. Bookstein FL. *Morphometric Tools for Landmark Data*. 1991.
- 471 9. Hasanzadeh, N., Majidi, M. R., Kianifar, H., and Eslami, N. Facial soft-tissue
472 morphology of adolescent patients with nonsyndromic bilateral cleft lip and
473 palate. *Journal of Craniofacial Surgery*. 2014; 25(1):314-7.

- 474 10. Hopkins, Dr Will G. Measures of Reliability in Sports Medicine and Science.
475 Sports Medicine. 2000; 30(1):1.
- 476 11. Liu R., Wamalwa P, Lu DW, Li CH, Hu HK, Zou S.. Soft-Tissue Characteristics of
477 Operated Unilateral Complete Cleft Lip and Palate Patients in Mixed Dentition.
478 Journal of Craniofacial Surgery. 2011; 22(4):1275-9.
- 479 12. Liu, R. , Lu, D. , Wamalwa, P. , Li, C. , Hu, H. , and Zou, S.. Craniofacial morphology
480 characteristics of operated unilateral complete cleft lip and palate patients in
481 mixed dentition. Journal of Craniofacial Surgery. 2011;112(6):e16-25.
- 482 13. Hansen L, Skovgaard LT, Nolting D, Hansen BF, Kjaer I. Human prenatal nasal
483 bone lengths: normal standards and length values in fetuses with cleft lip and
484 cleft palate. The Cleft palate-craniofacial journal : official publication of the
485 American Cleft Palate-Craniofacial Association. 2005;42(2):165-70.
- 486 14. Nielsen BW, Mølsted K, Skovgaard LT, Kjaer I. Cross-sectional study of the
487 length of the nasal bone in cleft lip and palate subjects. The Cleft palate-
488 craniofacial journal : official publication of the American Cleft Palate-
489 Craniofacial Association. 2005;42(4):417.
- 490 15. Miyamoto J, Nakajima T. Anthropometric evaluation of complete unilateral
491 cleft lip nose with cone beam CT in early childhood. Journal of Plastic
492 Reconstructive & Aesthetic Surgery. 2009;63(1):9-14.
- 493 16. Goyenc, Y. B., H. G. Gurel, and B Memili. Craniofacial morphology in children
494 with operated complete unilateral cleft lip and palate. Journal of Craniofacial

Surgery. 2008;19(5):1396-401.

17. Moreira I, Suri S, Ross B, Tompson B, Fisher D, Lou W. Soft-tissue profile growth in patients with repaired complete unilateral cleft lip and palate: A cephalometric comparison with normal controls at ages 7, 11, and 18 years. American Journal of Orthodontics and Dentofacial Orthopedics. 2014;145(3):341-58.

18. Ferrario VF, Chiarella S, Claudia D, Laura V, Armando C. Three-dimensional nasal morphology in cleft lip and palate operated adult patients. Ann Plast Surg. 2003;51(4):390-7.

19. Buschang, P. H., La Cruz R De, A. D. Viazis, and A Demirjian. Longitudinal shape changes of the nasal dorsum. American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics. 1993; 104(6):539-543.

Fig.1

Fig.2

Control

CL

CLP

bioRxiv preprint doi: <https://doi.org/10.1101/806406>; this version posted October 15, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Fig.3