Stimulus-specific neural encoding of a persistent, internal defensive state in the hypothalamus

Ann Kennedy^{1*}, Prabhat S. Kunwar^{1*†}, Lingyun Li¹, Daniel Wagenaar¹, and David J. Anderson^{1,2,3}

¹Division of Biology and Biological Engineering 156-29, Tianqiao and Chrissy Chen Institute for Neuroscience, California Institute of Technology, Pasadena, California 91125, USA

²Howard Hughes Medical Institute, California Institute of Technology, 1200 East California Blvd, Pasadena, California 91125 USA

³Author for correspondence

*These authors contributed equally to this work.

[†]Present address: Biogen, 225 Binney St. Cambridge, MA 02142 USA

14 Summary

15 Persistent neural activity has been described in cortical, hippocampal, and motor networks as
 16 mediating short-term working memory of transiently encountered stimuli¹⁻⁴. Internal emotion states
 17 such as fear also exhibit persistence following exposure to an inciting stimulus^{5,6}, but such
 18 persistence is typically attributed to circulating stress hormones⁷⁻⁹; whether persistent neural activity
 19 also plays a role has not been established. SF1⁺/Nr5a1⁺ neurons in the dorsomedial and central
 20 subdivision of the ventromedial hypothalamus (VMHdm/c) are necessary for innate and learned
 21 defensive responses to predators¹⁰⁻¹³. Optogenetic activation of VMHdm^{SF1} neurons elicits defensive
 22 behaviors that can outlast stimulation^{11,14}, suggesting it induces a persistent internal state of fear or
 23 anxiety. Here we show that VMHdm^{SF1} neurons exhibit persistent activity lasting tens of seconds, in
 24 response to naturalistic threatening stimuli. This persistent activity was correlated with, and
 25 required for, persistent thigmotactic (anxiety-like) behavior in an open-field assay. Microendoscopic
 26 imaging of VMHdm^{SF1} neurons revealed that persistence reflects dynamic temporal changes in
 27 population activity, rather than simply synchronous, slow decay of simultaneously activated
 28 neurons. Unexpectedly, distinct but overlapping VMHdm^{SF1} subpopulations were persistently
 29 activated by different classes of threatening stimuli. Computational modeling suggested that
 30 recurrent neural networks (RNNs) incorporating slow excitation and a modest degree of
 31 neurochemical or spatial bias can account for persistent activity that maintains stimulus identity,
 32 without invoking genetically determined “labeled lines”¹⁵. Our results provide causal evidence that
 33 persistent neural activity, in addition to well-established neuroendocrine mechanisms, can
 34 contribute to the ability of emotion states to outlast their inciting stimuli, and suggest a mechanism

35 that could prevent over-generalization of defensive responses without the need to evolve hardwired
36 circuits specific for each type of threat. (267 words)

37 Main Text

38 We performed fiber photometry¹⁶ in VMHdm^{SF1} neurons expressing GCaMP6s¹⁷ in freely behaving mice
39 during a 10s presentation of a predator (an anesthetized rat¹⁸) (**Fig. 1a-c**). We observed a rapid
40 increase in the bulk calcium signal at the onset of rat presentation (**Fig. 1d, e**). Remarkably, this activity
41 persisted for over a minute following rat removal, decaying exponentially with time constant $\tau_{\text{decay}} =$
42 26.7 ± 2.2 seconds (**Fig. 1d-e, h-i; Supplemental Video 1**). In contrast, a toy rat presented in the same
43 manner produced a far weaker and less persistent response (**Fig. 1d, e**).

44 To better control the timing and location of stimulus presentation, we repeated this experiment
45 using a head-fixed preparation. The magnitude, decay constants and specificity of VMHdm^{SF1} persistent
46 responses were comparable to those measured in freely behaving mice (**Fig. 1f-i; Extended Data. 1**). As
47 discussed below, this persistence is unlikely to be due simply to slow GCaMP decay kinetics (**Fig. 2 and**
48 **Extended Data Fig. 5**). VMHdm^{SF1} neurons also responded to rat urine alone (**Extended Data Fig. 2**),
49 consistent with earlier studies^{19,20}.

50 To investigate whether VMHdm^{SF1} neurons were also active during persistent defensive
51 behaviors, we devised a novel rat exposure assay in an open arena. Following a ten-minute period of
52 habituation to the arena by the mouse, an awake rat in a cage was presented to the mouse for 15
53 seconds, and then removed. After rat exposure, mice exhibited thigmotaxis, an indication of increased
54 anxiety²¹, lasting for minutes (**Fig. 1l, m; control; blue line**). Thigmotaxis was not observed if the mouse
55 was introduced to the arena following, rather than before, rat presentation, indicating that it is unlikely

56 to be due to residual rat-derived odors (**Extended Data Fig. 3a-c**). Persistent thigmotaxis could also be
 57 evoked by brief optogenetic stimulation of VMHdm^{SF1} neurons (ref ¹¹ and **Extended Data Fig. 3d, e**)).
 58 Fiber photometry confirmed that VMHdm^{SF1} neurons were strongly and persistently activated for
 59 minutes following rat presentation in the arena (**Extended Data Fig. 4c-e**), with kinetics well correlated
 60 with behavior (**Extended Data Fig. 4f-i**).

61 We next tested a requirement for VMHdm^{SF1} neurons in the rat-evoked increase in thigmotactic
 62 behavior, using the light-gated chloride channel iC++²² to reversibly silence these cells. First, we
 63 confirmed that iC++ inhibition of VMHdm^{SF1} neurons in a home cage rat exposure assay increased the
 64 time the mice spent in close proximity to the rat (**Fig. 1j, k**), phenocopying genetic ablation of these
 65 neurons¹¹. Next, we repeated the open field rat exposure test while photo-inhibiting VMHdm^{SF1}
 66 neurons continuously for a 3 min period, either starting five seconds prior to rat introduction, or
 67 immediately following rat removal (**Fig. 1m; “light on,” red vs. green bars, respectively**).

68 When silencing was initiated five seconds prior to rat introduction, no significant rat-induced
 69 increase in thigmotaxis was observed (**Fig. 1m-o; red plots**). Importantly, when iC++ photostimulation
 70 was initiated only after rat removal, mice showed an initial increase in thigmotaxis, but quickly
 71 returned to their pre-rat baseline behavior (**Fig. 1m-o, green plots; Extended Data Fig. 4a**). These data
 72 indicate that VMHdm^{SF1} neuronal activity is essential for maintaining a persistent defensive behavioral
 73 response to a predator.

74 To investigate the neural dynamics underlying the rat-evoked persistent state, we next
 75 performed microendoscopic imaging^{23,24} of VMHdm^{SF1} neurons (**Fig. 2a-b**). Head-fixed mice were
 76 presented for ten seconds each with a pseudorandomized set of stimuli including a rat, toy rat, and

77 conspecific male, on each of three days of imaging ($n=5$ mice, 187.3 ± 8.1 cells imaged per day, 78 cells
78 tracked across days.) While VMHdm^{SF1} neurons, as a population, showed persistent activation
79 following rat presentation (**Fig. 2c-d**), individual VMHdm^{SF1} neurons showed diverse but reproducible
80 stimulus-evoked dynamics (**Fig. 2e-g**). Although many cells showed activation from stimulus onset
81 followed by slow decay, other cells reached their peak activation only after stimulus removal. Thus the
82 slow, monotonic decay of the population response reflects a diverse, time-evolving pattern of activity
83 at the level of individual cells (**Fig. 2j-p**). Application of a spike-deconvolution algorithm²⁵ indicated that
84 the persistent $\Delta F/F$ signal likely reflects persistent underlying spiking activity (**Extended Data Fig. 5**).
85 Strikingly, the rat and conspecific activated distinct subpopulations of VMHdm^{SF1} neurons with only
86 moderate overlap (**Fig. 2h, i**).

87 In rodents, nonvolatile odor cues can activate neurons of the vomeronasal organ (VNO) for
88 several seconds following inhalation²⁶. It was therefore possible that the persistent activity we
89 observed in VMHdm^{SF1} neurons reflected persistence of the stimulus signal. If this were the case, we
90 would expect VMHdm^{SF1} neurons to show only transient responses to time-resolved stimuli, such as
91 purely visual or auditory stimuli. VMHdm neurons are not activated by (**Extended Data Fig. 6**), or
92 required for¹¹ defensive responses to, an overhead visual threat stimulus²⁷, and have not previously
93 been shown to respond to auditory stimuli. We therefore imaged VMHdm^{SF1} neuron activity in
94 response to an auditory stimulus that evokes defensive behaviors in mice^{28,29}: a series of ultrasonic
95 sweeps (USS) in the frequency range typical of rat distress vocalizations³⁰ (Methods).

96 The USS strongly activated VMHdm^{SF1} neurons, and this activation persisted on a similar time
97 scale as that following rat exposure, after stimulus termination (**Fig. 3a-b**). Underlying this population
98 response, individual USS-responsive VMHdm^{SF1} neurons showed diverse dynamics, as was observed for

the rat and mouse stimuli (**Fig. 3c-f**). The USS activated VMHdm^{SF1} neurons were spatially intermingled with the rat (and mouse) -responsive populations, and overlapped them at a frequency equal to that expected by chance (**Fig. 3g-j; Extended Data Fig. 7**). This suggests that these populations are randomly distributed, but reproducibly activated by a given stimulus. Of the neurons showing a significant response to at least one of our five test stimuli (74.4%), 43.6% responded to only one stimulus, and over 70% of cells responded to ≤ 2 of the five tested stimuli (**Fig. 3k**). Most stimulus-responsive VMHdm^{SF1} neurons were excited, although a small number showed stimulus-evoked inhibition (**Fig. 3c-f, l**). Principal component analysis (PCA) indicated that in addition to stimulus identity, VMHdm^{SF1} neurons may encode an additional feature common to multiple stimuli (**Fig. 3o**, y-axis). Since this feature has higher values on initial vs. later trials, it may relate to novelty or salience. A 5-way Naïve Bayes decoder was able to predict stimulus identity in held-out trials across three days of imaging with above-chance accuracy (**Fig. 3m**). Importantly, despite the gradual decay in population activity following stimulus offset, stimulus identity could still be decoded with above chance accuracy for tens of seconds (**Fig. 3n, Extended Data Fig. 8**). Thus, the VMHdm^{SF1} population response can encode the identity of the presented stimulus, even after its removal.

We next used computational modeling to investigate the space of mechanisms that can account for persistent activity in VMHdm^{SF1} neurons, incorporating features that have been proposed in other systems^{2,3,31-36}. We evaluated four classes of models (**Fig 4a1-4**) in terms of their ability to capture two main features of VMHdm^{SF1} neural activity: it changes over time, and it is stimulus-specific. To achieve persistent activity with time-evolving dynamics that aligned with our experimental observations, it was necessary to combine two elements: inhibition-stabilized recurrent connectivity and slow excitation on a time scale of several seconds (mediated by, e.g., peptidergic transmission³⁷);

we call the resulting class of models peptidergic recurrent neural networks (pRNNs) (**Fig. 4a3**). The importance of feedback inhibition in our model suggests the existence of an inhibitory population that is activated by threatening stimuli; a likely candidate are GABAergic neurons in the surrounding DMH^{38,39}.

To compare the time-evolving population code of VMHdm^{SF1} neurons (**Fig4b**) with the dynamics of these models, we computed the autocorrelation matrix of stimulus-evoked activity for all values of t and t' between 0 and 45 s after stimulus onset(**Fig 4c**), and summarized it by computing the mean correlation over all neurons as a function of Δt ($t - t'$) (**Fig 4d**). Several versions of our pRNN models showed autocorrelation dynamics similar to observed dynamics (**Fig 4e**). Importantly, models that used slow peptidergic transmission alone (**Fig 4a1**), or RNNs with NMDA-mediated transmission⁴⁰ (**Fig 4a2**) could not match the diverse responses we observed in VMHdm^{SF1} neurons (**Fig. 4e1-2**).

The stimulus specificity of VMHdm^{SF1} responses (**Fig 4f**) was quantified as the time-evolving Pearson's correlation between rat- vs. USS-evoked activity (**Fig 4g**). Maintaining stimulus specific representations during persistence could only be achieved in the pRNN by increasing the gain (strength) of excitatory synapses^{41,42} (**Fig. 4a3, h3-4**), but this came at the cost of failing to match the autocorrelation dynamics of observed activity (**Fig e3-4**). However, both main features of the data could be matched if the synaptic time constant of peptidergic excitation was further increased from 6 to 20 seconds (**Fig 4a3,h5,e5**). Alternatively, both features could be matched by assuming that the probability of synapse formation between a pair of model neurons decreased slightly with the distance between them, and imposing a mild differential spatial bias on the responses to different stimuli (rat vs. USS) (**Fig 4a4,h6,e6, ED Fig 9**). Imposing further network structure produced too much segregation between stimulus representations, compared to the overlap observed (**ED Fig 9**). We summarized the

143 performance of our models by creating a pair of “data similarity scores” quantifying model similarity to
144 data in terms of time-evolving dynamics and stimulus specificity (**Fig 4i-k**).

145 Persistent defensive states are typically attributed to neuroendocrine mechanisms, such as
146 activation of the HPA axis⁷⁻⁹. Here we provide the first evidence that persistent neural activity can
147 contribute causally to such persistent internal states. VMHdm^{SF1} activity may also activate longer-
148 lasting neuroendocrine processes, as stimulation of VMHdm^{SF1} neurons elevates serum cortisol
149 levels¹¹. However unlike circulating hormones, persistent activity in VMHdm^{SF1} neurons is stimulus-
150 specific, and may thereby prevent over-generalization of defensive responses. The observed persistent
151 activity can be modeled best by recurrent excitatory networks incorporating fast feedback inhibition
152 and slow peptidergic transmission⁴³. VMHdm^{SF1} neurons are exclusively glutamatergic, densely
153 interconnected⁴⁴, and express several neuropeptides as well as neuropeptide receptors⁴⁵, properties
154 consistent with optimal model features. While our data do not exclude a role for interconnected
155 structures^{10,46} in establishing persistent activity in VMHdm, they demonstrate that hypothalamic
156 neuronal population dynamics contribute to the persistence of emotional behaviors.

ACKNOWLEDGMENTS

We thank H. Inagaki, M. Meister, L.F. Abbott, U. Rutishauser, and members of the Anderson lab for helpful comments on the manuscript, T. Davidson and K. Deisseroth for help implementing fiber photometry, R. Remedios for help with miniscope imaging, X. Da, J.S. Chang and C. Kim for technical assistance, X. Da and C. Chiu for laboratory management and G. Mancuso for administrative support. This work was supported by NIH Grant R01 MH112593. K99 MH117264 to A.K. and a Helen Hay Whitney Foundation Postdoctoral Fellowship to L. L. D.J.A. is an Investigator of the Howard Hughes Medical Institute.

WORKS CITED

- 1 Aksay, E. *et al.* Functional dissection of circuitry in a neural integrator. *Nat Neurosci* **10**, 494-504, doi:10.1038/nn1877 (2007).
- 2 Major, G. & Tank, D. Persistent neural activity: prevalence and mechanisms. *Current opinion in neurobiology* **14**, 675-684, doi:10.1016/j.conb.2004.10.017 (2004).
- 3 McCormick, D. A. Brain calculus: neural integration and persistent activity. *Nat Neurosci* **4**, 113-114, doi:10.1038/83917 (2001).
- 4 Curtis, C. E. & Lee, D. Beyond working memory: the role of persistent activity in decision making. *Trends Cogn Sci* **14**, 216-222, doi:10.1016/j.tics.2010.03.006 (2010).
- 5 Anderson, D. J. & Adolphs, R. *The Neuroscience of Emotion: A New Synthesis*. (Princeton University Press, 2018).
- 6 Anderson, D. J. & Adolphs, R. A framework for studying emotions across species. *Cell* **157**, 187-200, doi:10.1016/j.cell.2014.03.003 (2014).
- 7 Chen, A. Genetic Dissection of the Neuroendocrine and Behavioral Responses to Stressful Challenges in *Stem Cells in Neuroendocrinology* (eds D. Pfaff & Y. Christen) (2016).
- 8 McEwen, B. S. *et al.* Mechanisms of stress in the brain. *Nat Neurosci* **18**, 1353-1363, doi:10.1038/nn.4086 (2015).
- 9 Sapolsky, R. M., Romero, L. M. & Munck, A. U. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine reviews* **21**, 55-89, doi:10.1210/edrv.21.1.0389 (2000).
- 10 Gross, C. T. & Canteras, N. S. The many paths to fear. *Nature reviews. Neuroscience* **13**, 651-658, doi:10.1038/nrn3301 (2012).
- 11 Kunwar, P. S. *et al.* Ventromedial hypothalamic neurons control a defensive emotion state. *Elife* **4**, doi:10.7554/eLife.06633 (2015).
- 12 Silva, B. A. *et al.* Independent hypothalamic circuits for social and predator fear. *Nature neuroscience* **16**, 1731-1733, doi:10.1038/nn.3573 (2013).

- 13 Silva, B. A. *et al.* The ventromedial hypothalamus mediates predator fear memory. *Eur J Neurosci* **43**, 1431-1439, doi:10.1111/ejn.13239 (2016).
- 14 Wang, L., Chen, I. Z. & Lin, D. Collateral Pathways from the Ventromedial Hypothalamus Mediate Defensive Behaviors. *Neuron*, doi:10.1016/j.neuron.2014.12.025 (2015).
- 15 Ishii, K. K. *et al.* A Labeled-Line Neural Circuit for Pheromone-Mediated Sexual Behaviors in Mice. *Neuron* **95**, 123-137 e128, doi:10.1016/j.neuron.2017.05.038 (2017).
- 16 Gunaydin, L. A. *et al.* Natural neural projection dynamics underlying social behavior. *Cell* **157**, 1535-1551, doi:10.1016/j.cell.2014.05.017 (2014).
- 17 Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**, 295-300, doi:10.1038/nature12354 (2013).
- 18 Blanchard, D. C., Griebel, G. & Blanchard, R. J. in *Eur J Pharmacol* Vol. 463 97-116 (2003).
- 19 Papes, F., Logan, D. W. & Stowers, L. The vomeronasal organ mediates interspecies defensive behaviors through detection of protein pheromone homologs. *Cell* **141**, 692-703, doi:10.1016/j.cell.2010.03.037 (2010).
- 20 Perez-Gomez, A. *et al.* Innate Predator Odor Aversion Driven by Parallel Olfactory Subsystems that Converge in the Ventromedial Hypothalamus. *Current biology : CB* **25**, 1340-1346, doi:10.1016/j.cub.2015.03.026 (2015).
- 21 Simon, P., Dupris, R. & Costentin, J. Thigmotaxis as an index of anxiety in mice: influence of dopaminergic transmissions. *Behavioral brain research* **61**, 59-64 (1994).
- 22 Berndt, A. *et al.* Structural foundations of optogenetics: Determinants of channelrhodopsin ion selectivity. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 822-829, doi:10.1073/pnas.1523341113 (2016).
- 23 Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nat Neurosci* **16**, 264-266, doi:10.1038/nn.3329 (2013).
- 24 Remedios, R. *et al.* Social behaviour shapes hypothalamic neural ensemble representations of conspecific sex. *Nature* **550**, 388-392, doi:10.1038/nature23885 (2017).
- 25 Pnevmatikakis, E. A. *et al.* Simultaneous Denoising, Deconvolution, and Demixing of Calcium Imaging Data. *Neuron* **89**, 285-299, doi:10.1016/j.neuron.2015.11.037 (2016).
- 26 Ben-Shaul, Y., Katz, L. C., Mooney, R. & Dulac, C. In vivo vomeronasal stimulation reveals sensory encoding of conspecific and allospecific cues by the mouse accessory olfactory bulb. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 5172-5177, doi:10.1073/pnas.0915147107 (2010).
- 27 Yilmaz, M. & Meister, M. in *Current biology : CB* Vol. 23 2011-2015 (Elsevier Ltd, 2013).
- 28 Mongeau, R., Miller, G. A., Chiang, E. & Anderson, D. J. Neural correlates of competing fear behaviors evoked by an innately aversive stimulus. *J Neurosci* **23**, 3855-3868 (2003).
- 29 Evans, D. A. *et al.* A synaptic threshold mechanism for computing escape decisions. *Nature* **558**, 590-594, doi:10.1038/s41586-018-0244-6 (2018).
- 30 Brudzynski, S. Pharmacological and behavioral characteristics of 22 kHz alarm calls in rats. *Neurosci Biobehav R* **25**, 611-617 (2001).
- 31 Guo, Z. V. *et al.* Maintenance of persistent activity in a frontal thalamocortical loop. *Nature* **545**, 181-186, doi:10.1038/nature22324 (2017).
- 32 Inagaki, H. K., Fontolan, L., Romani, S. & Svoboda, K. Discrete attractor dynamics underlies persistent activity in the frontal cortex. *Nature* **566**, 212-217, doi:10.1038/s41586-019-0919-7 (2019).
- 33 Brody, C. D., Romo, R. & Kepecs, A. Basic mechanisms for graded persistent activity: discrete attractors, continuous attractors, and dynamic representations. *Current opinion in neurobiology* **13**, 204-211, doi:10.1016/s0959-4388(03)00050-3 (2003).
- 34 Goldman, M., Compte, A. & Wang, X.-J. Neural Integrators: recurrent mechanisms and models. *New Encyclopedia of Neuroscience*, 1-26 (2007).

- 35 Wang, X.-J. Synaptic reverberation underlying mnemonic persistent activity. *Trends in Neuroscience* **24**, 455-463 (2001).
- 36 Durstewitz, D., Seamans, J. K. & Sejnowski, T. J. Neurocomputational models of working memory. *Nat Neurosci* **3**, 1184 (2000).
- 37 Jan, Y. N., Jan, L. Y. & Kuffler, S. W. Further evidence for peptidergic transmission in sympathetic ganglia. *Proceedings of the National Academy of Sciences of the United States of America* **77**, 5008-5012, doi:10.1073/pnas.77.8.5008 (1980).
- 38 Canteras, N. S., Chiavegatto, S., Ribeiro do Valle, L. E. & Swanson, L. W. Severe reduction of rat defensive behavior to a predator by discrete hypothalamic chemical lesions. *Brain Res Bull* **44**, 297-305 (1997).
- 39 Zelikowsky, M. *et al.* The Neuropeptide Tac2 Controls a Distributed Brain State Induced by Chronic Social Isolation Stress. *Cell* **173**, 1265-1279 e1219, doi:10.1016/j.cell.2018.03.037 (2018).
- 40 Wang, X. J. Synaptic basis of cortical persistent activity: the importance of NMDA receptors to working memory. *J Neurosci* **19**, 9587-9603 (1999).
- 41 Sompolinsky, H., Crisanti, A. & Sommers, H.-J. Chaos in random neural networks. *Physical review letters* **61**, 259 (1988).
- 42 Toyozumi, T. & Abbott, L. F. Beyond the edge of chaos: amplification and temporal integration by recurrent networks in the chaotic regime. *Phys. Rev. E* **84**, 051908(051908), doi:10.1103/PhysRevE.84.051908 (2011).
- 43 Jan, Y. N., Jan, L. Y. & Kuffler, S. W. A peptide as a possible transmitter in sympathetic ganglia of the frog. *Proceedings of the National Academy of Sciences of the United States of America* **76**, 1501-1505, doi:10.1073/pnas.76.3.1501 (1979).
- 44 Lo, L. *et al.* Connectional architecture of a mouse hypothalamic circuit node controlling social behavior. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 7503-7512, doi:10.1073/pnas.1817503116 (2019).
- 45 Kim, D.-W. *et al.* Multimodal analysis of cell types in a hypothalamic node controlling social behavior. *Cell in press* (2019).
- 46 Canteras, N. S. The medial hypothalamic defensive system: hodological organization and functional implications. *Pharmacology, biochemistry, and behavior* **71**, 481-491 (2002).

Figure 1. Persistent activity in VMH SF1+ neurons evoked by predatory and conspecific cues.

a, Circuits for innate and learned fear. Abbreviations below. **b**, Site of fiber photometry in VMHdm/c (green). **c**, GCaMP6s expression in VMHdmSF1 neurons. **d**, Activity of SF1+ neurons in freely moving mice exposed to a live or toy rat for 10 seconds (gray shading). (n = 4 mice; mean $\pm$ SEM). **e**, Peak activity from (d) (n = 4 mice; mean $\pm$ SEM). **f**, Responses of VMHdmSF1 neurons in head-fixed mice to rat, mouse, or toy rat (n=8 mice; mean $\pm$ SEM). **g**, Peak activity from (f) (n = 8 mice; mean $\pm$ SEM). **h**, **i**, Decay constants (h) or time to half-peak (i) of activity in freely moving ("home cage") or head fixed mice (mean $\pm$ SEM; home cage n = 4; head-fixed n = 8). **j**, Home cage rat exposure assay. **k**, Percent time in zone 1 during 3-minute rat presentation (PS; n=7 control mice; n=7 iC++ mice; mean $\pm$ SEM). **l**, Tracking of mouse in open field rat exposure assay, blue line marks "edge zone". **m**, Fraction of time spent in edge zone. Colored horizontal bars denote PS periods (n = 12 control mice, n = 6 iC++ mice with PS during+after rat; n = 6 iC++ mice with PS after rat only. * p < 0.05; ** p < 0.01, repeated measures ANOVA test. Data are mean $\pm$ SEM.) **n**, Mean time in edge zone, times defined in Methods. **o**, Expression of anxiety behaviors (see Methods) before vs after rat exposure. (mean $\pm$ SEM).

VNO - vomeronasal organ, AOB - accessory olfactory bulb, MeApv -posteroventral medial amygdala, BNSTif - interfascicular part of bed nucleus of the stria terminalis, VMHdm - doromedial ventromedial hypothalamus, AHN - anterior hypothalamic nucleus, PMd - dorsal premammillary nucleus, PAG - periaqueductal gray, Thal - thalamus, LA - lateral amygdala, BS - brain stem, BLA - basolateral amygdala, CEA - central amygdala.

Figure 2. Microendoscopic imaging reveals persistent activity at the single cell level.

a, Microendoscopic imaging in VMHdm/c. **b**, Field of view in an imaged mouse. **c**, Mean population response of imaged neurons to each stimulus (n = 2 trials/mouse from 5 mice, mean $\pm$ SEM). **d**, Fit decay constants of population response to rat and mouse (n=5 mice, mean $\pm$ SEM). **e**, Rat- and mouse-responsive neuron responses (from n=5 mice, mean over 2 trials). **f**, Example cells responding to rat in one imaged mouse on two repeated trials. **g**, Example cells responding to mouse on two repeated trials (same mouse as (f)). **h**, Example spatial map of cells responsive to rat, mouse, or both (white). **i**, Histogram of cell tuning preference for rat vs. mouse. Cells at ± 1 respond exclusively to rat or mouse, respectively; cells at 0 ("both") respond equally to both stimuli (n = 219 cells from 5 mice across 3 days of imaging). **j**, Scatterplot comparing cell responses at 2 vs 20 seconds after rat introduction, in one example mouse. **k**, Peak time for rat-responsive cells (n = 202 rat-responsive cells from 5 mice across 3 days of imaging). **l**, Peak time for mouse-responsive cells (n = 160 mouse-responsive cells from 5 mice across 3 days of imaging). **m**, Fraction of cells with peak after time T. **n**, Half-peak time for rat-responsive cells (n same as k). **o**, Half-peak time for mouse-responsive cells (n same as l). **p**, Fraction of cells with half-peak later than time T, legend as in (m).

Figure 3. SF1+ neurons respond to a threatening auditory stimulus, and encode stimulus identity.

a, Mean VMHdmSF1 population response to USS and tone (n = 5 mice, mean $\pm$ SEM). **b**, Fit decay constants of population response to USS (rat reproduced from Fig2d for comparison). **c**, Example cells responding to USS in one imaged mouse, on two repeated stimulus presentations (dark trace: 1st presentation; light trace: 2nd presentation). Black = excited cells, red = inhibited cells. **d**, Response peak time of USS-responsive cells (n = 133 USS-responsive cells from 5 mice across 3 days of imaging). **e**, Response half-peak time for USS-responsive cells (n same as (d)). **f**, Example spatial map of cells responsive to rat, USS, or both (white). **g**, Histogram of cell tuning preference, rat vs USS (n = 216 cells from 5 mice across 3 days of imaging). **h**, Example spatial map of cells responsive to mouse and USS. **i**, Histogram of cell tuning preference, mouse vs USS (n = 219 cells from 5 mice across 3 days of imaging). **j**, Percent of cells responding to zero to five out of five stimuli. **k**, Percent of cells excited or inhibited by each stimulus. **l**, Accuracy of a 5-way Naïve Bayesian decoder for stimulus identity, trained on six trials across three days of imaging (n=5 mice; mean $\pm$ SEM). **m**, Decoder accuracy as a function of time. **n**, Principal component analysis (PCA) of time-averaged population responses, pooled across mice, from 5 trials per stimulus across 3 days of imaging.

Figure 4. Data constrains set of possible mechanisms for persistent neural activity.

a Six tested models of persistent activity. **a1** slow-acting neuropeptide activation, no connectivity between model neurons. **a3** recurrent excitation in a randomly connected network with fast inhibitory feedback; persistence maintained via NMDA channels, time constant of NMDA excitation is $\tau_s = 200\text{ms}$. **a3** as in **a2**, replacing NMDA with slow peptidergic excitation (light/dark blue, $\tau_s = 6\text{sec}$; black, $\tau_s = 20\text{sec}$), tested for different strengths of recurrent synapses (“gain scaling”, light/dark blue). **a4** as in **a3**, but with “local connectivity” in which probability of a synapse $p(\text{syn})$ between neurons decreased with distance (inset). **b** Trial-averaged, normalized $\Delta F/F$ traces from USS-responsive neurons, sorted by time of response peak. **c** Autocorrelation matrix of USS-evoked population activity. **d** Time-averaged autocorrelation of USS-evoked population activity ($n=5$ mice, mean $\pm$ SEM). **e** Autocorrelation as in **d**, for each model; model identity indicated by line color (legend in panel **h**), dashed line = data ($n=10$ repeat simulations, mean $\pm$ SEM). **f** Trial-averaged $\Delta F/F$ traces from rat- or USS-responsive neurons, sorted by projection on first principal component. **g** Pearson’s correlation between rat- vs USS-evoked population activity as a function of time. **h** Pearson’s correlation between simulated “rat” and “USS” inputs to each model, dashed line = data ($n=10$ repeat simulations, mean $\pm$ SEM). **i** “Similarity score” (see Methods) of models vs. data, summarizing plots in (**e**). ($n=5$ mice, mean $\pm$ SEM). **j** Similarity score of model vs data, summarizing plots in **h**. ($n=5$ mice, mean $\pm$ SEM). **k** scatter plot of **i-j**.

METHODS

Animals. All experimental procedures involving the use of live animals or their tissues were performed in accordance with the NIH guidelines and approved by the Institutional Animal Care and Use Committee (IACUC) and the Institutional Biosafety Committee at the California Institute of Technology (Caltech). SF1-Cre mice were obtained from Dr. Brad Lowell¹ and maintained as heterozygotes in the Caltech animal facility as described previously; the SF1-Cre line is also available from the Jackson Laboratory (Stock No: 012462). An account of the specificity of SF1-Cre expression within VMH and characterization of neurons labeled by Cre-expression can be found in². Male mice, heterozygotes or their wild-type littermates aged between 8 to 20 weeks were used in this study. All mice were housed in ventilated micro-isolator cages in a temperature- and humidity- controlled environment under a reversed 12 hour dark-light cycle, and had free access to food and water. Mouse cages were changed weekly on a fixed day on which experiments were not performed. Long-Evans rats (for use as predators) were obtained from Charles River at 2-3 months of age, and raised to 5-10 months in the Caltech animal facilities.

Virus. AAV1.Syn.Flex.GCaMP6s.WPRE.SV40 (CS1113) was obtained from the Penn Vector Core. AAV5.EF1a.DIO.iC++eYFP and AAV2.EF1a.DIO.hChR2.eYFP.WPRE.pA were obtained from the University of North Carolina Vectors Core.

Surgery. Mice 8-20 weeks old were anesthetized with 5% isoflurane and mounted in a stereotaxic apparatus (Kopf Instruments). 1% - 1.5% isoflurane was used to maintain the anesthesia throughout the surgery procedure. An incision was made to exposure the skull and small craniotomies were made dorsal to each injection site with a stereotaxic mounted drill. Virus suspension (~600 nl) was injected to the VMHdm/c (ML +/- 0.5, AP -4.65, DV -5.6) at a rate of 60 nl/minute using a pulled glass capillary (~40 µm inner diameter at tip) mounted in a nanoliter injector (Nanoliter 2000, World Precision Instruments) controlled by a four channel micro controller (Micro4, World Precision Instruments). Capillaries were kept in place for 10 minutes following injections to allow the adequate diffusion of virus solution and to reduce the virus backflow during capillary withdraw.

For fiber photometry, a custom-made unilateral fiber cannula (400 µm in core diameter, 0.48 NA, Doric Lenses) was implanted after virus injection (ML +/-0.4, AP -4.65, DV -5.4). Metabond (Parkell) and dental cement (Bosworth) were applied to secure the implanted ferrule and cover the exposed skull. For optogenetics, a custom-made bilateral fiber cannula aimed 500 µm above each injection site (200 µm in core diameter, 0.37 NA, Doric Lenses) was implanted and held in place with Metabond and dental cement.

Surgery for microendoscopic imaging was performed as previously described³. Briefly, we first performed a series of titration experiments of the original viral stock, to determine the virus concentration at which the brightest cytoplasmic but non-nuclear GCaMP6s expression could be observed in slices of fixed brain tissue of the injected mice 4 weeks after injection. The optimal viral dilution was then used to inject mice for *in vivo* imaging as described above. 2-3 weeks after viral injections, mice were implanted with a graded-index (GRIN) lens (diameter - 0.5 mm, length - 8.4 mm, catalogue #1050-002212, Inscopix) using a supporting device (Proview Implant Kit, cat# 1050-002334, Inscopix). The implantation depth of the lens was determined based on the live visualization of (anesthetized) neural activity as the lens was inserted. Metabond was used to stabilize the lens, and Kwik-Sil sealant (World Precision Instruments) was used to cover the lens surface. After another 2-3 weeks, mice were anesthetized for placement of a microendoscope baseplate (cat# 1050-002192, Inscopix) and a baseplate cover (catalogue #1050-002193,

Inscopix) was used to protect the lens when not in use. Five out of twenty implanted animals were selected for *in vivo* imaging studies based on clarity of cytoplasmic GCaMP6s expression.

Stimuli Presentation. Stimuli were presented either in the mouse's home cage or in a head-fixation set up. In the home cage, a hand-held anesthetized rat weighing 400-600 g was brought in close proximity to the mouse. A stuffed toy rat of approximately the same size as the live rat was used as a control. For the head-fixed preparation, the mouse was placed on a plastic running wheel (15.5 cm diameter) and stabilized by the head-plate (World Precision Instruments, Catalogue #503617) with a custom made tethering system. Animals were habituated to the head-fixation setup for 1 hour each day for 2-3 days before experiments began. Physical stimuli (an awake behaving rat, a conspecific BALB/6 male mouse or a toy rat) were each presented inside a small wire mesh cage, which was held by the experimenter in front of the experimental mouse. Auditory stimuli were presented at 85 dB SPL from above the animal. The ultrasound stimulus (USS) consists of repeated 100 ms frequency sweeps from 17-20 kHz, as described previously⁴. A pure tone of 2 kHz was used as a control. Rat urine was collected in-house and kept at 4°C for up to two weeks. A cotton swab soaked with 100 µl of rat urine or water was presented in front of the experimental mouse. 500 ms looming stimulus was displayed on an overhead screen above the mouse home cage 10 times with 500 ms inter stimulus interval. All stimuli were pseudo-randomized and presented for 10 seconds unless otherwise clarified, with an inter-trial interval of at least five minutes. For microendoscopic imaging, two trials for each stimulus were presented on each of three consecutive days.

Optogenetic manipulation. Optogenetic experiments were performed as described in². Animals were briefly anaesthetized by isoflurane to connect the fiberoptic patch cord to the bilateral implanted optic cannula (Doric Lenses). Mice were then allowed to recover for at least 15-20 minutes in their home cage before being transferred to the behavioral testing room. Light for both iC++ and ChR2 activation was delivered via a 473nm laser (Shanghai Laser) controlled by a signal generator (A-M systems, isolated pulse stimulator). Laser intensity was calibrated at the distance of 0.5 mm below the implanted fiber tip. 3 minutes continuous light was used for iC++ activation; 10 seconds (20 Hz, 20 ms pulse width) pulse trains was used for ChR2 activation.

Home cage rat exposure assay. The mouse home cage was placed into a custom made testing apparatus (35 x 40 x 40 cm), and video of behavior was collected from a side-view camera. After a 6 minute baseline, a predator rat in a cage with a mesh wall (10 x 20 x 35 cm) was placed at one end of the mouse home cage. Ethovision XT software was used to track mouse position and quantify time spent in proximity to the rat.

Open field rat exposure assay. The mouse was placed in a plastic open top arena (50 x 50 cm, 30 cm walls), with behavior captured using an overhead mounted camera. Following a 10 minute baseline, a rat held in a cage with a mesh wall was held in close proximity to the mouse for 15 seconds, and then removed. Behavior of the mouse was then recorded for an additional 6 minutes. For behavior quantification, Ethovision tracking data was segmented into 30-second chunks, and percent of time spent in the "edge zone" (within 4cm of arena walls) was quantified. For bar graphs in **Fig 1n**, we define before rat = average over a window from -1 to 0 min relative to rat presentation, after rat = average from 0-1 min after the rat was removed, and after PS off = average from 3-4 min after rat was removed. Anxiety behaviors for **Fig 1o** were defined as thigmotaxis, immobility, and jumping (escape attempts) and were manually annotated

at 30Hz. Pre and post windows correspond to -3 to 0 and 0 to 3 min, respectively, relative to rat presentation.

Fiber photometry data acquisition and processing. Fiber photometry was performed as described in ⁵. Briefly, two LEDs modulated at different frequencies (490 nm and 405 nm, Thorlabs) were used to excite GCaMP6s-expressing neurons via implanted optical fiber. Excitation light at 490 nm activates GCaMP6s in a calcium-dependent manner, while excitation at 405 nm activates GCaMP6s in a calcium-independent manner, thus the 405nm signal can be used to control for bleaching and movement artifacts in the 490nm channel. A photometer (Newport Femtowatt Photoreceiver) received GCaMP6s fluorescent signals, and custom-designed software separated the signals generated by the two LEDs. The output power of both LED was set between 30-50 μ W at the fiber tip to obtain an optimal baseline fluorescence without photobleaching.

To calculate $\Delta F/F$ of the 490nm signal, we normalized it to the 405nm baseline as in ⁵. The 405nm signal was scaled to match the amplitude of the 490nm signal using linear regression, and $\Delta F/F$ computed as (490nm signal – scaled 405nm signal) / (scaled 405 nm signal).

Microendoscopic imaging data acquisition and processing. We used a head-mounted miniaturized microscope (nVista, Inscopix) for calcium imaging. Pilot experiments were done to identify imaging parameters that produced the clearest signal to noise ratio while limiting photobleaching. All mice except one were recorded at 11 Hz with 90.0ms exposure time, 10-20% LED illumination and 1.5 - 2.5x gain; the remaining mouse was imaged at 20Hz with 50ms exposure time. A custom-built system was used to synchronize the cameras for behavioral recordings and devices for neural recordings and stimuli delivery.

Imaging frames were spatially downsampled by a factor of two in the X and Y dimensions, and spatially high-pass filtered with a cutoff spatial frequency of 40 μ m. All frames collected over the course of a single day were then concatenated into a single stack and registered to each other to correct for motion artifacts using a rigid-body transformation (TurboReg plugin for ImageJ). Single cell Ca^{2+} activity traces and spatial filters were extracted from the registered movie using CNMF-E⁶. Extracted traces and ROIs were manually screened to remove neuropil or other non-neuronal signals. The cleaned set of cells were then registered across three consecutive days of imaging as described in³. Briefly, all extracted spatial filters from a given day of imaging were added to create a cell map, and intensity-based image registration was used to identify a pair of rigid-body transformations to align the day 1 and 3 maps to the day 2 maps. Overlapping triplets of spatial filters for the three days were identified by finding cells on day 1 and 3 with the smallest Euclidean distance to each day 2 cell. All identified triplets were then manually screened for accuracy. Roughly half of all cells could be registered across all three days of imaging.

Spike inference. Spiking of Sf1⁺ neurons was estimated from extracted GCaMP fluorescence traces using the constrained deconvolution approach of ⁷. Calcium transients were modeled as a first-order autoregressive (AR) process, and with the exception of the AR coefficient γ (GCaMP decay rate), model parameters were fit as in ⁷ to all data from each cell on a given day of imaging. The value of γ was selected to give a GCaMP half-decay time of two seconds, following the expected decay kinetics of GCaMP6s in high firing rate conditions⁸. This is a conservative estimate (other reported half decay times of AAV-GCaMP6s range from 510ms to 1.8sec⁹) to obtain an approximate upper limit on the contribution of slow GCaMP dynamics to the observed persistence of SF1⁺ population activity.

Units of neuronal activation. Stimulus-evoked responses of SF1⁺ neurons are reported in units of baseline standard deviation, σ , defined as the standard deviation of observed fluorescence in a 30-second pre-stimulus baseline.

Fitting decay constants in fiber photometry and population average microendoscope data. The stimulus-evoked response of the SF1⁺ population could be well fit by a difference of exponentials of the form $K(t) = A \cdot (\tau_{\text{decay}} - \tau_{\text{rise}})^{-1} (e^{-t/\tau_{\text{rise}}} - e^{-t/\tau_{\text{decay}}})$, where t is time in seconds, and A , τ_{decay} , and τ_{rise} are fit parameters characterizing the amplitude and kinetics of the response. Values of τ_{decay} and τ_{rise} were fit for each trial to minimize the mean squared error between $K(t)$ and the SF1⁺ population response over a 30 second window following the start of stimulus presentation, using the *fminunc* function in Matlab.

Identifying stimulus-responsive cells. Some analyses, such as calculation of time to peak or half-peak strength of stimulus preference, were performed only on cells that showed a significant response to the stimulus or stimuli in question. For these analyses, we defined a pre-stimulus baseline as the $\Delta F/F$ in a ten-second window prior to stimulus presentation, and defined responsive cells as those neurons for which the average $\Delta F/F$ value for any one-second window in the first 30 seconds after stimulus presentation was more than four standard deviations above the mean of baseline activity on that trial. Only cells that passed this criterion on both trials within a day were included for analysis.

Finding peak and decay to half-peak times. The time of the peak population response was defined as the first time (relative to start of stimulus presentation) the population $\Delta F/F$ passed 95% of its maximum observed value on a given trial. The time to decay to half-peak was defined as the last time the population $\Delta F/F$ was above 50% of its maximum value relative to pre-stimulus baseline. This analysis was performed separately for each cell on each day of imaging, using only stimulus-responsive cells (see above); values were averaged across two repeated stimulus presentations.

Strength of cell preference. The strength of cell preference for either of a pair of stimuli (**Fig 2g** and **Fig 3i,k**) was defined as $|s_a - s_b| / (|s_a| + |s_b|)$, where s_a and s_b are the average $\Delta F/F$ of that cell in a 45-second window following stimulus onset, for stimulus pair a vs b (eg rat vs USS). This analysis was performed separately for each cell on each day of imaging, using only cells that responded to either one (or both) of the two stimuli, as defined above; values of s_a and s_b were averaged across two repeated stimulus presentations.

Decoder analysis. Stimulus identity was decoded from the activity of all SF1⁺ neurons that could be reliably tracked across three days of imaging, using a two-class or five-class cross-validated Naïve Bayes decoder (*fitcnb* in Matlab). Bar plots of decoder accuracy (**Fig 3n**) and confusion matrix (**ED 7a**) were generated using held-out test data for a five-class Naïve Bayes decoder trained on the time-averaged responses of imaged neurons in a window from 30 seconds before to approximately 60 seconds after stimulus presentation. Time-evolving plots of decoder accuracy (**Fig 3o**, **ED 8b-c**) were constructed by training a separate cross-validated decoder on the time-averaged activity of imaged neurons in a one-second window, for each one-second window from 10 seconds before to 30 seconds after stimulus presentation. Decoder performance is reported as the average prediction accuracy on held-out test data; chance accuracy is 1/2 for the two-class decoder and 1/5 for the five-class decoder.

Stimulus-evoked autocorrelation. We constructed the standard correlation matrix C of VMHdm^{SF1} cell activity, defined as the pairwise correlation coefficient between all columns of a neurons x time matrix of

imaged activity. Values in C were averaged across each trial for a given stimulus over three days of imaging, and then averaged across $n=5$ imaged mice, for all imaging frames from zero to 45 seconds relative to the onset of stimulus presentation (imaging framerate was 11Hz). The mean correlation for lag Δt was computed by averaging $C(x, x+\Delta t)$ for all x between 0 and $45-\Delta t$ seconds.

The same calculation was used for simulated data, with correlations computed every 10 simulation timesteps (10ms). To make values comparable to the experimental data, model cell spikes were convolved with a pair of exponential filters with time constants of 0.5 seconds and 1.5 seconds, simulating the kinetics of the GCaMP6s response.

Rat/USS Pearson's correlation. The Pearson's correlation between rat and USS responses was computed for each mouse using the trial-averaged response of all neurons on the first day of imaging. Pearson's correlation was computed between the vectors of population activity from 10 seconds before to 30 seconds after stimulus onset, sampled at 11Hz (acquisition frequency).

For simulated data, the "rat" and "USS" inputs were assumed to be excitatory inputs to a randomly selected fraction of neurons in the model (temporal structure of stimulus and percent of neurons receiving input specified below for each model). Pearson's correlation between these two stimuli was computed across all model cells that fired 10 or more spikes across the two stimuli. GCaMP6s kinetics were simulated as for the stimulus-evoked autocorrelation analysis.

Neuropeptide model. For this model we assumed that VMHdm^{SF1} neuron dynamics were determined entirely by long-lasting peptidergic input, and that there were no recurrent connections between neurons within VMHdm. Given a model population of $N = 1000$ neurons, we assumed that a random 10% of neurons received peptidergic input for any given stimulus. For cells receiving stimulus-evoked input, we modeled the firing rate $r_i(t)$ of neuron i as $r_i(t) = g \cdot p_i(t)$, where $g \sim U(0,60)$ sets the strength of excitatory peptidergic input onto neuron i and $p_i(t)$ is a stimulus-evoked peptide-mediated excitatory current. Dynamics of $p_i(t)$ evolve as $\tau_p \frac{dp_i}{dt} = -p_i(t) + \delta(t)$, where $\delta(t)$ is the delta function and $\tau_p = 25$ sec, the decay time constant that sets the duration of peptidergic excitation, was set to match the observed decay time constant of the population average VMHdm^{SF1} response. To simulate spiking, the firing rate $r_i(t)$ was used to set the instantaneous rate constant of a non-homogeneous Poisson process with a simulation timestep of $dt = 1$ ms.

Spiking recurrent neural network model + NMDA. We constructed a model population of $N = 1000$ standard current-based leaky integrate-and-fire neurons, in which each neuron has membrane potential x_i characterized by dynamics $\tau_m \frac{dx_i}{dt} = -x_i(t) + I(t)$, where $\tau_m = 20$ ms is a membrane time constant and I (specified below) is a combination of external and recurrent inputs. To model spiking, we set a threshold θ (typically $\theta=0.1$) such that when the membrane potential $x_i(t) > \theta$, $x_i(t)$ is reset to 0 and instantaneous spiking rate $r_i(t)$ is set to 1. Spiking-evoked input to postsynaptic neurons was modeled as a synaptic current with dynamics $\tau_E \frac{dp_i}{dt} = -p_i(t) + r_i(t)$, where τ_E is the decay time constant of excitatory currents. To simulate the slow excitatory currents produced by NMDA receptors, we set $\tau_E = 200$ ms.

We next added recurrent connectivity between model units. Connectivity between model units is random and sparse, with $p = 10\%$ probability of a synapse forming between any two neurons, and weights of

existing synapses sampled from a uniform distribution: $W_{ij} \sim U(0, 1/\sqrt{N \cdot p})$. We also defined a gain parameter g that scales the strength of all synapses in the network.

To reduce finite-size effects in this model, we modeled recurrent inhibition by a single graded input I_{inh} representing an inhibitory population that receives equal input from, and provides equal input to, all excitatory units; dynamics of I_{inh} thus evolve as $\tau_I \frac{dI_{inh}}{dt} = -I_{inh}(t) + \frac{1}{N} \sum_{n=1}^N r_N(t)$, where $\tau_I = 50$ ms is the decay time constant of inhibitory currents.

Each modeled “stimulus” input to the network was modeled with the same dynamics, with a high initial firing rate that decayed to a much lower sustained firing rate, and dropped to zero ten seconds after stimulus onset: specifically, in our model this input took the form $s(t) = \mathbb{I}(t < 10) \int_{-\infty}^t \mathbb{I}(0 < t' < 2)e^{-\frac{t-t'}{2}} dt'$ where $\mathbb{I}$ is the indicator function. Each stimulus drove a random 50% of excitatory units in the network with input strength $w_i \sim g \cdot U(0, 1)$.

Thus, outside of spiking events, the membrane potential of neuron i evolves as $\tau_m \frac{dx_i}{dt} = -x_i(t) + g \left(\sum_{j=1}^N J_{ij} p_j(t) - g_{inh} I_{inh}(t) \right) + w_i s(t)$. Model dynamics were simulated in discrete time using first-order Euler’s method with a timestep of $dt = 1$ ms; a small Gaussian noise term $\eta_i \sim \mathcal{N}(0, 1)/5$ was added at each timestep. We explored model dynamics over a range of values of g and g_{inh} , by selecting a value of g and performing a grid search over g_{inh} until the desired degree of persistence was achieved. Figures in the paper correspond to $g = 1$, $g_{inh} = 3.8$.

Spiking recurrent neural network model + peptidergic excitation (pRNN). In experimenting with the RNN+NMNDA model described above, we found that we could achieve diverse temporal dynamics of spiking neurons if the time constant of excitation (τ_E) was further increased, causing excitation to be much slower than inhibition. This allows model neurons to act as leaky integrators of their excitatory inputs, and start spiking when the population average activity (reflected by inhibitory input) drops below the integrated excitation. Because VMHdm^{SF1} neurons are known to express both glutamate and a variety of neuropeptides + neuropeptide receptors, we further modified the model by replacing the excitatory current with a mix of fast (glutamatergic) and slow (peptidergic) excitatory neurotransmission, although similar results are obtained in a model with just the slow component of excitatory neurotransmission.

We modeled fast excitatory currents as in the prior model, with dynamics $\tau_{E_{fast}} \frac{dp_{fast_i}}{dt} = -p_{fast_i}(t) + r_i(t)$, however we set $\tau_{E_{fast}} = 50$ ms to better match the decay time constant of glutamatergic excitation. To model slow peptidergic excitation, we assumed that when a neuron spiked, peptide release was contingent on the recent firing rate history of that neuron, with peptide release only occurring if the average number of spikes in the last second exceeded a threshold T (typically $T = 20$, although performance was not strongly dependent on this parameter.) That is, the spiking of neuron i evoked peptide release if $\int_{t-1}^t r_i(\tau) d\tau > T$. Dynamics of peptide-mediated excitation were otherwise modeled as before, thus giving $\tau_{E_{slow}} \frac{dp_{slow_i}}{dt} = -p_{slow_i}(t) + r_i(t) \cdot \mathbb{I} \left(\int_{t-1}^t r_i(\tau) d\tau > T \right)$, where $\mathbb{I}$ is the indicator function. We used $\tau_{E_{slow}} = 6$ sec for all versions of the pRNN except for the third (black traces in Fig 4), for which $\tau_{E_{slow}} = 20$ sec ($\tau_{E_{slow}}$ is abbreviated as τ_S in Fig 4).

For simplicity we assumed the synaptic weight matrix J was the same for both fast and slow components of excitation. Membrane potential dynamics in this model are therefore given by $\tau_m \frac{dx_i}{dt} = -x_i + g \sum_{j=1}^N J_{ij} p_{\text{fast}j}(t) + g \sum_{j=1}^N J_{ij} p_{\text{slow}j}(t) - g_{\text{inh}} I_{\text{inh}}(t) + w_i s(t)$. We present three versions of this model in **Fig 4**: in the “low gain” model, $g = 1$, $g_{\text{inh}} = 8.8$, $\tau_{E_{\text{slow}}} = 6$ sec; in the “high gain” model, $g = 6$, $g_{\text{inh}} = 7.8$, $\tau_{E_{\text{slow}}} = 6$ sec; in the “high τ_s ” model, $g = 2.5$, $g_{\text{inh}} = 4.25$, $\tau_{E_{\text{slow}}} = 20$ sec. Simulation was performed as for the NMDA-RNN model, and as above parameters were fit by fixing the value of g (and $\tau_{E_{\text{slow}}}$) and performing a grid search over values of g_{inh} to achieve the desired degree of persistence.

PRNN + local connectivity. The locally connected version of the pRNN model was created by adding a “distance dependence” on the probability of a pair of neurons forming a synaptic connection. Model neurons were numbered between 1 and N , and for neurons i and j the probability of forming a synapse was defined as $p_{ij} = p_0 e^{-(i-j)^2/\sigma}$, where $p_0 = 0.1$ is the baseline degree of connectivity in the network, and σ sets the rate at which connectivity falls off with distance (here distance is defined as $|i - j|$). We found that broad connectivity was necessary to match the stimulus representation overlap seen in the data; plots in **Fig 4** and the illustration of distance-dependent connectivity in **ED Fig 9a-b** were constructed using $\sigma = 0.7N$.

As in the pRNN, each stimulus in the local connectivity model provided input to 50% of model neurons. To match the observed Pearson’s correlation of the data, we found that it was necessary for stimulus inputs to reflect the structure of the model network, by targeting separate but still overlapping portions of the band of model neurons. Specifically, we found that the data was well fit when the middle 50% of model neurons in the band could receive input from both rat and USS stimuli, while the outermost 25% could only receive rat or USS input (see **ED Fig 9a**).

Data similarity score, time-evolving dynamics. We constructed a data similarity score to quantify the degree of similarity between the plotted curves in **Fig 4e**, thus capturing how much the time-evolving dynamics of model neurons looked like that of the data. For each model and each mouse, we computed the Mean Correlation as defined above, which we will call $MC_{\text{model}}(t)$ for a given model and $MC_{\text{mouse } i}(t)$ for a given mouse. MC is a function of time-- thus to quantify the mean similarity between the data and a given model over time, we considered the value of $MC_{\text{model}}(t)$ and $MC_{\text{mouse } i}(t)$ for all imaging frames (acquired at 11Hz) from 0 to 45 seconds relative to stimulus onset, which we reference using a frame index $t = 1 \dots T$ (so $t=1$ corresponds to a time of 0sec and $t=T$ corresponds to a time of 45sec). Given these definitions, we define the data similarity score of the model dynamics as:

$$\text{Similarity Score}_{\text{dynamics}} = \frac{1}{N} \sum_{i=1}^N 1 - \left(\sum_{t=1}^T |MC_{\text{mouse } i}(t) - MC_{\text{model}}(t)| \right) / \left(\sum_{t=1}^T |MC_{\text{mouse } i}(t)| \right)$$

This can be simply interpreted as akin to the area between the data/model curves for each plot in **Fig 4e**. Note that the MC for the data here was computed from the USS-evoked neural activity, however MC for other stimuli gave similar results, as we found little difference between the MC for different stimuli.

Data similarity score, stimulus specificity. This data similarity score quantifies the degree of similarity between the plotted curves in **Fig 4h**, ie how much the Pearson’s correlation between rat- and USS-evoked activity in each model looked like that observed in the data. We computed the Pearson’s correlation (as defined above) for each model and each mouse, which we call $PC_{\text{model}}(t)$ for a given model and

$PC_{\text{mouse } i}(t)$ for a given mouse. We define frames $t=1\dots T$ as all imaging frames from times 0 to 45 seconds relative to stimulus onset (same as for the similarity score of dynamics). We then define the data similarity score of model stimulus-specific activity as:

$$\text{Similarity Score}_{\text{specificity}} = \frac{1}{N} \sum_{i=1}^N 1 - \frac{1}{T} \left(\sum_{t=1}^T |PC_{\text{mouse } i}(t) - PC_{\text{model}}(t)| \right)$$

Like the similarity score of the dynamics, this can be interpreted as the area between the data/model curves for each plot in **Fig 4h**.

- 1 Dhillon, H. *et al.* Leptin directly activates SF1 neurons in the VMH, and this action by leptin is required for normal body-weight homeostasis. *Neuron* **49**, 191-203, doi:10.1016/j.neuron.2005.12.021 (2006).
- 2 Kunwar, P. S. *et al.* Ventromedial hypothalamic neurons control a defensive emotion state. *Elife* **4**, doi:10.7554/eLife.06633 (2015).
- 3 Remedios, R. *et al.* Social behaviour shapes hypothalamic neural ensemble representations of conspecific sex. *Nature* **550**, 388-392, doi:10.1038/nature23885 (2017).
- 4 Mongeau, R., Miller, G. A., Chiang, E. & Anderson, D. J. Neural correlates of competing fear behaviors evoked by an innately aversive stimulus. *J Neurosci* **23**, 3855-3868 (2003).
- 5 Lerner, T. N. *et al.* Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* **162**, 635-647, doi:10.1016/j.cell.2015.07.014 (2015).
- 6 Zhou, P. *et al.* Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data. *Elife* **7**, doi:10.7554/eLife.28728 (2018).
- 7 Pnevmatikakis, E. A. *et al.* Simultaneous Denoising, Deconvolution, and Demixing of Calcium Imaging Data. *Neuron* **89**, 285-299, doi:10.1016/j.neuron.2015.11.037 (2016).
- 8 Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**, 295-300, doi:10.1038/nature12354 (2013).
- 9 Dana, H. *et al.* Thy1-GCaMP6 transgenic mice for neuronal population imaging in vivo. *PLoS One* **9**, e108697, doi:10.1371/journal.pone.0108697 (2014).

Table 1: Statistical significance testing

All t-tests are two-sided unless otherwise stated. All tests from distinct samples unless otherwise stated.

Figure	Panel	Identifier	Sample size	p-value	Statistical test	Notes
1	e	rat vs toy	4 mice	0.0078	paired t-test	
	g	rat vs mouse	8 mice	0.0001	paired t-test	
		rat vs toy	8 mice	0.0005	paired t-test	
		mouse vs toy	8 mice	0.0015	paired t-test	
	h	rat vs mouse - headfixed	4 mice home cage, 8 mice head fixed	0.0052	paired t-test	
	i	rat vs mouse - headfixed	4 mice home cage, 8 mice head fixed	0.0110	paired t-test	
	k	rat control vs rat iC++	7 mice control, 7 mice iC++	0.0289	t-test	
	m	during+after vs control	12 mice control, 6 mice during+after	0.0078	rep. meas. ANOVA	covariates tested: group vs time
		after only vs control	12 mice control, 6 mice after-only	0.0321	rep. meas. ANOVA	covariates tested: group vs time
	n	control before vs after rat	12 mice	0.0006	paired t-test	
		control before rat vs after PS off	12 mice	0.0074	paired t-test	
		after rat, control vs during+after	12 mice control, 6 mice during+after	0.0023	t-test	
		after PS off, control vs during+after	12 mice control, 6 mice during+after	0.0223	t-test	
		after PS off, control vs after only	12 mice control, 6 mice after-only	0.0062	t-test	
	o	control vs during+after	12 mice control, 6 mice during+after	0.0003	rep. meas. ANOVA	covariates tested: group vs time
		control vs after only	12 mice control, 6 mice after-only	0.0044	rep. meas. ANOVA	covariates tested: group vs time
2	d	*	5 mice	0.0117	paired t-test

Extended Data 1. Comparison of SF1+ neurons' response to rat in mouse home cage and head-fixed set-up.

a, Peak $\Delta F/F$ activity in response to rat in home cage and a head-fixed set-up. (home cage group $n = 4$; head-fixed group $n = 8$). **b**, Response time measured at 10% of peak. **c**, Rise time constant measured as time elapsed to reach $1/e$ of peak amplitude. **d**, Decay time constant measured as time elapsed from peak to $1/e$ of peak.

Extended Data 2. SF1+ neurons' response to rat urine.

a, Urine presentation to head-fixed fiber mouse. **b**, Averaged $\Delta F/F$ activity traces of SF1+ neurons in response to rat urine and water. $n = 6$. **c**, Peak $\Delta F/F$ activity triggered by rat urine and water. Paired t-test, $n = 6$. **d**, Decay time constant for rat and rat urine response, $n = 6$.

Extended Data 3. No change in mouse behavior due to potential lingering odor from rat.

a, Schematic plot showing the experiment protocol: top, a live rat or toy rat (control) was brought to the open field arena in a wire mesh cage for 15 seconds; bottom, mouse was introduced to arena afterwards immediately. **b**, Fraction of time spent in center zone (as shown by red dashed line in a) for rat group and control group, $n = 6$ for each group. **c**, Distance from mouse body center to arena center. **d**, Schematic plot showing the optogenetic activation experiment protocol: mice expressing ChR2 was brought to the open field arena. After a 5 minutes habituation period, a 10 seconds light or mock stimulation was delivered to the mice. **e**, Fraction of time spent in the edge zone, $n = 4$ mice for each group.

Extended Data 4. Fiber photometry and SF1+ neuron silencing in open field rat exposure assay.

a, Distance from mouse body center to arena center during three different time periods: before rat, after rat and after photo stimulation offset, corresponding to -1~0, 0~1 and 3~4 minute in Fig.1m. **b**, Mean velocity was not altered by photo stimulation for iC++ and control mice. Velocity was measured in mouse home cage and averaged during a three-minute period for light off and light on sessions. Paired t test, $n = 12$. **c**, ΔF/F activity traces (mean ± SEM) of SF1+ neurons in response to rat in open field arena. Shaded gray bar denotes the presentation of rat. $n = 9$. **d**, Peak ΔF/F activity triggered by rat in open field arena ($n = 9$) and head-fixed set up ($n = 8$) (mean ± SEM). **e**, Decay constants of ΔF/F activity in open field arena and head-fixed set up. **f**, Traces (mean ± SEM) for ΔF/F activity and the distance from mouse body center to arena center, aligned to rat removal. $n = 9$. Distance to center is plotted as 30 seconds moving average. **g**, Decay time measured as the time elapsed to reach 50% of the peak for linearly fitted data. $n = 9$, paired t test. **h**, Scatter plot of ΔF/F activity and the distance from mouse to arena center with a linear regression fitting for 2 example mice. Top, $r = 0.958$, $p < 0.0001$; bottom, $r = 0.808$, $p < 0.0001$. **i**, Pearson's correlation coefficient between ΔF/F activity and the distance to center. $n = 9$.

Extended Data 5. Inferred spiking of SF1+ neurons.

a, Calcium traces (black) and spiking events inferred using constrained deconvolution1 (blue) for microendoscopic imaging of an example SF1+ neuron in response to five rat presentation trials in a head-fixed mouse. **b**, Response to one presentation of the rat stimulus in microendoscopic imaging experiments, averaged across cells ($n=5$ mice, mean $\pm$ SEM). **c**, Spiking events inferred from all individual SF1+ cells from the data in (b), averaged across cells and smoothed with a 1-second sliding window ($n=5$ mice, mean $\pm$ SEM). **d-e**, As in b-c, but for the USS stimulus.

Extended Data 6. SF1+ population shows minimal response to the looming disk stimulus.

Peak $\Delta F/F$ in response to rat, toy, or looming disk stimuli presented for 10 seconds in the animal's home cage (points are individual mice).

Extended Data 7. Additional Pearson's correlations between stimulus pairs.

Pearson's correlation between SF1+ population activity evoked by all possible pairs of stimuli, as a function of time.

Extended Data 8. Additional decoder analysis of SF1+ population activity.

a, Confusion matrix of the five-way Naïve Bayes decoder shown in Fig 3n, showing predicted stimulus identity for each stimulus class. Matrix is normalized so rows sum to 100%. **b**, Accuracy of the time-dependent five-way Naïve Bayes decoder shown in Fig 3o, as a function of time, for each tested stimulus. **c**, Accuracy of time-dependent binary Naïve Bayes decoders trained on all possible pairs of stimuli. The pair of stimuli being decoded for each plot is specified by the labels on the left and top. All plots show mean $\pm$ SEM across 5 imaged mice.

Extended Data 9. Locally connected model networks.

a, Probability of synapse formation between neuron pairs decreases moderately as a function of "distance" (neuron number) in the locally connected sRNN model. Segments of the model targeted by rat and USS model input are also shown (blue/purple lines.) **b**, Example synaptic weight matrix generated from probability matrix shown in **a**; for visibility every 10th model neuron is shown. **c**, Example of a more highly structured model network, in which largely separate populations of neurons respond to the rat vs USS model inputs. **d**, Pearson's correlation and stimulus-evoked autocorrelation for a network model such as that in (**c**), in which network structure results in no overlap between rat and USS representations.