Dalbavancin and wound healing: new evidences/insights in a mouse model of skin infection.

Oriana Simonetti^{1*}, Guendalina Lucarini^{2*}, Gianluca Morroni³, Fiorenza Orlando⁴, Raffaella Lazzarini², Antonio Zizzi⁵, Lucia Brescini², Mauro Provinciali⁴, Andrea Giacometti³, Annamaria Offidani¹, Oscar Cirioni³.

* both the authors equally contributed to this work.

¹ Clinic of Dermatology, Department of Clinical and Molecular Sciences, Polytechnic University of Marche, Ancona, Italy; ² Histology, Dept. of Clinical and Molecular Sciences, Polytechnic University of Marche, Ancona, Italy; ³ Clinic of Infectious Diseases, Dept. of Biomedical Sciences and Public Health, Polytechnic University of Marche, Ancona, Italy; ⁴ Experimental Animal Models for Aging Unit, Scientific Technological Area, I.N.R.C.A. I.R.R.C.S., Ancona, Italy; ⁵ Pathologic Anatomy and Histopathology, Department of Biomedical Sciences and Public Health, Polytechnic University of Marche, Ancona, Italy.

Correspondence to: Oriana Simonetti, MD, PhD

Clinic of Dermatology, Department of Clinical and Molecular Sciences, Polytechnic University of Marche, c/o Ospedali Riuniti. Via Conca 71, 60126 Ancona AN

Telephone number: +39 071 5963494 Fax number: +39 071 5963446

E.mail: o.simonetti@univpm.it

Running title: **Dalbavancin and wound healing**

ABSTRACT

Dalbavancin is an effective antibiotic widely used to treat skin infection. Our aim was to determinate the effects of dalbavancin administration on wound healing compared to vancomycin, and to elucidate if EGFR, MMP-1, MMP-9 and VEGF could be involved in its therapeutic mechanism.

A mouse model of MRSA skin infection was established. Mice were treated daily with vancomycin (10mg/kg) and weekly with dalbavancin, at day 1 (20 mg/kg) and day 8 (10 mg/kg). After 14 days wounds were excised and bacterial counts were performed. Wound healing was assessed by histological and immunohistochemical staining, followed by protein extraction and immunoblotting. Our microbiological results confirmed that both dalbavancin and vancomycin are effective in reducing the bacterial load in wounds. Dalbavancin had a strong effect compared with infected untreated animals and vancomycin treated group. The wounds treated with dalbavancin showed robust epidermal coverage with a reconstitution of the regular and keratinized epidermal lining and a well-organized granulation tissue with numerous blood vessels, although slightly less than in the uninfected group, while in vancomycin treated group the epithelium appeared in general still hypertrophic, the granulation tissue appears even less organized.

We observed elevated EGFR and VEGF expression in both treated groups, although it was higher in dalbavancin treated mice. MMP-1 and MMP-9 were decreased in uninfected and in both treated tissue when compared with untreated infected wounds.

This study showed faster healing with dalbavancin treatment that might be associated with a higher EGFR and VEGF levels.

INTRODUCTION

It has recently proposed a new classification of skin and soft tissue infections namely acute bacterial skin and skin structure infections (ABSSSIs) where Superficial incisional Surgical Site Infections or SSIs (involving only skin or subcutaneous tissue of the incision) are recognized a subgroup ABSSSIs (1). In particular SSIs constitute a significant burden for the healthcare system, because of the associated morbidity, mortality, length of hospital stay, a higher probability of intensive care unit (ICU) admission with overall direct and indirect costs (2).

The common pathogens that are responsible of SSI are *Staphylococcus aureus*, followed by Coagulase-negative staphylococci and *Enterococcus* species. In particular methicillin-resistant *Staphylococcus aureus* (MRSA) is considered a leading cause of human nosocomial infections (3) as well as community-associated MRSA (CA-MRSA) is associated to deadly infection (4) and livestock-associated MRSA (LA-MRSA) represent major reservoir of infection (5).

In the last decades the epidemiological aspects of nosocomial infection as well as the growing prevalence of CA-MRSA have been increasingly studied to counteract this worldwide public health problem. Particular attention has been paid to the problem of LA-MRSA. Several studies demonstrated that this superbug can be transmitted to human by either healthy and disease animals, usually cow, horses and companion animals (6-7).

Taking into account these considerations, the respect of the principles of a good antibiotic stewardship is mandatory for fighting antimicrobial resistance and aim, next to optimization of clinical outcomes, to the prevention of the spread of antibiotic-resistant strains (8). Dalbavancin is a novel antimicrobial agent, belonging to the lipoglycopeptides family (a subclass of glycopeptides), active against *S. aureus* (including MRSA), *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus anginosus* group, *Enterococcus faecalis* and *Enterococcus faecium* (including ampicillin-resistant isolates) and approved for the treatment of ABSSSIs caused by susceptible Gram-positive organisms (9). Moreover, it has a bactericidal effect and an extremely extended half-

70 life that make it a promising alternative to conventional antibacterials such as vancomycin, that is
 71 considered the standard of care for treatment of gram-positive skin and soft-tissue infections (10).
 72 Wound healing is a highly complex physiological process involving ordered events where
 73 growth factors are critically important for coordinating cell-cell and cell-matrix interactions during
 74 normal injury repair. In particular Epidermal keratinocytes (KCs) respond to injury by becoming
 75 hyperproliferative, migratory and pro-inflammatory in processes regulated by several factors such
 76 as epidermal growth factor receptor (EGFR), a 170-kDa transmembrane protein belonging to ErbB
 77 family receptors, which includes three other members (ErbB2/HER-2, ErbB3/HER-3, ErbB4/HER-
 78 4). EGFR signaling plays key roles in skin regeneration and wound healing by stimulating KCs
 79 proliferation (11). Among the many proteins, that are essential for the restoration of tissue integrity,
 80 the metalloproteinase family is one of the most important. Matrix metalloproteinases (MMPs)
 81 comprise a large family of extracellular proteases, which are up-regulated during wound healing by
 82 epidermal, dermal, fibroblast, or blood cells in mammals (12). MMPs contribute significantly in
 83 reorganization of the extracellular matrix for numerous physiological functions including
 84 morphogenesis, tissue remodeling and wound healing (13). In particular MMP-1 and MMP-9
 85 reflect the correlation between high protease activity and poor healing outcomes (12).
 86 An essential feature of normal wound repair is the formation of granulation tissue, i.e. fibrovascular
 87 tissue containing fibroblasts, collagen and blood vessels. Vascular endothelial growth factor (VEGF)
 88 has shown to be important in wound healing by promoting the early events in angiogenesis,
 89 particularly endothelial cell migration and proliferation (14). VEGF is produced by endothelial cells,
 90 keratinocytes, fibroblast, smooth muscle cells, platelets, neutrophils and macrophages¹⁵, and,
 91 although in normal skin is negligible, in injured skin is markedly up-regulated (14-16).
 92 This study aimed to explore the therapeutic effects of dalbavancin on wound healing in an infected
 93 MRSA mouse model, using histological, immunohistochemical and western blot analysis, and to
 94 elucidate if EGFR, MMP-1, MMP-9 and VEGF could be involved in its therapeutic mechanisms.

RESULTS

Bacterial Counts

Overall, data analysis showed that inhibition of bacterial growth was achieved in all treated groups. In details, mean bacterial numbers in untreated group ($7,51 \times 10^8 \pm 5,05 \times 10^7$ CFU/ml) were significantly higher than those recovered from all treatment groups (Fig. 1). Treatment with vancomycin reduced the mean bacterial numbers to $8,04 \times 10^6 \pm 7,96 \times 10^6$ CFU/ml, while dalbavancin had a strong effect compared with control infected untreated animals and vancomycin treated group ($8,71 \times 10^5 \pm 9,02 \times 10^5$ CFU/ml).

Histological evaluation

To investigate the effects of the different treatments on wound healing, we analyzed the area of the wound, in particular reepithelialization, granulation tissue, collagen deposition and inflammatory cells according to wound healing scores²⁰ summarized in Table1. The statistical analysis was reported in Table 2 and the morphological features was shown in Figure 2.

The extent of newly formed epithelium and collagen deposition, the increase in the thickness of granulation tissue and the decrease in the number of inflammatory cells were considered as progress in the healing process.

Wound healing was still incomplete in the untreated groups, as observed by the partial regeneration of the epidermis, poor organization of dermis layers, immature granulation tissue (many cells and some fibers), and abundant inflammatory infiltrate. These mice showed the worst degree of healing ($p < 0.05$).

From a microscopic examination, the neo-epithelium developed significantly different patterns of regeneration among the treated groups: interestingly, the re-epithelialization of the wounds treated with dalbavancin showed robust epidermal coverage with a reconstitution of the regular and keratinized epidermal lining, comparable to the re-epithelialization of the uninfected mice, except in a small central area where the epithelium appears to be hypertrophic. The wounds treated with

vancomycin showed a good re-epithelialization, although the epithelium appeared in general still hypertrophic when compared to dalbavancin and uninfected groups ($p < 0.05$).

The wounds treated with dalbavancin showed a well-organized granulation tissue with numerous blood vessels, although slightly less than in the uninfected group. In the wounds treated with vancomycin the granulation tissue appears even less organized.

The treated wounds presented a distinct pattern of collagen distribution and remodeling similar to that of uninfected skin; in addition, mild or moderate inflammation was observed in the underlying connective tissue.

Immunohistochemical evaluation

To investigate the influence of the dalbavancin and vancomycin treatment on the expression of VEGF, EGFR, MMP-9 and MMP-1 and in wound healing, an immunohistochemical evaluation was performed. The immunohistochemical staining is shown in Figures.3-4. and the data are summarized in Table.3.

VEGF expression was greater in the uninfected and treated infected groups respect to the untreated ($p < 0.05$), the wounds treated with dalbavancin showed the highest VEGF values.

EGFR expression was significantly higher in the uninfected and infected group treated with dalbavancin respect to the other two groups ($p < 0.05$), the wounds treated with dalbavancin showed the highest expression.

MMP-9 and MMP-1 expression was high in the untreated infected wounds, while a significant reduced expression was found in the other groups ($p < 0.05$), in particular the treated wounds showed the lowest values.

Analysis by Western blot of the Expression of EGFR, VEGF, MMP-1 and MMP-9

The expression of EGFR, VEGF, MMP-1 and MMP-9 was tested by Western blot followed by densitometric analysis (Fig. 5). All proteins were detectable in uninfected, untreated infected, infected treated with dalbavancin and infected treated with vancomycin tissue samples. The

146 expression of EGFR and VEGF was significantly higher in infected treated with dalbavancin tissue
 147 than all the other tissue samples ($p < 0.05$, Fig. 5). VEGF protein expression level was significantly
 148 lower in the untreated infected compared with uninfected and treated infected ($p < 0.05$ Fig. 5).
 149 MMP-1 and MMP-9 protein expression levels were significantly higher in the untreated infected
 150 compared with the other groups; in particular, the treatments with dalbavancin and Vamcomycin
 151 resulted in a significant decrease of protein expression levels ($p < 0.05$ Fig. 5).

152 **DISCUSSION**

153 MRSA has demonstrated to be prevalent in ABSSSIs and in SSIs. Indeed, the rates of MRSA
154 infection are increasing dramatically and MRSA has emerged as the most common cause of
155 surgical infections (18).

156 Vancomycin is the first-choice antibiotic for the treatment of MRSA infection (19), but several
157 papers described infections sustained by vancomycin-intermediate *S. aureus* (VISA) and
158 heterogeneous VISA (hVISA) strains. Different studies reported that the host immune system has
159 difficulty in eradicating VISA/hVISA, leading to chronic, recurrent, and persistent infections (20) .

160 Dalbavancin is an optimal alternative to conventional antibacterials for the treatment of ABSSSIs.
161 Like other lipoglycopeptides, dalbavancin binds the C-terminal acyl-d-Ala-d-Ala subunit of
162 peptidoglycan precursors. Furthermore, the positively charged C-terminal dimethylaminopropyl
163 group may interact with the negative phospholipid head groups of the bacterial membrane (21). The
164 fatty acyl group allows nonspecific protein binding that likely contributes to dalbavancin's long
165 plasma half-life and its ability to bind its target sites (21). This particular structure confers
166 exceptional pharmacokinetic properties to dalbavancin and allow the administration in a two-dose
167 regimen (9).

168 Dalbavancin demonstrated superior *in vitro* activity versus comparators, including vancomycin,
169 with dramatically lower MICs without evidence of resistance through direct selection or serial
170 passage (22,9).

171 Our microbiological results confirmed that both dalbanvancin and vancomycin are effective in
172 reducing the bacterial load in wounds. These results are not surprising, considering that efficacy of
173 these antibiotics was proven by the large clinical use and the numerous reports (23-25). Moreover,
174 due to the long biological half-life, daily treatment didn't result in better reduction of bacteria, as
175 also previously demonstrated by other authors (26).

176 But, till now there is no published data implicating the effect of dalbavancin in wound healing
177 comparing to vancomycin.

178 After external skin injury, such as surgical intervention, a rapid repair is needed to avoid
 179 dehydration and blood loss. Thus, a wound healing process involves a coordinated cascade of
 180 events comprising inflammation, formation of granulation tissue and dermis maturation (27-28).
 181 It has been shown that the bacteria are specifically able to alter stromal components and
 182 inflammatory cells and their responses. Infact, Human stromal cells demonstrated, to be susceptible
 183 to soluble factors of *S. aureus*, biofilms, decreasing cell differentiation, viability, migration and
 184 angiogenesis (29). Therefore *S. aureus* showed to have a dramatic effect on keratinocyte
 185 proliferation rates as well as an impedance to the migration of the keratinocytes, evidenced by
 186 delayed wound closure (30). Thus, we studied the effect of dalbanvancin vs vancomycin on the
 187 infected MRSA wound healing process through morphological studies besides an
 188 immunoistochemical evidences; moreover, to confirm our results we performed a western blot
 189 analysis.

190 As expected, in the infected but untreated groups, wound healing was still incomplete, with partial
 191 regeneration of the epidermis, poor organization of dermis layers, immature granulation tissue, and
 192 abundant inflammatory infiltrate. Both treatments showed to be efficacy on wound healing process.
 193 In particular, in dalbavancin treated group we observed a robust epidermal coverage with a
 194 reconstitution of the regular and keratinized epidermal lining, while epidermis in vancomycin
 195 treated group was still hypertrophic, revealing that epithelial healing problem is still occurring.
 196 Besides, in the present study we discovered elevated EGFR expression in the treated groups,
 197 although it was higher in dalbavancin treated mice. These immunohistochemical results correlated
 198 with the histological features and suggested that dalbavancin and vancomycin were able to induce
 199 EGFR expression and consequently to have an effect on wound healing. In particular, dalbavancin
 200 seemed to be more effective than vancomycin in inducing reepithelialization.

201 Epidermal keratinocytes are a rich source of EGFR ligands, and EGFR signaling has a major effect
 202 on the proliferation and differentiation of keratinocytes. Therefore, EGFR plays a key role in skin
 203 development and homeostasis (31) and its expression in the epithelium is a good indicator of the

204 healing potential as observed in previous studies (32,14). In our study MMP-1 and MMP-9 were
 205 decreased in uninfected and in both treated tissues, when compared with untreated infected wounds.
 206 It has demonstrated that these metalloproteinases play a pivotal role, with their inhibitors, in
 207 regulating extracellular matrix (ECM) degradation and deposition that is essential for wound
 208 reepithelialization (33).

209 While, in normal tissue, MMPs are expressed at basal levels, when tissue remodeling is required (as
 210 in wound healing), they can be rapidly expressed and activated. Therefore, MMPs are upregulated
 211 during wound healing in the very early stages in epidermal cells, dermal cells, fibroblasts and blood
 212 cells (12).

213 In fact, it has been shown that MMP-1 expression peaks at day 1 after wounding in migrating basal
 214 keratinocytes at the wound edge followed by a gradual decrease until re-epithelialization is
 215 complete. Downregulation of MMP-1 seems to be important for normal tissue remodeling as high
 216 levels of MMP-1 in chronic nonhealing wounds are described (33). Similarly migrating
 217 keratinocytes at the leading edges secrete MMP-9, that modulate keratinocyte motility by
 218 degradation of proteins involved in cell–cell and cell–matrix adhesion, allowing re-epithelialization
 219 to occur (33). In particular, MMP-9 knockout (KO) mice, and MMP-9-deficient mice showed
 220 delayed wound closure with inhibition of cell proliferation (34-35), whereas the persistence of high
 221 level of MMP-9 has been associated with poor healing in several studies (36-38). These findings
 222 highlighted the role of MMP-9 in wound healing. In addition, when the balance of MMPs activity
 223 and inhibition is disturbed the wound progress to a state of increased ECM degradation, disrupting
 224 the equilibrium of synthesis and remodeling of ECM components, determinant for successful skin
 225 repaired (39). It is well known that *MMPs* play also a significant role in stimulation of angiogenesis
 226 in the proximity of wounds to accelerate recovery (33). In particular, the role of MMP-9 in
 227 angiogenesis include VEGF release (40-41). Actually, VEGF and MMP seem to regulate each other,
 228 contributing to wound repair (42). Previous studies have evidenced that keratinocytes express
 229 elevated VEGF as early as 1 day after injury and eventually in those cells that migrate to cover the

defect. Epidermal labeling for VEGF mRNA reaches a peak after 2–3 days, and levels remain elevated during the granulation tissue formation and until epidermal coverage is complete (43). In this period, VEGF is upregulated to promote the angiogenesis (i.e., vascular dilation, permeability, migration, and proliferation), providing a conduit for nutrients and other mediators of the healing response (43). In our study we observed that VEGF expression was highly expressed in the uninfected and both treated infected groups, although the wounds treated with dalbavancin showed the highest VEGF values, underlining the positive effect of dalbavancin on angiogenesis.

In conclusion it can be speculated that both treatments may have positive effect not only on the bacterial load, but also on tissue healing after surgical infected skin wound by downregulation of MMP-1 and MMP-9 and increase of EGFR and VEGF expression. In particular this study showed faster healing with dalbavancin treatment that might be associated with a higher EGFR and VEGF levels. However, further studies are needed to clarify the additional effects of Dalbavacin on wound healing.

MATERIALS AND METHODS

Ethics

In vivo experiments were approved by the Institutional Animal Care Committee of the Ministry of Health and by the Animal Research Ethics Committee of IRCCS-INRCA (Istituto di Ricovero e Cura a Carattere Scientifico – Istituto nazionale di Riposo e Cura per Anziani) 767/2016 Pr 28/07/2016.

Bacterial strains and drugs

S. aureus ATCC 43300, the reference MRSA strain, was used in the mouse infection model. Vancomycin and dalbavancin were diluted in accordance with manufacturers' recommendations

yielding 10 mg/mL stock solutions. Solutions were made fresh on the day of assay or stored at -80°C in the dark for short periods.

Mouse infection model

6-month-old BALB/c mice weighting 28–30 g from the Specific Pathogen Free (SPF) animal facility of INRCA (Italian National Centre on Health and Science on Aging, Ancona, Italy) were used for all the studies. All the procedures involving animals were conducted in conformity with the institutional guidelines in compliance with the national (Legislative Decree n. 26, March 4, 2014; Authorization n.767/2016-PR issued July 28, 2016, by the Italian MoH) and international law and policies (EEC Council Directive 2010/63/EU). Mice were individually caged in visual, auditory and olfactory contact with other mice of the same experimental group (within ventilated cabinets cage systems) as well as in presence of environmental enrichments consisting of nesting materials and wooden toys. Mice were kept in a 12 h light dark cycle with food and drinking water ad libitum and allowed to equilibrate in the phenotyping area allocated in the Scientific and Technological Pole of INRCA for approximately 1 month before starting the experiments. After surgical intervention, all mice were housed individually under constant temperature (22 ± 2 °C) and humidity. The animals were fed with standard pellet food (Harlan Laboratories, Udine, Italy), and fresh daily tap water.

The study included a total of 48 animals divided into 4 groups (each composed of 12 mice): uninfected group (C0, sham control); infected but not treated group (C1); vancomycin group (infected and daily treated with vancomycin; C2); dalbavancin group (infected and weekly treated with dalbavancin ; C3).

The MRSA ATCC43300 were grown in brain hearth infusion and diluted in saline to a final concentration of 5×10^7 CFU/ml, prepared freshly at time of intervention.

Mice were anesthetized by an intramuscular injection of ketamine (50 mg/kg of body weight) and xylazine (8mg/kg of body weight), the hair on their back was shaved and the skin cleansed with 10% povidone-iodine solution (no animals dropped out due to infection or anesthetics). One full

278 thickness wound was established through the panniculus carnosus on the back subcutaneous tissue
 279 of each animal. Small gauze was placed over each wound and then inoculated with 200 µl of
 280 bacterial culture previously diluted; in the control group the gauze was soaked only with sterile
 281 saline solution. The pocket was closed by means of skin clips. This procedure resulted in a local
 282 abscess at 24 h. One wound was created per animal. The animals were returned to individual cages
 283 and thoroughly examined daily. After 24 h, the wound was opened and washed with saline, the
 284 gauze was removed, and treatment started (44). The group C0 did not received any treatment. The
 285 groups C1, C2 were treated with daily intraperitoneal administration of 200 µl of saline and
 286 intraperitoneal vancomycin (10mg/kg) respectively. Group C3 was treated with intraperitoneal
 287 injection of 200 µl of dalbavancin at day 1 (20 mg/kg) and day 8 (10 mg/kg), mimicking human
 288 regimens (45). After 14 days, animals, were euthanized and a 1x2 cm area of skin, including the
 289 wound, was excised aseptically for histological and western blot examination (see below) and for
 290 bacterial count. For the bacterial analysis the samples were weighted and then homogenized in 1 ml
 291 phosphate-buffered saline (PBS) using a stomacher. Quantitation of viable bacteria was performed
 292 by culturing serial dilutions of the bacterial suspension on mannitol-salt agar plates at 37°C for 24-
 293 48 hours. The limit of detection for this method was approximately 10 CFU/ml.

294 **Histological and immunohistochemical staining**

295 To understand the efficacy of the treatments in promoting wound healing, the re-epithelialized
 296 tissue of each group was investigated histologically. Five micrometer thick sections were stained
 297 with hematoxylin and eosin (H&E) according to standard protocol. The stained sections were
 298 observed under the light microscope (Nikon DS-Vi1, Nikon Instruments, EuropeBV, Kingston,
 299 Surrey, England) and scored according to wound repair fivetiered grading system based on
 300 epithelial presence, degree of stratification, degree of differentiation as well as maturational features
 301 of the granulation tissue, dermal tissue and presence of inflammatory cells (46) (Table.1).

302 In addition, identification of VEGF, MMP-1, MMP-9 and EGFR expression was carried out using
 303 immunohistochemistry of histological samples with a heat-mediated antigen retrieval process.
 304 Briefly, 5 µm paraffin sections were deparaffinized and rehydrated by sequential immersion in
 305 xylene, ethanol, and water. Antigen retrieval was carried out by heating the sections in 0.1 M citrate
 306 buffer solution (pH 6.0) at 98 °C for 5 min via microwave. Then they were incubated with
 307 monoclonal anti-mouse VEGF (clone C-1; Santa Cruz Biotechnologies, Santa Cruz, CA), MMP-1
 308 (clone H-300; Santa Cruz Biotechnologies, Santa Cruz, CA), MMP-9 (clone 2C3; Santa Cruz
 309 Biotechnologies, Santa Cruz, CA) and EGFR (clone 1005; Santa Cruz Biotechnologies, Santa Cruz,
 310 CA) at 4 °C overnight and immunostained using the streptavidin-biotin peroxidase technique
 311 (Envision universal peroxidase kit; Dako Cytomation, Milan, Italy). After incubation, the tissue
 312 sections were colored with 3,3-diaminobenzidine (DAB), stained with hematoxylin, and
 313 coverslipped with Eukitt mounting medium (Electron Microscopy Sciences, PA, USA).
 314 Two investigators (G.L., A.Z.) blinded to the mice outcome, performed all counts separately. We
 315 counted the number of marker positive cells by using a light microscope (Nikon Eclipse E 600,
 316 Nikon Instruments, Europe B.V., Amsterdam, Netherlands) at x200 magnification in at least ten
 317 fields/sample and the positive cells were expressed as mean value ± standard deviation (SD).
 318 Images were captured with a Nikon DS-Vi1 digital camera (Nikon Instruments) connected to a
 319 computer. The area of each field (0.07 mm²) was standardized using NIS Elements BR 3.22
 320 imaging software (Nikon Instruments).

321 **Protein extraction and immunoblotting**

322 Total proteins were extracted from wound tissue samples and immediately pulverized in liquid
 323 nitrogen. The pulverized skins were homogenized and lysed in ice-cold Radio immunoprecipitation
 324 assay buffer [150 mM NaCl, 10 mM Tris (pH 7.2), 0.1 % sodium dodecyl sulfate, 1.0 % Triton X-
 325 100, and 5 mM ethylenediaminetetraacetic acid (pH 8.0)] containing protease inhibitor cocktail
 326 (Roche Applied Science, Indianapolis, IN, USA) for 45 minutes. The lysates were centrifuged at 14

000 rpm for 30 minutes at 4°C. The supernatant was collected and the protein concentration was determined. Protein concentration was determined using Bradford reagent (Sigma-Aldrich). Total protein extracts (30 µg) were reduced in dithiothreitol (DDT, 0.5 M) for 10 min at 70 °C and samples run on a 4–12 % gradient precast NuPAGE Bis–Tris polyacrylamide gel for 1 h at 200 V. After electrophoresis, the proteins in the gel were transferred to a nitrocellulose membrane, the blots were blocked with 5% nonfat dry milk in PBS containing 0.1% Tween-20 for 1 hours at room temperature. Membranes were incubating overnight with primary antibodies anti EGFR, VEGF, MMP-1 and MMP-9, (Santa Cruz Biotechnology, CA, USA) and GAPDH ((Santa Cruz Biotechnology) as endogenous control, followed by incubation with a secondary antibody conjugated to horseradish peroxidase (Santa Cruz Biotechnology). The detection of the proteins on the membrane was performed using the Clarity Western ECL Substrate kit (*Bio-rad*). The signals were captured using the Alliance Mini (*UVITEC Cambridge*) system. The UVITEC software carried out the quantization of the bands and the intensity of each band of interest was normalized comparing it to the housekeeping GAPDH protein, used as loading control. Subsequently, the intensity of each tested band was compared to negative controls.

Statistical analysis

All results were expressed as mean values $\pm$ SD. Differences between the groups were analyzed by one-way analysis of variance (ANOVA) tests and p values were determined using the Bonferroni test. Statistical analyses were performed using the SPSS 16 package (SPSS Inc., Chicago, IL, USA). Significance was set at $p < 0.05$.

348 **Funding**

349 This work was supported by Progetto Scientifico di Ateneo 2016 – Università Politecnica delle
350 Marche “Study of new compounds and innovative strategies to control complicated bacterial skin
351 infections”.

352

353 **Trasparency declaration**

354 Nothing to declare.

REFERENCES

- 1) Russo A, Concia E, Cristini F, De Rosa FG, Esposito S, Menichetti F, Petrosillo N, Tumbarello M, Venditti M, Viale P, Viscoli C, Bassetti M. 2016. Current and future trends in antibiotic therapy of acute bacterial skin and skin-structure infections. Clin Microbiol Infect 22 Suppl 2:S27-36
- 2) Ray GT, Suaya JA, Baxter R. 2013. Incidence, microbiology, and patient characteristics of skin and soft-tissue infections in a U.S. population: a retrospective population-based study. BMC Infect Dis 13:252.
- 3) Boucher HW, Corey GR. 2008. Epidemiology of methicillin-resistant *Staphylococcus aureus*. Clin Infect Dis 46:S344–S349.
- 4) Khan A, Wilson B, Gould IM. 2018. Current and future treatment options for community-associated MRSA infection. Expert Opin Pharmacother 19:457–470.
- 5) Holmes MA, Zadoks RN. 2011. Methicillin resistant *S. aureus* in human and bovine mastitis. J Mammary Gland Biol 16:373–382.
- 6) Pexara A, Solomakos N, Govaris A. 2013. Prevalence of methicillin-resistant *Staphylococcus aureus* in milk and dairy products. J Hell Vet Med Soc 64:17–34.
- 7) Ferreira JP, Anderson KL, Correa MT, Lyman R, Ruffin F, Reller LB, Fowler VG Jr. 2011. Transmission of MRSA between companion animals and infected human patients presenting to outpatient medical care facilities. PLoS One 6:e26978.
- 8) Barlam TF, Cosgrove SE, Abbo LM, MacDougall C, Schuetz AN, Septimus EJ, Srinivasan A, Dellit TH, Falck-Ytter YT, Fishman NO, Hamilton CW, Jenkins TC, Lipsett PA, Malani PN, May LS, Moran GJ, Neuhauser MM, Newland JG, Ohl CA, Samore MH, Seo SK, Trivedi KK. 2016. Implementing an Antibiotic Stewardship Program: guidelines by the infectious diseases society of America and the society for healthcare epidemiology of America. Clin Infect Dis 62(10):e51–77.
- 9) Boucher HW, Wilcox M, Talbot G, Puttagunta S, Das AF, Dunne MW. 2014. Once-weekly dalbavancin versus daily conventional therapy for skin infection. N Engl J Med 370 (23):2169-79
- 10) Tsoulas C, Nathwani D. 2015. Review of meta-analyses of vancomycin compared with new treatments for Gram-positive skin and soft-tissue infections: Are we any clearer? Int J Antimicrob Agents 46 (1):1-7.
- 11) Doma E., Rupp C., Baccarini M. 2013. EGFR-ras-raf signaling in epidermal stem cells: roles in hair follicle development, regeneration, tissue remodeling and epidermal cancers. Int J Mol Sci 14:19361–19384.
- 12) Gill SE, Parks WC. 2008. Metalloproteinases and their inhibitors: regulators of wound healing. Int J Biochem Cell Biol 40(6–7):1334–1347.
- 13) Amar S, Smith L, Fields GB. 2017. Matrix metalloproteinase collagenolysis in health and disease Biochim Biophys Acta Mol Cell Res 1864(11 Pt A):1940-1951.
- 14) Barrientos S, Stojadinovic O, Golinko MS, Brem H, Tomic-Canic M. 2008. Growth factors and cytokines in wound healing. Wound Repair Regen 16(5):585-601.
- 15) Werner S, Grose R. 2003. Regulation of wound healing by growth factors and cytokines. Physiol Rev 83(3):835-70.
- 16) Simonetti O, Cirioni O, Ghiselli R, Goteri G, Orlando F, Monfregola L, De Luca S, Zizzi A, Silvestri C, Veglia G, Giacometti A, Guerrieri M, Offidani A, Scaloni A. 2012.

- Antimicrobial properties of distinctin in an experimental model of MRSA-infected wounds. *Eur J Clin Microbiol Infect Dis* 31(11):3047-55.
- 17) Simonetti O, Cirioni O, Goteri G, Ghiselli R, Kamysz W, Kamysz E, Silvestri C, Orlando F, Barucca C, Scalise A, Saba V, Scalise G, Giacometti A, Offidani A. 2008. Temporin A is effective in MRSA-infected wounds through bactericidal activity and acceleration of wound repair in a murine model. *Peptides* 29(4):520-8.
- 18) Sganga G, Tascini C, Sozio E, Colizza S. 2017. Early recognition of methicillin-resistant *Staphylococcus aureus* surgical site infections using risk and protective factors identified by a group of Italian surgeons through Delphi method. *World J Emerg Surg* 12:25.
- 19) Liu C, Bayer A, Cosgrove SE, Daum RS, Daum RS, Fridkin SK, Gorwitz RJ, Kaplan SL, Karchmer AW, Levine DP, Murray BE, J Rybak M, Talan DA, Chambers HF. 2011. Clinical practice guidelines by the Infectious Diseases Society of America for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children. *Clin Infect Dis* 52(3):e18–e55.
- 20) Howden BP, Peleg AY, Stinear TP. 2014. The evolution of vancomycin intermediate *Staphylococcus aureus* (VISA) and heterogenous-VISA. *Infect Genet Evol* 575-582.
- 21) Economou N J, Nahoum V, Weeks SD, Grasty KC, Zentner IJ, Townsend TM, Bhuiya MW, Cocklin S, Loll PJ. 2012. A carrier protein strategy yields the structure of dalbavancin. A carrier protein strategy yields the structure of dalbavancin. *J Am Chem Soc* 134(10): 4637–4645
- 22) Ramdeen S, Boucher HW. 2015. Dalbavancin for the treatment of acute bacterial skin and skin structure infections. *Expert Opin Pharmacother* 16(13):2073-81.
- 23) Bassetti M, Peghin M, Castaldo N, Giacobbe DR. 2019. The safety of treatment options for acute bacterial skin and skin structure infections. *Expert Opin Drug Saf* 18(8):635-650
- 24) Savoldi A, Azzini AM, Baur D, Tacconelli E. 2018. Is there still a role for vancomycin in skin and soft-tissue infections? *Curr Opin Infect Dis* 31(2):120-130
- 25) Mendes RE, Castanheira M, Farrell DJ, Flamm RK, Sader HS, Jones RN. 2016. Update on dalbavancin activity tested against Gram-positive clinical isolates responsible for documented skin and skin-structure infections in US and European hospitals (2011-13). *J Antimicrob Chemother* 71:276–8.
- 26) Malabarba A, Goldstein BP. 2005. Origin, structure, and activity in vitro and in vivo of dalbavancin. *J Antimicrob Chemother* 55 Suppl 2:ii15-20.
- 27) Chow O, Barbul A. 2014. Immunonutrition: role in wound healing and tissue regeneration. *Ad. Wound Care* 3:46–53.
- 28) Landén NX, Li D, Ståhle M. 2016. Transition from inflammation to proliferation: a critical step during wound healing. *Cell Mol Life Sci* 73: 3861–3885.
- 29) Ward CL, Sanchez CJ Jr, Pollot BE, Romano DR, Hardy SK, Becerra SC, Rathbone CR, Wenke JC. 2015. Soluble factors from biofilms of wound pathogens modulate human bone marrow-derived stromal cell differentiation, migration, angiogenesis, and cytokine secretion. *BMC Microbiol* 15: 75.
- 30) Chaney SB, Ganesh K, Mathew-Steiner S, Stromberg P, Roy S, Sen CK, Wozniak DJ. 2017. Histopathological comparisons of *Staphylococcus aureus* and *Pseudomonas aeruginosa* experimental infected porcine burn wounds. *Wound Repair Regen* 25(3):541-549
- 31) Zhang Z, Xiao C, Gibson, M, Bass SA, Khurana Hershey GK. 2014. EGFR signaling blunts allergen-induced IL-6 production and th17 responses in the skin and attenuates development and relapse of atopic dermatitis. *J Immunol* 192(3):859–866.

- 32) Simonetti O, Lucarini G, Orlando F, Pierpaoli E, Ghiselli R, Provinciali M, Castelli P, Guerrieri M, Di Primio R, Offidani A, Giacometti A, Cirioni O. 2017. Role of Daptomycin on Burn Wound Healing in an Animal Methicillin-Resistant *Staphylococcus aureus* Infection Model. *Antimicrob Agents Chemother* 24;61, pii: e00606-17.
- 33) Caley MP, Martins VL, O'Toole EA. 2015. Metalloproteinases and wound healing. *Adv Wound Care (New Rochelle)* 4(4):225–234.
- 34) Hattori N, Mochizuki S, Kishi K, Nakajima T, Takaishi H, D'Armiento J, Okada Y. 2009. MMP-13 plays a role in keratinocyte migration, angiogenesis, and contraction in mouse skin wound healing. *Am J Pathol* 175:533–546.
- 35) Mohan R, Chintala SK, Jung JC, Villar WV, McCabe F, Russo LA, Lee Y, McCarthy BE, Wollenberg KR, Jester JV, Wang M, Welgus HG, Shipley JM, Senior RM, Fini ME. 2002. Matrix metalloproteinase gelatinase B (MMP-9) coordinates and effects epithelial regeneration. *J Biol Chem* 277:2065–2072.
- 36) Lazaro JL, Izzo V, Meaume S, Davies AH, Lobmann R, Uccioli L. 2016. Elevated levels of matrix metalloproteinases and chronic wound healing: an updated review of clinical evidence. *J Wound Care* 25(5):277-87.
- 37) Simonetti O, Cirioni O, Lucarini G, Orlando F, Ghiselli R, Silvestri C, Brescini L, Rocchi M, Provinciali M, Guerrieri M, Di Primio R, Giacometti A, Offidani A. 2012. Tigecycline accelerates staphylococcal-infected burn wound healing through matrix metalloproteinase-9 modulation. *J Antimicrob Chemother* 67(1):191-201.
- 38) Simonetti O, Lucarini G, Cirioni O, Zizzi A, Orlando F, Provinciali M, Di Primio R, Giacometti A, Offidani A. 2014. Delayed wound healing in aged skin rat models after thermal injury is associated with an increased MMP-9, K6 and CD44 expression. Erratum in: *Burns*. 40(1):175 related to *Burns*. 2013 Jun;39(4):776-87.
- 39) McCarty SM, Percival SL. 2013. Proteases and Delayed Wound Healing. *ADV Wound Care (New Rochelle)* 2(8):438-447.
- 40) Bergers G, Brekken RA, McMahon G, Vu TH, Itoh T, Tamaki K, Tanzawa K, Thorpe P, Itohara S, Werb Z, Hanahan D. 2000. Matrix metalloproteinase-9 triggers the angiogenic switch during carcinogenesis. *Nat Cell Biol* 2:737–44.
- 41) Ardi VC, Van den Steen PE, Opdenakker G, Schweighofer B, Deryugina EI, Quigley JP. 2009. Neutrophil MMP-9 proenzyme, unencumbered by TIMP-1, undergoes efficient activation *in vivo* and catalytically induces angiogenesis via a basic fibroblast growth factor (FGF-2)/FGFR-2 pathway. *J Biol Chem* 284:25854–66.
- 42) Howard EW, Soo C, Hunter S, Zhang X, Shaw WW, Longaker MT. 1997. Differences in MMP and TIMP gene expression during wound repair. *Surg Forum* 48:688–91.
- 43) Bao P, Kodra A, Tomic-Canic M, Golinko MS, Ehrlich HP, Brem H. 2009. The role of vascular endothelial growth factor in wound healing. *J Surg Res* 153(2):347-58.
- 44) Silvestri C, Cirioni O, Arzeni D, Ghiselli R, Simonetti O, Orlando F, Ganzetti G, Staffolani S, Brescini L, Provinciali M, Offidani A, Guerrieri M, Giacometti A. *In vitro* activity and *in vivo* efficacy of tigecycline alone and in combination with daptomycin and rifampin against Gram-positive cocci isolated from surgical wound infection. *Eur J Clin Microbiol Infect Dis* 31(8):1759-64.
- 45) Seltzer E, Dorr MB, Goldstein BP, Perry M, Dowell JA, Henkel T; Dalbavancin Skin and Soft-Tissue Infection Study Group. 2003. Dalbavancin Skin and Soft-Tissue Infection Study Group. Once-weekly dalbavancin versus standard-of-care antimicrobial regimens for treatment of skin and soft-tissue infections. *Clin Infect Dis* 37(10):1298-303.
- 46) Simonetti O, Cirioni O, Cacciatore, Baldassarre L, Orlando F, Pierpaoli E, Lucarini G, Orsetti E, Provinciali M, Fornasari E, Di Stefano A, Giacometti A, Offidani A. 2016. Efficacy of the Quorum Sensing Inhibitor FS10 Alone and in Combination with

499 Tigecycline in an Animal Model of Staphylococcal Infected Wound. PLoS One
500 11(6):e0151956.

Figure Legend

Figure 1. Bacterial counts of excised wound. Counts are expressed as log of CFU/ml. The limit of detection for this method was approximately 10 CFU/ml. * represent $p < 0.05$ compared to infected but not treated group-

Figure 2. Representative images of hematoxylin- and eosin-stained histological sections of wound healing tissues from uninfected mice, infected rats treated with dalbavancin or vancomycin, and infected untreated mice on day 14 after injury. Wound healing was still incomplete in the untreated groups, both treated groups exhibited a good re-epithelialization; in particular, the wounds treated with Dalbavancin showed robust epidermal coverage and well organized granulation tissue (Original magnification 100x).

Figure 3. Immunohistochemical staining of VEGF and EGFr in wound healing tissues from uninfected mice, infected rats treated with dalbavancin or vancomycin, and infected untreated mice on day 14 after injury. VEGF and EGFr expression was greater in the uninfected and treated infected groups respect to the untreated, in particular the wounds treated with dalbavancin showed the highest value (immunoperoxidase; original magnification 200x).

Figure 4. Immunohistochemical staining of MMP-9 and MMP-1 in wound healing tissues from uninfected mice, infected mice treated with dalbavancin or vancomycin, and infected untreated mice on day 14 after injury. MMP-9 and MMP-1 expression was high in the untreated infected wounds, the treated wounds showed the lowest values (immunoperoxidase; original magnification 200x).

Figure 5. Analysis of the expression of EGFR, MMP-1, MMP-9 and VEGF. Western blot and densitometric analyses of EGFR, MMP-1, MMP-9 and VEGF, in a representative of the three

527 performed experiments. Densitometric analyses of the immunoreactive bands is quantified as ratio
 528 between band relative to EGFR, VEGF, MMP-1, MMP-9 and, and GAPDH in corresponding
 529 samples, revealed by arbitrary units. Data are expressed as a mean $\pm$ SD from analyses performed
 530 on uninfected, untreated infected, infected treated with vancomycin and infected treated with
 531 dalbavancin tissue samples. * infected treated with dalbavancin versus uninfected, untreated
 532 infected, infected treated with vancomycin $p < 0.05$. ** untreated infected versus uninfected,
 533 infected treated with dalbavancin, infected treated with vancomycin $p < 0.05$.

Table 1-A five-tiered grading system to evaluate wound healing.

Scores	Re-epithelialization	Granulation tissue	Dermis organization	Inflammatory infiltrate
0	Trace and modest keratinocyte migration	Trace	Absent	Absent
1	Evident keratinocyte migration	Hypocellular and none vessel	Trace	Modest
2	Differentiation	Many cells and few vessels	Modest	Good
3	Hypertrophic and partial stratum corneum	Many fibroblasts, some fibers and some vessels	Good	Evident
4	Complete and normal	More fibers, few cells	Evident	–

Table 2- Analysis of wound healing in the different mice groups according to the histological scores.

Model	Re-epithelialization (mean±SD)	Granulation tissue (mean±SD)	Dermis organization (mean±SD)	Inflammatory infiltrate (mean±SD)
Uninfected (C0)	3.85±0.25	3.52±0.22	3.85±0.50	1.02±0.02
Untreated infected (C1)	0.50±0.08	1.25±0.08	1.80±0.30	2.00±0.25
Infected treated with vancomycin (C2)	2.85±0.20	3.00±0.25	3.20±0.28	1.20±0.03
Infected treated with dalbavancin (C3)	3.35±0.10	3.20±0.15	3.75±0.18	1.2±0.08
ANOVA test Test di Bonferroni p<0.05	C0 vs C1, C2, C3 C1 vs C0, C2, C3 C3 vs C2	C0 vs C1, C2 C1 vs C2,C3,	C0,C3 vs C0, C1 C2 vs C1	C1vs C0,C2, C3

Table 3- Immunohistochemical expression of VEGF, MMP2, MMP9, MMP1 and EGFr in the different mice groups.

Model	VEGF (mean±SD)**	EGFr (mean±SD)	MMP-9 (mean±SD)	MMP-1 (mean±SD)
Uninfected (C0)	30.0±3.2	40.0±2.5	20.3±2.6	22.0±1.2
Untreated Infected (C1)	15.5±2.6*	31.0±2.5	37.0±1.6	45.0±5.2
Infected treated with vancomycin (C2)	31.6±3.2	30.0±2.0	16.2±1.8*	18.0±3.6*
Infected treated with dalbavancin (C3)	40.0±4.2*	50.0±2.2*	15.0±2.5*	20.0±3.0*
ANOVA test Test di Bonferroni p<0.05*	C3 vs C0, C1, C2, C1 vs C0, C2	C3 vs C0, C1, C2 C0, C3 vs C1, C2 C3 vs C0	C1 vs C0, C2, C3 C2,C3 vs C0	C1 vs C0, C2, C3

** percentage of positive cells

Untreated infected

Dalbavancin

Vancomycin

Uninfected

VEGF

EGFr

Uninfected

Dalbavancin

Vancomycin

Untreated

MMP9

MMP1

Uninfected

Dalbavancin

Vancomycin

Untreated