How long is long enough? Decreasing on plant extract lethal effects, over time, affecting *Aedes aegypti* larval mortality

Short title: Aedes aegypti mortality by plant extracts

Gilberto Dinis Cozzer^{1¶}; Renan de Souza Rezende^{1¶}; Junir Antônio Lutinski^{2&}; Walter Antônio Roman Junior^{2&}; Maria Assunta Busato^{2&}; Jacir Dal Magro; Daniel Albeny-Simões^{1¶*}

¹ Environmental Sciences Department, Community University of Chapecó Region, Chapecó, Santa Catarina, Brazil.

² Health Sciences Department, Community University of Chapecó Region, Chapecó, Santa Catarina, Brazil.

*Corresponding author:
E-mail: danielalbeny@unochapeco.edu.br (DAS)

¶ These authors contributed equally to this work

& These authors also contributed equally to this work

Abstract

The mosquito *Aedes aegypti* has overcome all kinds of human being mosquito control attempts over the last century. Strategies for vector population control resorts to the use of synthetic insecticides, which can lead to problems of intoxication in humans and environmental contamination. We evaluated the effects of *Bacillus thuringiensis* var. *israelensis* (Bti), *Ilex paraguariensis* (mate-herb) and *Ilex theezans* (caúna-herb) extracts against *A. aegypti* larvae mortality. The bioassays were conducted under controlled laboratory conditions of temperature (27±3°C) and photoperiod (12h). Hydroalcoholic *I. theezans* leaves extract displayed better residual effect compared to *I. paraguariensis* fruit aqueous extract. Variation in larval mortality was also observed in the exposure periods (low after a few weeks). Low mortality after a few weeks may mean increased the food for mosquito in a oppose effect over time. The residual effect of Bti was observed during the 56 days of the study duration (100% of mortality). The strongest residual effect of *I. theezans* was probably due to the presence of chemical on its leaves, such coumarins, hemolytic saponins and cyanogenic glucosides, absent in *I. paraguariensis*. On the other hand, alternative methods to vector control present risks in a long term scale by reversal of larvicide effect into food resource. Our results contributed to the prospection of natural insecticides and open the possibility for subsequent studies of the use of plant extracts in field situations in a short time scale.

Keywords: Dengue, vector control, inseticide, entomology, yerba mate.

Introduction

Over the last century the mosquito *Aedes aegypti* Linnaeus, 1762 has overcome all mosquito control attempts by human being. *Aedes* females are well known by their capacity of naturally and/or under laboratory conditions replicate and transmit over 100 kinds of viruses [1]. As an example, in Brazil dengue, chikungunya, zika and most recently mayaro [1] viruses are a real threat to public health [2]. It became necessary the *A. aegypti* population control in order to reduce viruses' transmission and consequently the epidemic status. Although several chemical and natural products have being extensively used on attempts to reduce adults and immature mosquito population [3,4], appropriated mosquito population reduction is not close to come true. Especially because the gene resistance selectivity due to chemicals and natural products application misuse [5]. Accordingly, the most effective disease prevention method still remains targeting the mosquito population by elimination of mosquito breeding places [6].

A very promising field for mosquito population reduction is focusing the mosquito control strategies on the immature aquatic stages, when the insect is more vulnerable [7,8]. For this purpose, the use of synthetic insecticides is well known as being quite effective causing mosquito larvae mortality [9–13]. However, those chemicals might affect humans causing intoxication and environmental contamination, affecting biodiversity [9,10,12,13]. For the environment, the continuous use of synthetic insecticides may present undesirable effects, such as the long-term residence in the environment, selection of resistant populations and the appearance of new pests [10,12,13]. Regarding to the human health, the presence of such synthetic chemicals on the environment can cause neurological damage, and is associated with a wide range of symptoms, with significant deficits in nervous system function [14].

Alternatively to the use of synthetic chemicals, biological control play an important role on mosquito control [15,16]. The use of bacteria spores as a mosquito larvicial has been highlighted among the several components that are part of the mosquito integrated management program [15]. In the last decades, the use of inactivated spores of the *Bacillus thuringiensis* var. *israelensis* (Bti) bacteria spread into the mosquito breeding places water, has accomplished desired results, reaching mortality rates above 99% [17,18]. Additionally, during the last few years, plant derived compounds have been extensively used as an alternative method for controlling mosquitoes, not only because it is a new insecticidal agent, but also because it has been described as being environmental friendly [11,19–21]. The use of natural insecticides has some advantages over traditional synthetic products, since natural products are potentially less toxic to the environment. Environmental friendly compounds are less concentrated, have faster degradation and are specific to certain insect groups, resulting in less occupational exposure and less environmental pollution [22].

Ilex paraguariensis A. St.-Hil (Aquifoliaceae), known as “Mate” is an abundant plant [23], native of South America, twenty meters tall, endowed with dense crown and very branched [23]. *I. paraguariensis* leaves, after processing, are traditionally used to prepare a regional tea known as “Chimarrão” in Argentina, Brazil, Paraguay and Uruguay [23]. *I. paraguariensis* is commercially interesting due to caffeine and theobromine presence, both, recognized by displaying a nervous and cardiocirculatory systems stimulant effect [24]. The described pharmacological activities for *I. paraguariensis* leaf extracts include antioxidant, hypolipidemic [25,26] and hypoglycemic effects [27]. Also, *Ilex theezans* Mart. Ex Reissek (Aquifoliaceae), popularly known as cauna, is commonly found at southern Brazil [23]. It is well known due to the physiological characteristics of its leaves as adulterant of *I. paraguariensis* [28]. It is an evergreen

tree, early secondary or late secondary species [23], 20 meters long by 70 cm diameter average [28]. Both, *I. paraguariensis* fruit extract and *I. theezans* hydroalcoholic leaves extracts have larvicidal effect against *A. aegypti* larvae [9,20].

There are studies showing that *Ilex* spp. leaves and fruit extracts kill *A. aegypti* larvae within a 24h observational time [9,20]. However, there are no studies on the current literature evaluating the effects of time on the bioinsecticides lethal activity. How long is long enough for the bioinsecticide to still keep the mosquito larvae killing activity? [29–31]. In this context, we evaluated the lethal residual effect of *B. thuringiensis* var. *israelensis*, and leaves and fruits extracts of *I. theezans*, and *I. paraguariensis* (respectively), on *A. aegypti* larvae survivorship. We hypothesized that time would positively affect *A. aegypti* larvae survival due to the plant extracts lethal compounds decay, but the mortality caused by *I. theezans* would be higher compared to *I. paraguariensis* due to the difference in physical and chemical characteristics.

Materials and Methods

Animal source

The *A. aegypti* larvae used in this experiment were provided by the Laboratório de Entomologia Ecológica (LABENT-Eco) mosquito colony. A filter paper holding about 300 eggs was placed in a plastic tray (30x20cm) holding 1L of tap dechlorinated water. After hatching the larvae were split among 3 equal size, above described, plastic tray and feed with 2g of fish food. The mosquito larvae were raised for about 4-5 days until reaching 3rd and 4rd instars.

Plant source and extracts preparation

Fruits and leaves were obtained from native trees located at the Marechal Bormann district (27 ° 19'05 "S; 52 ° 65'11"W), Chapecó (SC), in December, 2016. The plant parts were dehydrated at room temperature ($\pm 20^{\circ}\text{C}$), pulverized in a knife mill (Cielamb®, CE 430) and stored away from light and moisture. The plant extracts were done according to Busato *et al.* (2015) and Knakiewicz *et al.* (2016). We took a sample of twenty grams of *I. paraguariensis* dehydrated fruits and *I. theezans* leaves. Both samples were extracted by turbolysis using 200ml of distilled and deionized water and a hydroalcoholic solution (90% ethanol; 200 ml) as the solvent, respectively [32]. The extracts were filtered in Büchner, rotavapor concentrated under reduced pressure, lyophilized, weighed, identified and stored in a freezer at -20°C . Hydroalcoholic and aqueous extracts were prepared using *I. theezans* and *I. paraguariensis* leaves and fruits, respectively. We used leaves in a concentration of $1000\mu\text{g/ml}$ and fruits were diluted to $2000\mu\text{g/ml}$. *B. thuringiensis* var. *israelensis* (Bti) strain WG® was used in a concentration of 0.004g/L , lethal dose specified by the manufacturer.

Experimental microcosms and design

We used 300ml plastic cups holding 100ml of dechlorinated water plus the treatment proposed. In each individual microcosm we added twenty 3rd and 4th instar *A. aegypti* larvae. Each container was covered with a mosquito net held by a rubber elastic band. We tested for the effects of Bti spores, *I. theezans*, *I. paraguariensis* leaves and fruits, respectively and clean aged water (control) on the *A. aegypti* larval mortality after seven days exposition. Before running the mortality test, each experimental treatment was aged from one to eight weeks. We considered each week as one age block with each treatment replicated six times. By doing this experimental design we assure the independence of each set of treatments. The aged treatments were used to test for

larval survival in each experimental week. At the end of the seventh day the larval survival was recorded. Both pupae and emerged adults were considered survivals. The experiment was performed for eight weeks (56 days) and carried out between April and May 2017 at the LABENT-Eco mosquito colony room under controlled conditions of temperature and photoperiod ($27 \pm 3^\circ\text{C}$, 12h D:L).

Statistics

Because both negative (Bti) and positive (tap water) control survivor rate was 0.16% and 100%, respectively, we analyzed the data in both ways, with (complete model) and without (simple model) these two categories. To evaluate differences in percentage of larval mortality (response variables) in simple (*I. paraguariensis* and *I. theezans*) and complete model (only water, Bti spores, *I. paraguariensis* and *I. theezans*), week (1 to 8) and interaction between week and treatment (explicative variables) we used factorial GLM, with binomial correct to quasi-binomial (link= logit, test= Chi-square) distributions for larval mortality (response variable; [33]). All GLMs analyzed were corrected for cases of under- or overdispersion.

Differences among the categorical variables were assessed through a contrast analysis [33]. In this contrast analysis (orthogonal), the dependent variables of different treatment and weeks were ordered (increasingly) and tested pairwise (with the closest values). Sequentially these dates are adding to the model values with no differences and testing with the next in a steps model simplification (for more see also chapter 9 of Crawley, (2007)). All analyses were performed using R [34].

Results

The *A. aegypti* larvae mortality was not affected by water age (positive control). We observed no larva deaths independent of the water age used. Oppositely, the Bti lead to zero percent of larvae survivorship until the age of 7 weeks, with only 6,6% of larvae alive on the age of 8 weeks (negative control). In this way, due to these extreme results in controls (0% of mortality in positive control and 100% of mortality in negative control), the mortality data were also analyzed only between treatments (*I. paraguariensis* and *I. theezans*).

The treatment (*I. paraguariensis* and *I. theezans*), week (1 to 8) and interaction factor (week:treatment) was significantly different among *A. aegypti* larvae mortality for both GLMs models (With and Without positive and negative controls; Table 1). The higher *A. aegypti* larvae mortality was found in Bti treatment (negative control), followed to *I. theezans*, *I. paraguariensis* and positive control (Table 1; Fig 1a). Also, *I. theezans* hydroalcoholic leaves extract, independently of the extract age, significantly killed more *A. aegypti* larvae than the aqueous *I. paraguariensis* fruits extract (Table 1; Fig 1b).

Table 1: Generalized linear models (GLM), degrees of freedom (Df), Residual Deviance (total and in) and p values, comparing the percentage of *Aedes aegypti* larvae mortality after the exposure to treatments (water control, *Bacillus thuringiensis israelensis* - Bti, hydroalcoholic dried leaves extract of *Ilex theezans* and aqueous *Ilex paraguariensis* fruits extract), time (8 weeks) and interaction among treatments and weeks, under laboratory conditions.

GLM	Df	Resid.		Pr(>Chi)	Analysis of contrast
		Dev.	%		
		Resid. Dev.	%		
a. With positive and negative controls					
Treatments	3	132.9	74.1	< 0.001	Control < <i>Ilex paraguariensis</i> < <i>Ilex theezans</i> < Bti

Weeks	7	33.4	18.6	< 0.001	Week 8 = 7 < 5 < 4 = 6 < 3 = 2 < 1
Treatment:weeks	21	3.0	1.7	< 0.001	
Residual	160	10.0	5.6		

b. Without positive and negative controls

Treatments	1	3.1	6.6	< 0.001	<i>Ilex paraguariensis</i> < <i>Ilex theezans</i>
Weeks	7	32.3	68.7	< 0.001	Week 8 = 7 < 5 < 4 = 6 < 3 = 2 < 1
Treatment:weeks	7	2.4	5.1	0.002	
Residual	80	9.2	19.6		

177

178 We observed a positive relationship between *A. aegypti* larvae survivorship and plant

179 extract age. We can also say that in general, both *I. paraguariensis* and *I. theezans* kill less (mainly

180 after 7 weeks) mosquito larvae as the plant extracts age (Fig MS1). We found higher significant in

181 *A. aegypti* larvae mortality in 1 week followed to 2 and 3 weeks, 4 and 6 weeks, 5 weeks and 7

182 and 8 weeks (Figs 1b and c).

183

184 **Fig 1. *Aedes aegypti* larvae mortality among treatments (a), sample weeks with positive and**

185 **negative controls (b) and sample weeks without positive and negative controls (c) among.**

186 Different letters (“a”, “b”, “c”, “d” and “e”) indicate significant differences. Boxes represent the

187 quartiles, the bold line represents the median, horizontal dashed line the mean, the vertical dashed

188 line represents the upper and lower limits and circles the outliers.

189

190 The residual deviance (estimate to explain the variance of the tested variables) in GLM

191 with positive and negative controls, showed that differences in all treatments (74%) was the main

192 responsible for the *A. aegypti* larvae mortality followed by weeks (18%; Table 1). On the other

193 hand, the residual deviance in GLM without positive and negative controls, showed that

differences in all weeks (68%) was the main responsible for the *A. aegypti* larvae mortality followed by treatments (19%), and only later was explained by treatments (6%; Table 1).

Fig MS1: *A. aegypti* larvae survival mean as a function of plant extracts age.

The extracts age are represented by weeks. The circles represent, aqueous *I. paraguariensis* fruits extract (open) and *I. theezans* hydroalcoholic leaves extract (closed). The asterisks (*) represents significant statistical difference between *I. paraguariensis* and *I. theezans* affecting larvae survival.

Discussion

Reversal of larvicide effect into food resource over time

I. theezans and *I. paraguariensis* extracts are promising against mosquito larvae (if applied and monitored in the first weeks) and the possibility of using these plants extracts as larvicides for *A. aegypti* represents. As it is natural extracts and does not leave toxic waste in the environment, its use may be advantageous. This extracts are an abundant and accessible alternative in southern Brazil, where *A. aegypti* infestation and dengue cases have been observed in the last decade [9]. However, extracts age caution should be applied on mosquito control attempts. The main idea of this study is not discouraging the use of such alternative method. We are only concerned in bringing an advice that we need to be aware about the *Ilex* spp. plant and fruit extract age before using it for mosquito control purposes. Better results can also be obtained with the development of additional studies, evaluating the larvicidal activity of pure compounds isolated from these plants. Also, better results can also be obtained as well as evaluate if there is a supporting effect of more than one active principle with larvicidal action for *A. aegypti*.

It is possible that the plant extracts degrade as the time goes on and these organic compounds with previous larvicidal activity may become food for *A. aegypti* larvae (corroborate to increase of importance to residual deviance percentage in GLMs models). The transformation of larvicide effect into food, especially on those treatments seven and eight weeks old, which displayed the highest survivorship rate. Because of this apparently plant extract larvicidal effect loss. Therefore, we recommend that the plant extract age should be considered into attention, caution when using these extracts in time series, since they may serve as food for bacteria [35]. Finally, plant extract age will serve as food for the larvae, contributing to its development rather than combat it, in a negative effect to to mosquito population control.

A. aegypti larvae mortality between plant extracts

The plant extracts preparation age plays an important role on mosquito larvae mortality (mainly with positive and negative controls). Despite the plant species and parts and extraction method, mosquito larval mortality decrease as the plant extracts age. However, the extracts tested were high efficient in the first weeks of the environment (high mortality). The higher *A. aegypti* larvae mortality found on *I. theezans* extracts can be in partially explained by the use of solvents during the extraction process [36]. The hydroalcoholic extraction method, used for *I. theezans*, extracts reduced polarity chemical constituents from the plant tissues, and these molecules have a greater ability to penetrate the mosquito larvae cells and modify metabolic activities.

On the other hand, aqueous extraction, used for the *I. paraguariensis* fruits, preferentially removes high polarity chemical compounds which are not able to easily penetrate such cells [36]. Also, *A. aegypti* larvae susceptibility to *I. theezans* may be explained by the presence of secondary metabolites of the coumarin class and absence of alkaloids when compared to *I. paraguariensis*

[37]. Coumarins are part of several plants secondary metabolism being well known for displaying insecticidal activities, acting as an adult repellent, oviposition deterrence, feeding and growth inhibition, morphogenetic and hormonal system alterations, sexual behavior changes, adult sterilization, among others [38]. It is important to point out that in raw plant extracts the active constituents are usually found in small concentrations [22].

A. aegypti larvae mortality among week old

In both, *I. theezans* and *I. paraguariensis*, the *A. aegypti* larvae mortality exposed to one-week old plant extracts was reduced to 100%. However, the plant extracts for both plants' species decrease the mosquito larvae lethality as the plant extracts age. Those results pointed out the need to carefully select the right age for a *Ilex* spp. plant extract before using it for mosquito larvae control purpose. Especially because, regarding to the residual larvicidal power, a product is considered efficient for a pest population control when it reaches a population reduction above 80% of the individuals, otherwise there is the selection of resistance genes [39].

In addition, it is possible that beyond decaying the lethal chemical compounds, which are toxic for mosquito larvae on the first week. As the time goes on the organic compounds present on the plant extracts act as a profitable food source for mosquito larvae. In this way, since organic compounds are well known as being an important component of several larval habitats, forming the basis of many food webs [40,41]. Microorganisms, such as bacteria, play an important role in the cycling and breaking of large organic molecules [42]. Therefore, microorganisms may making them more easily assimilable to aquatic organisms such as mosquito larvae, especially those belonging to the Culicidae family [40]. As a result of this, decomposing microbial communities

present a relevant contribution to the diet of culicid larvae, which end up ingested with the organic remains over time [40,43–45].

Our Bti results lead to 93.4% mortality using the eighth week aged solution. The residual effect described in the manufacturer's technical manual is 30 days. Also, with the values obtained in this work being much higher than those described, with a lethality of 100% for 49 days and 6.6% of larval survival up to 56 days. In this way, it was possible to evaluate the effectiveness of the products that are already being used by the state programs to *A. aegypti* control and combat. Therefore, testing their effectiveness against the existing vector populations, since the information obtained could indicate if there is a need insecticides currently used. It should be emphasized here that the use of the methodology without water renewal in the experiment ends up resulting in a longer residual effect, not evaluating the impact of water renewal in reducing the residual effect [46]. It would be important in other studies to simulate the situation in the field of realities where deposits dominated by permanent emptying and replacement of water, which would probably contribute to reduction of the residual effect duration for all the treatments tested.

Conclusions

Our study reinforces the idea of the continuously need to implement alternative methods to vector control, but there are long term risk. Also, identifying potential alternative pathways for mosquito population control using natural products originated from the native flora in the short term. The above described plants have high larvicidal potential against the mosquito *A. aegypti*, allowing it's easy access by the local population, and contributing to the maintenance of the quality of life and well-being of the population. As well as reducing public expenditures with vector

control and in the treatment of confirmed cases of dengue. Finally, we found that time would positively affect *A. aegypti* larvae survival due to the plant extracts lethal compounds decay, corroborating our first hypothesis. We also found that the mortality by *B. thuringiensis* var. *israelensis* will be constant throughout the experimental period and *A. aegypti* larvae survival will be lower in the treatments of plant extracts compared to *B. thuringiensis* var. *israelensis*. The high mortality was observed in *I. theezans* compared to *I. paraguariensis*, corroborating our second hypothesis. The strongest residual effect of *I. theezans* was probably due to the presence of chemical on its leaves, such coumarins, hemolytic saponins and cyanogenic glucosides, absent in *I. paraguariensis*.

Acknowledgements

The authors thank Unochapecó for the availability of the laboratories use and for the Research Program for the Unified Health System.

References

1. Weaver SC, Reisen WK. Present and future arboviral threats. *Antiviral Res.* 2010;85(2):328–45.
2. Lopes N, Nozawa C, Linhares REC. Características gerais e epidemiologia dos arbovírus emergentes no Brasil. *Rev Pan-Amazônica Saúde.* 2014;5(3):55–64.
3. Liu N. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research Directions. *Annu Rev Entomol.* 2015;60(1):537–59.
4. Apaire-Marchais V, Ogliastro M, Chandre F, Pennetier C, Raymond V, Lapied B. Virus

- 306 and calcium: An unexpected tandem to optimize insecticide efficacy. *Environ Microbiol*
307 *Rep.* 2016;8(2):168–78.
- 308 5. Sun Y, Dong Y, Li J, Lai Z, Hao Y, Liu P, et al. Development of large-scale mosquito
309 densovirus production by in vivo methods. *Parasites and Vectors.* 2019;12(1):1–13.
- 310 6. Brasil. Programa Nacional de Controle da Dengue (PNCD). 1st ed. Brasília: Ministério da
311 Saúde; 2002. Available from: http://bvsmms.saude.gov.br/bvs/publicacoes/pncd_2002.pdf
- 312 7. Dengue: Instruções para Pessoal de Combate ao Vetor - Manual de Normas Técnicas. 3rd
313 ed. Brasília: Ministério da Saúde; 2001.
- 314 8. Crivelenti LZ, Guilherme LC, Morelli S, Borin S. Toxicidade do Inseticida
315 Organofosforado Abate® em Alevinos de *Poecilia reticulata*. *J Brazilian Soc Ecotoxicol.*
316 2010;5(2–3):1–13.
- 317 9. Busato MA, Vitorello J, Lutinski JA, Magro JD, Scapinello J. Potencial Larvicida De
318 *Melia Azedarach* L. E *Ilex Paraguariensis* St. Hil. No Controle De *Aedes Aegypti*
319 (Linnaeus, 1762) (Diptera: Culicidae). *Ciência e Nat.* 2015;37(2):277–82.
- 320 10. Tahir U, Khan UH, Zubair MS, Bahar-e-Mustafa. *Wolbachia pipientis*: A potential
321 candidate for combating and eradicating dengue epidemics in Pakistan. *Asian Pac J Trop*
322 *Med.* 2015;8(12):989–98.
- 323 11. Guarda C, Lutinski JA, Antônio W, Maria R, Busato A. Susceptibilidade ao inseticida
324 temefós no controle do *Aedes aegypti* (Diptera : Culicidae). *Interciencia.* 2016;41(4):243–
325 7.
- 326 12. Baskar K, Sudha V, Nattudurai G, Ignacimuthu S, Duraipandiyan V, Jayakumar M, et al.
327 Larvicidal and repellent activity of the essential oil from *Atalantia monophylla* on three
328 mosquito vectors of public health importance, with limited impact on non-target zebra

- 329 fish. *Physiol Mol Plant Pathol*. 2018;101:197–201.
- 330 13. Govindarajan M, Rajeswary M, Senthilmurugan S, Vijayan P, Alharbi NS, Kadaikunnan
331 S, et al. Larvicidal activity of the essential oil from *Amomum subulatum* Roxb.
332 (*Zingiberaceae*) against *Anopheles subpictus*, *Aedes albopictus* and *Culex*
333 *tritaeniorhynchus* (Diptera: Culicidae), and non-target impact on four mosquito natural
334 enemies. *Physiol Mol Plant Pathol*. 2018;101:219–24.
- 335 14. De Araújo AJ, De Lima JS, Moreira JC, Jacob SDC, Soares MDO, Monteiro MCM, et al.
336 Exposição múltipla a agrotóxicos e efeitos à saúde: Estudo transversal em amostra de 102
337 trabalhadores rurais, Nova Friburgo, RJ. *Cienc e Saude Coletiva*. 2007;12(1):115–30.
- 338 15. Zara AL de SA, Santos SM Dos, Fernandes-Oliveira ES, Carvalho RG, Coelho GE.
339 Estratégias de controle do *Aedes aegypti*: uma revisão. *Epidemiol e Serv saude Rev do*
340 *Sist Unico Saude do Bras*. 2016;25(2):391–404.
- 341 16. Coelho WMD, Coêlho J de CA, Bresciani KDS, Buzetti WAS. Biological control of
342 *Anopheles darlingi*, *Aedes aegypti* and *Culex quinquefasciatus* larvae using shrimps.
343 *Parasite Epidemiol Control*. 2017;2(3):91–6.
- 344 17. Nascimento AKS. O uso do *Bacillus thuringiensis* variedade israelensis , no controle das
345 larvas do *Aedes aegypti* . M. Sc. Thesis, Centro Universitário de Brasília. 2003. Available
346 from: [https://docplayer.com.br/15917402-O-uso-do-bacillus-thuringiensis-variedade-](https://docplayer.com.br/15917402-O-uso-do-bacillus-thuringiensis-variedade-israelensis-no-controle-das-larvas-do-aedes-aegypti.html)
347 [israelensis-no-controle-das-larvas-do-aedes-aegypti.html](https://docplayer.com.br/15917402-O-uso-do-bacillus-thuringiensis-variedade-israelensis-no-controle-das-larvas-do-aedes-aegypti.html)
- 348 18. Soares-da-Silva J, Queirós SG, de Aguiar JS, Viana JL, Neta M dos RAV, da Silva MC, et
349 al. Molecular characterization of the gene profile of *Bacillus thuringiensis* Berliner
350 isolated from Brazilian ecosystems and showing pathogenic activity against mosquito
351 larvae of medical importance. *Acta Trop*. 2017;176(June):197–205.

- 352 19. Gomes PRB, Silva ALS, Pinheiro HA, Carvalho LL, Lima HS, Silva EF, et al. Avaliação
353 da atividade larvívica do óleo essencial do *Zingiber officinale* Roscoe (Gengibre) frente ao
354 mosquito *Aedes aegypti*. *Rev Bras Plantas Med.* 2016;18(2):597–604.
- 355 20. Knakiewicz AC, Lutinski JA, Guarda C, Paris A, Belotti A, Busato MA, et al. Larval
356 susceptibility of *Aedes Aegypti* (L.) (Diptera: Culicidae) to extracts of *Ilex Paraguariensis*
357 and *Ilex Theezans*. *Rev Bras Ciências Ambient.* 2016;(42):113–20.
- 358 21. Rosa CS, Veras KS, Silva PR, Lopes Neto JJ, Cardoso HLM, Alves LPL, et al.
359 Composição química e toxicidade frente *Aedes aegypti* L. e *Artemia salina* Leach do óleo
360 essencial das folhas de *Myrcia sylvatica* (G. Mey.) DC. *Rev Bras Plantas Med.*
361 2016;18(1):19–26.
- 362 22. Krinski D, Massaroli A, Machado M. Potencial insetívica de plantas da família
363 Annonaceae. *Rev Bras Frutic.* 2014;36(edição especial):225–42.
- 364 23. Souza MFF. Chá mate (*Ilex paraguariensis*): compostos bioativos e relação com atividade
365 biológica. M. Sc. Thesis, Universidade de São Paulo; 2009. Available from:
366 <http://www.teses.usp.br/teses/disponiveis/6/6138/tde-21092009-102503/pt-br.php>
- 367 24. Castaldelli APA, Vieira LP, Przygodda F, Nunes Z, Jorge M. Efeito da erva mate (*Ilex*
368 *paraguariensis* A . St . -Hill) no comportamento e fisiologia de ratos Wistar. *Rev Bras*
369 *Biociências.* 2011;9(4):514–9.
- 370 25. Gao H, Liu Z, Wan W, Qu X, Chen M. Aqueous extract of yerba mate tea lowers
371 atherosclerotic risk factors in a rat hyperlipidemia model. *Phyther Res.* 2013;27(8):1225–
372 31.
- 373 26. Messina D, Soto C, Méndez A, Corte C, Kemnitz M, Avena V, et al. Efecto
374 hipolipemiante del consumo de mate en individuos dislipidémicos. *Nutr Hosp.*

- 2015;31(5):2131–9.
27. Conceição EPS, Kaezer AR, Peixoto-Silva N, Felzenszwalb I, De Oliveira E, Moura EG, et al. Effects of *Ilex paraguariensis* (yerba mate) on the hypothalamic signalling of insulin and leptin and liver dysfunction in adult rats overfed during lactation. *J Dev Orig Health Dis.* 2017;8(1):123–32.
28. Athayde ML, Schenkel EP, Gosmann G, Guillaume D. Triterpenoids from the leaves of *Ilex theezans* Martius ex Reiss. *Acta Farm Bonaer.* 1999;18(1):49–52.
29. Resende MC, Gama RA. Persistência e eficácia do regulador de crescimento pyriproxyfen em condições de laboratório para *Aedes aegypti*. *Rev Soc Bras Med Trop.* 2006;39(1):72–5.
30. dos Santos RLC, Fayal A da S, Aguiar AEF, Vieira DBR, Póvoa MM. Evaluation of the residual effect of pyrethroids on *Anopheles* in the Brazilian Amazon. *Rev Saude Publica.* 2007;41(2):276–83.
31. Guirado MM, Elly H, Campos M De. Alguns aspectos do controle populacional e da resistência a inseticidas em *Aedes aegypti* (*Diptera* , *Culicidae*). *Inst Biociências, Let e Ciências Exatas.* 2009;6(64):5–14.
32. Brasil. *Farmacopeia Brasileira.* 6th ed. Brasília: Anvisa; 2019.
33. Crawley MJ. *The R Book.* 1st ed. Chichester: John Wiley e Sons Inc.; 2007.
34. Venables WN, Smith DM, Team RC. *An Introduction to R.* N Venables e D M Smith Copyr. 2019;2:105. Available from: <https://cran.r-project.org/doc/manuals/r-release/R-intro.pdf>
35. Albeny-Simões D, Murrell EG, Vilela EF, Juliano SA. A multifaceted trophic cascade in a detritus-based system : *Ecosphere.* 2015;6(3):1–13.

36. Lee CC, Houghton P. Cytotoxicity of plants from Malaysia and Thailand used traditionally to treat cancer. *J Ethnopharmacol.* 2005;100(3):237–43.
37. Valduga E, Freitas RJS de, Reissmann CB, Nakashima T. Caracterização química da folha de *Ilex paraguariensis* St. Hil. (Erva-Mate) e de outras espécies utilizadas na adulteração do mate. *Bol do Cent Pesqui Process Aliment.* 1997;15(1):25–36.
38. Dietrich F, Aparecida A, Strohschoen G, Schultz G, Deise A. Utilização De Inseticidas Botânicos Na Agricultura Orgânica De Arroio Do Meio / RS. *Rev Bras Agrociências.* 2011;17(2–4):251–5.
39. Jagadeesan R, Collins PJ, Nayak MK, Schlipalius DI, Ebert PR. Genetic characterization of field-evolved resistance to phosphine in the rusty grain beetle, *Cryptolestes ferrugineus* (Laemophloeidae: Coleoptera). *Pestic Biochem Physiol.* 2016;127:67–75.
40. Merritt, R. W.; Dadd, R. H.; Walker ED. Feeding Behavior, Natural Food, And Nutritional Relationships Of Larval Mosquitos. *Annu Rev Entomol.* 1992;37(1):349–76.
41. Moore JC, Berlow EL, Coleman DC, De Suiter PC, Dong Q, Hastings A, et al. Detritus, trophic dynamics and biodiversity. *Ecol Lett.* 2004;7(7):584–600.
42. Sinsabaugh RL, Linkins AE. Enzymic and chemical analysis of particulate organic matter from a boreal river. *Freshw Biol.* 1990;23(2):301–9.
43. Cochran-Stafira DL, Von Ende CN. Integrating bacteria into food webs: Studies with *Sarracenia purpurea* inquilines. *Ecology.* 1998;79(3):880–98.
44. Eisenberg JNS, Washburn JO, Schreiber SJ. Generalist feeding behaviors of *Aedes sierrensis* larvae and their effects on protozoan populations. *Ecology.* 2000;81(4):921–35.
45. Kaufman MG, Walker ED, Smith TW, Merritt RW, Klug MJ. Effects of larval mosquitoes (*Aedes triseriatus*) and stemflow on microbial community dynamics in container habitats.

- 421 Appl Environ Microbiol. 1999;65(6):2661–73.
- 422 46. Pontes RJS, Regazzi ACF, Lima JWO, Kerr-Pontes LRS. Efeito residual de apresentações
- 423 comerciais dos larvicidas temefos e *Bacillus thuringiensis israelensis* sobre larvas de
- 424 *Aedes aegypti* em recipientes com renovação de água. Rev Soc Bras Med Trop.
- 425 2005;38(4):316–21.

Figure 1

Figure 2