Transduction of mechanical cellular oscillation by the plasma-membrane mechanosensitive channel MSL10

Daniel Tran¹, Tiffanie Girault¹, Marjorie Guichard¹, Sébastien Thomine¹, Nathalie Leblanc-Fournier^{2,3}, Bruno Moulia^{2,3}, Emmanuel de Langre⁴, Jean-Marc Allain^{5,6}, Jean-Marie Frachisse¹

¹ Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud, Université Paris-Saclay, 91198 Gif sur Yvette Cedex, France

² INRA, UMR 547 PIAF, F-63100 Clermont-Ferrand, France

³ Université Clermont Auvergne, UMR 547 PIAF, F-63100 Clermont-Ferrand, France

⁴ LadHyX, Ecole Polytechnique, Palaiseau, 91128 Cedex, France

⁵ LMS, Ecole Polytechnique, CNRS, Palaiseau, France

⁶ Inria, Université Paris-Saclay, Palaiseau, France

Key words:

Mechanosensitive channel, mechanosensing, mechanotransduction, oscillation, frequency, plant, Arabidopsis, wind

***Correspondence:**

Jean-Marie Frachisse

Institute for Integrative Biology of the Cell

Saclay Plant Sciences, CNRS

Avenue de la Terrasse, 91198 Gif-sur-Yvette, France

jean-marie.frachisse@i2bc.paris-saclay.fr

Abstract

Throughout their life, plants are submitted to recurrent cyclic mechanical loading due to wind. The resulting passive oscillation movements of stem and foliage is an important phenomenon for biological and ecological issues such as photosynthesis optimization¹⁻³ and thermal exchange⁴. The induced motions at plant scale are well described and analyzed, with oscillations at typically 1 to 3 Hz in trees⁵⁻¹⁰. However, the cellular perception and transduction of such recurring mechanical signals remains an open question. Multimeric protein complexes forming mechanosensitive (MS) channels embedded in the membrane provide an efficient system to rapidly convert mechanical tension into electrical signal¹¹. Here we show that the plasma membrane mechanosensitive channel MscS-LIKE 10 (MSL10) from the model plant *Arabidopsis thaliana* responds to pulsed membrane stretching with rapid activation and relaxation kinetics in the range of one second. Under sinusoidal membrane stretching MSL10 presents a greater activity than under static stimulation and behaves as a large bandpass oscillation “follower” without filtering the signal in the range of 0.3 to 3 Hz. With a localization in aerial organs naturally submitted to oscillations, our results suggest that the mechanosensitive channel MSL10 represents a molecular component of a universal system of oscillatory perception in plants.

Main

In animals, transduction of vibrational stimulation is achieved through MS channels in organs with specialized structures, such as the ear in which the different frequencies are spatially separated¹², or by the organ motion as in touch sensation¹³. In plants, such specialized features have not yet been reported, and it remains unclear whether and how MS channels participate in the perception of oscillatory stimuli. To investigate this question, we studied MSL family members, homologues of the Mechanosensitive channel of Small conductance (MscS) from *E.*

coli^{14,15}, as they are found in all land plant genomes¹⁶. We have focused our study on MSL10, the most widely expressed, plasma membrane-localized and functionally characterized member in *Arabidopsis thaliana*^{17,18}.

To characterize the response of *Arabidopsis* to wind mechanical stimulation, we examined the frequency of free oscillations of plant with young flowering stem subjected to a short air pulse (supplementary movie 1). Using the Vibration Phenotyping System (<https://vimeo.com/213665517>)¹⁹, we determined the image correlation coefficient depicting the pendulum movement of the stem on 6 plants and obtained a mean frequency of 2.8 ± 1.0 Hz (mean $\pm$ SD, n=131) (Fig. 1a). This frequency is inside the one excited by the wind²⁰. Then, to determine whether MSL10 localization is compatible with a function as oscillation sensor, we characterized its expression pattern on plants at a flowering stage. GUS reporter driven by *MSL10* promoter was present in stem and leaf vasculature and at the root tip (Fig. 1b, c)¹⁷. This expression pattern of *MSL10*, especially at the junction between roots and shoots, which experience the major tension induced by leaves and stem motion, is an expected location for probing the motion induced by the wind (Fig. 1d).

Channel kinetic properties are crucial for its ability to perceive oscillatory stimulation at various frequencies. In order to know how fast MSL10 responds to rapid variations in membrane mechanical tension, we characterized the kinetics of this channel using the patch-clamp technique. To specifically monitor MSL10 activity in its endogenous environment, we expressed the *MSL10* gene in protoplasts from a quintuple mutant ($\Delta 5$) lacking the activity of five *MSL*-encoding genes (*msl4;msl5;msl6;msl9;msl10*). This provides a low background to record mechanically-activated currents from *MSL10*-expressing protoplasts ($\Delta 5$ +*MSL10*)¹⁷ by

Figure 1 | Oscillatory movement and *MSL10* expression pattern in aerial part of *Arabidopsis* plants. **a**, Images of the oscillatory movement of the stem induced by an air pulse of 60 ms, the correlation coefficient curve visualize the oscillating and the damping of the stem movement (red arrow: air pulse), **b-d** Blue staining represents the β -glucuronidase (GUS) activity driven by the promoter of *MSL10*. The *MSL10* promoter drives expression of the reporter gene **a**, in the root tip (indicated by arrows in **c**) and throughout vasculature of the leaves and stem, **c**, at the bottom of leaf petioles and **d**, in the root-stem junction (**d** is the same view as **c**, with petioles entirely removed). Scale: **b**, 5mm; **c** and **d**, 500 μ m..

75 applying pulses of pressure whilst monitoring transmembrane currents at a constant voltage (-
76 186mV) on a membrane excised patch in outside-out configuration (Fig. 2a). At this
77 physiologically relevant membrane potential, opening of a single stretch-activated channel
78 caused a current variation of 19.4 ± 1.7 pA (Fig. 2a, n=14) as reported in root protoplasts
79 expressing *MSL10*^{17,21}. The sustained activity under the membrane tension of *MSL10* without
80 inactivation clearly distinguishes it from the plasma membrane rapidly activated calcium MS
81 channel activity (RMA), which displays rapid inactivation^{22,23}. We observed that the activation

82 of MSL10 current increased exponentially in response to pressure, with time constants τ_{act}
 83 ranging from 1000 ms at 30 mmHg to 200 ms at 100 mmHg (Fig. 2b and Supplementary Fig.
 84 1, $n = 15$). The current-pressure relationship representing the MSL10 channel sensitivity to
 85 membrane tension was well described by a Boltzmann function with a pressure for half-

Figure 2 | Gating kinetics and pressure dependence of MSL10 in native membrane. **a**, Quintuple mutant ($\Delta 5$) stimulated by increasing pressure steps in outside-out patch configuration shows no mechanically activated current. $\Delta 5$ mutant expressing MSL10 ($\Delta 5 + MSL10$) shows currents stimulated in outside-out configuration by increasing pressure steps with slow activation kinetics. The current amplitude varies from one patch to another depending on channel density (expression of the protein). Single channel amplitude shows current transitions of 19.4 ± 1.7 pA at -186 mV ($n = 14$ protoplasts). **b**, Pressure dependence of activation time constant for MSL10 channel in excised outside-out-patch configuration (see Extended Data Fig. 1). Results are normalized with the $P_{0.5}$ of each patch and data represent mean $\pm$ S.E.M ($n = 15$ protoplasts). **c**, I_{max} normalized current–pressure relationship of stretch-activated currents in excised outside-out-patch configuration in $\Delta 5 + MSL10$, fitted with a Boltzmann equation. $P_{0.5}$ of 49.3 ± 3.4 mmHg is the average value determined for individual cells. Data represent mean $\pm$ S.E.M ($n = 15$ protoplasts).

The membrane potential is clamped at -186 mV. MSL10 protein is transiently expressed in quintuple *msl4;msl5;msl6;msl9;msl10* mutant ($\Delta 5$) protoplasts. Ionic conditions are described in the Materials and Methods.

86 activation ($P_{0.5}$) of 49.3 ± 3.4 mmHg, an activation threshold of about 30 mmHg and a saturation
 87 pressure of about 70 mmHg (Fig.2c and Supplementary Fig. 2a-c).

We then examined the effect of oscillatory membrane tension on MSL10 channel activity. To do so, MSL10 activity was recorded under oscillating pressures at a wide range of frequencies from 0.3 to 30 Hz (supplementary movie 2, Fig. 3a). Whatever the frequency tested, opening events occurred almost exclusively during the upper phase of the period ($\geq 80\%$ of cases) (Fig. 3b, Supplementary Fig. 3). At low frequency (≤ 1 Hz), at least one opening transition of the channel was triggered during each period, (Fig. 3c, 100 % of cases), at 3 Hz 70 % of the periods triggered channel opening, while at 30 Hz only 20 % of the periods were efficient (Fig 3c and Supplementary Fig. 3). Thus, the channel does not open randomly in response to oscillatory stimulations, even when some pressure pulses at 3 Hz and beyond were not efficient to trigger the channel opening (Supplementary Fig. 4).

We then undertook a comparison between static and oscillatory stimuli for a same applied “mean-pressure” using a protocol alternating the two types of stimulations (see Fig. 3d). A static stimulation held at “mean-pressure” was applied for 1 min followed by a sinusoidal pressure stimulation of ± 15 mmHg from the mean-pressure baseline at a given frequency for 1 min (Fig. 3d). This protocol was repeated to sweep frequencies from 0.3 to 30 Hz and then from 30 to 0.3 Hz, always with the same mean-pressure, in order to determine the effect of frequencies on channel activity compared to that under prior static stimulation (Fig. 3d). Figure 3e-g shows the relative effect of frequencies (ratios oscillatory/static) on channels NP(o), τ_{open} and τ_{close} on at least 5 membrane patches.

A ratio (NP(o) osc/ NP(o) stat) above 1 indicates a greater activity of the channel under sinusoidal stimulation than under static stimulation. We observed that at each frequency, the NP(o) ratio is significantly greater than one, meaning that the mean open probability is significantly higher upon dynamic than static stimulation, while the pressure applied was on average the same (Fig. 3e; red asterisks, Mann-Whitney Rank Sum Test, $p \leq 0.05$). The highest ratios are observed at low frequency (0.3, 1 and 3 Hz) corresponding to the frequencies of plant

113 oscillation measured in Figure 1a (Fig. 3e, green asterisks, Mann-Whitney Rank Sum Test,

Figure 3 | Effect of oscillatory pressure stimulation on MSL10 channel characteristics.

a, Representative recording of single channel activity of MSL10 in excised outside-out-patch configuration, in response to oscillatory pressure stimulation at 0.3 Hz, 3 Hz and 30 Hz. An oscillatory pressure of +15/-15 mmHg from a mean pressure is applied. **b**, *Left*, Idealized MSL10 current used for event analysis in response to oscillatory pressure. *Right*, Distribution of close-open transitions (at least one) elicited at upper (red bars) or lower (light grey bars) pressure as a function of frequency for 15 seconds of loading. Results represent mean $\pm$ S.E.M ($n \geq 5$ protoplasts). **c**, Probability that MSL10 channel undergoes at least one close-open transition per period as a function of frequency. Results represent mean $\pm$ S.E.M ($n \geq 5$ protoplasts). **d**, Sequences of 1 min oscillatory pressure alternating with 1 min static pressure are performed on excised outside-out-patches over time. The oscillatory stimulation (same protocol for Supplementary Fig. 6a-c) is of 30 mmHg amplitude (+15/-15 mmHg from a mean-pressure level) with a sweep of frequencies from 0.3 to 30 Hz (--), while static stimulation is at mean-pressure (--). **e-g**, Relative effect of frequency (oscillatory/static mean-pressure) on **e**, open probability $NP(o)$, **f**, open state time constant and **g**, closed state time constant. The red dashed line represents the relative ratio static/static (=1). Each point represents each biological replicate ($n \geq 5$ for a given frequency); asterisk in red (*) indicates in **e**, **f** and **g** that mean value is significantly different from 1 (Mann-Whitney Rank Sum Test, $p < 0.05$), asterisk in green (•) in **e** indicates mean $NP(o)$ is significantly different from mean $NP(o)$ obtained at 30 Hz (Mann-Whitney Rank Sum Test, $p < 0.05$).

We have determined $NP(o)_{osc}$ for each oscillatory sinusoidal pressure frequency (0.3 to 30 Hz) and $NP(o)_{stat}$ for each static stimulation prior to frequency stimulation. In 3b we present the ratio $NP(o)_{osc} / NP(o)_{stat}$ called $NP(o)$ Ratio. The same principle is applied for T_{open} ratio and T_{close} ratio in 3c and d. Other conditions same as Fig. 2.

114 $p \leq 0.05$). The asymmetry observed in $NP(o)$ distribution for decreasing and increasing
 115 frequencies (Fig. 3e and supplementary Fig.5a) likely reflect the diminution of the active
 116 channels over time of the experiment. Supplementary Figure 5 exemplifies the effect of
 117 frequencies on $NP(o)$, τ_{open} and τ_{close} obtained for a representative recording. Under oscillatory
 118 stimulation $NP(o)$ increased, τ_{open} were unchanged while τ_{close} were decreased compared to the
 119 static stimulation. In order to further quantify the opening and closing oscillation dependency
 120 of MSL10, we compared open and close time constants obtained on five patches, either under
 121 static or dynamic conditions. We measured a mean open time constant in static condition of
 122 $\tau_{open} \text{ static} = 14.7 \pm 1.9 \text{ ms}$ ($n \geq 5$). This time constant is not or weakly affected by oscillatory
 123 stimulation with a τ_{open} oscillation relative to static above 1 (Fig. 3f). The mean close time
 124 constant τ_{close} decreased significantly from $\tau_{close} \text{ static} = 164.5 \pm 24.8 \text{ ms}$ to $\tau_{close} \text{ oscillation} =$
 125 $106.4 \pm 17.6 \text{ ms}$ (all frequencies, $n \geq 5$). This is reflected by the ratios $\tau_{close} \text{ osc} / \tau_{close} \text{ stat}$ lower

than 1 (Fig. 3g), pointing to the fact that MSL10 spends less time in the closed state due to an increase in their opening probability upon oscillatory stimuli.

Mammalian Piezo1 and Piezo2 have been reported as pronounced frequency filters¹³, thus allowing transduction of repetitive mechanical stimuli at a given frequency. This was attributed to their strong inactivation. In MSL10, we didn't observe a strong inactivation, but still we observed a clear oscillation dependence in a wide range of frequencies. We tested if this may come from the channel natural kinematic of opening and closing as a function of the tension. To do so, we implemented a two-states model (see Material & Methods), which fits well our data despite its lack of explicit frequency dependency (Fig. 4a). We observe an oscillatory dependence of the response, as the model predicts a higher NP(o) ratio between oscillatory and static pressure at low frequency than at high frequency.

Figure 4 | Modelling of MSL 10 channel as a classical double state system. The channel is modelled as an 2 states (open-close) system, with rates classically changing exponentially with the applied pressure. No specific frequency dependence is introduced. **(a)** Adjustment of the model to experimental data. Model predictions (blue circles) are superimposed over data from Fig. 4e. **(b)** Prediction of the NP(o) ratio between oscillatory/static mean-pressure for different initial mean-pressure, using the same parameters. Blue: parameters obtained from the experiment (mean-pressure 7.6mmHg below $P_{0.5}$). Red: mean-pressure taken as $P_{0.5}$. Black: mean-pressure increased by 7.6mmHg with respect to $P_{0.5}$.

The low frequency response of the channel can be explained physically by the non-linear response of the channel to static pressures (the Boltzmann curve). As the mean-pressure is below $P_{0.5}$, an increase in pressure will open more channels than the same decrease in pressure.

Thus, it is expected that at low frequencies the ratio is greater than one. Choosing an initial mean-pressure exactly at $P_{0.5}$ will have symmetrized the effects of the increase and decrease in pressure leading to a ratio of 1 (see Fig. 4b). Similarly, starting from a pressure above $P_{0.5}$ will have led to a decrease of the ratio. The reliability of this model raises two questions: at which frequencies the channel is solicited in vivo and what is the mean pressure applied to the membrane in vivo? The free oscillations of Arabidopsis around 3 Hz that we have measured (Fig. 1a) are within the range of low frequencies presented in the model for which oscillatory stimulations are more efficient than static one (Fig. 4b), and are well described by the Boltzman response. For the mean pressure applied in vivo, one should expect to have two contributions: a baseline due to the turgor pressure, and one due to the mean bending during the plant oscillation, proportional to the amplitude of the plant oscillations. Thus, for a plant in rest condition, not submitted to wind loading, the MSL10 channel is expected to be in the very low domain of solicitation of the Boltzmann curve (Fig. 2c) for which the channel is closed. At low amplitudes of oscillation, corresponding to a solicitation in the domain of the Boltzmann below $P_{0.5}$, the channel will be more active under movement than during an equivalent static bending and thus will transduce oscillation into cellular ion fluxes. At high amplitude of oscillation, corresponding to a domain above $P_{0.5}$ on the Boltzmann curve, the channel will be less active during oscillation than during a static bending. This might represent a homeostatic behavior amplifying channel activity at low amplitude but decreasing channel activity at large amplitude.

At high frequency, we predict ratio larger than 1, whatever the initial mean-pressure. This effect is due to higher pressure sensitivity for opening than for closing the channel, but is harder to explain intuitively. Interestingly, the characteristic frequency, at which the channel changes from one regime to the other one, doesn't seem to depend on the mean-pressure. This effect on the channel observed for mechanical stimulation at frequency higher than 10 Hz is difficult to rely to a cellular physiological function.

Our finding also raises the question on how oscillations occurring at the scale of the plant organ could be relayed at the scale of the cell membrane. We know that a mechanical stimulation, in order to be efficient (in term of physiological response), should produce a tissue/cell deformation^{24,25}. In previous study on Arabidopsis²⁶, sinusoidal sweep excitation, mimicking wind, combined with high-speed imaging allowed us to estimate several modal frequencies and the corresponding spatial localizations of deformation. The spatial localizations of the deformation are compatible with the localizations in the plant of MSL10 as measured here (Fig. 1b). Therefore, to link membrane and organ scales we propose a qualitative model in which tissue/cell deformation induced by mechanical oscillations would induce local membrane tension able to trigger MSL10 channel. However, a full assessment of this hypothesis requires working out the full “localization/distribution/intensity” of the membrane stretching or tensions.

Conclusion

In plants, the functions of plasma membrane-located MSLs are unknown, with the exception of MSL8 which was shown to be involved in pollen hydration²⁷. This is particularly surprising for MSL10 as it is the most studied member of the MSL family. Actually, MSL10 was shown to induce cell death, but this effect was found to be separable from its mechanosensitive ion channel activity²⁸. In the present study we provide compelling evidences supporting that MSL10 acts not only as a classical transducer of sustained force but also as a transducer able to translate mechanical oscillations. With a selectivity in favor of anions^{17,18} the stretch-activated channel MSL10 is a potent actor of the plasma membrane depolarization. Thus, solicitation of MSL10 via mechanical stress delivered as sustained or even more efficiently as repetitive load to the membrane is a favorable situation to initiate electrical signaling.

This study supports that MSL10 might represent a molecular component of a system of oscillatory perception in plants. Our findings open new avenues for studying the molecular mechanisms involved in perception of oscillations that allows environmental adaptation.

Methods

Histology

Transgenic *Arabidopsis* lines used for histochemical studies and carrying the *pMSL10::GUS* promoter-reporter gene fusion were obtained previously¹⁷. In order to perform detection of β -glucuronidase (GUS) activity on whole plant, plants were grown on agar plate. In this culture condition, mature plants with flowering stem have a reduced height of ~4 cm and are suitable for staining. Tissue was fixed for 30 min in ice-cold 90% acetone, then incubated overnight at 37°C in 0.5 μ g/mL 5-bromo-4-chloro-3-indoyl β -glucuronic acid, 100mM NaPO₄ (pH 7), 0.1% Triton X-100, 5 mM potassium ferricyanide, and 10mM EDTA. Samples were then dehydrated through an ethanol series and photographed with a camera or with a Nikon AZ100 MultiZoom macroscope (objective: AZ-Plan Apo 1X NA 0.1 WD 35 mm (Nikon)).

Callus initiation and maintenance

Arabidopsis thaliana (Col-0 accession) surface-sterilized seeds were sown on “initiation medium” containing 4.3 g/L Murashige and Skoog salts (MS, Sigma-Aldrich), 2% sucrose, 10 mg/L myo-inositol, 100 μ g/L nicotinic acid, 1 mg/L thiamine-HCl, 100 μ g/L pyridoxine-HCl, 400 μ g/L glycine, 0.23 μ M kinetin, 4.5 μ M 2,4-D, 1% Phytigel (pH 5.7). For callus generation, seeds were cultured in a growth chamber for 15 days. Calli were then transferred onto “maintenance medium” containing 4.3 g/L MS salts (Sigma-Aldrich), 2% sucrose, 10 mg/L myo-inositol, 100 μ g/L nicotinic acid, 1 mg/L thiamine-HCl, 100 μ g/L pyridoxine-

HCl, 400 µg/L glycine, 0.46 µM kinetin, 2.25 µM 2,4-D, 1% phytigel, (pH 5.7), and sub-cultured every 15 days onto fresh “maintenance medium”.

Protoplast isolation and transient transformation

Calli from *Arabidopsis* were digested for 15 min at 22 °C under hyperosmotic conditions (2 mM CaCl₂, 2 mM MgCl₂, 1 mM KCl, 10 mM MESs (pH 5.5), 0.2 % cellulysin (Calbochem), 0.2 % cellulase RS (Onozuka RS, Yakult Honsha Co.), 0.004 % pectolyase Y23 (Kikkoman Corporation), 0.35 % Bovine Serum Albumin (Sigma) and mannitol to 600 mOsmol. For enzyme removal, the preparation was washed twice with 2 mM CaCl₂, 2 mM MgCl₂, 10 mM MES (pH 5.5), and mannitol to 600 mOsmol. For protoplast liberation, the preparation was incubated with 2 mM CaCl₂, 2 mM MgCl₂, 10 mM MES (pH 5.5), and mannitol to 280 mOsmol. Filtering the suspension (through a 80 µm nylon mesh) yielded protoplasts. For transient expression, protoplasts were co-transformed as described by Haswell et al. (2008)¹⁷. Silent protoplasts obtained from quintuple mutant ($\Delta 5$) *Arabidopsis* calli were co-transformed with 2.5 µg 35Sp::GFP in the p327 vector and with 10 µg 35Sp::MSL10 in the pAlligator2 vector. We only used fluorescent protoplasts, indicating a co-transformation, for patch-clamp experiments. As controls for transfection, we tested patches from $\Delta 5$ cells transfected with soluble GFP alone (n= 5) and found no mechanically activated currents in the pressure range from 0 to 60 mmHg.

Electrophysiology

Patch-clamp experiments were performed as described at room temperature with a patch-clamp amplifier (model 200A, Axon Instruments, Foster City) and a Digidata 1322A interface (Axon Instruments). Currents were filtered at 1 kHz, digitized at 4 kHz, and analyzed with pCLAMP8.1 and Clampfit 10 software. During patch-clamp recordings, the membrane potential was clamped at -186 mV and the pressure was applied with a High Speed Pressure-Clamp system²⁹ (ALA Scientific Instrument), allowing the application of precise and controlled

either pressure pulses or continuous sinusoidal variations in the pipette¹¹. Media are designed in order to eliminate stretch-activated K⁺ currents whereas the Ca²⁺ current is negligible compared to that Cl⁻. Isolated protoplasts were maintained in bathing medium: 50 mM CaCl₂, 5 mM MgCl₂, 10 mM MES-Tris, and 0,25 mM LaCl₃ (pH 5.6). Membrane seal with low resistance (< 1 GΩ) and with unstable current after excision were rejected. The pipettes were filled with 150 mM CsCl, 2 mM MgCl₂, 5 mM EGTA, 4.2 mM CaCl₂, and 10 mM Tris-HEPES (pH 7.2), supplemented with 5 mM MgATP. We adjusted the osmolarity with mannitol to 450 mosmol for the bath solution and 460 mosmol for the pipette solution using an osmometer (Type 15, Löser Meßtechnik). Gigaohm resistance seals between pipettes (pipette resistance, 0.8-1.5 MΩ), coated with Sylgard (General Electric) and pulled from capillaries (Kimax-51, Kimble Glass), and the protoplast membranes were obtained with gentle suction leading to the whole-cell configuration, and then excised to an outside-out configuration. The current-pressure relationship data were fitted to a Boltzmann function

$$I = I_r + I_m \cdot \frac{1}{1 + e^{(P_{0.5}-P)/P_c}}$$

where I_r is the background current at zero pressure, I_m is the maximum steady state current intensity, P_c is the slope of the tangent at inflexion point and $P_{0.5}$ is the pressure of half activation.

The current activation kinetics were fitted with a mono-exponential function

$$F(t) = A \cdot e^{-t/\tau} + C$$

where A is current-scale coefficient, τ is the time constant and C maximum current intensity.

Mechanosensitive channels respond to the membrane tension, which itself depends on the pipette (and patch) geometry^{30,31}. Thus, as the membrane geometry is slightly different from one patch to another (Extended Data Fig. 2a and 2d), Boltzmann functions were determined for each patch individually, prior oscillatory pressure stimulation was applied (Extended Data Fig. 2a). This allows delivering the oscillatory pressure in the same zone of membrane tension

sensitivity. The amplitude of the oscillation was +15/-15 mmHg from a mean-pressure baseline that we choose slightly below (5 to 10 mmHg) the $P_{0.5}$ (Fig. 3a).

Statistical analysis

The data were analysed using Student's t -test and analysis of variance. Comparison of NP(o) at different frequencies (Fig.3 and Extended Fig.4) were analysed with Rank Sum test. NP(o)_{osc} was determined for each oscillatory sinusoidal pressure frequency (0.3 to 30 Hz) and NP(o)_{stat} for each static stimulation prior to frequency stimulation. In Fig. 3b we present the ratio NP(o)_{osc} / NP(o)_{stat} called NP(o) Ratio. The same logic is applied for T_{open} ratio and T_{close} ratio Fig. 3c and d.

Cloning and genetics

All plasmid constructs were made with Gateway technology (Life Technologies). The *MSL10* cDNA was cloned previously into pENTR/D-TOPO¹⁷. This pENTR construct was then used in recombination reactions with pAlligator³² to create the MSL10 protein overexpression construct (*p35S::MSL10*). This construct was used for transient expression in protoplasts obtained from the quintuple mutant *msl4;msl5;msl6; msl9;msl10* ($\Delta 5 + MSL10$).

Modeling

We modeled MSL10 as a 2 states channel: an open one (O), in which the channel is activated, and a closed one (C) in which the channel is completely closed. The equilibrium between the 2 states is given by the classical chemical reaction:

with k_o and k_c the opening and closing rates, respectively. Opening (resp. closing) rate increases (resp. decreases) exponentially with the applied pressure, describing the mechano-sensitivity in the Arrhenius framework. This model doesn't contain any specific oscillatory sensitivity, but

the reaction rates are affected by the changes in pressure. We then adjusted the four constants of the model to reproduce the experimental data (Fig. 4a).

Accession numbers

MSL4: At1g53470 (SALK_142497, msl4-1)
MSL5: At3g14810 (SALK_127784, msl5-2)
MSL6: At1g78610 (SALK_06711, msl6-1)
MSL9: At5g19520 (SALK_114626, msl9-1)
MSL10: At5g12080 (SALK_076254, msl10-1)

References

1. Roden, J. S. Modeling the light interception and carbon gain of individual fluttering aspen (*Populus tremuloides* Michx) leaves. *Trees* **17**, 117–126 (2003).
2. Roden, J. S. & Pearcy, R. W. Effect of leaf flutter on the light environment of poplars. *Oecologia* **93**, 201–207 (1993).
3. Roden, J. S. & Pearcy, R. W. Photosynthetic gas exchange response of poplars to steady-state and dynamic light environments. *Oecologia* **93**, 208–214 (1993).
4. Grace, J. The turbulent boundary layer over a flapping *Populus* leaf. *Plant Cell Environ.* **1**, 35–38 (1978).
5. Tadrist, L. *et al.* Foliage motion under wind, from leaf flutter to branch buffeting. *J. R. Soc. Interface* **15**, 20180010 (2018).
6. Jackson, T. *et al.* An architectural understanding of natural sway frequencies in trees. *J. R. Soc. Interface* **16**, 20190116 (2019).

7. Rodriguez, M., Langre, E. de & Moulia, B. A scaling law for the effects of architecture and allometry on tree vibration modes suggests a biological tuning to modal compartmentalization. *Am. J. Bot.* **95**, 1523–37 (2008).
8. Gardiner, B., Berry, P. & Moulia, B. Review: Wind impacts on plant growth, mechanics and damage. *Plant Sci.* **245**, 94–118 (2016).
9. de Langre, E. Effects of wind on plants. in *Annual Review of Fluid Mechanics* vol. 40 141–168 (Annual Reviews, 2008).
10. de Langre, E. Plant vibrations at all scales: a review. *J. Exp. Bot.* **70**, 3521–3531 (2019).
11. Peyronnet, R., Tran, D., Girault, T. & Frachisse, J.-M. Mechanosensitive channels: feeling tension in a world under pressure. *Front. Plant Sci.* **5**, 558 (2014).
12. Kazmierczak, P. & Müller, U. Sensing sound: molecules that orchestrate mechanotransduction by hair cells. *Trends Neurosci.* **35**, 220–229 (2012).
13. Lewis, A. H., Cui, A. F., McDonald, M. F. & Grandl, J. Transduction of Repetitive Mechanical Stimuli by Piezo1 and Piezo2 Ion Channels. *Cell Rep.* **19**, 2572–2585 (2017).
14. Booth, I. R. *et al.* Sensing bilayer tension: bacterial mechanosensitive channels and their gating mechanisms. *Biochem. Soc. Trans.* **39**, 733–740 (2011).
15. Kung, C., Martinac, B. & Sukharev, S. Mechanosensitive Channels in Microbes. *Annu. Rev. Microbiol.* **64**, 313–329 (2010).
16. Haswell, E. S. MscS-Like Proteins in Plants. in *Current Topics in Membranes* vol. 58 329–359 (Elsevier, 2007).
17. Haswell, E. S., Peyronnet, R., Barbier-Brygoo, H., Meyerowitz, E. M. & Frachisse, J.-M. Two MscS homologs provide mechanosensitive channel activities in the Arabidopsis root. *Curr. Biol.* **18**, 730–4 (2008).

18. MaksaeV, G. & Haswell, E. S. MscS-Like10 is a stretch-activated ion channel from *Arabidopsis thaliana* with a preference for anions. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 19015–20 (2012).
19. de Langre, E. *et al.* Nondestructive and Fast Vibration Phenotyping of Plants. *Plant Phenomics* **2019**, 1–10 (2019).
20. Stull, R. B. Mean Boundary Layer Characteristics. in *An Introduction to Boundary Layer Meteorology* (ed. Stull, R. B.) 1–27 (Springer Netherlands, 1988). doi:10.1007/978-94-009-3027-8_1.
21. Peyronnet, R., Haswell, E. S., Barbier-Brygoo, H. & Frachisse, J.-M. Sensors of plasma membrane tension in *Arabidopsis* roots. *Plant Signal. Behav.* **3**, 726–729 (2008).
22. Tran, D. *et al.* A mechanosensitive Ca²⁺ channel activity is dependent on the developmental regulator DEK1. *Nat. Commun.* **8**, 1009 (2017).
23. Guerringue, Y., Thomine, S. & Frachisse, J.-M. Sensing and transducing forces in plants with MSL10 and DEK1 mechanosensors. *FEBS Lett.* **592**, 1968–1979 (2018).
24. Moulia, B., Coutand, C. & Julien, J.-L. Mechanosensitive control of plant growth: bearing the load, sensing, transducing, and responding. *Front. Plant Sci.* **6**, (2015).
25. Coutand, C., Julien, J. L., Moulia, B., Mauget, J. C. & Guitard, D. Biomechanical study of the effect of a controlled bending on tomato stem elongation: global mechanical analysis. *J. Exp. Bot.* **51**, 1813–1824 (2000).
26. Der Loughian, C. *et al.* Measuring local and global vibration modes in model plants. *Comptes Rendus Mécanique* **342**, 1–7 (2014).
27. Hamilton, E. S. *et al.* Mechanosensitive channel MSL8 regulates osmotic forces during pollen hydration and germination. *Science* **350**, 438–441 (2015).

28. Veley, K. M. *et al.* *Arabidopsis* MSL10 Has a Regulated Cell Death Signaling Activity That Is Separable from Its Mechanosensitive Ion Channel Activity. *Plant Cell* **26**, 3115–3131 (2014).
29. Besch, S. R., Suchyna, T. & Sachs, F. High-speed pressure clamp. *Pflüg. Arch. Eur. J. Physiol.* **445**, 161–6 (2002).
30. Suchyna, T. M., Markin, V. S. & Sachs, F. Biophysics and Structure of the Patch and the Gigaseal. *Biophys. J.* **97**, 738–747 (2009).
31. Lewis, A. H. & Grandl, J. Mechanical sensitivity of Piezo1 ion channels can be tuned by cellular membrane tension. *eLife* **4**, (2015).
32. Bensmihen, S. *et al.* Analysis of an activated ABI5 allele using a new selection method for transgenic *Arabidopsis* seeds. *FEBS Lett.* **561**, 127–131 (2004).

Acknowledgements

This work is supported by the grant ANR-09-BLAN-0245-03 from the Agence Nationale de la Recherche (ANR, project SENZO) and the grant ANR-11-BSV7-010-02 from the Agence Nationale de la Recherche (ANR, project CAROLS). We would like to thank Dr. Sébastien Thomine (I2BC, Gif-sur-Yvette, France) and Dr. Alexis De Angeli (I2BC, Gif-sur-Yvette, France) for their help and for insightful discussions, Dr. Elizabeth Haswell (Washington University, St. Louis, USA) for providing quintuple *msl* mutant and GUS reporter lines, Gwyneth C. Ingram for corrections on the manuscript (ENS, Lyon, France).

Author Contributions

D.T performed experiments; T.G generated MSL10 lines; M.G, N.L.F, B.M and E.dL were involved in study design; D.T, J.M.A and J.M.F designed the study; D.T, J.M.A and J.M.F

analyzed data; D.T and J.M.F wrote the paper. All authors discussed the results and commented on the manuscript.

Author information

The authors declare no competing financial interest. Correspondence and requests for materials should be addressed to J.M.F (jean-marie.frachisse@i2bc.paris-saclay.fr).

Figure legends

Figure 1 | Oscillatory movement and *MSL10* expression pattern in aerial part of *Arabidopsis* plants.

Figure 2 | Gating kinetics and pressure dependence of MSL10 in native membrane.

Figure 3 | Effect of oscillatory pressure stimulation on MSL10 channel characteristics.

Figure 4 | Modelling of MSL 10 channel as a classical double state system.

Additional information

Supplementary Figure 1 | MSL10 pressure dependence of activation time constants.

Supplementary Figure 2 | Determination of current-pressure relationship of MSL10 channel.

Supplementary Figure 3 | MSL10 channel opening occurred almost exclusively during the upper phase of the stimulation period.

Supplementary Figure 4 | Patch clamp recording of MSL10 elicited by oscillatory pressure of 0.1 Hz, 1 Hz, 3 Hz.

Supplementary Figure 5 | MSL10 channel characteristics obtained on an individual patch using a protocol alternating oscillatory and static stimulation.

Supplementary Movie 1 | Oscillatory movement of an *Arabidopsis thaliana* plant after elicitation by an air pulse. Slow motion X 16.

Supplementary Movie 2 | Activation of the mechanosensitive channel MSL10 elicited by oscillatory pressure of 0.1 Hz, 1 Hz, 3 Hz.