Cortisol-induced signatures of stress in the fish microbiome

Tamsyn M. Uren Webster¹, Deiene Rodriguez-Barreto¹, Sofia Consuegra¹, Carlos Garcia de Leaniz¹

¹Centre for Sustainable Aquatic Research, College of Science, Swansea University,
Swansea, SA2 8PP, UK

Abstract

Stress experienced in intensive aquaculture can compromise fish growth, condition and immunity. Microbiome disruption may contribute to these adverse health effects, but little is known about how stress affects fish microbial communities. Here, we specifically examined the effects of stress-induced cortisol production on the fish microbiome. We exposed juvenile Atlantic salmon to a mild confinement stressor for two weeks. We then measured cortisol in the plasma, skin-mucus and faeces, and characterised the skin and faecal microbiome. Faecal and skin cortisol concentrations increased in fish exposed to confinement stress, and were positively correlated with plasma cortisol. Elevated faecal cortisol was associated with pronounced changes in the diversity and structure of the faecal microbiome. In particular, we identified a marked decline in probiotic Lactobacillales (*Carnobacterium* sp.) and an increase in pro-inflammatory and pathogenic taxa within the classes Clostridia and Gammaproteobacteria. In contrast, skin-mucus cortisol concentrations were lower and not associated with any detectable changes in the skin microbiome. Our results demonstrate that cortisol disrupts the gut microbiome, which may, in turn, contribute to the adverse effects of stress on fish health. They also highlight the value of using non-invasive faecal samples to monitor stress, including simultaneous determination of cortisol and stress-responsive bacteria.

Keywords: Stress response, Microbiota, Glucocorticoid, *Salmo salar*, Lactic acid bacteria

33 Introduction

34 Aquaculture is the fastest growing food producing sector, and plays an increasingly
35 important role in global food security in the face of a growing human population, depletion of
36 capture fisheries and climate change (Teletchea & Fontaine 2014). However, the
37 intensification of aquaculture is often associated with an increase in stress, for example due
38 to crowding and handling, which can impact animal health and welfare, and threaten
39 aquaculture sustainability (Iwama et al. 2011). Stress response in fish includes activation of
40 the hypothalamus-pituitary-interrenal (HPI) axis, culminating in the release of glucocorticoids
41 from interrenal cells located in the head kidney (Barton 2002). As for mammals, cortisol is
42 the predominant glucocorticoid released as part of the primary stress response, and is
43 critical for mediating adaptive metabolic, physiological and behavioural adjustments
44 (Schreck & Tort 2016). However, prolonged elevation of cortisol is associated with adverse
45 health effects. In particular, stress-mediated impairment of immune function has been widely
46 described in cultured fish and has been associated with reduced disease resistance, which
47 is of critical importance for aquaculture sustainability (Ellison et al. 2018; Uren Webster et al.
48 2018b; Yada & Tort 2016).

49 Recent research has revealed the diverse influence of microbiota and their metabolites
50 on many aspects of host health and fitness, including digestion and nutrient uptake,
51 metabolism and immune development (Hooper et al. 2012; Rea et al. 2016). In mammals,
52 stress is well known to disrupt the diversity, structure and function of the microbiome which,
53 in turn, has been associated with long term health effects in the host, including metabolic
54 and immune impairment, and a range of diseases (Foster et al. 2017; Tetel et al. 2018). The
55 mechanisms by which stress impacts the microbiome are complex, and not fully understood.
56 Elevated plasma cortisol, resulting from social or psychological stress, has been associated
57 with alterations in the structure and/or diversity of the mammalian faecal or oral microbiome,
58 including the abundance of lactic acid bacteria and opportunistic pathogens (Galley et al.
59 2014; Jasarevic et al. 2015; Mudd et al. 2017). Direct administration of glucocorticoids has
60 also been shown to exert stimulatory and inhibitory effects on specific microbial taxa,
61 together with wider effects on host metabolism in some cases (Huang et al. 2015; Jentsch et
62 al. 2013; Petrosus et al. 2018; Wu et al. 2018). Additionally, while host stress response
63 influences the microbiome, the microbiome can also influence host stress response.
64 Microbiota and their metabolites are known to exert effects throughout the mammalian
65 hypothalamus-pituitary-adrenal (HPA) axis, influencing glucocorticoid synthesis, release and
66 signalling pathways (Burokas et al. 2017; de Weerth 2017; Simard et al. 2014; Vodicka et al.
67 2018).

68 Disruption of the microbiome is likely to represent an important mechanism by which
69 stress affects fish health, welfare and performance in aquaculture. There is some evidence

that environmental and social stressors disrupt microbial communities associated with the fish gut and skin (Boutin et al. 2013; Sylvain et al. 2016; Uren Webster et al. 2019; Zha et al. 2018), however, a potential role of cortisol in mediating these effects is unknown. We hypothesised that stress-induced cortisol production would directly disrupt the fish microbiome. To test this, we examined the effects of cortisol on the skin and gut microbiome of juvenile Atlantic salmon, following confinement, a mild aquaculture-relevant stressor.

Methods

Stress experiment

Prior to the start of the experiment 0+ Atlantic salmon fry (mean mass 3.92 ± 0.11 g; fork length 7.46 ± 0.07 cm), were housed in stock tanks (80 L) supplied with a constant flow of aerated, de-chlorinated tap water in a recirculation system with a temperature of $15^{\circ}\text{C} \pm 0.5^{\circ}\text{C}$ and photoperiod of 12L:12D. Water oxygen saturation was maintained above 90%, and ammonia (<0.02 mg/L), nitrite (<0.01 mg/L), nitrate (<15 mg/L) and pH (7.5 ± 0.2) were maintained within the optimal range for the species. Fish were fed with a commercial salmon feed (Skretting Nutra Parr) at a rate of 3% body weight per day.

Experimental fish were assigned at random to the control and confinement-stress treatment groups, with three replicate 20 L tanks per group, each containing 28 fish. Confinement stress consisted of slowly lowering the water volume in each tank (via draining) from 20 L to 5 L for one hour, and this was repeated every day at the same time (1100 hrs) for two weeks. All other husbandry conditions were as before. At the end of the experiment, fish were euthanised via an overdose of anaesthetic (Phenoxyethanol; 0.5 mg/L), followed by destruction of the brain according to UK Home Office regulations. The fish were measured (fork length), weighed (wet weight) and Fulton's condition factor was calculated as a measure of body condition. Blood samples were collected from the caudal vein using heparinised capillary tubes, centrifuged (5 min, $5000 \times g$) and the plasma supernatant removed and stored at -80°C prior to cortisol analysis. For each fish, a sample of skin-associated mucus for microbiome analysis was collected by swabbing the left side lateral line five times using Epicentre Catch-All™ Sample Collection Swabs (Cambio, Cambridge, UK). A sample of skin-associated mucus for cortisol analysis was collected by scraping mucus from a 2 cm^2 region of skin on the left-hand side of the fish, above the lateral line between the head and dorsal fin, using a scalpel blade. Faecal samples were collected from each fish by gently pressing along the length of the abdomen and collecting expelled faeces, which were then split evenly between samples for cortisol and microbiome analysis. All skin-mucus and faecal samples were directly frozen at -80°C prior to analysis.

Cortisol measurement

Quantification of cortisol concentration in plasma, skin mucus and faecal samples was performed using the DetectX Cortisol Enzyme Immunoassay Kit (Arbor Assays, Michigan, USA), according to the manufacturer's recommendations. Briefly, plasma samples were first pre-treated with dissociation reagent then diluted in assay buffer (1:50 final dilution) before cortisol measurement. Faecal samples were suspended in 100 µl ethanol, vortexed for 30 minutes, centrifuged (5 min, 5000 x g), then the supernatant was then diluted in assay buffer (1:20) before cortisol measurement. Skin-mucus samples were suspended in 100 µl 1M Tris-HCl, vortexed for 30 minutes and centrifuged (5 min, 5000 x g), then the supernatant was used directly in the assay without dilution.

Cortisol concentration was measured in the plasma, faeces and skin-mucus for a total of 60 individual fish (40 stressed fish and 20 controls; distributed evenly amongst replicate tanks). Each of these 180 samples was analysed in duplicate, across five 96-well plates. Cortisol concentration was calculated based on a standard curve run on each plate, and adjusted for dilution factor and initial sample volume (plasma) or weight (skin-mucus/faeces). Inter-assay variability, measured as the coefficient of variation (CV%) of four repeats across the five plates, was 4.62% and the lower limit of detection was 76.4 pg/ml. We removed one outlier of faecal cortisol (47.3 ng/g) from the stressed group (most likely resulting from an error in sample preparation) using Tukey's 1.5*IQR method, as it was 3.4x higher than the mean value (13.76 ng/g) and 1.6x higher than the next highest value in this group (29.1 ng/g).

16S rRNA amplicon sequencing

16S rRNA amplicon sequencing was performed using the faecal and skin mucus samples for the same 60 individual fish for which cortisol quantification was performed (40 stressed and 20 controls), as described previously (Uren Webster et al. 2018a). Briefly, DNA was extracted from all samples using the MoBio PowerSoil® DNA Isolation Kit (Qiagen) according to the manufacturer's instructions. Libraries were prepared amplifying the V4 hypervariable region of the bacterial 16S gene using the primers 341F and 785R, and sequenced across two lanes of an Illumina MiSeq. Raw sequence reads were quality filtered using Trimmomatic (Bolger et al. 2014), before analysis with mothur v1.39 (Kozich et al. 2013). Concatenated reads were aligned to the Silva seed reference database (version 128) (Quast et al. 2013), chimeric reads were removed using UCHIME (Edgar et al. 2011), and Bacteria and Archaea contigs were classified using the Silva reference taxonomy. Contigs were clustered into operational taxonomic units (OTUs) using mothur, based on 97% sequence similarity. Singleton OTUs were removed from the dataset then all faecal samples

were subsampled to an equal depth of 19,724 reads and all skin samples were subsampled to 10,133 reads. Measures of alpha diversity (Chao1 richness and Shannon diversity) for each sample were calculated in *mothur*.

Statistical analysis

All statistical analysis was performed in R v3.5.0. Firstly, we assessed whether non-invasive measurements of faecal and skin cortisol were indicative of cortisol in blood plasma by calculating the Pearson correlation coefficient. We then employed linear mixed effects models (LMM) using the *lme4* package to examine the effects of confinement stress and fish size on measured cortisol values in plasma, skin and faeces, using tank identity as a random factor. We also used linear mixed effects models to examine the effects of confinement stress, fish size and measured faeces/skin cortisol on faeces/skin microbial alpha diversity (Chao1 richness and Shannon diversity), including tank as a random factor. We used fish length as a covariate to control for size effects as it had a lower coefficient of variation (CV = 0.067) than fish mass (CV = 0.216). To achieve model simplification, we started with a model with all main effects and selected the model with the lowest AIC value via backward selection using the *step* and *drop1* functions and the *lmerTest* package (Kuznetsova et al. 2017). A minimal adequate model was then refitted via Restricted Maximum Likelihood, or as a linear model when the random component (tank identity) did not improve model fit compared to the fixed effects only model, as indicated by the Likelihood Ratio Test (LRT). We used the *VCA* package to estimate the amount of variability in cortisol due to confinement stress, tank effects, and differences among individual fish.

Analysis of microbial community structure (beta diversity) was performed within the *Vegan* package in R (Oksanen et al. 2017), using the Bray-Curtis dissimilarity index. Non-metric multidimensional scaling (NMDS) ordination of Bray-Curtis distances were visualised, including measured cortisol concentration as an environmental vector. Multivariate statistical analysis of microbial community separation in the faecal and skin samples was performed by PERMANOVA using *Adonis* in the *Vegan* package with confinement stress and measured faecal/skin cortisol as predictors. Statistical analysis of OTU abundance was performed using *DeSeq2* (Love et al. 2014). The effect of confinement stress and faecal/skin cortisol on relative abundance of faecal/skin OTUs was tested using a multifactorial design. Within the *DeSeq* model, low coverage OTUs were independently filtered to optimise power for identification of differentially abundant OTUs at a threshold of $\alpha=0.05$. Outlier detection and moderation of OTU level dispersion estimates were performed using default settings, and OTUs were considered significantly differentially abundant at FDR <0.05.

Results

Relation between plasma cortisol and non-invasive measures of cortisol in faeces and skin

Cortisol concentrations ranged from 2.9 to 65.8 ng/ml in blood plasma, 3.6 to 29.1 ng/g in faeces, and 0.14 to 9.45 ng/g in skin mucus across all samples. Significant positive correlations were found between plasma and faecal cortisol (Pearson's $r_{56}=0.615$, $P < 0.001$), between plasma and skin cortisol ($r_{56}=0.289$, $P = 0.028$), and between faecal and skin cortisol ($r_{57}=0.422$, $P < 0.001$; Figure 1a). Variance component analysis indicated that 82-85% of the variation in cortisol was due to variation between individuals, and 0-18% was due to variation between tanks.

Effects of confinement stress on cortisol

There was a significant increase in cortisol in the faeces and skin of stressed fish compared to unstressed controls (Welch two sample t-test; faeces, $t_{56.766} = 2.955$, $P = 0.004$; skin, $t_{52.917} = 2.819$, $P = 0.007$), but not in blood plasma (LMM stress effect, $t_{4.785} = 0.603$, $P = 0.574$; Figure 1b), which showed a significant tank effect (LRT $\chi^2 = 6.01$, $P = 0.014$). There was no association between cortisol and the length, weight or body condition of fish at the end of the experiment ($P > 0.2$ in all cases).

Effects of cortisol on microbial diversity

Faecal cortisol was negatively correlated with faecal Chao1 microbial richness (Chao1 Cortisol estimate: -79.17 ± 32.33 , $t_{1,57} = -2.449$, $P = 0.017$), but positively correlated with Shannon diversity (Shannon Cortisol estimate: 0.08 ± 0.024 , $t_{1,57} = 3.536$, $P < 0.001$; Figure 2). There was no effect of confinement stress, fish size or tank identity on faecal microbial diversity beyond that accounted by an increase in cortisol ($P > 0.4$ in all cases). For the skin, there was no significant effect of confinement stress, skin cortisol, or fish size on skin microbial Chao1 richness or Shannon diversity ($P > 0.1$ in all cases), although there were significant tank effects for skin Chao1 richness (LRT $\chi^2 = 15.53$, $P < 0.001$).

Microbial community structural diversity was performed based on the Bray-Curtis dissimilarity metric, and visualised using NMDS analysis (Figure S1). As for alpha diversity, there was a significant effect of faecal cortisol on faecal microbiome beta diversity (Cortisol: $F_{1,56} = 9.525$, $P = 0.001$), but the confinement stress had no additional effect beyond that caused by an increase in cortisol (Stress $F_{1,56} = 2.487$, $P = 0.064$). For the skin microbiome there was no detectable effect of stress or skin cortisol concentration on beta diversity (Stress: $F_{1,53} = 1.263$, $P=0.151$, Cortisol: $F_{1,53} = 0.960$, $P=0.475$).

Effects of cortisol on microbial composition

The effects of stress and cortisol concentration on OTU relative abundance was investigated using DeSeq2. For the faecal microbiome, the abundance of 44 OTUs (27 increased, 17 decreased) were significantly associated with faecal cortisol concentration, but only one OTU (*Vagococcus* sp.) was significantly elevated in the confinement stress group independently of cortisol (Figure 3; Table S1). Strikingly, of the 17 OTUs which were negatively associated with cortisol concentration, the vast majority (15) were classified as belonging to the genus *Carnobacterium* sp. in the order Lactobacillales, including the most abundant OTU overall. Of the OTUs that were positively associated with faecal cortisol concentration, 10 (37%) were from the class Gammaproteobacteria and, notably, two highly abundant OTUs from the family Clostridiaceae were also increased. In contrast, for the skin microbiome, no OTUs were significantly associated with either confinement stress or measured skin cortisol concentration.

Discussion

Our study indicates that the cortisol stress response modulates the intestinal, but not the skin, microbiome of juvenile Atlantic salmon. Although the cortisol stress response to confinement was variable amongst individuals, we identified a very distinctive relationship between cortisol and the diversity and structure of the faeces microbiome, including a clear inhibition of lactic acid-producing bacteria and the promotion of pro-inflammatory and pathogenic taxa.

Exposure to a routine, aquaculture-relevant stressor (confinement) increased the cortisol concentration in the skin and faeces of juvenile salmon compared to control fish, although there was considerable variation among individuals. This variation is consistent with the existence of low and high cortisol response fish (e.g. Pottinger & Carrick 1999; Samaras et al. 2016). We also identified a positive association between plasma and faecal cortisol, and, to a lesser extent, between plasma and skin-mucus cortisol. Plasma cortisol is typically used to measure the stress response in fish, but blood sampling is invasive and may require terminal sampling in the case of small fish (Sadoul & Geffroy 2019). Plasma cortisol concentration is also known to be influenced by acute spikes in glucocorticoid production, for example caused by handling stress prior to sampling, which may mask underlying stress levels (Bertotto et al. 2010). Our results add to that of other recent studies suggesting that faecal and skin sampling provide non-destructive alternatives to measuring plasma cortisol in Atlantic salmon (Bertotto et al. 2010; Cao et al. 2017; De Mercado et al. 2018; Lupica & Turner Jr 2009), which can also be linked directly to microbiome analysis.

Across all fish, we identified a strong association between cortisol in the faeces and both alpha and beta measures of gut microbiome diversity. Faecal cortisol was negatively associated with Chao1 richness, but positively associated with Shannon diversity,

suggesting that there were fewer, but more evenly distributed, bacterial taxa in the intestine of stressed fish. This is likely to reflect an inhibitory effect of cortisol on dominant OTUs normally present in non-stressed individuals. In particular, there was a striking decline in *Carnobacterium* sp. with increasing levels of faecal cortisol, including the most abundant OTU in non-stressed fish, together with +10 other OTUs assigned to this genus. *Carnobacterium* (order Lactobacillales, class Bacilli, phylum Firmicutes) is a genus of facultatively anaerobic, cold tolerant lactic acid bacteria comprising +12 species (Pikuta & Hoover 2014). This genus, particularly *C. pisciola*, *C. divergens* and *C. inhibens*, is commonly found in the intestinal communities of healthy fish, including Atlantic salmon (Ringø et al. 2001). *Carnobacterium* sp. are also widely used as probiotics in aquaculture, due to their beneficial effects on gut health and fish performance, and their ability to inhibit the growth of several common fish pathogens (Ringø 2008; Ringø et al. 2001).

Individuals that displayed a high cortisol response to confinement stress had a distinct faecal microbiome, that was very different from that of non-responsive fish, or from control fish that had low baseline cortisol levels. Alongside a marked decline in *Carnobacterium* sp., this structural change was characterised by a notable increase in the relative abundance of two Clostridiaceae OTUs. This family (class Clostridia, phylum Firmicutes) is commonly found in the gut of healthy mammals and fish, but also includes a number of opportunistic pathogens. An increased abundance of Clostridiaceae has been associated with microbial dysbiosis, intestinal inflammation and gastrointestinal diseases (Lopetuso et al. 2013; Muñoz Pedrego et al. 2018). Several OTUs within the class Gammaproteobacteria, including two *Yersinia* sp., *Pseudomonas* sp., *Acinetobacter* sp. and *Aeromonas* sp, were also particularly abundant in fish that had high levels of faecal cortisol. These genera include a range of opportunistic fish pathogens (Austin & Austin 2007), and tend to increase following exposure to different types of environmental stress in fish (Boutin et al. 2013; Uren Webster et al. 2019), suggesting they may represent a common signature of stress exposure. In mammals, experimental administration of cortisol results in a similar reduction in probiotic lactic acid-producing bacteria and an increase in pro-inflammatory microbiota (Huang et al. 2015; Petrosus et al. 2018; Wu et al. 2018), suggesting that these bacteria taxa could represent useful biomarkers of stress across vertebrates.

Our results demonstrate how an increase in cortisol can affect the diversity and structure of the salmon gut microbiome, which may in turn contribute to the adverse effects of stress on fish health. This is consistent with the results of previous studies that experimentally administered glucocorticoids to rats, mice and pigs (Huang et al. 2015; Petrosus et al. 2018; Wu et al. 2018). However, it is not clear exactly how cortisol may affect different taxa within complex host-associated microbial communities. Potential inhibitory mechanisms could include direct toxicity, metabolic impairment, disruption of ion-regulation, endocrine signalling

or nutrient depletion, similar to the effects of other chemical or physical stressors (Harms et al. 2016; Weber et al. 2014). At the same time, taxa more tolerant of cortisol may flourish in the absence of previous niche completion (Hibbing et al. 2010), and cortisol is also known to specifically promote the growth of certain oral pathogens *in vitro* (Jentsch et al. 2013). However, the overall relationship between stress response, cortisol and the microbiome is complex. The microbiome may be influenced by other stress hormones, by interactions amongst microbes or with the host immune system, while microbiota and/or their metabolites are also well known to influence host stress response signalling (Burokas et al. 2017; de Weerth 2017; Simard et al. 2014; Vodicka et al. 2018).

The impacts of cortisol on the microbiome are also likely to depend on the nature of the microbial community, and the stress response. In contrast to the faecal microbiome, we found no significant effects of confinement stress or skin cortisol on the diversity or structure of the skin microbiome. It is possible that skin-associated communities, which are dominated by Proteobacteria with much lower levels of Lactobacillales, are less sensitive to cortisol than faecal microbiota. On the other hand, cortisol concentrations in the skin mucus were much lower than those measured in faecal samples, which may also help explain the lack of observed effects of skin cortisol on the skin microbiome.

To conclude, our study shows, for the first time, that an increase in stress-related cortisol alters the fish intestinal microbiome, notably by decreasing the abundance of *Carnobacterium*, a Lactobacillales commonly used as a probiotic in aquaculture, and increasing pro-inflammatory and opportunistic bacterial pathogens. Given the fundamental influence of microbiota and their metabolites on many aspects of host health, this suggests that cortisol-mediated disruption of the intestinal microbiome is likely to contribute to the adverse effects of stress on immune function and disease resistance. These results have important implications for health and welfare of fish exposed to environmental stress, and more broadly, to research on stress-related diseases, such as metabolic syndrome, obesity and IBD, which have been associated with microbiome dysbiosis. Finally, our study demonstrates that both cortisol measurements and microbiome analysis can be performed simultaneously on faecal and skin samples collected non-invasively, which could represent a valuable screening tool for evaluating stress in fish.

Acknowledgements

We thank the staff at CSAR for help with fish husbandry, Nikita Berry for assistance with sample preparation and Matthew Hitchings for performing the Illumina sequencing.

Author contributions

TUW, DRB, CGL and SC designed the study; TUW and DRB performed the experiment; TUW and CGL analysed the data; TUW drafted the manuscript. All authors contributed to the final version of the manuscript.

Funding

This work was funded by a BBSRC-NERC Aquaculture grant (BB/M026469/1) to CGL and the Welsh Government and Higher Education Funding Council for Wales (HEFCW) through the Sêr Cymru National Research Network for Low Carbon Energy and Environment (NRN-LCEE) to SC.

Ethics

All experimental procedures were approved by Swansea Animal Welfare and Ethical Review Body (number IP-1415-2).

Data accessibility

All 16S rRNA sequence reads are available from the European Nucleotide Archive under accession PRJEB32276. Full metadata, and measured cortisol concentrations are available in the supporting information.

References

- Austin, B. & Austin, D. A. 2007 *Bacterial Fish Pathogens, Disease of Farmed and Wild Fish*. Chichester: Springer-Praxis.
- Barton, B. A. 2002 Stress in fishes: a diversity of responses with particular reference to changes in circulating corticosteroids. *Integrative and comparative biology* **42**, 517-525.
- Bertotto, D., Poltronieri, C., Negrato, E., Majolini, D., Radaelli, G. & Simontacchi, C. 2010 Alternative matrices for cortisol measurement in fish. *Aquaculture Research* **41**, 1261-1267.
- Bolger, A. M., Lohse, M. & Usadel, B. 2014 Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-20.

364 Boutin, S., Bernatchez, L., Audet, C. & Derôme, N. 2013 Network analysis highlights
365 complex interactions between pathogen, host and commensal microbiota. *PLoS ONE*
366 **8**, e84772.

367 Burokas, A., Arbolea, S., Moloney, R. D., Peterson, V. L., Murphy, K., Clarke, G., Stanton,
368 C., Dinan, T. G. & Cryan, J. F. 2017 Targeting the Microbiota-Gut-Brain Axis:
369 Prebiotics Have Anxiolytic and Antidepressant-like Effects and Reverse the Impact of
370 Chronic Stress in Mice. *Biological Psychiatry* **82**, 472-487.

371 Cao, Y., Tveten, A.-K. & Stene, A. 2017 Establishment of a non-invasive method for stress
372 evaluation in farmed salmon based on direct fecal corticoid metabolites
373 measurement. *Fish & Shellfish Immunology* **66**, 317-324.

374 De Mercado, E., Larrán, A. M., Pinedo, J. & Tomás-Almenar, C. 2018 Skin mucous: A new
375 approach to assess stress in rainbow trout. *Aquaculture* **484**, 90-97.

376 de Weerth, C. 2017 Do bacteria shape our development? Crosstalk between intestinal
377 microbiota and HPA axis. *Neuroscience & Biobehavioral Reviews* **83**, 458-471.

378 Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. & Knight, R. 2011 UCHIME improves
379 sensitivity and speed of chimera detection. *Bioinformatics* **27**, 2194-2200.

380 Ellison, A. R., Webster, T. M. U., Rey, O., de Leaniz, C. G., Consuegra, S., Orozco-
381 terWengel, P. & Cable, J. 2018 Transcriptomic response to parasite infection in Nile
382 tilapia (*Oreochromis niloticus*) depends on rearing density. *BMC genomics* **19**, 723.

383 Foster, J. A., Rinaman, L. & Cryan, J. F. 2017 Stress & the gut-brain axis: Regulation by the
384 microbiome. *Neurobiology of Stress* **7**, 124-136.

385 Galley, J. D., Nelson, M. C., Yu, Z., Dowd, S. E., Walter, J., Kumar, P. S., Lyte, M. & Bailey,
386 M. T. 2014 Exposure to a social stressor disrupts the community structure of the
387 colonic mucosa-associated microbiota. *BMC Microbiology* **14**, 189.

388 Harms, A., Maisonneuve, E. & Gerdes, K. 2016 Mechanisms of bacterial persistence during
389 stress and antibiotic exposure. *Science* **354**, aaf4268.

390 Hibbing, M. E., Fuqua, C., Parsek, M. R. & Peterson, S. B. 2010 Bacterial competition:
391 surviving and thriving in the microbial jungle. *Nature reviews. Microbiology* **8**, 15-25.

392 Hooper, L. V., Littman, D. R. & Macpherson, A. J. 2012 Interactions between the microbiota
393 and the immune system. *Science* **336**, 1268-1273.

394 Huang, E. Y., Inoue, T., Leone, V. A., Dalal, S., Touw, K., Wang, Y., Musch, M. W.,
395 Theriault, B., Higuchi, K., Donovan, S., Gilbert, J. & Chang, E. B. 2015 Using
396 corticosteroids to reshape the gut microbiome: implications for inflammatory bowel
397 diseases. *Inflamm Bowel Dis* **21**, 963-72.

398 Iwama, G. K., Pickering, A. & Sumpter, J. 2011 *Fish stress and health in aquaculture*:
399 Cambridge University Press.

400 Jasarevic, E., Howerton, C. L., Howard, C. D. & Bale, T. L. 2015 Alterations in the Vaginal
401 Microbiome by Maternal Stress Are Associated With Metabolic Reprogramming of
402 the Offspring Gut and Brain. *Endocrinology* **156**, 3265-76.

403 Jentsch, H. F., Marz, D. & Kruger, M. 2013 The effects of stress hormones on growth of
404 selected periodontitis related bacteria. *Anaerobe* **24**, 49-54.

405 Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D. 2013
406 Development of a dual-index sequencing strategy and curation pipeline for analyzing
407 amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ*
408 *Microbiol* **79**, 5112-20.

409 Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. 2017 lmerTest package: tests in
410 linear mixed effects models. *Journal of Statistical Software* **82**.

411 Lopetuso, L. R., Scaldaferri, F., Petito, V. & Gasbarrini, A. 2013 Commensal Clostridia:
412 leading players in the maintenance of gut homeostasis. *Gut pathogens* **5**, 23-23.

413 Love, M. I., Huber, W. & Anders, S. 2014 Moderated estimation of fold change and
414 dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550-555.

415 Lupica, S. J. & Turner Jr, J. W. 2009 Validation of enzyme-linked immunosorbent assay for
416 measurement of faecal cortisol in fish. *Aquaculture Research* **40**, 437-441.

417 Mudd, A. T., Berding, K., Wang, M., Donovan, S. M. & Dilger, R. N. 2017 Serum cortisol
418 mediates the relationship between fecal Ruminococcus and brain N-acetylaspartate
419 in the young pig. *Gut Microbes*, 1-12.

420 Muñiz Pedrego, D. A., Chen, J., Hillmann, B., Jeraldo, P., Al-Ghalith, G., Taneja, V., Davis,
421 J. M., Knights, D., Nelson, H., Faubion, W. A., Raffals, L. & Kashyap, P. C. 2018 An
422 Increased Abundance of Clostridiaceae Characterizes Arthritis in Inflammatory Bowel
423 Disease and Rheumatoid Arthritis: A Cross-sectional Study. *Inflammatory bowel*
424 *diseases*.

425 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.
426 R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. &
427 Wagner, H. 2017 vegan: Community Ecology Package: [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
428 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan).

429 Petrosus, E., Lay, D., Jr., Silva, E. B. & Eicher, S. D. 2018 Effects of orally administered
430 cortisol and norepinephrine on weanling piglet gut microbial populations and
431 Salmonella passage1. *Journal of Animal Science* **96**, 4543-4551.

432 Pikuta, E. V. & Hoover, R. B. 2014 The genus Carnobacterium. In *Lactic Acid Bacteria:*
433 *Biodiversity and Taxonomy* (ed. W. H. Holzapfel & B. J. Wood): John Wiley & Sons,
434 Ltd.

435 Pottinger, T. G. & Carrick, T. R. 1999 A comparison of plasma glucose and plasma cortisol
436 as selection markers for high and low stress-responsiveness in female rainbow trout.
437 *Aquaculture* **175**, 351-363.

438 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. &
439 Glöckner, F. O. 2013 The SILVA ribosomal RNA gene database project: improved
440 data processing and web-based tools. *Nucleic Acids Research* **41**, D590-D596.

441 Rea, K., Dinan, T. G. & Cryan, J. F. 2016 The microbiome: a key regulator of stress and
442 neuroinflammation. *Neurobiology of stress* **4**, 23-33.

443 Ringø, E. 2008 The ability of carnobacteria isolated from fish intestine to inhibit growth of fish
444 pathogenic bacteria: a screening study. *Aquaculture Research* **39**, 171-180.

445 Ringø, E., Wesmajervi, M. S., Bendiksen, H. R., Berg, A., Olsen, R. E., Johnsen, T.,
446 Mikkelsen, H., Seppola, M., Strøm, E. & Holzapfel, W. 2001 Identification and
447 Characterization of Carnobacteria Isolated from Fish Intestine. *Systematic and*
448 *Applied Microbiology* **24**, 183-191.

449 Sadoul, B. & Geffroy, B. 2019 Measuring cortisol, the major stress hormone in fishes.
450 *Journal of Fish Biology*.

451 Samaras, A., Dimitroglou, A., Sarropoulou, E., Papaharisis, L., Kottaras, L. & Pavlidis, M.
452 2016 Repeatability of cortisol stress response in the European sea bass
453 (*Dicentrarchus labrax*) and transcription differences between individuals with
454 divergent responses. *Scientific Reports* **6**, 34858.

455 Schreck, C. B. & Tort, L. 2016 1 - The Concept of Stress in Fish. In *Fish Physiology*, vol. 35
456 (ed. C. B. Schreck, L. Tort, A. P. Farrell & C. J. Brauner), pp. 1-34: Academic Press.

457 Simard, M., Hill, L. A., Underhill, C. M., Keller, B. O., Villanueva, I., Hancock, R. E. &
458 Hammond, G. L. 2014 *Pseudomonas aeruginosa* elastase disrupts the cortisol-
459 binding activity of corticosteroid-binding globulin. *Endocrinology* **155**, 2900-8.

460 Sylvain, F.-É., Cheaib, B., Llewellyn, M., Correia, T. G., Fagundes, D. B., Val, A. L. &
461 Derome, N. 2016 pH drop impacts differentially skin and gut microbiota of the
462 Amazonian fish tambaqui (*Colossoma macropomum*). *Scientific Reports* **6**, 32032.

463 Teletchea, F. & Fontaine, P. 2014 Levels of domestication in fish: implications for the
464 sustainable future of aquaculture. *Fish and fisheries* **15**, 181-195.

465 Tetel, M. J., de Vries, G. J., Melcangi, R. C., Panzica, G. & O'Mahony, S. M. 2018 Steroids,
466 stress and the gut microbiome-brain axis. *Journal of Neuroendocrinology* **30**, e12548.

467 Uren Webster, T., Consuegra, S. & Garcia de Leaniz, C. 2019 Early life stress causes
468 lasting impacts on the microbiome of Atlantic salmon. *BioRxiv*.

469 Uren Webster, T. M., Consuegra, S., Hitchings, M. & Garcia de Leaniz, C. 2018a Inter-
470 population variation in the Atlantic salmon microbiome reflects environmental and
471 genetic diversity. *Appl Environ Microbiol* **84**, e0061-18.

- Uren Webster, T. M., Rodriguez-Barreto, D., Martin, S. A. M., Van Oosterhout, C., Orozco-terWengel, P., Cable, J., Hamilton, A., Garcia De Leaniz, C. & Consuegra, S. 2018b Contrasting effects of acute and chronic stress on the transcriptome, epigenome, and immune response of Atlantic salmon. *Epigenetics* **13**, 1191-1207.
- Vodicka, M., Ergang, P., Hrnčir, T., Mikulecka, A., Kvapilova, P., Vagnerova, K., Sestakova, B., Fajstova, A., Hermanova, P., Hudcovic, T., Kozakova, H. & Pacha, J. 2018 Microbiota affects the expression of genes involved in HPA axis regulation and local metabolism of glucocorticoids in chronic psychosocial stress. *Brain Behav Immun* **73**, 615-624.
- Weber, M. F., Poxleitner, G., Hebisch, E., Frey, E. & Opitz, M. 2014 Chemical warfare and survival strategies in bacterial range expansions. *Journal of The Royal Society Interface* **11**, 20140172.
- Wu, T., Yang, L., Jiang, J., Ni, Y., Zhu, J., Zheng, X., Wang, Q., Lu, X. & Fu, Z. 2018 Chronic glucocorticoid treatment induced circadian clock disorder leads to lipid metabolism and gut microbiota alterations in rats. *Life sciences* **192**, 173-182.
- Yada, T. & Tort, L. 2016 Stress and disease resistance: immune system and immunoendocrine interactions. In *Fish Physiology*, vol. 35, pp. 365-403: Elsevier.
- Zha, Y., Eiler, A., Johansson, F. & Svanbäck, R. 2018 Effects of predation stress and food ration on perch gut microbiota. *Microbiome* **6**, 28.

Figure Legends

Figure 1. A) Relation between measured cortisol in the plasma, skin and faeces across individual fish, and B) measured cortisol in the plasma, faeces and skin of Atlantic salmon exposed to confinement stress compared to control fish.

Figure 2. Relationship between measured cortisol and microbial alpha diversity (Chao1 richness and Shannon diversity) in the faeces.

Figure 3. Genus-level composition of the faecal microbiome, and measured faecal cortisol concentrations in individual fish. (C: control, S: confinement stress).

Figure 1. A) Relation between measured cortisol in the plasma, skin and faeces across individual fish, and **B)** measured cortisol in the plasma, faeces and skin of Atlantic salmon exposed to confinement stress compared to control fish.

Figure 2. Relationship between measured cortisol and microbial alpha diversity (Chao1 richness and Shannon diversity) in the faeces.

Figure 3. Genus-level composition of the faecal microbiome, and measured faecal cortisol concentrations in individual fish (C: control, S: confinement stress).