Rainer Waadt et al. – Coordination of signaling compound fluxes in roots

AUTHOR NAMES AND AFFILIATIONS

Rainer Waadt,^{a,1} Philipp Köster,^{b,d} Zaida Andrés,^a Christian Waadt,^c Gabriele Bradamante,^{a,e} Konstantinos Lampou,^{a,f} Jörg Kudla,^b and Karin Schumacher^a

^a Entwicklungsbiologie der Pflanzen, Centre for Organismal Studies, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany.

^b Molekulare Genetik und Zellbiologie der Pflanzen, Institut für Biologie und Biotechnologie der Pflanzen, Westfälische Wilhelms-Universität Münster, Münster Germany.

^c Not affiliated to a Research Institution.

^d Present address: Molecular and Cellular Plant Physiology, Department of Plant and Microbial Biology, University of Zurich, Zurich, Switzerland.

^e Present address: Epigenetic Changes in Plants, Gregor Mendel Institute of Molecular Plant Biology, Vienna, Austria.

^f Present address: Entwicklungsbiologie, Institut für Pflanzenwissenschaften und Mikrobiologie, Universität Hamburg, Germany.

¹ Address correspondence to rainer.waadt@gmail.com.

The authors responsible for distribution of materials integral to the findings presented in this article in accordance with policy described in the Instructions for Authors (www.plantcell.org) are: Rainer Waadt (rainer.waadt@gmail.com) and Karin Schumacher (karin.schumacher@cos.uni-heidelberg.de).

TITLE

Dual-sensing genetically encoded fluorescent indicators resolve the spatiotemporal coordination of cytosolic abscisic acid and second messenger dynamics in Arabidopsis

SHORT TITLE

Coordination of signaling compound fluxes in roots

ABSTRACT

Deciphering signal transduction processes is crucial for understanding how plants sense and respond to environmental changes. Various chemical compounds function as central messengers within deeply intertwined signaling networks. How such compounds act in concert remains to be elucidated. We have developed dual-sensing genetically encoded fluorescent indicators (2-In-1-GEFIs) for multiparametric in vivo analyses of the phytohormone abscisic acid (ABA), Ca^{2+} , protons (H^+), chloride (anions), the glutathione redox potential (E_{GSH}) and hydrogen peroxide (H_2O_2). Simultaneous analyses of two signaling compounds in *Arabidopsis* (*Arabidopsis thaliana*) roots revealed that ABA treatment and uptake did not trigger rapid cytosolic Ca^{2+} or H^+ fluxes. Glutamate, ATP, Arabidopsis PLANT ELICITOR PEPTIDE (AtPEP1) and glutathione disulfide (GSSG) treatments induced rapid spatiotemporally overlapping cytosolic Ca^{2+} , H^+ and anion fluxes, but except for GSSG only weakly affected the cytosolic redox state. Overall, 2-In-1-GEFIs enable complementary high-resolution in vivo analyses of signaling compound dynamics and facilitate an advanced understanding of the spatiotemporal coordination of signal transduction processes in *Arabidopsis*.

INTRODUCTION

Understanding how plants sense and respond to environmental and extracellular fluctuations is key for our strategic progressions to limit the consequences of climate change on plant growth and crop productivity. Plants have evolved complex signal transduction networks that enable the sensing and integration of extracellular signals, and the processing and transduction of the underlying information into physiological-, growth- and developmental responses. Within such signaling networks, spatiotemporal concentration changes of hormones, the divalent cation Ca^{2+} and reactive oxygen

species (ROS) mediate various downstream responses (Dodd et al., 2010; Kudla et al., 2010; Shan et al., 2012; Vanstraelen and Benková, 2012; Mittler et al., 2017; Waszczak et al., 2018; Smirnov and Arnaud, 2019). Among the plant hormones, abscisic acid (ABA) functions as a central regulator of the plant water status (Cutler et al., 2010; Finkelstein et al., 2013; Yoshida et al., 2019). Dynamic concentration changes of signaling compounds require inter- and intracellular transport, including long-distance transport and signaling, that often depend on proton (H^+) and electrochemical gradients across membranes (Schumacher, 2014; Choi et al., 2016; Sze and Chanroj, 2018). In addition, environmental and cellular H^+ concentration (pH) can affect plant growth, development and molecular properties (Shavrukov and Hirai, 2016). Therefore, H^+ may also function in signaling (Sze and Chanroj, 2018).

In plants, hormonal, Ca^{2+} , ROS and pH signaling processes are intertwined on several levels (Hauser et al., 2011; Vanstraelen and Benková, 2012; Gilroy et al., 2014; Steinhorst and Kudla, 2014; Edel and Kudla, 2016), for example to regulate stomatal movements or root hair and pollen tube growth (Munemasa et al., 2015; Mangano et al., 2016; Hauser et al., 2017; Michard et al., 2017). In response to the growth hormone auxin, extracellular ATP, touch or wounding, Ca^{2+} signals are accompanied by an apoplastic alkalization and/or cytosolic acidification (Monshausen et al., 2009, 2011; Behera et al., 2018). However, except for the auxin response (Shih et al., 2015; Dindas et al., 2018), the underlying mechanisms are not well understood. Current models suggest that cytosolic Ca^{2+} and extracellular ROS signals are important for cell-to-cell communication and long-distance signaling (Gilroy et al., 2014; Steinhorst and Kudla, 2014). Although some components, such as the ROS producing Arabidopsis (*Arabidopsis thaliana*) RESPIRATORY BURST OXIDASE HOMOLOG D (AtRBOHD), the ion channel TWO PORE CHANNEL1 (TPC1), GLUTAMATE RECEPTOR-LIKE CHANNELS (GLRs), Ca^{2+} -DEPENDENT PROTEIN KINASES (CDPKs/CPKs) and CALCINEURIN B-LIKE (CBL) proteins together with CBL-INTERACTING PROTEIN KINASES (CIPKs) have been linked to such processes (Choi et al., 2016; Waszczak et al., 2018), the underlying mechanisms remain unclear.

In order to decipher the coordination and interdependence of signaling processes, it is important to monitor the spatiotemporal dynamics of signaling

compounds. Genetically encoded fluorescent indicators (GEFIs) are currently the state-of-the-art technology for high resolution in vivo monitoring of biological processes (Grossmann et al., 2018; Hilleary et al., 2018; Walia et al., 2018). Although the number of GEFIs is steadily increasing, only a fraction has been introduced into plants, and less have been used in simultaneous multiparametric analyses (Okumoto et al., 2012; Kostyuk et al., 2018; Walia et al., 2018). To facilitate the use of GEFIs in multiparametric analyses, we introduce here the concept of dual-sensing genetically encoded fluorescent indicators (2-In-1-GEFIs) that enable the in vivo monitoring of at least two signaling compounds simultaneously. Through the genetic fusion of two distinct indicators, seven 2-In-1-GEFIs were generated that enable time-efficient and complementary in vivo analyses of ABA, Ca^{2+} , H^+ , Cl^- , H_2O_2 and the glutathione redox potential (E_{GSH}) at unprecedented spatiotemporal resolution. Microscopic analyses of these 2-In-1-GEFIs in Arabidopsis roots confirmed their functionality and revealed that extracellularly applied ABA was rapidly taken up, but without discernible effect on cytosolic Ca^{2+} and pH levels. In contrast, treatments with glutamate, ATP, AtPEP1 and GSSG induced spatiotemporally overlapping fluxes of Ca^{2+} , H^+ and Cl^- , without noticeable rapid effect on the cytosolic redox state.

RESULTS

Approaches to optimize FRET-based ABA indicators

Currently available ABA indicators are ABACUS and ABAleon (Jones et al., 2014; Waadt et al., 2014). Because their expression in plants has an impact on ABA signaling and because they exhibit a relatively small signal-to-noise ratio (Waadt et al., 2015), we aimed to optimize these indicators before utilizing them for multiparametric analyses. As testing system, we chose human embryonic kidney (HEK293T) cells that allow for efficient medium throughput plate reader-based screens. Compared to ABAleon2.15 and ABACUS1-2 μ used as positive control and non-responsive ABAleon2.15nr as negative control, the initial screening aimed to evaluate deletion variants of ABAleon2.15 and various combinations of five fluorescent protein Förster

Resonance Energy Transfer (FRET)-pairs and three sensory domains (SDs; Figure 1A, Supplemental Figures 1A to 1D). 60 min after application of 0 or 100 μ M ABA, ABAleon2.15 deletion variants (d1-d3), as well as ABACUS1-2 μ and SD2 variants exhibited emission ratio changes even in control conditions. When compared to ABAleon2.15, two indicators (PmTurquoise-SD1-Venus and PmTurquoise-SD1-cpVenus173) exhibited neglectable responses to control treatments, but increased negative emission ratio changes in response to ABA (Figure 1A). Because PmTurquoise-SD1-cpVenus173 (ABALeonSD1-3) differed from ABAleon2.15 only in the sequences that link the sensory domain with the attached fluorescent proteins (Supplemental Figures 1A and 1D), an additional ABALeonSD1-3 linker screening was performed (Figure 1B). This led to the identification of ABALeonSD1-3L21 (linkers LD and T; Supplemental Figure 1E) with similar properties as ABALeonSD1-3 (Figure 1B).

To corroborate these findings, ABAleon2.15, ABALeonSD1-3 and ABALeonSD1-3L21 proteins were purified from *Escherichia coli* (*E. coli*) and characterized in vitro. Although all three ABALeons were functional (Figures 1C to 1G), their properties in vitro were markedly different from results obtained in HEK293T cells (Figures 1G to 1I). Basal emission ratios (at 0 μ M ABA) were not the same in both assay systems (Figures 1G to 1H). ABAleon2.15 exhibited a higher ABA-induced emission ratio change in vitro, while ABALeonSD1-3 responses were larger in the HEK293T cell system (Figure 1I). ABALeonSD1-3L21 responded similar in vitro and in HEK293T cells (Figure 1I). In vitro calibrations revealed that ABALeonSD1-3 responded much weaker to ABA (maximum emission ratio change $\Delta R_{(\max)}/R_0 = -0.141$) compared to ABAleon2.15 and ABALeonSD1-3L21 ($\Delta R_{(\max)}/R_0 = -0.185$ and -0.167) (Figures 1C to 1E). Apparent ABA affinities of ABALeonSD1-3 (954 nM) and ABALeonSD1-3L21 (938 nM) were slightly lower compared to ABAleon2.15 (817 nM) (Figure 1F). Altogether, the ABA indicator screening led to the identification of ABALeonSD1-3L21. Its advanced properties in HEK293T cells requires therefore further validation in planta.

Evaluation of FRET-based ABA indicators in Arabidopsis

5-day-old Arabidopsis seedlings expressing ABAleon2.15, ABALeonSD1-3L21 or ABACUS1-2 μ in the cytosol and nucleus were compared for their ABA responses in

roots. Therefore, spatiotemporal vertical response profiles of emission ratios (R; Figure 2 top) or emission ratio changes normalized to 4 min average baseline recordings ($\Delta R/R_0$; Figure 2 middle) and overall emission ratio changes (Figure 2 bottom) were acquired in response to 10 μ M ABA treatments. ABAleon2.15 and ABAleonSD1-3L21 responded similar to ABA with a sigmoidal emission ratio decrease at a half response time of $t_{1/2} \sim 15$ min (Figures 2A and 2B). Note that the calyptra exhibited a much weaker response to ABA compared to the other tissues, likely because of the high cytosolic ABA concentration ($[ABA]_{\text{cyt}}$) there (depicted in dark blue; Figures 2A top and 2B top). The root elongation zone however, exhibited the lowest $[ABA]_{\text{cyt}}$ (depicted in white). ABACUS1-2 μ did not resolve this ABA concentration gradient, and responded slower ($t_{1/2} \sim 29$ min), but with larger increasing emission ratio changes that were more pronounced in the meristematic- and early elongation zone (Figure 2C, Supplemental Movie 1). None of the indicators exhibited emission ratio changes in response to control treatments (Supplemental Figure 2). Altogether, we conclude that ABAleons are more suitable for ABA analyses in tissues with low $[ABA]_{\text{cyt}}$, whereas ABACUS1-2 μ should be preferably used in tissues with high $[ABA]_{\text{cyt}}$. Because both ABAleons exhibited similar ABA response patterns, we decided to employ the latest version ABAleonSD1-3L21 for our multiparameter imaging approach.

Concept and design of dual-sensing genetically encoded fluorescent indicators (2-In-1-GEFIs)

Multiparametric analyses of signaling compounds requires the generation of transgenic plants that express several GEFIs simultaneously. Because the generation of transgenic plants is time-consuming and the insertion of multiple transgenes into the Arabidopsis genome often results in epigenetic silencing effects, we aimed to express two GEFIs from one mRNA. Therefore, GEFIs were genetically fused via sequences encoding for a 14 amino acid ASGGSGGTSGGGGS-linker (GSL), or the self-cleaving 22 amino acid P2A linker that enables the expression of two separate polypeptides (Kim et al., 2011). Six GEFIs: ABAleonSD1-3L21 (ABA), R-GECO1 (Ca^{2+} ; Zhao et al., 2011), Arabidopsis codon optimized red-fluorescing (P)A-17 (H^+ ; Shen et al., 2014), E^2GFP (H^+ and Cl^- ; Bizzarri et al., 2006), Grx1-roGFP2 (E_{GSH} ; Gutscher et al., 2008) and roGFP2-

Orp1 (H_2O_2 ; Gutscher et al., 2009) were combined for the generation of seven 2-In-1-GEFIs: ABAleonSD1-3L21-P2A-R-GECO1 (ABA and Ca^{2+}), PA-17-P2A-ABALEONSD1-3L21 (H^+ and ABA), R-GECO1-GSL- E^2GFP (Ca^{2+} , H^+ and Cl^-), PA-17-P2A-Grx1-roGFP2 (H^+ and E_{GSH}), PA-17-P2A-roGFP2-Orp1 (H^+ and H_2O_2), R-GECO1-P2A-Grx1-roGFP2 (Ca^{2+} and E_{GSH}) and R-GECO1-P2A-roGFP2-Orp1 (Ca^{2+} and H_2O_2). See Supplemental Data Sets 1B and 1C for information about constructs and transgenic Arabidopsis plants. In the following, we will describe the application of these 2-In-1-GEFIs in Arabidopsis and highlight the resulting biological findings.

ABA does not trigger rapid Ca^{2+} or pH changes in roots

To test the functionality of the 2-In-1-GEFIs, we first studied the interrelation of ABA with Ca^{2+} in roots. Therefore, ABAleonSD1-3L21-P2A-R-GECO1 seedlings were monitored in response to 10 μM ABA, which induced a typical ABAleonSD1-3L21 emission ratio decrease (Figure 3A left). The Ca^{2+} indicator R-GECO1 did not respond to this treatment (Figure 3A right). However, subsequent 1 μM indole-3-acetic-acid (IAA; auxin) treatment at 30 min induced a biphasic Ca^{2+} signal that initiated in the root elongation zone and spread from there to neighboring regions (Figure 3A right, Supplemental Movie 2), as observed before (Waadt et al., 2017).

To investigate the effect of ABA on cytosolic pH, PA-17-P2A-ABALEONSD1-3L21 seedlings were treated with 10 μM ABA. From 0-30 min after ABA treatment PA-17 fluorescence remained unchanged (Figure 3B left). However, in response to 1 mM ATP PA-17 fluorescence emission decreased, indicating a rapid and transient cytosolic acidification with a maximum pH drop in the root meristematic zone that also spread to the elongation zone (Figure 3B left). In this experiment, ABAleonSD1-3L21 exhibited a typical ABA response pattern that was not affected by ATP (Figure 3B right, Supplemental Movie 3). These experiments established the functionality of both 2-In-1-GEFIs and revealed that ABA does not trigger rapid cytosolic Ca^{2+} or pH changes in roots.

Auxin, ATP and glutamate treatments trigger spatiotemporally overlapping fluxes of Ca^{2+} , H^+ and Cl^-

Next, we used R-GECO1-GSL-E²GFP to simultaneously monitor Ca²⁺, H⁺ and Cl⁻ fluxes. Because anions, such as Cl⁻, quench the fluorescence of E²GFP and because its excitation ratiometric pH readout is Cl⁻ independent, E²GFP provides a means to simultaneously assess cytosolic H⁺ and Cl⁻ changes (Bizzarri et al., 2006; Arosio et al., 2010). In response to 1 μM IAA, R-GECO1 reported biphasic Ca²⁺ signals in the root elongation zone that travelled to neighboring regions. Subsequent 1 mM ATP treatments triggered Ca²⁺ signals in the calyptra and meristematic zone that proceeded shootward (Figure 4A left). Interestingly, both Ca²⁺ signals coincided with a cytosolic acidification reported by E²GFP (Figure 4A middle, Supplemental Movie 4). IAA also induced a Cl⁻ influx, indicated by a E²GFP fluorescence emission decrease in the entire imaged root, with a maximum decrease in the meristematic zone (Figure 4A right). ATP however induced Cl⁻ influx in the upper elongation zone and above, but Cl⁻ efflux in the lower elongation- and meristematic zone (Figure 4A right). Correlation analyses of the initial 15 min during the IAA response indicated a remarkable spatiotemporal overlap of Ca²⁺, H⁺ and Cl⁻ influx in the meristematic and elongation zone (Figure 4B).

In additional experiments, the effect of 1 mM glutamate was assessed. Compared to IAA, glutamate treatments triggered a more expanded and rapid Ca²⁺ transient that arrived in a wave-like shape from upper root regions (Figure 5A left, Supplemental Movie 5). H⁺ and Cl⁻ also displayed a rapid and transient initial influx with a maximum acidification in the meristematic zone, followed by a weak transient alkalization in the early elongation zone and a prolonged H⁺ and Cl⁻ influx (Figure 5A left, Supplemental Movie 5). During the initial 10 min of the glutamate response, Ca²⁺ and H⁺ influx exhibited a noticeable spatiotemporal overlap/correlation in the meristematic- and early elongation zone (Figure 5B left). Subsequent responses to ATP, used as positive control, were as observed before (Figures 4A and 5A). Correlation analyses indicated a remarkable coincidence of Ca²⁺ and H⁺ influx and Cl⁻ efflux in the meristematic zone (Figure 5B right).

In order to increase the spatial resolution for pH measurements, R-GECO1-P2A-E²GFP was fused to the N-terminus of LOW TEMPERATURE INDUCED PROTEIN 6B (LTI6b) or VESICLE TRANSPORT V-SNARE 11 (VTI11). This enabled the targeting of E²GFP to the cytosolic side of the plasma membrane (LTI6b; Cutler et al., 2000) or the

tonoplast (VTI11; Takemoto et al., 2018), while R-GECO1 remained in the cytosol and the nucleus (Supplemental Figures 3A and 4A). Compared to previous analyses (Figure 5), these indicators reported very similar Ca^{2+} and pH response patterns, irrespective of the subcellular localization of E^2GFP . However, it appeared that Cl^- responses were more variable with noticeably higher Cl^- fluxes at the tonoplast (Supplemental Figures 3B and 4B). Note that R-GECO1-P2A- E^2GFP -LTI6b expression induced more severe growth defects in Arabidopsis compared to the other GEFI lines (Supplemental Figure 5). Therefore, results obtained with this GEFI should be interpreted with caution. Taken together, R-GECO1-GSL- E^2GFP enables the simultaneous monitoring of Ca^{2+} , H^+ and Cl^- fluxes, that in response to IAA, ATP and glutamate exhibited a remarkably high spatiotemporal overlap.

Glutamate treatment induces cytosolic acidification without noticeable impact on the cytosolic redox state

To test whether glutamate has an impact on the cytosolic redox state, Arabidopsis seedlings expressing PA-17-P2A-Grx1-roGFP2 (pH and E_{GSH}) or PA-17-P2A-roGFP2-Orp1 (pH and H_2O_2) were exposed to 1 mM glutamate and 100 μM H_2O_2 treatments as positive control. As observed before, glutamate triggered a biphasic cytosolic acidification, that prolonged during the 100 μM H_2O_2 response (Figures 6A left and 6B left, Supplemental Movies 6 and 7). Application of glutamate did not induce cytosolic redox changes. Whereas, 100 μM H_2O_2 treatments triggered a steep cytosolic oxidation that remained high for longer than 30 min (Figures 6A right and 6B right). During this response, the roGFPs indicated a cytosolic oxidation predominantly in epidermis and cortex cells of the elongation zone and above, with faster responses in upper root regions. Except of the epidermis, cells of the meristematic zone only slightly increased their redox state in response to H_2O_2 (Supplemental Movies 6 and 7). Altogether, these data indicate that 100 μM H_2O_2 treatments rapidly induce cytosolic oxidation, and that the root meristematic zone is less sensitive to this oxidative stress.

H_2O_2 treatment triggers spatiotemporally overlapping but also distinct patterns of cytosolic oxidation and Ca^{2+} fluxes

Current models propose an interdependence of Ca^{2+} - and ROS signaling (Gilroy et al., 2014; Steinhorst and Kudla, 2014). To investigate the spatiotemporal relationships of cytosolic Ca^{2+} and ROS signals, we first treated Arabidopsis seedlings expressing R-GECO1-P2A-Grx1-roGFP2 (Ca^{2+} and E_{GSH}) or R-GECO1-P2A-roGFP2-Orp1 (Ca^{2+} and H_2O_2) with 20 and 100 μM H_2O_2 . The E_{GSH} and H_2O_2 indicators responded to both treatments with similar patterns, albeit with increased signal changes in response to 100 μM H_2O_2 (Figures 7A right and 7B right, Supplemental Movies 8 and 9). Although 20 μM H_2O_2 treatments induced a detectable cytosolic oxidation, discernible Ca^{2+} signals were not observed (Figure 7, Supplemental Movies 8 and 9). In response to 100 μM H_2O_2 treatments, cytosolic oxidation preceded detectable Ca^{2+} signals. Although, both signals appeared to arrive from upper root regions, Ca^{2+} signals in the elongation zone exhibited a maximum response in the vasculature, whereas cytosolic oxidation was more pronounced in epidermis and cortex cells. Both signals exhibited a minimum response in the meristematic zone (Figure 7, Supplemental Movies 8 and 9). We conclude that these 2-In-1-GEFIs exhibit sufficient sensitivity for resolving the interrelation of cytosolic Ca^{2+} and ROS signals, which, in response to 100 μM H_2O_2 treatments, exhibit overlapping but not similar spatiotemporal response patterns. In addition, the roGFPs facilitate the detection of cytosolic oxidation in response to H_2O_2 below the threshold of Ca^{2+} channel activation.

ATP and AtPEP1 treatments trigger Ca^{2+} , H^+ and Cl^- fluxes, and a weak cytosolic oxidation

Extracellular ATP and the signaling peptide AtPEP1 function as damage-associated elicitors that trigger Ca^{2+} signals and ROS production (Song et al., 2006; Demidchik et al., 2009; Ma et al., 2014). To investigate the spatiotemporal relationships of these processes, Arabidopsis seedlings expressing R-GECO1 and roGFP2-Orp1 or Grx1-roGFP2 from individual expression cassettes located on one T-DNA, were subjected to 1 mM ATP and consecutive 100 μM H_2O_2 treatments as positive control. ATP triggered typical Ca^{2+} responses, but its effect on the cytosolic redox state was rather weak (Figure 8A, Supplemental Figure 6A). 100 μM H_2O_2 treatments induced Ca^{2+} fluxes and cytosolic oxidation as observed before (Supplemental Movies 10 and

12). Experiments using R-GECO1-P2A-roGFP2-Orp1 and R-GECO1-P2A-Grx1-roGFP2 revealed that 1 μ M (At)PEP1 treatments induced Ca^{2+} signals that initiated in epidermis cells, followed by an overall Ca^{2+} burst, after which Ca^{2+} oscillations in the meristem appeared that proceeded to the vasculature and further shootward. However, PEP1 treatments had only weak effects on the cytosolic redox state (Figure 8B, Supplemental Figure 6B, Supplemental Movies 11 and 13). To better resolve the roGFP responses, the initial 30 min signal change response profiles were extracted from original data sets and calibrated to the same color scale. The data indicate a detectable cytosolic oxidation in response to glutamate, ATP and PEP1 that was however low compared to the 20 μ M H_2O_2 response (Supplemental Figure 7). We also investigated the effect of PEP1 using R-GECO1-GSL- E^2GFP . 1 μ M PEP1 triggered a transient Ca^{2+} , H^+ and Cl^- influx that, during the initial 20 min of the response, exhibited a spatiotemporal overlap/correlation mainly in the meristematic- and elongation zone (Figures 9A and 9B, Supplemental Movie 14). Altogether, these experiments established that PEP1 triggers spatiotemporally overlapping Ca^{2+} , H^+ and Cl^- fluxes in roots. Whereas, the effect of PEP1 and ATP on the cytosolic redox state was below the threshold of ROS-induced Ca^{2+} signaling (Supplemental Figure 7).

Glutathione disulfide (GSSG) treatments trigger rapid Ca^{2+} , H^+ and Cl^- fluxes that precede a slow-progressing cytosolic oxidation

GSSG is known to trigger cytosolic Ca^{2+} signals (Gomez et al., 2004) and to directly oxidize Grx1-roGFP2 (Gutscher et al., 2008). We sought to resolve the spatiotemporal relations of these responses in Arabidopsis seedlings expressing R-GECO1-P2A-Grx1-roGFP2. Although 1 mM GSSG-induced Ca^{2+} signals were variable, they appeared to arrive from upper root regions and travelled towards the root tip, followed by a second Ca^{2+} burst in the vasculature and oscillations in the meristematic- and elongation zone (Figure 10A left, Supplemental Movie 15). After the initial Ca^{2+} signal reached the root tip, in this region a cytosolic oxidation was observed that slowly progressed towards the elongation zone, where oscillation became visible (Figure 10A right, Supplemental Movie 15). Note that the Ca^{2+} and E_{GSH} oscillations were shifted in phase (Supplemental Movie 15). Additional experiments using R-GECO1-GSL- E^2GFP

revealed that GSSG treatments also induced H^+ and Cl^- influx, exhibiting the most noticeable spatiotemporal overlap with Ca^{2+} signals in the meristematic- and early elongation zone during the initial 20 min of the GSSG response (Figures 10B and 10C, Supplemental Movie 16). In summary, GSSG treatments trigger Ca^{2+} signals, cytosolic acidification and Cl^- influx that precede a cytosolic oxidation.

DISCUSSION

Optimization of ABA indicators in HEK293T cells

Optimization procedures of FRET-based indicators usually comprise the testing of FRET-pair-, sensory domain- and linker variants (Okumoto et al., 2012; Hochreiter et al., 2015). Although such testing has been performed on ABACUS, the linkers between the sensory domain and attached fluorescent proteins remained invariant (Jones et al., 2014). Early optimizations of ABAleon focused on sensory domain modifications that led to the development of ABAleon2.15 with improved stereospecificity for (+)-ABA (Waadt et al., 2014). Using the HEK293T cell system, we have developed ABAleonSD1-3L21 that exhibits an improved signal-to-noise ratio compared to ABAleon2.15 (Figure 1B). HEK293T cells are a convenient system for GEFI screenings, because they can be easily transfected, cultivated and analyzed in a plate reader (Tian et al., 2009). HEK293T cells contain neglectable ABA levels, and are therefore well suited for ABA indicator screenings with a potential to facilitate the heterologous characterization of ABA transporters using ABA indicators. Successful characterizations of plasma membrane proteins in HEK293T cells has been demonstrated for RBOHs and CYCLIC NUCLEOTIDE-GATED ION CHANNEL (CNGC)-type Ca^{2+} channels (Gao et al., 2016; Han et al., 2019). The differences of ABAleon characteristics between HEK293T cell and in vitro analyses might be due to a lower stability of ABAleon2.15 in HEK293T cells. Similar issues have been reported for Ca^{2+} indicators (Tian et al., 2009). However, in vitro characterizations of ABA indicators are time-consuming, and screenings using *E. coli* are not practical due to a likely even lower protein stability in this system (Jones et al., 2014; Waadt et al., 2014). Note that previously measured properties of ABAleon2.15

($\Delta R_{(\max)}/R_0 \sim -0.10$ and $k'_d \sim 500$ nM; Waadt et al., 2014) were different compared to results in Figures 1C and 1F. Here we used a faster sandwich-tag purification procedure with subsequent characterization of freshly purified proteins that might give more reliable results.

ABAlcon2.15 and ABAlconSD1-3L21 exhibited similar ABA responses in Arabidopsis. ABACUS1-2 μ responded slower to ABA and did not resolve the ABA gradient in roots (Figure 2, Supplemental Figure 2), probably due to the lower ABA affinity (Jones et al., 2014). However, this indicator might have advantages for the analyses of ABA dynamics in the root tip, the root-hypocotyl junction and in guard cells, where ABAlcons are close to saturation (Figure 2; Waadt et al., 2015). In the future, optimization of ABA indicators will require the development of alternative sensory domains and the investigation of alternative biosensor designs.

2-In-1-GEFIs facilitate simultaneous multiparametric analyses

Multiplexed live imaging in plants has been performed via the combination of GEFIs with fluorescent dyes, the use of GEFIs in parallel experiments, or through dual-expression of Ca^{2+} indicators (Monshausen et al., 2007, 2009, 2011; Loro et al., 2012; Schwarzländer et al., 2012; Ngo et al., 2014; Keinath et al., 2015; Behera et al., 2018; Kelner et al., 2018; Wagner et al., 2019). However, GEFI-based simultaneous analyses of two signaling compounds has been established in Arabidopsis only for Ca^{2+} and ABA (Waadt et al., 2017). Because most GEFIs are FRET- or green fluorescent protein-based, simultaneous multiparametric analyses have become possible through the development of red fluorescent protein-based indicators for Ca^{2+} , redox/ H_2O_2 and pH (Bilan and Belousov, 2017; Martynov et al., 2018; Walia et al., 2018). Yet, except for R-GECO1, their application in plants is rare. Here, we introduced the intensimetric red fluorescing pH indicator (P)A-17 (Shen et al., 2014), which is well suited for pH analyses in Arabidopsis with similar responsiveness compared to the ratiometric E^2GFP (Figures 3B left, 4A middle and 5A middle).

As the generation of stable transgenic organisms is time-consuming, simultaneous expression of GEFIs, or the generation of dual-sensing GEFIs, is advantageous. Moreover, the latter approach minimizes epigenetic silencing effects,

often observed in lines carrying multiple transgenes. Dual-sensing GEFIs have been developed for pH and Cl^- (ClopHensor; Arosio et al., 2010) and for phosphatidylinositol 3,4,5-trisphosphate localization and H_2O_2 concentration (PIP-SHOW; Mishina et al., 2012). For the generation of our 2-In-1-GEFIs we were inspired by ClopHensor and the incorporated E^2GFP that we fused with R-GECO1 in analogy to R-GECO1-GSL-mTurquoise (Waadt et al., 2017). Note that recent studies indicated that ClopHensor/ E^2GFP might also respond to NO_3^- (<https://doi.org/10.1101/716050>). Because our imaging buffer contained 5 mM Cl^- and the microscope-dish agarose media contained 4 mM NO_3^- , the observed E^2GFP responses likely depended on both anion species. In contrast to R-GECO1-GSL- E^2GFP , the other 2-In-1-GEFIs have been linked via the self-cleaving P2A-peptide, which enables efficient cleavage in Arabidopsis (Burén et al., 2012; Supplemental Figures 3 and 4). In addition, P2A-based 2-In-1-GEFIs exhibit similar responses compared to indicators expressed from separate expression cassettes (Figure 8, Supplemental Figure 6). Because only one expression cassette is used, P2A-linked GEFIs can be more easily screened at the microscope and are less prone to unwanted silencing effects. Our work established several 2-In-1-GEFIs based on the P2A-linkage, which is applicable to any eukaryotic system (Kim et al., 2011).

Ca^{2+} , H^+ and anion fluxes exhibit a high spatiotemporal overlap

Previous work established that mechanical stimulation, wounding, ATP and auxin simultaneously induce Ca^{2+} and H^+ fluxes (Monshausen et al., 2009, 2011; Behera et al., 2018). We found that, in addition to auxin and ATP, also glutamate, PEP1 and GSSG trigger Ca^{2+} , H^+ and anion fluxes with high spatiotemporal overlap (Figures 4, 5, 9 and 10). The linkage of Ca^{2+} and H^+ fluxes may depend on H^+ pumps and $\text{Ca}^{2+}/\text{H}^+$ -coupled transport via CATION/PROTON EXCHANGERS (CAXs) or Ca^{2+} -ATPases (Bonza and De Michelis, 2011; Pittman and Hirschi, 2016). However, knowledge about their role in Ca^{2+} signaling is fragmentary, probably due to functional overlap of gene family members (Behera et al., 2018). Simultaneous Ca^{2+} and H^+ fluxes in response to auxin are mediated by the auxin/ H^+ -symporter AUXIN RESISTANT 1 (AUX1) and the Ca^{2+} channel CNGC14 that are functionally coupled (Shih et al., 2015; Dindas et al.,

2018). Since the activation of plasma membrane H^+ -ATPases is coupled to AUX1 (Inoue et al., 2016), this could explain the subsequent H^+ -efflux.

Mechanical stimulation-induced Ca^{2+} and H^+ fluxes depend on the RECEPTOR-LIKE KINASE (RLK) FERONIA, which acts as a receptor for RAPID ALKALIZATION FACTOR (RALF) peptides (Haruta et al., 2014; Shih et al., 2014; Stegmann et al., 2017). Several RLKs, including the ATP receptor DOES NOT RESPOND TO NUCLEOTIDES 1 (DORN1) and PEP RECEPTORS (PEPRs), can induce Ca^{2+} signals, apoplastic alkalization (coupled to cytosolic acidification), and ROS production (Qi et al., 2010; Choi et al., 2014; Ma et al., 2014; Seybold et al., 2014; Haruta et al., 2015; Chen et al., 2017; Kimura et al., 2017). The effect of DORN1 and PEPRs on anion efflux was observed in guard cells during stomatal closure (Chen et al., 2017; Zheng et al., 2018). Our analyses revealed that PEP1 induced a transient anion influx along the entire imaged root axis. Extracellular ATP triggered anion influx in the differentiation- and elongation zone, but efflux in the meristematic zone (Figure 9). Early research revealed that cytosolic but not extracellular ATP is required for adenine nucleotide activation of R-type anion channels (Hedrich et al., 1990) and protein kinase-mediated activation of S-type anion channels (Schmidt et al., 1995). It is likely that ATP-triggered Ca^{2+} signals activate Ca^{2+} -dependent protein kinases required for the activation of anion channels (Mori et al., 2006). The differences in PEP1- and ATP-induced anion fluxes in roots might be due to the distinct Ca^{2+} signatures observed in the meristematic zone (Figure 9). In the future, it will be interesting to discriminate the differences in anion-flux regulation between roots and guard cells.

On the interdependence of Ca^{2+} and ROS signaling

The interdependence of Ca^{2+} and ROS signaling has been extensively discussed (Gilroy et al., 2014; Steinhorst and Kudla, 2014). In the context of long-distance and systemic signaling, current models propose that Ca^{2+} signals trigger the ion channel TPC1 for signal amplification. Ca^{2+} signal propagation occurs via plasmodesmata or Ca^{2+} -dependent activation of RBOHs. RBOH-derived extracellular ROS propagate to adjacent cells to activate plasma membrane localized Ca^{2+} channels (Evans et al., 2016; Choi et al., 2016). Ca^{2+} -dependent activation of RBOHs is well established

(Kadota et al., 2015; Han et al., 2019). However, whether RBOH-dependent ROS contribute to Ca^{2+} channel activation, has only been inferred from pharmacological- and genetic analyses (Kwak et al., 2003; Evans et al., 2016). In *Arabidopsis* guard cells, hyperpolarization-activated Ca^{2+} -permeable channels can be activated by 50 μM H_2O_2 (Pei et al., 2000). In *Vicia faba* guard cells such channels exhibit a half response at 76 μM H_2O_2 (Köhler et al., 2003). Analyses in root epidermis cells revealed a Ca^{2+} channel activation by 10 μM H_2O_2 in the elongation zone and by 1 mM H_2O_2 in the maturation zone (Demidchik et al., 2007). The threshold concentrations of ROS required to activate Ca^{2+} channels may depend on the cell type, the location (apoplast or cytosol) and the chemical nature of ROS (Demidchik et al., 2007).

Our analyses revealed that in *Arabidopsis* roots 20 μM extracellular H_2O_2 triggered a detectable cytosolic oxidation, but no considerable Ca^{2+} signals (Figure 7). On the other hand, glutamate, ATP and PEP1, which efficiently trigger Ca^{2+} signals, induced a cytosolic oxidation rather below this threshold (Supplemental Figure 7). These data are consistent with a slow progressing cytosolic oxidation in response to the pathogen-associated molecular pattern flagellin fragment flg22 (Nietzel et al., 2019). Whether such cytosolic oxidation is Ca^{2+} dependent, requires further experimentation. However, compared to 20 μM H_2O_2 responses, our data suggest that glutamate-, ATP- and PEP1-induced cytosolic oxidation is not sufficient to trigger root Ca^{2+} channels (Supplemental Figure 7). We hypothesize that the observed ROS dependence of Ca^{2+} signaling may be indirectly linked to the impact of ROS on the cell wall, which binds considerable amounts of Ca^{2+} in Ca^{2+} -pectate cross-linked complexes (Hepler et al., 2010; Peaucelle et al., 2012; Kärkönen and Kuchitsu, 2015). Such a model would be consistent with a rather slow H_2O_2 activation of Ca^{2+} channels in patch clamp analyses (20-60 min; Demidchik et al., 2007). Another possibility would be that a signaling component triggers both, Ca^{2+} and ROS signaling. The BOTRYTIS-INDUCED KINASE 1 (BIK1) could be a good candidate for such a mechanism (Kadota et al., 2014; Li et al., 2014; Kimura et al., 2017; Tian et al., 2019).

Conclusions

Our work established 2-In-1-GEFI-based simultaneous multiparametric in vivo analyses of signaling compounds in Arabidopsis. Using the 2-In-1-GEFIs, we observed that in roots ABA does not trigger rapid Ca^{2+} or pH changes. Whereas, auxin, glutamate, ATP, PEP1 and GSSG induce Ca^{2+} , H^+ and anion fluxes with high spatiotemporal overlap (Figure 11, Supplemental Figure 8). These results suggest an interdependence and coordination of ion fluxes that need to be dissected in future research. Findings that glutamate-, ATP- and PEP1-induced cytosolic oxidation is below the threshold required for triggering Ca^{2+} channels argue against the current model of a ROS-assisted Ca^{2+} signal propagation mechanism (Evans et al., 2016). We hypothesize that ROS may have an indirect effect on Ca^{2+} signaling. Overall, 2-In-1-GEFI-based imaging will allow for high resolution in vivo analyses of signaling processes beyond the model plant Arabidopsis.

METHODS

Generation of plasmids

Oligonucleotides used for cloning procedures (Supplemental Data Set 1A) were obtained from Eurofinsgenomics. Plasmids (Supplemental Data Set 1B) were constructed using classical cloning procedures and the GreenGate system (Lampropoulos et al., 2013) utilizing enzymes from Thermo Fisher Scientific. Arabidopsis codon-optimized DNA fragments of PmTurquoise and PA-17 were designed using GeneArtTM gene synthesis (Thermo Fisher Scientific). Expression of GEFIs in *Arabidopsis thaliana* Col-0 was carried out utilizing the promoter of a ubiquitous and highly expressed reference gene *ASPARTIC PROTEASE A1* (*APA1*, AT1G11910) that was chosen based on searches using Genevestigator (Hruz et al., 2008). The expression cassette also included the terminator of the *HEAT SHOCK PROTEIN 18.2* (*HSP18.2*) gene (AT5G59720; Nagaya et al., 2010; Waadt et al., 2014).

Optimization of ABA indicators in HEK293T cells

Transformation and cultivation of HEK293T cells was performed as described previously (Ogasawara et al., 2008; Zhang et al., 2018). Spectral characteristics of ABA indicators were recorded in Greiner flat bottom 96-well plates (Greiner BIO-ONE) using a TECAN Safire plate reader (TECAN) operated by the XFLUOR4.51 software with the following parameters: fluorescence emission scan bottom mode; excitation wavelength 440 nm, bandwidth 12.5 nm; emission wavelength scan from 460-600 nm, bandwidth 10 nm; gain 100-115; flashes 10; integration time 40-60 μ s; temperature 37 °C. HEK293T cells were kept in Hanks Balanced Salt Solution (HBSS; Thermo Fisher Scientific) and fluorescence emission spectra were recorded before (t_0) and 60 min (t_{60}) after exchange of solution to either HBSS with 100 μ M ($\pm$)-ABA (Merck) and 0.1 % EtOH (treatment) or HBSS with 0.1 % EtOH (solvent control). ABA indicator emission ratios (R) were calculated as average emission at 518-538 nm divided by average emission at 470-490 nm after subtraction of the non-transfected HEK293T cell background emission spectrum. Emission ratio change ($\Delta R/R_0$) was calculated as $[R(t_{60})-R(t_0)]/R(t_0)$. Experiments were performed in triplicates.

Purification and in vitro characterization of ABAleons

BL21-CodonPlus (DE3)-RIL cells transformed with pET28-6xHis-ABAlleon-(P)StreptII constructs were shaken at 150 rpm and 37 °C in 2x 1 L Luria Broth (LB) media supplemented with 50 μ g mL⁻¹ kanamycin and 30 μ g mL⁻¹ chloramphenicol. At an optical density (OD₆₀₀) of 0.5, 1 mM Isopropyl β -D-1-thiogalactopyranoside (IPTG; Carl Roth) was added and protein expression was conducted in a shaking incubator at 24 °C for 6 h. Cultured cells were harvested by several centrifugation steps at 4 °C and 4000 rpm and bacterial pellet was flash frozen in liquid N₂ and stored at - 80 °C.

The bacterial pellet was thawed on ice and resuspended in 30 mL lysis buffer (1x PBS [137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 1.8 mM KH₂PO₄], 10 mM imidazole [Merck], 1x Roche protease inhibitor EDTA-free, 1 mM Phenylmethylsulfonyl fluoride [PMSF; Carl Roth] and 1 mg mL⁻¹ lysozym [VWR], pH 7.4). After 40-60 min incubation on ice, cells were disrupted through microtip-based sonication (25 % amplitude, 21x 20 s) and cell debris were removed by centrifugation (2x 30 min, 20000 g, 4 °C) and filtering through 0.45 μ m syringe filters (Merck).

6x-His purification was conducted in 20 mL gravity columns (VWR) loaded with 4 mL HisPur™ Ni-NTA resin (Thermo Fisher Scientific). After binding of proteins to the Ni-NTA resin, columns were washed 5x with 10 mL His-wash buffer (1x PBS, 25 mM imidazole, pH 7.4) and proteins were eluted in 3x 2 mL His-elution buffer (1x PBS, 250 mM imidazole, pH 7.4). Purified proteins were then loaded onto a 20 mL gravity column supplemented with 3 mL 50 % Strep Tactin Superflow (IBA). After 4x washing with 7.5 mL SII-wash buffer 1 (30 mM Tris/HCl pH 7.4, 250 mM NaCl) and 3x washing with 7.5 mL SII-wash buffer 2 (30 mM Tris/HCl pH 7.4, 250 mM NaCl, 10 mM MgCl₂, 1 mM MnCl₂), proteins were eluted in 3x 1.5 mL SII-elution buffer (30 mM Tris/HCl pH 7.4, 250 mM NaCl, 10 mM MgCl₂, 1 mM MnCl₂, 2.5 mM desthiobiotin [IBA]) and concentrated to a final volume of about 1 mL using Amicon Ultra-4 30 K filters (Merck). Purity of proteins was analyzed by SDS-PAGE using 10 % Mini-PROTEAN® TGX™ Precast Gels (BioRad) and InstantBlue staining (Expedeon). In a similar procedure, protein yield was calculated according to a bovine serum albumin standard curve.

For vitro calibration, a 100 mM (+)-ABA (TCI) stock solution dissolved in 100 % EtOH was used for an ABA dilution series in SII-wash buffer 2 and 0.2 % EtOH. 10 µL of each ABA dilution were added to 3 wells of black flat bottom µclear® 96-well plates (Greiner BIO-ONE) containing 90 µL of ~ 1.1 µM ABAleon protein, diluted in SII-wash buffer 2, or to 90 µL SII-wash buffer 2 alone as background control. Fluorescence emission spectra were recorded after 20 min incubation at room temperature in the dark using a TECAN Infinite M1000 plate reader (TECAN) operated by the i-control 1.10.4.0 software with the following parameters: fluorescence emission scan bottom mode; excitation wavelength 440 nm, bandwidth 10 nm; emission wavelength scan from 460-650 nm, bandwidth 10 nm; gain 98-104, flashes 10 at 100 Hz, integration time 60 µs, temperature 21-22 °C. ABA-dependent ABAleon emission ratios (R) were calculated as described above. Maximum emission ratio change ($\Delta R_{(max)}/R_0$) was calculated as $[R(\text{at } 20 \mu\text{M ABA}) - R(\text{at } 0 \mu\text{M ABA})]/R(\text{at } 0 \mu\text{M ABA})$. Apparent ABA affinities (k'_d ; EC50) of ABAleons were calculated by fitting the emission ratio values of all three replicates to a 4-parameter logistic function using OriginPro 2018 (OriginLab Corporation).

Generation of transgenic Arabidopsis plants expressing GEFIs

Agrobacterium strain ASE containing the pSOUP helper plasmid and the respective plant expression vectors (Supplemental Data Set 1B) were used for transformation of *Arabidopsis thaliana* ecotype Col-0 by floral dip (Clough & Bent, 1998) to generate the transgenic lines listed in Supplemental Data Set 1C. Seeds of transformed plants were surface sterilized for 10-15 min in 70 % EtOH, washed three times with 100 % EtOH and sowed on half-strength Murashige & Skoog (0.5 MS) media (Duchefa) supplemented with 5 mM MES-KOH pH 5.8, 0.8 % phytoagar (Duchefa) and 10 µg mL⁻¹ Glufosinate-ammonium or 25 µg mL⁻¹ hygromycin B (Merck) for herbicide selection. After 3-6 days of stratification in the dark at 4 °C, transgenic plants were grown for six-days in a growth room (16 h day/8 h night, 22 °C, 65 % relative humidity, photon fluence rate 100 µmol m⁻² s⁻¹). Positive transformants were then transferred to herbicide-free 0.5 MS media-containing petri dishes. After one day recovery, GEFI expression was confirmed by visual inspection at a Zeiss Discovery.V20 fluorescence stereo microscope equipped with GFP, YFP and RFP filters and a Plan S 1.0x FWD 81 mm lens. Approximately 40 herbicide resistant and fluorescing seedlings were then transferred to round 7 cm pots containing classic soil (Einheitserde) and grown until seed ripening in the growth room. ABAleon expressing plants were covered with a plastic lid and grown in a Conviron CMP6010 growth chamber (16 h day/8 h night, 20 °C, 65 % relative humidity, photon fluence rate 150 µmol m⁻² s⁻¹). To confirm proper GEFI expression, compare GEFI fluorescence emissions and avoid silencing effects in next generations, one leaf of each individual about three-week-old plant was examined at a confocal laser scanning microscope Leica SP5 II equipped with a HCX PL APO CS 20.0 x 0.7 IMM UV objective (Leica Microsystems) using emission and excitation settings listed in Supplemental Data Set 1D. For each construct, at least two transgenic lines with highest GEFI expression, proper 3:1 segregation in the 2nd generation and least silencing were selected for further propagation. One line, indicated with (microscope) in Supplemental Data Set 1C, was used for microscopic experiments.

Phenotypic characterization of GEFI lines

Seeds were surface sterilized and sown on 0.5 MS media supplemented with 5 mM MES-KOH pH 5.8 and 0.8 % phytoagar. Seven-day-old seedlings grown in the

growth room (16 h day/8 h night, 22 °C, 65 % relative humidity, photon fluence rate 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$) were transferred to soil in single pots and further grown until 28-days-old. Pictures from 9-12 plants per genotype were acquired from the top and rosette area values were obtained using the Rosette Tracker Fiji plugin (De Vylder et al., 2012).

Microscopic analyses

Seeds of GEFI expressing lines were surface sterilized and sown in four horizontal rows on square petri dishes containing LAK media (Barragán et al., 2012; 1 mM KH_2PO_4 , 2 mM $\text{Ca}(\text{NO}_3)_2$, 1 mM MgSO_4 , 30 μM H_3BO_3 , 10 μM MnSO_4 , 1 μM ZnSO_4 , 1 μM CuSO_4 , 0.03 μM $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$, 50 μM FeNaEDTA) supplemented with 10 mM MES-Tris pH 5.6 and 0.8 % phytoagar. After six days of stratification in the dark at 4 °C, seedlings were grown vertically in a Conviron CMP 6010 growth chamber (16 h day/8 h night, 22 °C, 65 % relative humidity, photon fluence rate 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$). After four days, seedlings were transferred to microscope dishes (MatTek) containing 200 μL polymerized LAK media supplemented with 10 mM MES-Tris pH 5.6 and 0.7 % low melting point (LMP) agarose (Carl Roth). Seedlings were incubated vertically overnight in the growth chamber until the microscopic experiments were conducted.

Before microscopic analyses, seedlings on microscope dishes were placed horizontally and topped with 90 μL imaging buffer (Allen et al., 2001; 5 mM KCl, 50 μM CaCl_2 , 10 mM MES-Tris pH 5.6). Using a 200 μL pipet tip, seedlings were gently attached back to the LAK media-LMP agarose bed and incubated for 10-50 min for recovery until the GEFI fluorescence emission baseline was stable. Microscopic analyses were performed at Leica SP5 II and Leica SP8 confocal laser scanning microscopes using a 10x air objective and photomultiplier tube detectors (Leica Microsystems). Microscope settings were as follows: image format 1024x178 pixels (1536x256 pixels for RW300 experiment); bidirectional scanning at 400 Hz; zoom 0.75 (SP8) or 1 (SP5 II and RW300 experiment at SP8); pinhole 5 AU; line accumulation 2 (SP5 II) and 1 (SP8); line average 1 (SP5 II) and 2 (SP8); offset -0.4 % for blue, cyan, green and yellow emissions and -0.2 % for red emissions; frame rate 6 sec. Laser intensities and gain settings were optimized for each GEFI and kept stable for all experimental replicates. Emission and excitation settings for each GEFI are listed in

Supplemental Data Set 1D. After 4 min baseline recording, chemical treatments were performed by dropping 10 μ L of 10-fold concentrated stock solutions (Supplemental Data Set 1E) close to the imaged region.

Image processing and analyses were conducted using Fiji (Schindelin et al., 2012). Image processing included background subtraction (2-4), gaussian blur (1), median (1), 32-bit conversion, thresholding of background noise (2-5) and ratio image calculation for ratiometric GEFIs. Normalized fluorescence intensity ($\Delta F/F_0$) and emission ratio ($\Delta R/R_0$) analyses, and root tip localized time-dependent vertical response profiles were generated using a custom build Fiji plugin (will be uploaded to github after article acceptance) that utilizes additional plugins, such as VectorGraphics2D-0.13 (<https://github.com/eseifert/vectorgraphics2d>) and xchart-3.5.2 (<https://github.com/knowm/XChart>). Fluorescence emissions (F) and emission ratios (R) were measured as average values from each entire processed movie frame and signal changes ($\Delta F/F_0$ and $\Delta R/R_0$) were calculated relative to the average value of a 4 min baseline recording as $[F(t)-F(\text{baseline})]/F(\text{baseline})$ and $[R(t)-R(\text{baseline})]/R(\text{baseline})$. Means and SD of experimental replicates were calculated using Excel (Microsoft). For time-dependent vertical response profiles, root tips were detected within each movie frame and regions of interest (ROIs) were drawn to cover the entire x-axis and a defined area above the root tip. Vertical response profiles were calculated from each movie frame ROI as average of all x-axis pixel values within each y-axis pixel line (similar to the Plot Profile command in Fiji) and plotted in a time-dependent manner. Time-dependent signal change vertical response profiles were calculated using the raw response profiles as a basis and applying the signal change formulas to each y-axis pixel line. Average time-dependent vertical response profiles of multiple experimental replicates were generated using the average Z projection command in Fiji.

Statistical Analysis

For phenotypic analyses presented in Supplemental Figure 5, Box plot- and statistical analyses using pairwise Tukey test comparisons relative to Col-0 wild type were conducted using OriginPro 2018 (OriginLab Corporation).

Accession Numbers

The Arabidopsis Genome Initiative locus numbers for the genes used in this article are as follows: *ABI1* (AT4G26080), *APA1* (AT1G11910), *AtPEP1* (AT5G64900), *HSP18.2* (AT5G59720), *LTI6b* (AT3G05890), *PYL1* (AT5G46790), *PYR1* (AT4G17870), *VTI11* (AT5G39510).

Supplemental Data

Supplemental Figure 1. Topologies of ABA indicators.

Supplemental Figure 2. Solvent control experiments of ABA indicators in Arabidopsis.

Supplemental Figure 3. Targeting of E²GFP to the plasma membrane.

Supplemental Figure 4. Targeting of E²GFP to the tonoplast.

Supplemental Figure 5. Targeting of E²GFP to the plasma membrane induces plant growth defects.

Supplemental Figure 6. Cytosolic E_{GSH} is only weakly affected by ATP and PEP1 treatments.

Supplemental Figure 7. Glutamate-, ATP- and PEP1-dependent cytosolic oxidation is below the threshold of ROS-induced Ca²⁺ signaling.

Supplemental Figure 8. Ca²⁺, H⁺ and anion fluxes exhibit a high spatiotemporal overlap.

Supplemental Data Set 1. Lists of materials, imaging settings and chemicals used in this work.

Supplemental Movie 1. ABA indicator ABA responses in Arabidopsis.

Supplemental Movie 2. ABAleonSD1-3L21-P2A-R-GECO1 in response to ABA and IAA.

672 **Supplemental Movie 3.** PA-17-P2A-ABALeonSD1-3L21 in response to ABA and
673 ATP.

674 **Supplemental Movie 4.** R-GECO1-GSL-E²GFP in response to IAA and ATP.

675 **Supplemental Movie 5.** R-GECO1-GSL-E²GFP in response to glutamate and ATP.

676 **Supplemental Movie 6.** PA-17-P2A-Grx1-roGFP2 in response to glutamate and
677 H₂O₂.

678 **Supplemental Movie 7.** PA-17-P2A-roGFP2-Orp1 in response to glutamate and
679 H₂O₂.

680 **Supplemental Movie 8.** R-GECO1-P2A-Grx1-roGFP2 in response to H₂O₂.

681 **Supplemental Movie 9.** R-GECO1-P2A-roGFP2-Orp1 in response to H₂O₂.

682 **Supplemental Movie 10.** R-GECO1 and roGFP2-Orp1 in response to ATP and
683 H₂O₂.

684 **Supplemental Movie 11.** R-GECO1-P2A-roGFP2-Orp1 in response to PEP1 and
685 H₂O₂.

686 **Supplemental Movie 12.** R-GECO1 and Grx1-roGFP2 in response to ATP and
687 H₂O₂.

688 **Supplemental Movie 13.** R-GECO1-P2A-Grx1-roGFP2 in response to PEP1 and
689 H₂O₂.

690 **Supplemental Movie 14.** R-GECO1-GSL-E²GFP in response to PEP1 and ATP.

691 **Supplemental Movie 15.** R-GECO1-P2A-Grx1-roGFP2 in response to GSSG.

692 **Supplemental Movie 16.** R-GECO1-GSL-E²GFP in response to GSSG.

693 **Supplemental Movie Legends**

ACKNOWLEDGEMENTS

We thank the groups at COS (Heidelberg) for generous access to equipment and GreenGate modules, Dr. Jana Hakenjos for initial help with ABAleon purifications, Dr. Andreas Meyer (University of Bonn) for roGFP2 PCR templates, Dr. Eugenia Russinova (VIB Gent) for providing the AtPEP1 peptide and Dr. Shintaro Munemasa (Okayama University) for helpful discussions. This work was supported by the Deutsche Forschungsgemeinschaft (DFG WA 3768/1-1) to R.W., (DFG AN 1323/1-1) to Z.A. and (DFG KU 931/14-1) to J.K..

AUTHOR CONTRIBUTIONS

R.W. conceived the project, generated most of the plasmids and transgenic plants, conducted the in vitro characterization of ABAleons, performed all microscopic analyses and wrote the manuscript. P.K. conducted the characterization of ABA indicators in HEK293T cells and revised the manuscript. Z.A. generated and characterized the transgenic lines RW307 and RW308 and revised the manuscript. C.W. developed the GEFI analyzer Fiji plugin and revised the manuscript. G.B. generated the plasmids indicated with GB and conducted preliminary ABA indicator characterizations. K.L. introduced A-17 and generated PA-17 GreenGate modules. J.K. supervised the ABA indicator characterization in HEK293T cells and revised the manuscript. K.S. supervised and hosted the project.

REFERENCES

- Allen, G.J., Chu, S.P., Harrington, C.L., Schumacher, K., Hoffmann, T., Tang, Y.Y., Grill, E., and Schroeder, J.I.** (2001). A defined range of guard cell calcium oscillation parameters encodes stomatal movements. *Nature* **411**, 1053-1057.
- Arosio, D., Ricci, F., Marchetti, L., Gualdani, R., Albertazzi, L., and Beltram, F.**

- (2010). Simultaneous intracellular chloride and pH measurements using a GFP-based sensor. *Nat Methods* **7**, 516-518.
- Barragán, V., Leidi, E.O., Andrés, Z., Rubio, L., De Luca, A., Fernández, J.A., Cubero, B., and Pardo, J.M.** (2012). Ion exchangers NHX1 and NHX2 mediate active potassium uptake into vacuoles to regulate cell turgor and stomatal function in *Arabidopsis*. *Plant Cell* **24**, 1127-1142.
- Behera, S., Zhaolong, X., Luoni, L., Bonza, M.C., Doccu, F.G., De Michelis, M.I., Morris, R.J., Schwarzländer, M., and Costa, A.** (2018). Cellular Ca^{2+} Signals Generate Defined pH Signatures in Plants. *Plant Cell* **30**, 2704-2719.
- Bilan, D.S., and Belousov, V.V.** (2017). New tools for redox biology: From imaging to manipulation. *Free Radic Biol Med* **109**, 167-188.
- Bizzarri, R., Arcangeli, C., Arosio, D., Ricci, F., Faraci, P., Cardarelli, F., and Beltram, F.** (2006). Development of a novel GFP-based ratiometric excitation and emission pH indicator for intracellular studies. *Biophys J* **90**, 3300-3314.
- Bonza, M.C., and De Michelis, M.I.** (2011). The plant Ca^{2+} -ATPase repertoire: biochemical features and physiological functions. *Plant Biol (Stuttg)* **13**, 421-430.
- Burén, S., Ortega-Villasante, C., Otvös, K., Samuelsson, G., Bakó, L., and Villarejo, A.** (2012). Use of the foot-and-mouth disease virus 2A peptide co-expression system to study intracellular protein trafficking in *Arabidopsis*. *PLoS One* **7**, e51973.
- Chen, D., Cao, Y., Li, H., Kim, D., Ahsan, N., Thelen, J., and Stacey, G.** (2017). Extracellular ATP elicits DORN1-mediated RBOHD phosphorylation to regulate stomatal aperture. *Nat Commun* **8**, 2265.
- Choi, J., Tanaka, K., Cao, Y., Qi, Y., Qiu, J., Liang, Y., Lee, S.Y., and Stacey, G.** (2014). Identification of a plant receptor for extracellular ATP. *Science* **343**, 290-294.
- Choi, W.G., Hilleary, R., Swanson, S.J., Kim, S.H., and Gilroy, S.** (2016). Rapid, Long-Distance Electrical and Calcium Signaling in Plants. *Annu Rev Plant Biol* **67**, 287-307.
- Clough, S.J., and Bent, A.F.** (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**, 735-743.
- Cutler, S.R., Rodriguez, P.L., Finkelstein, R.R., and Abrams, S.R.** (2010). Absciscic

- acid: emergence of a core signaling network. *Annu Rev Plant Biol* **61**, 651-679.
- De Vyllder, J., Vandenbussche, F., Hu, Y., Philips, W., and Van Der Straeten, D.** (2012). Rosette tracker: an open source image analysis tool for automatic quantification of genotype effects. *Plant Physiol* **160**, 1149-1159.
- Demidchik, V., Shabala, S.N., and Davies, J.M.** (2007). Spatial variation in H₂O₂ response of *Arabidopsis thaliana* root epidermal Ca²⁺ flux and plasma membrane Ca²⁺ channels. *Plant J* **49**, 377-386.
- Demidchik, V., Shang, Z., Shin, R., Thompson, E., Rubio, L., Laohavisit, A., Mortimer, J.C., Chivasa, S., Slabas, A.R., Glover, B.J., Schachtman, D.P., Shabala, S.N., and Davies, J.M.** (2009). Plant extracellular ATP signalling by plasma membrane NADPH oxidase and Ca²⁺ channels. *Plant J* **58**, 903-913.
- Dindas, J., Scherzer, S., Roelfsema, M.R.G., von Meyer, K., Müller, H.M., Al-Rasheid, K.A.S., Palme, K., Dietrich, P., Becker, D., Bennett, M.J., and Hedrich, R.** (2018). AUX1-mediated root hair auxin influx governs SCF^{TIR1/AFB}-type Ca²⁺ signaling. *Nat Commun* **9**, 1174.
- Dodd, A.N., Kudla, J., and Sanders, D.** (2010). The language of calcium signaling. *Annu Rev Plant Biol* **61**, 593-620.
- Edel, K.H., and Kudla, J.** (2016). Integration of calcium and ABA signaling. *Curr Opin Plant Biol* **33**, 83-91.
- Evans, M.J., Choi, W.G., Gilroy, S., and Morris, R.J.** (2016). A ROS-Assisted Calcium Wave Dependent on the AtRBOHD NADPH Oxidase and TPC1 Cation Channel Propagates the Systemic Response to Salt Stress. *Plant Physiol* **171**, 1771-1784.
- Fendrych, M., Leung, J., and Friml, J.** (2016). TIR1/AFB-Aux/IAA auxin perception mediates rapid cell wall acidification and growth of *Arabidopsis* hypocotyls. *Elife* **5**, e19048.
- Finkelstein, R.** (2013). Absciscic Acid synthesis and response. *Arabidopsis Book* **11**, e0166.
- Gao, Q.F., Gu, L.L., Wang, H.Q., Fei, C.F., Fang, X., Hussain, J., Sun, S.J., Dong, J.Y., Liu, H., and Wang, Y.F.** (2016). Cyclic nucleotide-gated channel 18 is an essential Ca²⁺ channel in pollen tube tips for pollen tube guidance to ovules in *Arabidopsis*. *Proc Natl Acad Sci U S A* **113**, 3096-3101.

- Gilroy, S., Suzuki, N., Miller, G., Choi, W.G., Toyota, M., Devireddy, A.R., and Mittler, R.** (2014). A tidal wave of signals: calcium and ROS at the forefront of rapid systemic signaling. *Trends Plant Sci* **19**, 623-630.
- Gomez, L.D., Noctor, G., Knight, M.R., and Foyer, C.H.** (2004). Regulation of calcium signalling and gene expression by glutathione. *J Exp Bot* **55**, 1851-1859.
- Grossmann, G., Krebs, M., Maizel, A., Stahl, Y., Vermeer, J.E.M., and Ott, T.** (2018). Green light for quantitative live-cell imaging in plants. *J Cell Sci* **131**.
- Gutscher, M., Pauleau, A.L., Marty, L., Brach, T., Wabnitz, G.H., Samstag, Y., Meyer, A.J., and Dick, T.P.** (2008). Real-time imaging of the intracellular glutathione redox potential. *Nat Methods* **5**, 553-559.
- Gutscher, M., Sobotta, M.C., Wabnitz, G.H., Ballikaya, S., Meyer, A.J., Samstag, Y., and Dick, T.P.** (2009). Proximity-based protein thiol oxidation by H₂O₂-scavenging peroxidases. *J Biol Chem* **284**, 31532-31540.
- Han, J.P., Köster, P., Drerup, M.M., Scholz, M., Li, S., Edel, K.H., Hashimoto, K., Kuchitsu, K., Hippler, M., and Kudla, J.** (2019). Fine tuning of RBOHF activity is achieved by differential phosphorylation and Ca²⁺ binding. *New Phytol* **221**, 1935-1949.
- Haruta, M., Sabat, G., Stecker, K., Minkoff, B.B., and Sussman, M.R.** (2014). A peptide hormone and its receptor protein kinase regulate plant cell expansion. *Science* **343**, 408-411.
- Haruta, M., Gray, W.M., and Sussman, M.R.** (2015). Regulation of the plasma membrane proton pump (H(+)-ATPase) by phosphorylation. *Curr Opin Plant Biol* **28**, 68-75.
- Hauser, F., Waadt, R., and Schroeder, J.I.** (2011). Evolution of abscisic acid synthesis and signaling mechanisms. *Curr Biol* **21**, R346-355.
- Hauser, F., Li, Z., Waadt, R., and Schroeder, J.I.** (2017). SnapShot: Absciscic Acid Signaling. *Cell* **171**, 1708-1708.e1700.
- Hedrich, R., Busch, H., and Raschke, K.** (1990). Ca²⁺ and nucleotide dependent regulation of voltage dependent anion channels in the plasma membrane of guard cells. *EMBO J* **9**, 3889-3892.
- Hepler, P.K., and Winship, L.J.** (2010). Calcium at the cell wall-cytoplasm interface. *J*

- Integr Plant Biol **52**, 147-160.
- Hilleary, R., Choi, W.G., Kim, S.H., Lim, S.D., and Gilroy, S.** (2018). Sense and sensibility: the use of fluorescent protein-based genetically encoded biosensors in plants. Curr Opin Plant Biol **46**, 32-38.
- Hochreiter, B., Garcia, A.P., and Schmid, J.A.** (2015). Fluorescent proteins as genetically encoded FRET biosensors in life sciences. Sensors (Basel) **15**, 26281-26314.
- Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W., and Zimmermann, P.** (2008). Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. Adv Bioinformatics **2008**, 420747.
- Inoue, S.I., Takahashi, K., Okumura-Noda, H., and Kinoshita, T.** (2016). Auxin Influx Carrier AUX1 Confers Acid Resistance for Arabidopsis Root Elongation Through the Regulation of Plasma Membrane H⁺-ATPase. Plant Cell Physiol **57**, 2194-2201.
- Jones, A.M., Danielson, J.A., Manojkumar, S.N., Lanquar, V., Grossmann, G., and Frommer, W.B.** (2014). Absciscic acid dynamics in roots detected with genetically encoded FRET sensors. Elife **3**, e01741.
- Kadota, Y., Sklenar, J., Derbyshire, P., Stransfeld, L., Asai, S., Ntoukakis, V., Jones, J.D., Shirasu, K., Menke, F., Jones, A., and Zipfel, C.** (2014). Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase BIK1 during plant immunity. Mol Cell **54**, 43-55.
- Kadota, Y., Shirasu, K., and Zipfel, C.** (2015). Regulation of the NADPH Oxidase RBOHD During Plant Immunity. Plant Cell Physiol **56**, 1472-1480.
- Kärkönen, A., and Kuchitsu, K.** (2015). Reactive oxygen species in cell wall metabolism and development in plants. Phytochemistry **112**, 22-32.
- Keinath, N.F., Waadt, R., Brugman, R., Schroeder, J.I., Grossmann, G., Schumacher, K., and Krebs, M.** (2015). Live Cell Imaging with R-GECO1 Sheds Light on flg22- and Chitin-Induced Transient [Ca(2+)]_{cyt} Patterns in Arabidopsis. Mol Plant **8**, 1188-1200.
- Kelner, A., Leitão, N., Chabaud, M., Charpentier, M., and de Carvalho-Niebel, F.** (2018). Dual Color Sensors for Simultaneous Analysis of Calcium Signal Dynamics

in the Nuclear and Cytoplasmic Compartments of Plant Cells. *Front Plant Sci* **9**,
245.

**Kim, J.H., Lee, S.R., Li, L.H., Park, H.J., Park, J.H., Lee, K.Y., Kim, M.K., Shin, B.A.,
and Choi, S.Y.** (2011). High cleavage efficiency of a 2A peptide derived from
porcine teschovirus-1 in human cell lines, zebrafish and mice. *PLoS One* **6**, e18556.

Kimura, S., Waszczak, C., Hunter, K., and Wrzaczek, M. (2017). Bound by Fate: The
Role of Reactive Oxygen Species in Receptor-Like Kinase Signaling. *Plant Cell* **29**,
638-654.

Köhler, B., Hills, A., and Blatt, M.R. (2003). Control of guard cell ion channels by
hydrogen peroxide and abscisic acid indicates their action through alternate
signaling pathways. *Plant Physiol* **131**, 385-388.

Kostyuk, A.I., Panova, A.S., Bilan, D.S., and Belousov, V.V. (2018). Redox
biosensors in a context of multiparameter imaging. *Free Radic Biol Med* **128**, 23-39.

Kudla, J., Batistic, O., and Hashimoto, K. (2010). Calcium signals: the lead currency
of plant information processing. *Plant Cell* **22**, 541-563.

**Kwak, J.M., Mori, I.C., Pei, Z.M., Leonhardt, N., Torres, M.A., Dangl, J.L., Bloom,
R.E., Bodde, S., Jones, J.D., and Schroeder, J.I.** (2003). NADPH oxidase
AtrbohD and *AtrbohF* genes function in ROS-dependent ABA signaling in
Arabidopsis. *EMBO J* **22**, 2623-2633.

**Lampropoulos, A., Sutikovic, Z., Wenzl, C., Maegele, I., Lohmann, J.U., and
Forner, J.** (2013). GreenGate---a novel, versatile, and efficient cloning system for
plant transgenesis. *PLoS One* **8**, e83043.

**Li, L., Li, M., Yu, L., Zhou, Z., Liang, X., Liu, Z., Cai, G., Gao, L., Zhang, X., Wang,
Y., Chen, S., and Zhou, J.M.** (2014). The FLS2-associated kinase BIK1 directly
phosphorylates the NADPH oxidase RbohD to control plant immunity. *Cell Host
Microbe* **15**, 329-338.

Liu, Z., Wu, Y., Yang, F., Zhang, Y., Chen, S., Xie, Q., Tian, X., and Zhou, J.M.
(2013). BIK1 interacts with PEPRs to mediate ethylene-induced immunity. *Proc Natl
Acad Sci U S A* **110**, 6205-6210.

Loro, G., Drago, I., Pozzan, T., Schiavo, F.L., Zottini, M., and Costa, A. (2012).
Targeting of Cameleons to various subcellular compartments reveals a strict

cytoplasmic/mitochondrial Ca^{2+} handling relationship in plant cells. *Plant J* **71**, 1-13.

Ma, C., Guo, J., Kang, Y., Doman, K., Bryan, A.C., Tax, F.E., Yamaguchi, Y., and Qi, Z. (2014). AtPEPTIDE RECEPTOR2 mediates the AtPEPTIDE1-induced cytosolic Ca^{2+} rise, which is required for the suppression of Glutamine Dumper gene expression in Arabidopsis roots. *J Integr Plant Biol* **56**, 684-694.

Mangano, S., Juárez, S.P., and Estevez, J.M. (2016). ROS Regulation of Polar Growth in Plant Cells. *Plant Physiol* **171**, 1593-1605.

Martynov, V.I., Pakhomov, A.A., Deyev, I.E., and Petrenko, A.G. (2018). Genetically encoded fluorescent indicators for live cell pH imaging. *Biochim Biophys Acta Gen Subj* **1862**, 2924-2939.

Michard, E., Simon, A.A., Tavares, B., Wudick, M.M., and Feijó, J.A. (2017). Signaling with Ions: The Keystone for Apical Cell Growth and Morphogenesis in Pollen Tubes. *Plant Physiol* **173**, 91-111.

Mishina, N.M., Bogeski, I., Bolotin, D.A., Hoth, M., Niemeyer, B.A., Schultz, C., Zagaynova, E.V., Lukyanov, S., and Belousov, V.V. (2012). Can we see PIP(3) and hydrogen peroxide with a single probe? *Antioxid Redox Signal* **17**, 505-512.

Mittler, R. (2017). ROS Are Good. *Trends Plant Sci* **22**, 11-19.

Monshausen, G.B., Bibikova, T.N., Messerli, M.A., Shi, C., and Gilroy, S. (2007). Oscillations in extracellular pH and reactive oxygen species modulate tip growth of Arabidopsis root hairs. *Proc Natl Acad Sci U S A* **104**, 20996-21001.

Monshausen, G.B., Bibikova, T.N., Weisenseel, M.H., and Gilroy, S. (2009). Ca^{2+} regulates reactive oxygen species production and pH during mechanosensing in Arabidopsis roots. *Plant Cell* **21**, 2341-2356.

Monshausen, G.B., Miller, N.D., Murphy, A.S., and Gilroy, S. (2011). Dynamics of auxin-dependent Ca^{2+} and pH signaling in root growth revealed by integrating high-resolution imaging with automated computer vision-based analysis. *Plant J* **65**, 309-318.

Mori, I.C., Murata, Y., Yang, Y., Munemasa, S., Wang, Y.F., Andreoli, S., Tiriack, H., Alonso, J.M., Harper, J.F., Ecker, J.R., Kwak, J.M., and Schroeder, J.I. (2006). CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca^{2+} -permeable channels and stomatal closure. *PLoS Biol* **4**, e327.

- Munemasa, S., Hauser, F., Park, J., Waadt, R., Brandt, B., and Schroeder, J.I.** (2015). Mechanisms of abscisic acid-mediated control of stomatal aperture. *Curr Opin Plant Biol* **28**, 154-162.
- Nagaya, S., Kawamura, K., Shinmyo, A., and Kato, K.** (2010). The HSP terminator of *Arabidopsis thaliana* increases gene expression in plant cells. *Plant Cell Physiol* **51**, 328-332.
- Ngo, Q.A., Vogler, H., Lituiev, D.S., Nestorova, A., and Grossniklaus, U.** (2014). A calcium dialog mediated by the FERONIA signal transduction pathway controls plant sperm delivery. *Dev Cell* **29**, 491-500.
- Nietzel, T., Elsässer, M., Ruberti, C., Steinbeck, J., Ugalde, J.M., Fuchs, P., Wagner, S., Ostermann, L., Moseler, A., Lemke, P., Fricker, M.D., Müller-Schüssele, S.J., Moerschbacher, B.M., Costa, A., Meyer, A.J., and Schwarzländer, M.** (2019). The fluorescent protein sensor roGFP2-Orp1 monitors *in vivo* H₂O₂ and thiol redox integration and elucidates intracellular H₂O₂ dynamics during elicitor-induced oxidative burst in Arabidopsis. *New Phytol* **221**, 1649-1664.
- Ogasawara, Y., Kaya, H., Hiraoka, G., Yumoto, F., Kimura, S., Kadota, Y., Hishinuma, H., Senzaki, E., Yamagoe, S., Nagata, K., Nara, M., Suzuki, K., Tanokura, M., and Kuchitsu, K.** (2008). Synergistic activation of the Arabidopsis NADPH oxidase AtrbohD by Ca²⁺ and phosphorylation. *J Biol Chem* **283**, 8885-8892.
- Okumoto, S., Jones, A., and Frommer, W.B.** (2012). Quantitative imaging with fluorescent biosensors. *Annu Rev Plant Biol* **63**, 663-706.
- Peaucelle, A., Braybrook, S., and Höfte, H.** (2012). Cell wall mechanics and growth control in plants: the role of pectins revisited. *Front Plant Sci* **3**, 121.
- Pei, Z.M., Murata, Y., Benning, G., Thomine, S., Klüsener, B., Allen, G.J., Grill, E., and Schroeder, J.I.** (2000). Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* **406**, 731-734.
- Pittman, J.K., and Hirschi, K.D.** (2016). CAX-ing a wide net: Cation/H(+) transporters in metal remediation and abiotic stress signalling. *Plant Biol (Stuttg)* **18**, 741-749.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J.,**

- Hartenstein, V., Eliceiri, K., Tomancak, P., and Cardona, A.** (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682.
- Schmidt, C., Schelle, I., Liao, Y.J., and Schroeder, J.I.** (1995). Strong regulation of slow anion channels and abscisic acid signaling in guard cells by phosphorylation and dephosphorylation events. *Proc Natl Acad Sci U S A* **92**, 9535-9539.
- Schumacher, K.** (2014). pH in the plant endomembrane system - an import and export business. *Curr Opin Plant Biol* **22**, 71-76.
- Schwarzländer, M., Logan, D.C., Johnston, I.G., Jones, N.S., Meyer, A.J., Fricker, M.D., and Sweetlove, L.J.** (2012). Pulsing of membrane potential in individual mitochondria: a stress-induced mechanism to regulate respiratory bioenergetics in *Arabidopsis*. *Plant Cell* **24**, 1188-1201.
- Seybold, H., Trempel, F., Ranf, S., Scheel, D., Romeis, T., and Lee, J.** (2014). Ca^{2+} signalling in plant immune response: from pattern recognition receptors to Ca^{2+} decoding mechanisms. *New Phytol* **204**, 782-790.
- Shan, X., Yan, J., and Xie, D.** (2012). Comparison of phytohormone signaling mechanisms. *Curr Opin Plant Biol* **15**, 84-91.
- Shavrukov, Y., and Hirai, Y.** (2016). Good and bad protons: genetic aspects of acidity stress responses in plants. *J Exp Bot* **67**, 15-30.
- Shen, Y., Rosendale, M., Campbell, R.E., and Perrais, D.** (2014). pHuji, a pH-sensitive red fluorescent protein for imaging of exo- and endocytosis. *J Cell Biol* **207**, 419-432.
- Shih, H.W., Miller, N.D., Dai, C., Spalding, E.P., and Monshausen, G.B.** (2014). The receptor-like kinase FERONIA is required for mechanical signal transduction in *Arabidopsis* seedlings. *Curr Biol* **24**, 1887-1892.
- Shih, H.W., DePew, C.L., Miller, N.D., and Monshausen, G.B.** (2015). The Cyclic Nucleotide-Gated Channel CNGC14 Regulates Root Gravitropism in *Arabidopsis thaliana*. *Curr Biol* **25**, 3119-3125.
- Smirnoff, N., and Arnaud, D.** (2019). Hydrogen peroxide metabolism and functions in plants. *New Phytol* **221**, 1197-1214.
- Song, C.J., Steinebrunner, I., Wang, X., Stout, S.C., and Roux, S.J.** (2006). Extracellular ATP induces the accumulation of superoxide via NADPH oxidases in

Arabidopsis. *Plant Physiol* **140**, 1222-1232.

Stegmann, M., Monaghan, J., Smakowska-Luzan, E., Rovenich, H., Lehner, A., Holton, N., Belkhadir, Y., and Zipfel, C. (2017). The receptor kinase FER is a RALF-regulated scaffold controlling plant immune signaling. *Science* **355**, 287-289.

Steinhorst, L., and Kudla, J. (2014). Signaling in cells and organisms - calcium holds the line. *Curr Opin Plant Biol* **22**, 14-21.

Sze, H., and Chanroj, S. (2018). Plant Endomembrane Dynamics: Studies of K⁺/H⁺ Antiporters Provide Insights on the Effects of pH and Ion Homeostasis. *Plant Physiol* **177**, 875-895.

Takemoto, K., Ebine, K., Askani, J.C., Krüger, F., Gonzalez, Z.A., Ito, E., Goh, T., Schumacher, K., Nakano, A., and Ueda, T. (2018). Distinct sets of tethering complexes, SNARE complexes, and Rab GTPases mediate membrane fusion at the vacuole in Arabidopsis. *Proc Natl Acad Sci U S A* **115**, E2457-E2466.

Tanaka, K., Swanson, S.J., Gilroy, S., and Stacey, G. (2010). Extracellular nucleotides elicit cytosolic free calcium oscillations in Arabidopsis. *Plant Physiol* **154**, 705-719.

Tian, L., Hires, S.A., Mao, T., Huber, D., Chiappe, M.E., Chalasani, S.H., Petreanu, L., Akerboom, J., McKinney, S.A., Schreiter, E.R., Bargmann, C.I., Jayaraman, V., Svoboda, K., and Looger, L.L. (2009). Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators. *Nat Methods* **6**, 875-881.

Tian, W., Hou, C., Ren, Z., Wang, C., Zhao, F., Dahlbeck, D., Hu, S., Zhang, L., Niu, Q., Li, L., Staskawicz, B.J., and Luan, S. (2019). A calmodulin-gated calcium channel links pathogen patterns to plant immunity. *Nature* **572**, 131-135.

Vanstraelen, M., and Benková, E. (2012). Hormonal interactions in the regulation of plant development. *Annu Rev Cell Dev Biol* **28**, 463-487.

Waadt, R., Hitomi, K., Nishimura, N., Hitomi, C., Adams, S.R., Getzoff, E.D., and Schroeder, J.I. (2014). FRET-based reporters for the direct visualization of abscisic acid concentration changes and distribution in Arabidopsis. *Elife* **3**, e01739.

Waadt, R., Hsu, P.K., and Schroeder, J.I. (2015). Absciscic acid and other plant hormones: Methods to visualize distribution and signaling. *Bioessays* **37**, 1338-1349.

- Waadt, R., Krebs, M., Kudla, J., and Schumacher, K.** (2017). Multiparameter imaging of calcium and abscisic acid and high-resolution quantitative calcium measurements using R-GECO1-mTurquoise in Arabidopsis. *New Phytol* **216**, 303-320.
- Wagner, S., Steinbeck, J., Fuchs, P., Lichtenauer, S., Elsässer, M., Schippers, J.H.M., Nietzel, T., Ruberti, C., Van Aken, O., Meyer, A.J., Van Dongen, J.T., Schmidt, R.R., and Schwarzländer, M.** (2019). Multiparametric real-time sensing of cytosolic physiology links hypoxia responses to mitochondrial electron transport. *New Phytol* **224**, 1668-1684.
- Walia, A., Waadt, R., and Jones, A.M.** (2018). Genetically Encoded Biosensors in Plants: Pathways to Discovery. *Annu Rev Plant Biol* **69**, 497-524.
- Waszczak, C., Carmody, M., and Kangasjärvi, J.** (2018). Reactive Oxygen Species in Plant Signaling. *Annu Rev Plant Biol* **69**, 209-236.
- Yoshida, T., Christmann, A., Yamaguchi-Shinozaki, K., Grill, E., and Fernie, A.R.** (2019). Revisiting the Basal Role of ABA - Roles Outside of Stress. *Trends Plant Sci* **24**, 625-635.
- Zhang, X., Köster, P., Schlücking, K., Balcerowicz, D., Hashimoto, K., Kuchitsu, K., Vissenberg, K., and Kudla, J.** (2018). CBL1-CIPK26-mediated phosphorylation enhances activity of the NADPH oxidase RBOHC, but is dispensable for root hair growth. *FEBS Lett* **592**, 2582-2593.
- Zhao, Y., Araki, S., Wu, J., Teramoto, T., Chang, Y.F., Nakano, M., Abdelfattah, A.S., Fujiwara, M., Ishihara, T., Nagai, T., and Campbell, R.E.** (2011). An expanded palette of genetically encoded Ca²⁺ indicators. *Science* **333**, 1888-1891.
- Zheng, X., Kang, S., Jing, Y., Ren, Z., Li, L., Zhou, J.M., Berkowitz, G., Shi, J., Fu, A., Lan, W., Zhao, F., and Luan, S.** (2018). Danger-Associated Peptides Close Stomata by OST1-Independent Activation of Anion Channels in Guard Cells. *Plant Cell* **30**, 1132-1146.

FIGURE LEGENDS

Figure 1. Development of ABAleonSD1-3L21. **(A)** FRET-pair and sensory domain, and **(B)** linker screening of ABA indicator variants after expression in HEK293T cells. Shown are emission ratio changes in response to 60 min treatments with 0 and 100 μ M ABA. Reference indicators are shown in cyan and new candidates in red. Information on ABA indicator topologies is given in Supplemental Figure 1. **(C to E)** Representative ABA-dependent normalized in vitro emission spectra of ABAleons with indicated maximum emission ratio change ($\Delta R_{\text{max}}/R_0$). **(F)** ABA-dependent in vitro emission ratios and apparent ABA affinities (k'_d) of ABAleons. **(G and H)** Comparison of ABA-dependent ABAleon emission ratios: **(G)** in vitro and **(H)** in HEK293T cells. **(I)** In vitro and HEK293T cell comparison of ABA-induced maximum emission ratio change. All data are shown as mean $\pm$ SD, $n = 3$.

Figure 2. Comparison of ABA indicator ABA responses in Arabidopsis. Five-day-old roots of Arabidopsis expressing **(A)** ABAleon2.15 ($n = 9$), **(B)** ABAleonSD1-3L21 ($n = 8$) and **(C)** ABACUS1-2 μ ($n = 6$) were imaged for 64 min at a frame rate of 10 min^{-1} and treated with 10 μ M ABA at $t = 0$ min. Shown are average vertical response profiles of (top) emission ratios (R) and (middle) emission ratio changes ($\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image average emission ratio changes (mean $\pm$ SD) with indicated half response times ($t_{1/2}$). A representative experiment is provided as Supplemental Movie 1. Data of 0 μ M ABA control experiments are shown in Supplemental Figure 2.

Figure 3. Application of ABA does not trigger rapid changes of cytosolic Ca^{2+} or pH in Arabidopsis roots. Analyses of five-day-old roots of Arabidopsis expressing **(A)** ABAleonSD1-3L21-P2A-R-GECO1 (ABA and Ca^{2+} ; $n = 5$) in response to 10 μ M ABA ($t = 0$ min) and 1 μ M IAA ($t = 30$ min), and **(B)** PA-17-P2A-ABALEONSD1-3L21 (pH and ABA; $n = 8$) in response to 10 μ M ABA ($t = 0$ min) and 1 mM ATP ($t = 30$ min). Images were acquired for 64 min at a frame rate of 10 min^{-1} . Shown are average vertical response profiles of (top) emission ratios (R) or fluorescence emissions (F) and (middle) signal changes ($\Delta R/R_0$ or $\Delta F/F_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom)

Full image signal changes (mean $\pm$ SD). Representative experiments are provided as Supplemental Movies 2 and 3.

Figure 4. Application of auxin and ATP triggers cytosolic Ca^{2+} , H^+ and Cl^- fluxes. Analyses of five-day-old roots of Arabidopsis expressing R-GECO1-GSL-E²GFP (Ca^{2+} , pH and Cl^-) in response to 1 μM IAA ($t = 0$ min) and 1 mM ATP ($t = 40$ min; $n = 6$). Images were acquired for 84 min at a frame rate of 10 min^{-1} . **(A)** Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R), (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings, and (bottom) full image signal changes (mean $\pm$ SD). **(B)** Spatiotemporal Pearson correlation analyses (mean $\pm$ SD) of indicated GEFI responses during the application of IAA ($t = -4$ -15 min). An adjacent representative bright field (BF) root image is shown for orientation. A representative experiment is provided as Supplemental Movie 4.

Figure 5. Application of glutamate triggers cytosolic Ca^{2+} , H^+ and Cl^- fluxes. Analyses of five-day-old roots of Arabidopsis expressing R-GECO1-GSL-E²GFP (Ca^{2+} , pH and Cl^-) in response to 1 mM glutamate (Glu; $t = 0$ min) and 1 mM ATP ($t = 30$ min; $n = 7$). Images were acquired for 64 min at a frame rate of 10 min^{-1} . **(A)** Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R), (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings, and (bottom) full image signal changes (mean $\pm$ SD). **(B)** Spatiotemporal Pearson correlation analyses (mean $\pm$ SD) of indicated GEFI responses during the application of glutamate (left; $t = -4$ -10 min) or ATP (right; $t = 26$ -60 min). An adjacent representative bright field (BF) root image is shown for orientation. A representative experiment is provided as Supplemental Movie 5. See also Supplemental Figures 3 and 4 for related experiments.

Figure 6. Application of Glutamate triggers a rapid cytosolic acidification without noticeable impact on the cytosolic redox state. Analyses of five-day-old roots of Arabidopsis expressing **(A)** PA-17-P2A-Grx1-roGFP2 (pH and E_{GSH} ; $n = 6$) and **(B)** PA-17-P2A-roGFP2-Orp1 (pH and H_2O_2 ; $n = 8$) in response to 1 mM glutamate (Glu; $t = 0$ min) and 100 μM H_2O_2 ($t = 30$ min). Images were acquired for 64 min at a frame rate of

10 min⁻¹. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Note that experiments in **(A)** and **(B)** were acquired at different magnifications. Representative experiments are provided as Supplemental Movies 6 and 7.

Figure 7. Application of H₂O₂ triggers overlapping but also distinct patterns of cytosolic oxidation and Ca²⁺ fluxes. Analyses of five-day-old roots of Arabidopsis expressing **(A)** R-GECO1-P2A-Grx1-roGFP2 (Ca²⁺ and E_{GSH}; n = 8) and **(B)** R-GECO1-P2A-roGFP2-Orp1 (Ca²⁺ and H₂O₂; n = 8) in response to 20 μ M H₂O₂ (t = 0 min) and 100 μ M H₂O₂ (t = 30 min). Images were acquired for 64 min at a frame rate of 10 min⁻¹. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Representative experiments are provided as Supplemental Movies 8 and 9.

Figure 8. Cytosolic oxidation is only weakly affected by ATP and PEP1. Analyses of five-day-old roots of Arabidopsis expressing **(A)** R-GECO1 and roGFP2-Orp1 (Ca²⁺ and H₂O₂; n = 7) in response to 1 mM ATP and 100 μ M H₂O₂, and **(B)** R-GECO1-P2A-roGFP2-Orp1 (Ca²⁺ and H₂O₂; n = 7) in response to 1 μ M PEP1 and 100 μ M H₂O₂. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Representative experiments are provided as Supplemental Movies 10 and 11. See also Supplemental Figures 6 and 7 for related experiments.

Figure 9. Application of PEP1 triggers Ca²⁺, H⁺ and Cl⁻ fluxes with high spatiotemporal overlap. **(A)** Analyses of five-day-old roots of Arabidopsis expressing R-GECO1-GSL-E²GFP (Ca²⁺, pH and Cl⁻; n = 6) in response to 1 μ M PEP1 (t = 0 min) and 1 mM ATP (t

= 30 min). Images were acquired for 64 min at a frame rate of 10 min⁻¹. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). **(B)** Spatiotemporal Pearson correlation analyses (mean $\pm$ SD) of indicated GEFI responses during the application of PEP1 (t = -4-20 min). A representative experiment is provided as Supplemental Movie 14.

Figure 10. GSSG-triggered Ca²⁺-, H⁺- and Cl⁻ influx precedes cytosolic oxidation. Analyses of five-day-old roots of Arabidopsis expressing **(A)** R-GECO1-P2A-Grx1-roGFP2 (Ca²⁺ and E_{GSH} ; n = 5) and **(B)** R-GECO1-GSL-E²GFP (Ca²⁺, pH and Cl⁻; n = 6) in response to 1 mM GSSG (t = 0 min). Images were acquired for 64 min at a frame rate of 10 min⁻¹. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). **(C)** Spatiotemporal Pearson correlation analyses (mean $\pm$ SD) of indicated GEFI responses during the application of GSSG (t = -4-20 min; data from **(B)**). Representative experiments are provided as Supplemental Movies 15 and 16.

Figure 11. Schematic model of observed ABA, Ca²⁺, H⁺, Cl⁻ and redox changes in roots. ABA treatment and uptake did not induce rapid fluxes of Ca²⁺ or H⁺. Whereas, IAA, glutamate (Glu), ATP, PEP1 and GSSG triggered Ca²⁺, H⁺ and Cl⁻ fluxes with high spatiotemporal overlap. For comparison of the obtained data see also Supplemental Figure 8. Compared to 20 μ M H₂O₂ and 1 mM GSSG, redox changes in response to glutamate, ATP and PEP1 were very low and below the threshold required to trigger ROS-induced Ca²⁺ signaling. Regions with highest response are color-coded according to the adjacent flux curves that were taken from the corresponding main figures (ABA, H₂O₂ and E_{GSH} , black, Ca²⁺, magenta; H⁺, yellow; Cl⁻, cyan). For a better illustration of H⁺ and Cl⁻ influx the PA-17 and E²GFP response curves were inverted. Arrows indicate the direction of the corresponding fluxes. RC, root cap; MEZ, meristematic zone; EZ, elongation zone; MZ, maturation zone.

Rainer Waadt et al. – Coordination of signaling compound fluxes in roots

1146

Figure 1. Development of ABAlleonSD1-3L21. **(A)** FRET-pair and sensory domain, and **(B)** linker screening of ABA indicator variants after expression in HEK293T cells. Shown are emission ratio changes in response to 60 min treatments with 0 and 100 μ M ABA. Reference indicators are shown in cyan and new candidates in red. Information on ABA indicator topologies is given in Supplemental Figure 1. **(C to E)** Representative ABA-dependent normalized in vitro emission spectra of ABAlleons with indicated maximum emission ratio change ($\Delta R_{(max)}/R_0$). **(F)** ABA-dependent in vitro emission ratios and apparent ABA affinities (k_d) of ABAlleons. **(G and H)** Comparison of ABA-dependent ABAlleon emission ratios: **(G)** in vitro and **(H)** in HEK293T cells. **(I)** In vitro and HEK293T cell comparison of ABA-induced maximum emission ratio change. All data are shown as mean $\pm$ SD, $n = 3$.

Figure 2. Comparison of ABA indicator ABA responses in Arabidopsis. Five-day-old roots of Arabidopsis expressing (A) ABAleon2.15 (n = 9), (B) ABAleonSD1-3L21 (n = 8) and (C) ABACUS1-2μ (n = 6) were imaged for 64 min at a frame rate of 10 min⁻¹ and treated with 10 μM ABA at t = 0 min. Shown are average vertical response profiles of (top) emission ratios (R) and (middle) emission ratio changes (ΔR/R₀) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image average emission ratio changes (mean ± SD) with indicated half response times (t_{1/2}). A representative experiment is provided as Supplemental Movie 1. Data of 0 μM ABA control experiments are shown in Supplemental Figure 2.

Figure 3. Application of ABA does not trigger rapid changes of cytosolic Ca^{2+} or pH in Arabidopsis roots. Analyses of five-day-old roots of Arabidopsis expressing **(A)** ABAleonSD1-3L21-P2A-R-GECO1 (ABA and Ca^{2+} ; $n = 5$) in response to 10 μM ABA ($t = 0$ min) and 1 μM IAA ($t = 30$ min), and **(B)** PA-17-P2A-ABALEonSD1-3L21 (pH and ABA; $n = 8$) in response to 10 μM ABA ($t = 0$ min) and 1 mM ATP ($t = 30$ min). Images were acquired for 64 min at a frame rate of 10 min^{-1} . Shown are average vertical response profiles of (top) emission ratios (R) or fluorescence emissions (F) and (middle) signal changes ($\Delta R/R_0$ or $\Delta F/F_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Representative experiments are provided as Supplemental Movies 2 and 3.

Figure 4. Application of auxin and ATP triggers cytosolic Ca^{2+} , H^+ and Cl^- fluxes. Analyses of five-day-old roots of *Arabidopsis* expressing R-GECO1-GSL-E²GFP (Ca^{2+} , pH and Cl^-) in response to 1 μM IAA (t = 0 min) and 1 mM ATP (t = 40 min; n = 6). Images were acquired for 84 min at a frame rate of 10 min⁻¹. **(A)** Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R), (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings, and (bottom) full image signal changes (mean $\pm$ SD). **(B)** Spatiotemporal Pearson correlation analyses (mean $\pm$ SD) of indicated GEFI responses during the application of IAA (t = -4-15 min). An adjacent representative bright field (BF) root image is shown for orientation. A representative experiment is provided as Supplemental Movie 4.

Figure 5. Application of glutamate triggers cytosolic Ca^{2+} , H^+ and Cl^- fluxes. Analyses of five-day-old roots of *Arabidopsis* expressing R-GECO1-GSL-E²GFP (Ca^{2+} , pH and Cl^-) in response to 1 mM glutamate (Glu; t = 0 min) and 1 mM ATP (t = 30 min; n = 7). Images were acquired for 64 min at a frame rate of 10 min⁻¹. **(A)** Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R), (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings, and (bottom) full image signal changes (mean $\pm$ SD). **(B)** Spatiotemporal Pearson correlation analyses (mean $\pm$ SD) of indicated GEF responses during the application of glutamate (left; t = -4-10 min) or ATP (right; t = 26-60 min). An adjacent representative bright field (BF) root image is shown for orientation. A representative experiment is provided as Supplemental Movie 5. See also Supplemental Figures 3 and 4 for related experiments.

Figure 6. Application of Glutamate triggers a rapid cytosolic acidification without noticeable impact on the cytosolic redox state. Analyses of five-day-old roots of *Arabidopsis* expressing **(A)** PA-17-P2A-Grx1-roGFP2 (pH and E_{GSH} ; $n = 6$) and **(B)** PA-17-P2A-roGFP2-Orp1 (pH and H_2O_2 ; $n = 8$) in response to 1 mM glutamate (Glu; $t = 0$ min) and 100 μM H_2O_2 ($t = 30$ min). Images were acquired for 64 min at a frame rate of 10 min⁻¹. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Note that experiments in **(A)** and **(B)** were acquired at different magnifications. Representative experiments are provided as Supplemental Movies 6 and 7.

Figure 7. Application of H_2O_2 triggers overlapping but also distinct patterns of cytosolic oxidation and Ca^{2+} fluxes. Analyses of five-day-old roots of *Arabidopsis* expressing **(A)** R-GECO1-P2A-Grx1-roGFP2 (Ca^{2+} and E_{GSH} ; $n = 8$) and **(B)** R-GECO1-P2A-roGFP2-Orp1 (Ca^{2+} and H_2O_2 ; $n = 8$) in response to 20 μM H_2O_2 ($t = 0$ min) and 100 μM H_2O_2 ($t = 30$ min). Images were acquired for 64 min at a frame rate of 10 min^{-1} . Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Representative experiments are provided as Supplemental Movies 8 and 9.

Figure 8. Cytosolic oxidation is only weakly affected by ATP and PEP1. Analyses of five-day-old roots of Arabidopsis expressing (A) R-GECO1 and roGFP2-Orp1 (Ca^{2+} and H_2O_2 ; $n = 7$) in response to 1 mM ATP and 100 μM H_2O_2 and (B) R-GECO1-P2A-roGFP2-Orp1 (Ca^{2+} and H_2O_2 ; $n = 7$) in response to 1 μM PEP1 and 100 μM H_2O_2 . Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Representative experiments are provided as Supplemental Movies 10 and 11. See also Supplemental Figures 6 and 7 for related experiments.

Figure 9. Application of PEP1 triggers Ca²⁺, H⁺ and Cl⁻ fluxes with high spatiotemporal overlap. **(A)** Analyses of five-day-old roots of Arabidopsis expressing R-GECO1-GSL-E²GFP (Ca²⁺, pH and Cl⁻; n = 6) in response to 1 μM PEP1 (t = 0 min) and 1 mM ATP (t = 30 min). Images were acquired for 64 min at a frame rate of 10 min⁻¹. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes (ΔF/F₀ or ΔR/R₀) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean ± SD). **(B)** Spatiotemporal Pearson correlation analyses (mean ± SD) of indicated GEFI responses during the application of PEP1 (t = -4-20 min). A representative experiments is provided as Supplemental Movie 14.

Figure 10. GSSG-triggered Ca^{2+} -, H^+ - and Cl^- influx precedes cytosolic oxidation. Analyses of five-day-old roots of *Arabidopsis* expressing **(A)** R-GECO1-P2A-Grx1-roGFP2 (Ca^{2+} and E_{GSH} ; $n = 5$) and **(B)** R-GECO1-GSL-E²GFP (Ca^{2+} , pH and Cl^- ; $n = 6$) in response to 1 mM GSSG ($t = 0$ min). Images were acquired for 64 min at a frame rate of 10 min^{-1} . Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). **(C)** Spatiotemporal Pearson correlation analyses (mean $\pm$ SD) of indicated GEFI responses during the application of GSSG ($t = -4$ -20 min; data from **(B)**). Representative experiments are provided as Supplemental Movies 15 and 16.

Figure 11. Schematic model of observed ABA, Ca^{2+} , H^+ , Cl^- and redox changes in roots. ABA treatment and uptake did not induce rapid fluxes of Ca^{2+} or H^+ . Whereas, IAA, glutamate (Glu), ATP, PEP1 and GSSG triggered Ca^{2+} , H^+ and Cl^- fluxes with high spatiotemporal overlap. For comparison of the obtained data see also Supplemental Figure 8. Compared to 20 μM H_2O_2 and 1 mM GSSG, redox changes in response to glutamate, ATP and PEP1 were very low and below the threshold required to trigger ROS-induced Ca^{2+} signaling. Regions with highest response are color-coded according to the adjacent flux curves that were taken from the corresponding main figures (ABA, H_2O_2 and E_{GSH} , black; Ca^{2+} , magenta; H^+ , yellow; Cl^- , cyan). For a better illustration of H^+ and Cl^- influx the PA-17 and $E^2\text{GFP}$ response curves were inverted. Arrows indicate the direction of the corresponding fluxes. RC, root cap; MEZ, meristematic zone; EZ, elongation zone; MZ, maturation zone.

Supplemental Figure 1. Topologies of ABA indicators. Indicated are fluorescent protein FRET-pairs (cyan and yellow), linkers (black), PYR1 and PYL1 moieties (green) and ABI1 moieties (orange). Point mutations and incorporated amino acids are given. **(A)** ABAleon2.15 and ABACUS1-2μ. **(B)** Non-responsive ABAleon2.15nr that contains two mutations PYR1_{K59Q} and ABI1_{W300A} to prevent ABA-binding. **(C)** ABAleon2.15 deletion variants. **(D)** FRET-pair and sensory domain (SD) variants. Arabidopsis codon-optimized (P)mTurquoise, PmTurquoise_{Δ11} with a C-terminal deletion of 11 amino acids or enhanced dimeric PedTurquoise were used as FRET donor. Venus, circularly permuted Venus (cpVenus173) or enhanced dimeric (ed)Venus were used as FRET acceptor. The sensory domains SD1 and SD3 derived from ABAleon2.15, with SD3 harboring a longer linker between the PYR1 and ABI1 moieties. SD2 derived from ABACUS1-2μ. **(E)** ABAleonSD1-3 linker variants. **(D and E)** Red color indicates the topology of ABAleonSD1-3 **(D)** and the linkers in ABAleonSD1-3L21 **(E)**.

Supplemental Figure 2. Solvent control experiments of ABA indicators in Arabidopsis. Five-day-old roots of Arabidopsis expressing (A) ABAleon2.15 (n = 6), (B) ABAleonSD1-3L21 (n = 7) and (C) ABACUS1-2μ (n = 6) were imaged for 64 min at a frame rate of 10 min⁻¹ and treated with 0.01 % EtOH (solvent control for ABA) at t = 0 min. Shown are average vertical response profiles of (top) emission ratios (R) and (middle) emission ratio changes (ΔR/R₀) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image average emission ratio changes (mean ± SD). Color scales of response profiles are identical to the scales in Figure 2.

Supplemental Figure 3. Targeting of E²GFP to the plasma membrane. **(A)** Subcellular localization of R-GECO1-P2A-E²GFP-LTI6b fluorescence emission in epidermis cells of three-week-old Arabidopsis leaves. **(B)** Analyses of five-day-old roots of Arabidopsis expressing R-GECO1-P2A-E²GFP-LTI6b in response to 1 mM glutamate (Glu; t = 0 min) and 1 mM ATP (t = 30 min; n = 8). Images were acquired for 64 min at a frame rate of 10 min⁻¹. Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). These data are related to experiments presented in Figure 5.

Supplemental Figure 5. Targeting of E²GFP to the plasma membrane induces plant growth defects. **(A)** Representative images of 28-day-old *Arabidopsis* lines expressing the indicated 2-In-1-GEFIs. **(B)** Rosette area quantification of 28-day-old 2-In-1-GEFI lines (square dots, mean; central lines in boxes, median; boxes, 25th and 75th percentiles, whiskers, $\pm$ 1.5 interquartile range; diamond dots, outliers). Asterisks indicate statistically significant differences relative to Col-0 wild type in pairwise Tukey test comparisons ($p < 0.05$; $n = 9-12$). These data are related to Figure 5 and Supplemental Figures 3 and 4.

Supplemental Figure 6. Cytosolic E_{GSH} is only weakly affected by ATP and PEP1 treatments. Analyses of five-day-old roots of Arabidopsis expressing **(A)** R-GECO1 and Grx1-roGFP2 (Ca^{2+} and E_{GSH} ; $n = 8$) in response to 1 mM ATP and 100 μM H_2O_2 , and **(B)** R-GECO1-P2A-Grx1-roGFP2 (Ca^{2+} and E_{GSH} ; $n = 7$) in response to 1 μM PEP1 and 100 μM H_2O_2 . Average vertical response profiles of (top) fluorescence emissions (F) or emission ratios (R) and (middle) signal changes ($\Delta F/F_0$ or $\Delta R/R_0$) normalized to 4 min average baseline recordings. An adjacent representative bright field (BF) root image is shown for orientation. (bottom) Full image signal changes (mean $\pm$ SD). Representative experiments are provided as Supplemental Movies 12 and 13. These data are related to experiments presented in Figure 8.

Supplemental Figure 7. Glutamate-, ATP- and PEP1-dependent cytosolic oxidation is below the threshold of ROS-induced Ca^{2+} signaling. 30 min signal change ($\Delta F/F_0$ or $\Delta R/R_0$) vertical response profiles of **(A)** R-GECO1 (Ca^{2+}), **(B)** roGFP2-Orp1 (H_2O_2) and **(C)** Grx1-roGFP2 (E_{GSH}) after treatments with 20 μM H_2O_2 , 1 mM glutamate (Glu), 1 mM ATP and 1 μM PEP1. These data were taken from analyses presented in Figure 7 (H_2O_2), Figures 5A and 6B (Glu) and Figure 8 and Supplemental Figure 6 (ATP and PEP1) and scaled to the adjacent color scale.

Supplemental Figure 8. Ca^{2+} , H^+ and anion fluxes exhibit a high spatiotemporal overlap. 30 min signal change ($\Delta F/F_0$ or $\Delta R/R_0$) vertical response profiles of (A) R-GECO1 (Ca^{2+}), (B) E²GFP (H^+) and (C) E²GFP (Cl^- /anions) after treatments with 1 μ M IAA, 1 mM glutamate (Glu), 1 mM ATP, 1 μ M PEP1 and 1 mM GSSG. These data were taken from analyses presented in Figure 4 (IAA), Figure 5 (Glu and ATP) Figure 9 (PEP1) and Figure 10 (GSSG) and scaled to the adjacent color scale.

Parsed Citations

Allen, G.J., Chu, S.P., Harrington, C.L., Schumacher, K., Hoffmann, T., Tang, Y.Y., Grill, E., and Schroeder, J.I. (2001). A defined range of guard cell calcium oscillation parameters encodes stomatal movements. *Nature* 411, 1053-1057.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Arosio, D., Ricci, F., Marchetti, L., Gualdani, R., Albertazzi, L., and Beltram, F. (2010). Simultaneous intracellular chloride and pH measurements using a GFP-based sensor. *Nat Methods* 7, 516-518.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Barragán, V., Leidi, E.O., Andrés, Z., Rubio, L., De Luca, A., Fernández, J.A., Cubero, B., and Pardo, J.M. (2012). Ion exchangers NHX1 and NHX2 mediate active potassium uptake into vacuoles to regulate cell turgor and stomatal function in *Arabidopsis*. *Plant Cell* 24, 1127-1142.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Behera, S., Zhaolong, X., Luoni, L., Bonza, M.C., Doccua, F.G., De Michelis, M.I., Morris, R.J., Schwarzländer, M., and Costa, A. (2018). Cellular Ca²⁺ Signals Generate Defined pH Signatures in Plants. *Plant Cell* 30, 2704-2719.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Bilan, D.S., and Belousov, V.V. (2017). New tools for redox biology: From imaging to manipulation. *Free Radic Biol Med* 109, 167-188.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Bizzarri, R., Arcangeli, C., Arosio, D., Ricci, F., Faraci, P., Cardarelli, F., and Beltram, F. (2006). Development of a novel GFP-based ratiometric excitation and emission pH indicator for intracellular studies. *Biophys J* 90, 3300-3314.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Bonza, M.C., and De Michelis, M.I. (2011). The plant Ca²⁺-ATPase repertoire: biochemical features and physiological functions. *Plant Biol (Stuttg)* 13, 421-430.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Burén, S., Ortega-Villasante, C., Otvös, K., Samuelsson, G., Bakó, L., and Villarejo, A. (2012). Use of the foot-and-mouth disease virus 2A peptide co-expression system to study intracellular protein trafficking in *Arabidopsis*. *PLoS One* 7, e51973.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Chen, D., Cao, Y., Li, H., Kim, D., Ahsan, N., Thelen, J., and Stacey, G. (2017). Extracellular ATP elicits DORN1-mediated RBOHD phosphorylation to regulate stomatal aperture. *Nat Commun* 8, 2265.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Choi, J., Tanaka, K., Cao, Y., Qi, Y., Qiu, J., Liang, Y., Lee, S.Y., and Stacey, G. (2014). Identification of a plant receptor for extracellular ATP. *Science* 343, 290-294.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Choi, W.G., Hilleary, R., Swanson, S.J., Kim, S.H., and Gilroy, S. (2016). Rapid, Long-Distance Electrical and Calcium Signaling in Plants. *Annu Rev Plant Biol* 67, 287-307.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16, 735-743.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Cutler, S.R., Rodriguez, P.L., Finkelstein, R.R., and Abrams, S.R. (2010). Absciscic acid: emergence of a core signaling network. *Annu Rev Plant Biol* 61, 651-679.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

De Vylder, J., Vandenbussche, F., Hu, Y., Philips, W., and Van Der Straeten, D. (2012). Rosette tracker: an open source image analysis tool for automatic quantification of genotype effects. *Plant Physiol* 160, 1149-1159.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Dermidchik, V., Shabala, S.N., and Davies, J.M. (2007). Spatial variation in H₂O₂ response of *Arabidopsis thaliana* root epidermal Ca²⁺ flux and plasma membrane Ca²⁺ channels. *Plant J* 49, 377-386.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Dennidchik, V., Shang, Z., Shin, R., Thompson, E., Rubio, L., Laohavisit, A., Mortimer, J.C., Chivasa, S., Slabas, A.R., Glover, B.J., Schachtman, D.P., Shabala, S.N., and Davies, J.M. (2009). Plant extracellular ATP signalling by plasma membrane NADPH oxidase and Ca²⁺ channels. *Plant J* 58, 903-913.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Dindas, J., Scherzer, S., Roelfsema, M.R.G., von Meyer, K., Müller, H.M., Al-Rasheid, K.A.S., Palme, K., Dietrich, P., Becker, D., Bennett, M.J., and Hedrich, R. (2018). AUX1-mediated root hair auxin influx governs SCFTIR1/AFB-type Ca²⁺ signaling. *Nat Commun* 9, 1174.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Dodd, A.N., Kudla, J., and Sanders, D. (2010). The language of calcium signaling. *Annu Rev Plant Biol* 61, 593-620.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Edel, K.H., and Kudla, J. (2016). Integration of calcium and ABA signaling. *Curr Opin Plant Biol* 33, 83-91.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Evans, M.J., Choi, W.G., Gilroy, S., and Morris, R.J. (2016). AROS-Assisted Calcium Wave Dependent on the AtRBOHD NADPH Oxidase and TPC1 Cation Channel Propagates the Systemic Response to Salt Stress. *Plant Physiol* 171, 1771-1784.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Fendrych, M., Leung, J., and Friml, J. (2016). TIR1/AFB-Aux/IAA auxin perception mediates rapid cell wall acidification and growth of *Arabidopsis hypocotyls*. *Elife* 5, e19048.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Finkelstein, R. (2013). Absciscic Acid synthesis and response. *Arabidopsis Book* 11, e0166.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Gao, Q.F., Gu, L.L., Wang, H.Q., Fei, C.F., Fang, X., Hussain, J., Sun, S.J., Dong, J.Y., Liu, H., and Wang, Y.F. (2016). Cyclic nucleotide-gated channel 18 is an essential Ca²⁺ channel in pollen tube tips for pollen tube guidance to ovules in *Arabidopsis*. *Proc Natl Acad Sci U S A* 113, 3096-3101.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Gilroy, S., Suzuki, N., Miller, G., Choi, W.G., Toyota, M., Devireddy, A.R., and Mittler, R. (2014). A tidal wave of signals: calcium and ROS at the forefront of rapid systemic signaling. *Trends Plant Sci* 19, 623-630.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Gomez, L.D., Noctor, G., Knight, M.R., and Foyer, C.H. (2004). Regulation of calcium signalling and gene expression by glutathione. *J Exp Bot* 55, 1851-1859.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Grossmann, G., Krebs, M., Maizel, A., Stahl, Y., Vermeer, J.E.M., and Ott, T. (2018). Green light for quantitative live-cell imaging in plants. *J Cell Sci* 131.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Gutscher, M., Pauleau, A.L., Marty, L., Brach, T., Wabnitz, G.H., Samstag, Y., Meyer, A.J., and Dick, T.P. (2008). Real-time imaging of the intracellular glutathione redox potential. *Nat Methods* 5, 553-559.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Gutscher, M., Sobotta, M.C., Wabnitz, G.H., Ballikaya, S., Meyer, A.J., Samstag, Y., and Dick, T.P. (2009). Proximity-based protein thiol oxidation by H₂O₂-scavenging peroxidases. *J Biol Chem* 284, 31532-31540.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Han, J.P., Köster, P., Drerup, M.M., Scholz, M., Li, S., Edel, K.H., Hashimoto, K., Kuchitsu, K., Hippler, M., and Kudla, J. (2019). Fine tuning of RBOHF activity is achieved by differential phosphorylation and Ca²⁺ binding. *New Phytol* 221, 1935-1949.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Haruta, M., Sabat, G., Stecker, K., Minkoff, B.B., and Sussman, M.R. (2014). A peptide hormone and its receptor protein kinase regulate plant cell expansion. *Science* 343, 408-411.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haruta, M., Gray, W.M., and Sussman, M.R. (2015). Regulation of the plasma membrane proton pump (H(+)-ATPase) by phosphorylation. Curr Opin Plant Biol 28, 68-75.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hauser, F., Waadt, R., and Schroeder, J.I. (2011). Evolution of abscisic acid synthesis and signaling mechanisms. Curr Biol 21, R346-355.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hauser, F., Li, Z., Waadt, R., and Schroeder, J.I. (2017). SnapShot: Absciscic Acid Signaling. Cell 171, 1708-1708.e1700.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hedrich, R., Busch, H., and Raschke, K. (1990). Ca²⁺ and nucleotide dependent regulation of voltage dependent anion channels in the plasma membrane of guard cells. EMBO J 9, 3889-3892.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Heppler, P.K., and Winship, L.J. (2010). Calcium at the cell wall-cytoplasm interface. J Integr Plant Biol 52, 147-160.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hilleary, R., Choi, W.G., Kim, S.H., Lim, S.D., and Gilroy, S. (2018). Sense and sensibility: the use of fluorescent protein-based genetically encoded biosensors in plants. Curr Opin Plant Biol 46, 32-38.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hochreiter, B., Garcia, A.P., and Schmid, J.A. (2015). Fluorescent proteins as genetically encoded FRET biosensors in life sciences. Sensors (Basel) 15, 26281-26314.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W., and Zimmermann, P. (2008). Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. Adv Bioinformatics 2008, 420747.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Inoue, S.I., Takahashi, K., Okumura-Noda, H., and Kinoshita, T. (2016). Auxin Influx Carrier AUX1 Confers Acid Resistance for Arabidopsis Root Elongation Through the Regulation of Plasma Membrane H⁺-ATPase. Plant Cell Physiol 57, 2194-2201.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jones, A.M., Danielson, J.A., Manojkumar, S.N., Lanquar, V., Grossmann, G., and Frommer, W.B. (2014). Absciscic acid dynamics in roots detected with genetically encoded FRET sensors. Elife 3, e01741.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kadota, Y., Sklenar, J., Derbyshire, P., Stransfeld, L., Asai, S., Ntoukakis, V., Jones, J.D., Shirasu, K., Menke, F., Jones, A., and Zipfel, C. (2014). Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase BIK1 during plant immunity. Mol Cell 54, 43-55.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kadota, Y., Shirasu, K., and Zipfel, C. (2015). Regulation of the NADPH Oxidase RBOHD During Plant Immunity. Plant Cell Physiol 56, 1472-1480.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kärkönen, A., and Kuchitsu, K. (2015). Reactive oxygen species in cell wall metabolism and development in plants. Phytochemistry 112, 22-32.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Keinath, N.F., Waadt, R., Brugman, R., Schroeder, J.I., Grossmann, G., Schumacher, K., and Krebs, M. (2015). Live Cell Imaging with R-GECO1 Sheds Light on flg22- and Chitin-Induced Transient [Ca²⁺]_{cyt} Patterns in Arabidopsis. Mol Plant 8, 1188-1200.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kelner, A., Leitão, N., Chabaud, M., Charpentier, M., and de Carvalho-Niebel, F. (2018). Dual Color Sensors for Simultaneous Analysis of Calcium Signal Dynamics in the Nuclear and Cytoplasmic Compartments of Plant Cells. Front Plant Sci 9, 245.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kim, J.H., Lee, S.R., Li, L.H., Park, H.J., Park, J.H., Lee, K.Y., Kim, M.K., Shin, B.A., and Choi, S.Y. (2011). High cleavage efficiency of a 2Apeptide derived from porcine teschovirus-1 in human cell lines, zebrafish and mice. PLoS One 6, e18556.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kimura, S., Waszczak, C., Hunter, K., and Wrzaczek, M. (2017). Bound by Fate: The Role of Reactive Oxygen Species in Receptor-Like Kinase Signaling. Plant Cell 29, 638-654.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Köhler, B., Hills, A., and Blatt, M.R. (2003). Control of guard cell ion channels by hydrogen peroxide and abscisic acid indicates their action through alternate signaling pathways. Plant Physiol 131, 385-388.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kostyuk, A.I., Panova, A.S., Bilan, D.S., and Belousov, V.V. (2018). Redox biosensors in a context of multiparameter imaging. Free Radic Biol Med 128, 23-39.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kudla, J., Batistic, O., and Hashimoto, K. (2010). Calcium signals: the lead currency of plant information processing. Plant Cell 22, 541-563.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kwak, J.M., Mori, I.C., Pei, Z.M., Leonhardt, N., Torres, M.A., Dangl, J.L., Bloom, R.E., Bodde, S., Jones, J.D., and Schroeder, J.I. (2003). NADPH oxidase *AtrbohD* and *AtrbohF* genes function in ROS-dependent ABA signaling in Arabidopsis. EMBO J 22, 2623-2633.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lampropoulos, A., Sutikovic, Z., Wenzl, C., Maegele, I., Lohmann, J.U., and Forner, J. (2013). GreenGate—a novel, versatile, and efficient cloning system for plant transgenesis. PLoS One 8, e83043.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Li, L., Li, M., Yu, L., Zhou, Z., Liang, X., Liu, Z., Cai, G., Gao, L., Zhang, X., Wang, Y., Chen, S., and Zhou, J.M. (2014). The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD to control plant immunity. Cell Host Microbe 15, 329-338.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Liu, Z., Wu, Y., Yang, F., Zhang, Y., Chen, S., Xie, Q., Tian, X., and Zhou, J.M. (2013). BIK1 interacts with PEPRs to mediate ethylene-induced immunity. Proc Natl Acad Sci U S A 110, 6205-6210.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Loro, G., Drago, I., Pozzan, T., Schiavo, F.L., Zottini, M., and Costa, A. (2012). Targeting of Cameleons to various subcellular compartments reveals a strict cytoplasmic/mitochondrial Ca²⁺ handling relationship in plant cells. Plant J 71, 1-13.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ma, C., Guo, J., Kang, Y., Doman, K., Bryan, A.C., Tax, F.E., Yamaguchi, Y., and Qi, Z. (2014). *At*PEPTIDE RECEPTOR2 mediates the *At*PEPTIDE1-induced cytosolic Ca(2+) rise, which is required for the suppression of Glutamine Dumper gene expression in Arabidopsis roots. J Integr Plant Biol 56, 684-694.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Mangano, S., Juárez, S.P., and Estevez, J.M. (2016). ROS Regulation of Polar Growth in Plant Cells. Plant Physiol 171, 1593-1605.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Martynov, V.I., Pakhomov, A.A., Deyev, I.E., and Petrenko, A.G. (2018). Genetically encoded fluorescent indicators for live cell pH imaging. Biochim Biophys Acta Gen Subj 1862, 2924-2939.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Michard, E., Simon, A.A., Tavares, B., Wudick, M.M., and Feijó, J.A. (2017). Signaling with Ions: The Keystone for Apical Cell Growth and Morphogenesis in Pollen Tubes. Plant Physiol 173, 91-111.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Mishina, N.M., Bogeski, I., Bolotin, D.A., Hoth, M., Niemeyer, B.A., Schultz, C., Zagaynova, E.V., Lukyanov, S., and Belousov, V.V. (2012). Can we see PIP(3) and hydrogen peroxide with a single probe? Antioxid Redox Signal 17, 505-512.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Mittler, R. (2017). ROS Are Good. Trends Plant Sci 22, 11-19.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Monshausen, G.B., Bibikova, T.N., Messerli, M.A., Shi, C., and Gilroy, S. (2007). Oscillations in extracellular pH and reactive oxygen species modulate tip growth of Arabidopsis root hairs. Proc Natl Acad Sci U S A 104, 20996-21001.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Monshausen, G.B., Bibikova, T.N., Weisenseel, M.H., and Gilroy, S. (2009). Ca²⁺ regulates reactive oxygen species production and pH during mechanosensing in Arabidopsis roots. Plant Cell 21, 2341-2356.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Monshausen, G.B., Miller, N.D., Murphy, A.S., and Gilroy, S. (2011). Dynamics of auxin-dependent Ca²⁺ and pH signaling in root growth revealed by integrating high-resolution imaging with automated computer vision-based analysis. Plant J 65, 309-318.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Mori, I.C., Murata, Y., Yang, Y., Munemasa, S., Wang, Y.F., Andreoli, S., Tiriach, H., Alonso, J.M., Harper, J.F., Ecker, J.R., Kwak, J.M., and Schroeder, J.I. (2006). CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca(2+)-permeable channels and stomatal closure. PLoS Biol 4, e327.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Munemasa, S., Hauser, F., Park, J., Waadt, R., Brandt, B., and Schroeder, J.I. (2015). Mechanisms of abscisic acid-mediated control of stomatal aperture. Curr Opin Plant Biol 28, 154-162.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Nagaya, S., Kawamura, K., Shinmyo, A., and Kato, K. (2010). The HSP terminator of Arabidopsis thaliana increases gene expression in plant cells. Plant Cell Physiol 51, 328-332.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ngo, Q.A., Vogler, H., Lituiev, D.S., Nestorova, A., and Grossniklaus, U. (2014). A calcium dialog mediated by the FERONIA signal transduction pathway controls plant sperm delivery. Dev Cell 29, 491-500.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Nietzel, T., Elsässer, M., Ruberti, C., Steinbeck, J., Ugalde, J.M., Fuchs, P., Wagner, S., Ostermann, L., Moseler, A., Lemke, P., Fricker, M.D., Müller-Schüssele, S.J., Moerschbacher, B.M., Costa, A., Meyer, A.J., and Schwarzländer, M. (2019). The fluorescent protein sensor roGFP2-Orp1 monitors in vivo H₂O₂ and thiol redox integration and elucidates intracellular H₂O₂ dynamics during elicitor-induced oxidative burst in Arabidopsis. New Phytol 221, 1649-1664.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ogasawara, Y., Kaya, H., Hiraoka, G., Yumoto, F., Kimura, S., Kadota, Y., Hishinuma, H., Senzaki, E., Yamagoe, S., Nagata, K., Nara, M., Suzuki, K., Tanokura, M., and Kuchitsu, K. (2008). Synergistic activation of the Arabidopsis NADPH oxidase AtrbohD by Ca²⁺ and phosphorylation. J Biol Chem 283, 8885-8892.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Okumoto, S., Jones, A., and Frommer, W.B. (2012). Quantitative imaging with fluorescent biosensors. Annu Rev Plant Biol 63, 663-706.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Peaucelle, A., Braybrook, S., and Höfte, H. (2012). Cell wall mechanics and growth control in plants: the role of pectins revisited. Front Plant Sci 3, 121.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Pei, Z.M., Murata, Y., Benning, G., Thomine, S., Klüsener, B., Allen, G.J., Grill, E., and Schroeder, J.I. (2000). Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. Nature 406, 731-734.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Pittman, J.K., and Hirschi, K.D. (2016). CAX-ing a wide net: Cation/H(+) transporters in metal remediation and abiotic stress signalling. Plant Biol (Stuttg) 18, 741-749.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., and Cardona, A. (2012). Fiji: an open-source platform for biological-

image analysis. Nat Methods 9, 676-682.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Schmidt, C., Schelle, I., Liao, Y.J., and Schroeder, J.I. (1995). Strong regulation of slow anion channels and abscisic acid signaling in guard cells by phosphorylation and dephosphorylation events. Proc Natl Acad Sci U S A 92, 9535-9539.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Schumacher, K. (2014). pH in the plant endomembrane system - an import and export business. Curr Opin Plant Biol 22, 71-76.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Schwarzländer, M., Logan, D.C., Johnston, I.G., Jones, N.S., Meyer, A.J., Fricker, M.D., and Sweetlove, L.J. (2012). Pulsing of membrane potential in individual mitochondria: a stress-induced mechanism to regulate respiratory bioenergetics in Arabidopsis. Plant Cell 24, 1188-1201.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Seybold, H., Trempel, F., Ranf, S., Scheel, D., Romeis, T., and Lee, J. (2014). Ca²⁺ signalling in plant immune response: from pattern recognition receptors to Ca²⁺ decoding mechanisms. New Phytol 204, 782-790.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Shan, X., Yan, J., and Xie, D. (2012). Comparison of phytohormone signaling mechanisms. Curr Opin Plant Biol 15, 84-91.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Shavruk, Y., and Hirai, Y. (2016). Good and bad protons: genetic aspects of acidity stress responses in plants. J Exp Bot 67, 15-30.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Shen, Y., Rosendale, M., Campbell, R.E., and Perrais, D. (2014). pHuji, a pH-sensitive red fluorescent protein for imaging of exo- and endocytosis. J Cell Biol 207, 419-432.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Shih, H.W., Miller, N.D., Dai, C., Spalding, E.P., and Monshausen, G.B. (2014). The receptor-like kinase FERONIA is required for mechanical signal transduction in Arabidopsis seedlings. Curr Biol 24, 1887-1892.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Shih, H.W., DePew, C.L., Miller, N.D., and Monshausen, G.B. (2015). The Cyclic Nucleotide-Gated Channel CNGC14 Regulates Root Gravitropism in Arabidopsis thaliana. Curr Biol 25, 3119-3125.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Smirnoff, N., and Arnaud, D. (2019). Hydrogen peroxide metabolism and functions in plants. New Phytol 221, 1197-1214.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Song, C.J., Steinebrunner, I., Wang, X., Stout, S.C., and Roux, S.J. (2006). Extracellular ATP induces the accumulation of superoxide via NADPH oxidases in Arabidopsis. Plant Physiol 140, 1222-1232.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Stegmann, M., Monaghan, J., Smakowska-Luzan, E., Rovenich, H., Lehner, A., Holton, N., Belkadir, Y., and Zipfel, C. (2017). The receptor kinase FER is a RALF-regulated scaffold controlling plant immune signaling. Science 355, 287-289.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Steinhorst, L., and Kudla, J. (2014). Signaling in cells and organisms - calcium holds the line. Curr Opin Plant Biol 22, 14-21.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Sze, H., and Chanroj, S. (2018). Plant Endomembrane Dynamics: Studies of K⁺/H⁺ Antiporters Provide Insights on the Effects of pH and Ion Homeostasis. Plant Physiol 177, 875-895.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Takemoto, K., Ebine, K., Askani, J.C., Krüger, F., Gonzalez, Z.A., Ito, E., Goh, T., Schumacher, K., Nakano, A., and Ueda, T. (2018). Distinct sets of tethering complexes, SNARE complexes, and Rab GTPases mediate membrane fusion at the vacuole in Arabidopsis. Proc Natl Acad Sci U S A 115, E2457-E2466.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tanaka, K., Swanson, S.J., Gilroy, S., and Stacey, G. (2010). Extracellular nucleotides elicit cytosolic free calcium oscillations in Arabidopsis. Plant Physiol 154, 705-719.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tian, L., Hires, S.A., Mao, T., Huber, D., Chiappe, M.E., Chalasani, S.H., Petreanu, L., Akerboom, J., McKinney, S.A., Schreiter, E.R., Bargmann, C.I., Jayaraman, V., Svoboda, K., and Looger, L.L. (2009). Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators. Nat Methods 6, 875-881.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tian, W., Hou, C., Ren, Z., Wang, C., Zhao, F., Dahlbeck, D., Hu, S., Zhang, L., Niu, Q., Li, L., Staskawicz, B.J., and Luan, S. (2019). A calmodulin-gated calcium channel links pathogen patterns to plant immunity. Nature 572, 131-135.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Vanstraelen, M., and Benková, E. (2012). Hormonal interactions in the regulation of plant development. Annu Rev Cell Dev Biol 28, 463-487.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Waadt, R., Hitomi, K., Nishimura, N., Hitomi, C., Adams, S.R., Getzoff, E.D., and Schroeder, J.I. (2014). FRET-based reporters for the direct visualization of abscisic acid concentration changes and distribution in Arabidopsis. Elife 3, e01739.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Waadt, R., Hsu, P.K., and Schroeder, J.I. (2015). Abscisic acid and other plant hormones: Methods to visualize distribution and signaling. Bioessays 37, 1338-1349.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Waadt, R., Krebs, M., Kudla, J., and Schumacher, K. (2017). Multiparameter imaging of calcium and abscisic acid and high-resolution quantitative calcium measurements using R-GECO1-mTurquoise in Arabidopsis. New Phytol 216, 303-320.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wagner, S., Steinbeck, J., Fuchs, P., Lichtenauer, S., Elsässer, M., Schippers, J.H.M., Nietzel, T., Ruberti, C., Van Aken, O., Meyer, A.J., Van Dongen, J.T., Schmidt, R.R., and Schwarzländer, M. (2019). Multiparametric real-time sensing of cytosolic physiology links hypoxia responses to mitochondrial electron transport. New Phytol 224, 1668-1684.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Walia, A., Waadt, R., and Jones, A.M. (2018). Genetically Encoded Biosensors in Plants: Pathways to Discovery. Annu Rev Plant Biol 69, 497-524.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Waszczak, C., Carmody, M., and Kangasjärvi, J. (2018). Reactive Oxygen Species in Plant Signaling. Annu Rev Plant Biol 69, 209-236.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Yoshida, T., Christmann, A., Yamaguchi-Shinozaki, K., Grill, E., and Fernie, A.R. (2019). Revisiting the Basal Role of ABA - Roles Outside of Stress. Trends Plant Sci 24, 625-635.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhang, X., Köster, P., Schlücking, K., Balcerowicz, D., Hashimoto, K., Kuchitsu, K., Vissenberg, K., and Kudla, J. (2018). CBL1-CIPK26-mediated phosphorylation enhances activity of the NADPH oxidase RBOHC, but is dispensable for root hair growth. FEBS Lett 592, 2582-2593.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhao, Y., Araki, S., Wu, J., Teramoto, T., Chang, Y.F., Nakano, M., Abdelfattah, A.S., Fujiwara, M., Ishihara, T., Nagai, T., and Campbell, R.E. (2011). An expanded palette of genetically encoded Ca²⁺ indicators. Science 333, 1888-1891.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zheng, X., Kang, S., Jing, Y., Ren, Z., Li, L., Zhou, J.M., Berkowitz, G., Shi, J., Fu, A., Lan, W., Zhao, F., and Luan, S. (2018). Danger-Associated Peptides Close Stomata by OST1-Independent Activation of Anion Channels in Guard Cells. Plant Cell 30, 1132-1146.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)