Genetic liability for internalizing versus externalizing behavior manifests in the developing and adult hippocampus: Insight from a meta-analysis of transcriptional profiling studies in a selectively-bred rat model

Isabelle A. Birt^{1*}, Megan H. Hagenauer, Ph.D.^{1*}, Sarah M. Clinton, Ph.D.², Cigdem Aydin, Ph.D.¹, Peter Blandino, Jr., Ph.D.¹, John D. H. Stead, Ph.D.³, Kathryn L. Hilde, Ph.D.¹, Fan Meng, Ph.D.¹, Robert C. Thompson, Ph.D.¹, Huzefa Khalil, Ph.D.¹, Alex Stefanov¹, Pamela Maras, Ph.D.¹, Zhifeng Zhou, Ph.D.⁴, Elaine K. Hebda-Bauer, Ph.D.¹, David Goldman, Ph.D.⁴, Stanley J. Watson, Jr., Ph.D.¹ and Huda Akil, Ph.D.¹

¹University of Michigan, Ann Arbor, MI, USA; ²Virginia Tech University, Blacksburg, VA, USA;

³Carleton University, Ottawa, Ontario, CA; ⁴ National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Bethesda, MD, USA

*Shared first authorship

Correspondence:

Megan Hastings Hagenauer, Ph.D.

e-mail: hagenaue@umich.edu

Molecular Behavioral Neuroscience Institute (MBNI)

205 Zina Pitcher Pl.

Ann Arbor, MI 48109

Abstract

Background: For over 16 years, we have selectively bred rats to show either high or low levels of exploratory activity within a novel environment. These “bred High Responder” (bHR) and “bred Low Responder” (bLR) rats serve as a model for temperamental extremes, exhibiting large differences in many internalizing and externalizing behaviors relevant to mood and substance abuse disorders.

Methods: Our study elucidated persistent differences in gene expression related to bHR/bLR phenotype across development and adulthood in the hippocampus, a region critical for emotional regulation. We meta-analyzed eight transcriptional profiling datasets (microarray, RNA-Seq) spanning 43 generations of selective breeding (adult: $n=46$, P7: $n=22$, P14: $n=49$, P21: $n=21$; all male). We cross-referenced these results with exome sequencing performed on our colony to pinpoint candidates likely to mediate the effect of selective breeding on behavioral phenotype.

Results: Genetic and transcriptional profiling results converged to implicate two genes with previous associations with metabolism and mood: Thyrotropin releasing hormone receptor and Uncoupling protein 2. Our results also highlighted bHR/bLR functional differences in the hippocampus, including a network essential for neurodevelopmental programming, proliferation, and differentiation, containing hub genes Bone morphogenetic protein 4 and Marker of proliferation ki-67. Finally, we observed differential expression related to microglial activation, which is important for synaptic pruning, including two genes within implicated chromosomal regions: Complement C1q A chain and Milk fat globule-EGF factor 8.

Conclusion: These candidate genes and functional pathways have the capability to direct bHR/bLR rats along divergent developmental trajectories and promote a widely different reactivity to the environment.

27

28 **Keywords:** Anxiety, Depression, Hyperactivity, Impulsivity, Limbic, Neonatal, Postnatal, Trhr,

29 Ucp2, Ncan, Bmp4, Mki67, C1qa, Mfge8, Etv4

30

31

Introduction

The strong pattern of comorbidity amongst psychiatric disorders is believed to be generated by a spectrum of latent liability (1), arising from a complex interplay of genetic risk and environmental factors, such as stress and childhood adversity (1–3). At one end of this spectrum are internalizing disorders, which are associated with neuroticism, anxiety, and depression. At the other end of the spectrum are externalizing disorders, which are associated with risk-taking and novelty-seeking, as seen in mania, substance abuse, and impulse-control disorders (1).

We model the genetic contributions underlying both extremes of this spectrum by selectively breeding rats that react differently to a novel environment. “Bred High Responder” (bHR) rats are highly exploratory with a disinhibited, novelty-seeking temperament, including hyperactivity, aggression, and drug-seeking. “Bred Low Responder” (bLR) rats are highly-inhibited, exhibiting reduced locomotor activity and anxious and depressive-like behavior (4–13). These behavioral propensities are robust and stable, beginning early in development (14,15) similar to temperament in humans (16).

This highly-differentiated phenotype makes bHR/bLR rats ideal for observing the developmental programming and adult manifestation of neurological factors underlying internalizing and externalizing tendencies (8,11,17). This study focused on the hippocampus, a region important for emotional regulation, behavioral inhibition (18–20), and reactivity to the environment (18), including stress-related glucocorticoid release (21,22). In the bHR/bLR model, we previously observed differences in hippocampal glucocorticoid receptor and growth factor expression, histone methylation, cell proliferation, survival, and overall volume (5,9,14,23,24).

Our current study characterized hippocampal gene expression in bHR/bLR rats across development and adulthood using a meta-analysis of eight transcriptional profiling datasets spanning 43 generations of selective breeding. Concurrently, we discovered chromosomal

regions containing bHR/bLR segregating variants that are likely to contribute to exploratory locomotor phenotype (25). By comparing across these studies, we identified differentially-expressed (DE) genes situated within implicated chromosomal regions and hippocampal functional pathways essential for mood and development (**Fig 1**). These genes are promising candidates for mediating the influence of selective breeding on behavioral phenotype.

[Figure 1]

Methods

Full methods for the individual experiments and analyses are provided in the supplement. The associated datasets have been released on GEO (**Table 1**) or FigShare (<https://doi.org/10.1101/774034>). Analysis code (R Studio v.1.0.153, R v.3.2.2) is available at <https://github.com/isabellie4/PhenotypeProject> and https://github.com/hagenaue/bHRbLR_MetaAnalysisProject.

[Table 1]

The bHR/bLR Rat Colony

All experiments were approved by the local University Committee on the Use and Care of Animals, in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Selective Breeding: We began selectively-breeding bHR/bLR rats in the Molecular Behavioral Neuroscience Institute (MBNI) at the University of Michigan in 2003 (protocol: (11)).

Later, a second colony was begun at the University of Alabama-Birmingham using generation F30 bHR/bLR rats from MBNI. Our meta-analyses use datasets derived from male bHR/bLR rats spanning generations F4-F43. We refer to these datasets according to their respective institution, transcriptional profiling platform, and generation (**Table 1**). *MBNI_RNASeq_F37* also included a bHRxbLR cross (“Intermediate Responder” (bIR) rats)).

Behavioral Testing: For each generation, locomotor response to a novel environment was assessed between P50–75 (11). For *MBNI_RNASeq_F37*, we measured anxiety-like behavior in adulthood (bHR/bLR: P160-P167; bIR: P65-75) using the percent time spent in the open arms of an Elevated Plus Maze (EPM; 5 min test, protocol: (26)). For *MBNI_RNASeq_F43*, we measured social interaction in adulthood (P92, protocol: (4)) after 15 min exposure to the anxiogenic open arms of the EPM (protocol: (27)).

Hippocampal Gene Expression Analyses

Broad Overview of the Datasets: Our meta-analyses included eight datasets from bHR/bLR rats aged P7, P14, P21, and adult (**Table 1**). The rats were housed in standard conditions with minimal intervention besides behavioral testing or saline injections, and sacrificed by rapid decapitation without anesthesia. The whole hippocampus was dissected, except in *Alabama_Nimblegen_F34*, where dorsal hippocampal tissue punches were performed on sliced frozen tissue (28). The extracted RNA was profiled using microarray (earlier generations: F4, F6, F15, F34) or RNA-Seq (later generations: F29, F37, F43).

Broad Overview of the Data Preprocessing: The pre-processing steps for each study varied according to platform, but included common steps, including re-annotation, normalization to reduce technical variation, and quality control. Microarray data were typically summarized into

log2-transformed expression sets using Robust Multi-Array Average (RMA: (29)). Gene-level RNA-Seq read count summaries were converted to log(2) fragments per million (FPM). When applicable, transcript data were averaged by gene symbol to obtain a single expression value per sample per gene.

Meta-Analyses: Within each dataset, we calculated the effect size (Cohen's d and variance of d) of bHR/bLR phenotype on the expression data for each gene within each age group. This output was aligned across datasets using official gene symbol. The meta-analysis for each age group was performed using *rma.mv()* (package *metafor* (30)) and corrected for false discovery rate (FDR; Benjamini-Hochberg method in *multtest*: (31)). Including generation as a co-variate provided little additional insight (*generation*: all genes FDR>0.3).

Overlap with Previously-Identified Genetic Variants: Our exome sequencing study identified bHR/bLR segregating variants (single-nucleotide polymorphisms, or SNPs), and used a sampling of those variants to pinpoint quantitative trait loci (QTLs) for exploratory locomotion using an bHRxbLR F2 intercross (25). In our current study, we identified all DE genes nearby (+/- 1 MB) the segregating variants and QTL peaks (LOD>3) and determined overlap with additional QTLs relevant to bHR/bLR behavioral phenotype from Rat Genome Database ((32), accessed 08/08/2019, keywords: "Anxiety", "Stress", and "Despair").

Positional Gene Enrichment Analysis: To explore which genes might be either co-regulated or in linkage disequilibrium with a causal genetic variant, we evaluated the clustering of top bHR/bLR DE genes within chromosomal regions using Positional Gene Enrichment analysis (*PGE*, <http://silico.biotoul.fr/pge/>, (33)) and the top results from the P14 and adult meta-analyses (nominal p<0.01, removing duplicated EntrezIDs).

Gene Set Enrichment Analysis (GSEA) & Protein-Protein Interaction (PPI)

Networks: To elucidate overall functional trends within the largest sets of results (P14 and Adult) we used GSEA ((34,35); package *fgsea* (36)) and gene set matrix files (.gmt) containing standard gene ontology for rats (go2msig.org; (37)), or customized hippocampal-specific gene sets (**Table S1**) including previously-identified co-expression modules (38,39) and expression specific to hippocampal neuronal subtypes or subregions (*Hipposeq*: (40)). We further explored top DE genes (adult meta-analysis FDR<0.10) and implicated hippocampal gene sets using predicted PPI networks (string-db.org (41)).

Cell Type Data Deconvolution: To interrogate the relative cell type composition of our samples, we used the *BrainInABlender* method (42). Data for genes previously identified as having cell type specific expression was extracted, normalized, and averaged to produce a “cell type index”. For this analysis, we excluded the small MBNI_RNASeq_F29 dataset (n=2/group). We then performed a meta-analysis of the effects of bHR/bLR phenotype on these cell type indices using aforementioned methods.

qPCR Validation: Hippocampal tissue from 6 bHR and 6 bLR males was collected at ages P14 (generation F55) and P90 (generation F51; **Fig S1**). Following cDNA synthesis, Bone morphogenetic protein 4 (*Bmp4*) was quantified using qPCR and custom-designed primers, using the Livak method (43) and glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*) as reference. Group differences in $-\Delta C_q$ at each age were assessed using Welch’s two-sample t-test (44).

Results

Selective Breeding Amplifies the Propensity for Internalizing vs. Externalizing Behavior

The divergence of bHR/bLR exploratory activity in response to selective breeding happened rapidly (**Fig 2A**), implying oligogenic inheritance (25). This divergence was accompanied by a general amplification of internalizing and externalizing tendencies (11,14,45,46). For example, in the behavioral data accompanying our transcriptional profiling datasets, bLRs showed more anxiety-like behavior than bHRs (**Fig 2B**), and spent less time interacting socially following a stressful challenge (**Fig 2C**). Therefore, we expected that examining gene expression across bHR/bLR generations would reveal a convergence of effects within implicated chromosomal regions and pathways essential to affective behavior and reactivity to the environment.

[FIGURE 2]

Selective Breeding for Exploratory Locomotion Alters Hippocampal Gene Expression

Between generations F4-43, we conducted eight exploratory studies transcriptionally profiling the hippocampus of bHR/bLR rats at four ages (P7, P14, P21, and adult). These small studies individually produced few reliable results (**Figs S2-S3**). Nevertheless, a formal meta-analysis revealed multiple genes with consistent DE across generations (**Fig 3, Table S2**).

These results can be explored interactively at <https://mbni.org/dashboard/huzefak/hrlrMetaAnalysis/>.

[FIGURE 3]

Adulthood: The effect of bHR/bLR phenotype on gene expression was significant for 74 genes (FDR<0.05, out of 16,269; **Fig 3**). In general, the estimated effect sizes (β 's) tended to be more extreme for genes exclusively represented in RNA-Seq data from later generations (**Fig**

S4), but due to smaller sample size, these effects were not overrepresented in the top results. To rule out bias due to platform, we ran a meta-analysis using only recent RNA-Seq data (F37/F43) and confirmed that similar DE genes and pathways were identified (**Fig S5**). In the following discussion we emphasize candidate genes for which there is evidence that expression diverged during the earliest generations.

Development: The developmental meta-analyses were more dependent on data from earlier generations and produced less robust results. However, the top genes had consistent effects across age groups. Within the P7 meta-analysis, only one gene (out of 3,257), Neurocan (Ncan), was differentially expressed, with higher expression in bLRs than bHRs since an early generation (**Fig 4B-C**), in a manner that nominally persisted at P14 (**Fig 4D**). Within the P14 and P21 meta-analysis, none of the effects survived multiple comparison correction (for 15,682 and 3,257 genes, respectively). However, the top gene at P14, Bmp4, was consistently expressed at higher levels in bLRs than bHRs within both the P14 (**Fig 5A**) and adult datasets since the F4 generation (**Fig 5B**). We confirmed DE at these ages using qPCR (**Fig 5C, Fig S1**).

[FIGURE 4, FIGURE 5]

Many bHR/bLR Differentially Expressed Genes Are Located Within Implicated Chromosomal Regions

Our exome sequencing study identified bHR/bLR segregating genetic variants, and then used a sampling of those variants to identify chromosomal regions that are likely to contribute to exploratory locomotor phenotype (QTL peaks). These implicated chromosomal regions overlapped extensively with QTLs relevant to internalizing/externalizing behavior (32), including

anxiety (47–52), stress response (53–55), and behavioral despair (56). In our current study, 68% of the genes DE in adulthood (FDR<0.05) were within +/-1 MB of a bHR/bLR segregating variant, and 21% were within +/-100 kB of a highly segregating variant (Bonferonni-corrected $\alpha=5.00E-05$), a 4.7x enrichment compared to non-DE genes (**Fig 6A**, Fischer's exact test: $p=3.50E-06$). DE genes were also 4.5x more likely to be located within a QTL for exploratory locomotion (LOD>4; **Fig 6B**). This overlap included two genes previously associated with internalizing and externalizing behaviors (Thyrotropin Releasing Hormone Receptor (Trhr), Uncoupling Protein 2 (Ucp2), **Fig 6C**, (57–62)). These results fit our expectation that the influence of bHR/bLR segregating genetic variants on exploratory locomotion is at least partially mediated by effects on gene expression within the hippocampus.

[FIGURE 6]

Positional Gene Enrichment Analysis Specifies Narrower Chromosomal Regions Contributing to bHR/bLR Phenotype

Positional Gene Enrichment (PGE) identified 132 chromosomal regions with a significant enrichment (FDR<0.05) of DE genes within the P14 and adult meta-analyses. We focused on the top regions (FDR<0.001; **Figure 6D**). We confirmed that most of these top loci (10/13) could be identified using recent RNA-Seq data (F37/F43), ruling out bias towards regions overrepresented on older microarray platforms. These 13 top loci were narrow chromosomal regions (measured in KB), but overlapped a strikingly high percentage of QTLs relevant to bHR/bLR phenotype. This overlap included three QTLs for exploratory activity (25), encompassing two DE genes associated with internalizing and externalizing behavior (Ucp2, Trhr; **Fig 7&8**), as well as 13% of the QTLs for anxiety (6/45, (47–50)), 21% of the QTLs for stress-related responses (8/38, (53–55,63–65)), and 23% of the QTLs for behavioral despair (3/13, (56)) in the Rat Genome Database (32). Therefore, these enriched loci could contain

genetic variants contributing to internalizing/externalizing aspects of the bHR/bLR behavioral phenotype beyond exploratory locomotion.

[FIGURE 7, FIGURE 8]

bHR/bLR Differential Expression is Enriched within Hippocampal Functional Pathways

bHR/bLR Phenotype is Associated with Proliferation and Differentiation: Eight functional ontology gene sets (out of 2,761) showed an enrichment of DE within the P14 meta-analysis (FDR<0.05), all of which were upregulated in bLRs (**Fig 9A, Table S3, Fig S6**) and predominantly related to neurogenesis, differentiation, and brain development. Within the adult meta-analysis, 2 of the 4 top gene sets enriched with DE (FDR<0.1, out of 2,761 total) were similarly related to proliferation and development, but upregulated in bHRs. This pattern was confirmed using recent RNA-Seq data (F37/F43, **Table S3**), ruling out bias towards gene families overrepresented on microarray platforms.

A PPI network constructed using the top genes from the adult meta-analysis (FDR<0.10: 192 genes) had a dominant subnetwork highlighting many of these same genes (**Fig 9B**), including hubs *Bmp4* (*discussed above*) and the canonical marker of proliferation *Mki67* (**Fig 9C-E**). Literature review confirmed the PPI interactions within this subnetwork and their role in proliferation and differentiation in the brain (66–78).

[FIGURE 9]

bHR/bLR Phenotype is Associated with the Dentate Gyrus (DG): Similarly, when performing GSEA using 69 gene sets custom-designed to reflect hippocampal-specific cell types and networks (**Table S1**), we observed an enrichment of DE related to the DG (**Fig 8E**,

Fig S7, Table S3), the location of neural proliferation within the hippocampus. At P14, bLRs showed an upregulation of genes with enriched expression in the DG as compared to the Cornu Ammonis (CA) regions (40) (FDR<0.05). In adulthood, bHRs showed an upregulation of genes with enriched expression in the ventral compared to the dorsal DG (40) (FDR<0.05), including *Trhr* (FDR<0.05, *discussed above*).

bHR/bLR Phenotype is Associated with Hippocampal Co-expression Networks

Related to Synaptic Signaling: Co-expression modules can capture regionally-important cell types and functions that remain undocumented in traditional ontology databases (79). We observed an enrichment of bHR/bLR effects within six previously identified hippocampal co-expression modules within the P14 meta-analysis (FDR<0.05) and five within the adult meta-analysis (FDR<0.05, **Fig 8E, Fig S7, Table S3**). Two of these modules included genes within implicated chromosomal regions.

The first was a large co-expression module (695 genes) previously identified in the mouse hippocampus ((39), *“lightcyan”*) which showed elevated expression in bLRs relative to bHRs at P14 (FDR<0.05) and adulthood (FDR<0.10). One of the top DE genes in this module, ETS Variant 4 (*Etv4*, FDR<0.05, bHR>bLR), is a transcription factor required for proper hippocampal dendrite development (80) located within an implicated chromosomal region (**Fig S8**). A PPI network constructed using the DE genes in this module (n=74, adult p<0.05), was enriched with genes related to cell projections, neurons, synapses, and cation binding (FDR<0.05).

The second was a small co-expression module (39 genes) previously identified in the mouse hippocampus ((39), *sienna3*), which showed elevated expression in bHRs in adulthood (FDR<0.05). The top gene in this module was *Trhr* (*discussed above*). A PPI network constructed using all 39 genes in this module centered on *Trhr* and its ligand, thyrotropin releasing hormone (*Trh*; **Fig 8G**), and included many reward-related signaling molecules,

including CART prepropeptide (Cartpt (81)), Oxytocin/Neurophysin I Prepropeptide (Oxt (82)), and Dopamine Receptor D1a (Drd1a (83,84)).

bHR/bLR Phenotype is Associated with Microglial and Endothelial-Specific Gene Expression: Our results suggested that bHR/bLR rats might have differences in hippocampal cell type composition. A cell type deconvolution analysis focused on well-characterized non-neuronal cell type categories revealed that bLRs had greater microglial-specific expression than bHRs at P14 and adulthood (**Fig 10A,C-D**). At P14, bLRs also showed greater endothelial-specific expression (**Fig 10B**). These effects could reflect differences in cell type composition or activation state. Notably, two of the top DE genes that were located within implicated chromosomal regions are regulators of microglial state: Milk fat globule-EGF factor 8 (Mfge8, **Fig 10E-H**) promotes alternative (M2) activation (85), and Complement component C1q A Chain (C1qa, **Fig 10I-L**) promotes classical activation (86). To interrogate less well-characterized hippocampal cell types, we compared our meta-analysis results to the new mousebrain.org database (87), and found that the top DE genes (FDR<0.10) were highly expressed in a variety of cell types, including neuronal subcategories (**Fig S9**), mirroring the diversity of hippocampal functions implicated in bHR/bLR phenotype.

[FIGURE 10]

Discussion

By selectively-breeding rats for over 16 years, we have produced a robust, genetic model of the co-occurrence of common internalizing and externalizing behaviors. Such large differences in behavior would be expected to be accompanied by similarly strong differences in gene expression in affective circuitry. By performing a formal meta-analysis across small,

exploratory datasets, we provide insight into bHR/bLR differences in hippocampal gene expression across development and adulthood. Further, by cross-referencing these results with our concurrent genetic study (25), we pinpoint strong candidates for mediating the influence of selective breeding on hippocampal function and internalizing/externalizing behavior.

Transcriptional profiling converges with genetic results to identify two strong candidates for contributing to bHR/bLR behavioral phenotype: Trhr and Ucp2

Our exome sequencing study offered a first glimpse of the genetic factors contributing to our selectively-bred phenotypes (25). However, the implicated chromosomal regions encompass several hundred genes, making their specific effects on gene expression and function difficult to predict. By cross-referencing these findings with our DE results, we discovered two strong candidates: Trhr and Ucp2. These candidates are located near bHR/bLR fully-segregating genetic variants (25) within narrow chromosomal regions highly enriched for DE genes, and overlap QTLs for exploratory activity (25), anxiety (47–49), and stress response (53,54). Trhr was also the top gene within a bHR-upregulated gene set associated with the ventral DG (40), a region important for proliferation and emotional regulation (18), and was a hub in a bHR-upregulated hippocampal network containing reward-related signaling molecules.

Trhr and Ucp2 are both important for energy metabolism and extensively linked to internalizing and externalizing behavior (88–90). Knocking out Ucp2 produces a bLR-like phenotype: higher anxiety-like behavior, lower locomotor activity, and reduced stress resilience (57,58,60,61). Trhr is an important component of the hypothalamic-pituitary-thyroid (HPT) axis and regulates anxiety and depressive-like behavior (59,62,91). In our exome-sequencing study, the variants associated with Trhr and Ucp2 explained a moderate portion of exploratory locomotor behavior (<10%, approximately 200 locomotor counts, (25)) – a magnitude akin to the bHR/bLR difference in locomotor score present in the F1 generation. Altogether, this evidence suggests that bHR/bLR segregating genetic variants are driving DE of Ucp2 and Trhr in a

manner that could meaningfully contribute to bHR/bLR differences in hippocampal function and internalizing/externalizing behavior.

The top genes identified in the developmental meta-analyses suggest that bHR/bLR differences in hippocampal structure arise early in development: Ncan and Bmp4

The different propensity of bHR/bLR rats towards externalizing or internalizing behavior is evident at a young age (14,15), paralleling hippocampal development (14). Our meta-analyses encompassed three postnatal ages (P7, P14, and P21) to provide insight into this neurodevelopmental trajectory. These meta-analyses depended on data from earlier generations and older transcriptional profiling platforms yet identified two strong candidates with clear associations with internalizing/externalizing behavior and hippocampal development.

The top P7 result was Ncan, exhibiting a strikingly large effect size (bHR<bLR) as early as generation F6. Ncan is located adjacent to a bHR/bLR segregating variant (25), overlapping a QTL for despair-related behavior (56). As part of the extracellular matrix, Ncan is upregulated during early brain development (92,93) and modulates cell adhesion, migration, and growth factor binding (94). Ncan has been linked to Bipolar Disorder (92), and knocking-out Ncan enhances locomotor activity, risk-taking, hedonia, and amphetamine hypersensitivity (93). Therefore, lower levels of Ncan in early development in bHRs could promote externalizing behavior as well as divergent hippocampal development.

The top P14 result was Bmp4, which was elevated in bLRs since the earliest generations in a manner that appeared to persist into adulthood. As a regulator of development, Bmp4 is initially important for neural induction (95,96), but later suppresses neurogenesis (97–101) and promotes other cell fates (78,95,102). Bmp4 was an important driver in bHR/bLR-enriched gene sets related to proliferation and differentiation at P14 and a hub in a related PPI network constructed from top adult DE genes. In the hippocampus, Bmp signaling promotes

dorsal cell type identity and is essential for DG formation (103), matching our results indicating bLR-enrichment of gene expression related to the DG at P14 and adulthood.

Moreover, blocking Bmp signaling can produce a bHR-like phenotype, reducing anxiety, fear conditioning, and depressive-like behavior (98,103). Therefore, Bmp4 is a strong candidate for driving long-term structural differences in the hippocampus capable of producing stable differences in temperament. However, Bmp4 was not located near a bHR/bLR segregating variant in our exome sequencing study (25), implying that impactful variation may be located in a nearby non-coding region or within a gene upstream in Bmp4's signaling pathway.

Functional analyses implicate hippocampal proliferation and differentiation in bHR/bLR phenotype

One of the most prominent themes in our results were functions related to cell proliferation and differentiation. Indeed, we found that Mki67 itself contained two bHR/bLR segregating genetic variants, and was more highly expressed in bLRs in adulthood (FDR<0.05), matching the upregulation in bLRs observed histologically in development (14) and maybe adulthood (9,104). These findings confirm that the relationship between internalizing behavior and cell proliferation in our model is unlikely to be as simple as a general stunting of growth-related processes, as suggested by the neurotrophic model of stress-related mood disorders (105).

Many of our top DE genes were also important regulators of cell fate. Bmp4, SRY-box 9 (Sox9), SRY-box 2 (Sox2), Hes Family BHLH Transcription Factor 5 (Hes5), CD24 Molecule (Cd24), and TEK Receptor Tyrosine Kinase (Tek) regulate functions such as the developmental progression of neural differentiation, gliogenesis, and endothelial proliferation (66,74,75,77,106–108). Their role in adulthood includes growth and plasticity in response to neural activity and injury (69,109–111). Therefore, these results could explain previous morphological findings indicating that cell differentiation progressed differently in the adult hippocampus in bHRs and

bLRs under conditions of mild stress (9). Together, these findings raise the interesting possibility that DE within neurodevelopmental programming pathways could provide a general mechanism by which environmental stimuli, such as stress or drugs, produces divergent changes in hippocampal structure in bHR/bLR animals.

Functional analyses implicate microglial activation in bHR/bLR phenotype

Microglial-specific gene expression was upregulated at both P14 and adulthood in bLRs, suggesting either an increased proportion of microglia cells or microglial activation. Several top candidate genes were important regulators of microglial function. bLRs had greater expression of C1qa than bHRs since the earliest generations. The C1q genes promote classical microglial activation (86) and are implicated in phagocytosis-driven synaptic pruning (86,112,113). In contrast, Mfge8 was more expressed in bHRs. Mfge8 is associated with reduced pro-inflammatory factors (114) as well as alternative (M2) microglial activation (85), playing an important role in phagocytosis (85,115,116). Ucp2, discussed previously, has anti-inflammatory function (57,58,60,117,118) and was more highly expressed in bHRs than bLRs. Both Mfge8 and Ucp2 contain bHR/bLR segregating genetic variants within probable QTLs for exploratory activity (25), suggesting that genetic variation could contribute to their DE in the hippocampus.

Together, these results seem to fit pro-inflammatory theories of internalizing disorders (119). However, we found little evidence of bHR/bLR differences in the expression of traditional inflammatory markers. Therefore, it seems more likely that bHR/bLR differences in microglial activation genes are tied to non-immune roles for microglia within the brain, including the regulation of neurogenesis, cell survival (120), and synaptic pruning (113,121) in response to neuronal activity (113). Therefore, microglial phagocytosis could be serving as a multi-faceted tool to tailor plasticity either during development, or in response to environmental stimuli like stress or drugs of abuse.

Conclusion and Future Directions

By comparing exome sequencing findings to hippocampal differential expression patterns during development and adulthood across many generations of selective breeding in our bHR/bLR colony, we implicate a diverse and compelling array of genes whose effects may converge to promote internalizing and externalizing behavior. Due to a dependence of our results on older platforms and exclusively male rats, we cannot claim to have identified all relevant candidates, nor have we highlighted all promising results in our text. However, we implicate two functional pathways with the capability to guide bHRs and bLRs along a divergent developmental trajectory and set the stage for a widely different reactivity to the environment. These findings will inspire new avenues of research (122–124), including cell type specific morphological analyses and the manipulation of candidate pathways to assess relevance to behavioral and neurological phenotype in our model.

Acknowledgements

The work that was performed at the Molecular Behavioral Neuroscience Institute was supported by the following grants: Hope for Depression Research Foundation (HDRF; RGA# DTF Phase II (D) (McEwen- PI)), Pritzker Neuropsychiatric Disorders Research Consortium, National Institute on Drug Abuse Grants 5P01 DA021633-02, NIDA (5RO1 DA013386), Office of Naval Research Grant N00014-12-1-0366, Office of Naval Research (ONR) N00014-09-1-0598, Office of Naval Research (ONR) N00014-02-1-0879, NIMH Conte Center Grant #L99MH60398, and NIMH Program Project Grant #P01MH42251-01. The University of Alabama study was funded by NIMH 4R00MH085859-02 (S.M.C.). Research training for I.B. was supported by the Undergraduate Research Opportunity Program at the University of Michigan. We would also

like to acknowledge everyone involved in the studies included in the meta-analysis: James Stewart, Angela Koelsch, Sharon Burke, Mary Hoverstein, Jennifer Fitzpatrick, Hui Li, Fei Li, and Amy Tang, the University of Michigan Advanced Genomics Core, and HudsonAlpha Institute for Biotechnology Genomics Services Lab. Finally, we thank everyone who has provided feedback on the manuscript: Dr. Shelly Flagel, Alek Pankonin, Pouya Mandi Gholami, Dr. Aram Parsegian, and Leah Johnson.

Disclosures

These results were previously published in pre-print format (*bioRxiv* 774034; doi: <https://doi.org/10.1101/774034>). Several of the authors are members of the Pritzker Neuropsychiatric Disorders Research Consortium (MHH, RCT, FM, HA, SJW), which is supported by Pritzker Neuropsychiatric Disorders Research Fund, LLC. A shared intellectual property agreement exists between the academic and philanthropic entities of the consortium. The authors have declared that no competing interests exist.

Works Cited

1. Krueger RF, Markon KE (2006): Reinterpreting comorbidity: a model-based approach to understanding and classifying psychopathology. *Annu Rev Clin Psychol* 2: 111–133.
2. Smoller JW (2016): The Genetics of Stress-Related Disorders: PTSD, Depression, and Anxiety Disorders. *Neuropsychopharmacology* 41: 297–319.
3. Wray NR, Ripke S, Mattheisen M, Trzaskowski M, Byrne EM, Abdellaoui A, *et al.* (2018): Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet* 50: 668–681.
4. Aydin C, Frohmader K, Akil H (2015): Revealing a latent variable: individual differences in affective response to repeated injections. *Behav Neurosci* 129: 679–682.
5. Chaudhury S, Aurbach EL, Sharma V, Blandino P, Turner CA, Watson SJ, Akil H (2014): FGF2 is a target and a trigger of epigenetic mechanisms associated with differences in emotionality: Partnership with H3K9me3. *Proc Natl Acad Sci U S A* 111: 11834–11839.
6. Clinton SM, Vazquez DM, Kabbaj M, Kabbaj M-H, Watson SJ, Akil H (2007): Individual differences in novelty-seeking and emotional reactivity correlate with variation in maternal behavior. *Horm Behav* 51: 655–664.
7. Flagel SB, Robinson TE, Clark JJ, Clinton SM, Watson SJ, Seeman P, *et al.* (2010): An animal model of genetic vulnerability to behavioral disinhibition and responsiveness to reward-related cues: implications for addiction. *Neuropsychopharmacology* 35: 388–400.
8. Flagel SB, Waselus M, Clinton SM, Watson SJ, Akil H (2014): Antecedents and consequences of drug abuse in rats selectively bred for high and low response to novelty. *Neuropharmacology* 76 Pt B: 425–436.

9. Perez JA, Clinton SM, Turner CA, Watson SJ, Akil H (2009): A New Role for FGF2 as an Endogenous Inhibitor of Anxiety. *J Neurosci* 29: 6379–6387.
10. Prater KE, Aurbach EL, Larcinese HK, Smith TN, Turner CA, Blandino P, *et al.* (2017): Selectively Bred Rats Provide a Unique Model of Vulnerability to PTSD-Like Behavior and Respond Differentially to FGF2 Augmentation Early in Life. *Neuropsychopharmacology* 42: 1706–1714.
11. Stead JDH, Clinton S, Neal C, Schneider J, Jama A, Miller S, *et al.* (2006): Selective breeding for divergence in novelty-seeking traits: Heritability and enrichment in spontaneous anxiety-related behaviors. *Behav Genet* 36: 697–712.
12. Stedenfeld KA, Clinton SM, Kerman IA, Akil H, Watson SJ, Sved AF (2011): Novelty-seeking behavior predicts vulnerability in a rodent model of depression. *Physiol Behav* 103: 210–216.
13. Turner CA, Clinton SM, Thompson RC, Watson SJ, Akil H (2011): Fibroblast growth factor-2 (FGF2) augmentation early in life alters hippocampal development and rescues the anxiety phenotype in vulnerable animals. *Proc Natl Acad Sci U S A* 108: 8021–8025.
14. Clinton SM, Stead JDH, Miller S, Watson SJ, Akil H (2011): Developmental underpinnings of differences in rodent novelty-seeking and emotional reactivity. *Eur J Neurosci* 34: 994–1005.
15. Turner CA, Hagenauer MH, Aurbach EL, Maras PM, Fournier CL, Blandino P, *et al.* (2019): Effects of early-life FGF2 on ultrasonic vocalizations (USVs) and the mu-opioid receptor in male Sprague-Dawley rats selectively-bred for differences in their response to novelty. *Brain Res* 1715: 106–114.
16. Robinson JL, Kagan J, Reznick JS, Corley R (1992): The heritability of inhibited and uninhibited behavior: A twin study. *Developmental Psychology* 28: 1030–1037.
17. Turner CA, Fligel SB, Blandino P, Watson SJ, Akil H (2017): Utilizing a unique animal model to better understand human temperament. *Curr Opin Behav Sci* 14: 108–114.

18. Fanselow MS, Dong H-W (2010): Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* 65: 7–19.
19. Gray JA (1982): *The Neuropsychology of Anxiety: An Enquiry into the Functions of the Septo-Hippocampal System*. New York, NY, US: Clarendon Press/Oxford University Press.
20. Papez JW (1937): A proposed mechanism of emotion. *Archives of Neurology & Psychiatry* 38: 725–743.
21. Campbell S, Macqueen G (2004): The role of the hippocampus in the pathophysiology of major depression. *J Psychiatry Neurosci* 29: 417–426.
22. Gerlach JL, McEwen BS (1972): Rat brain binds adrenal steroid hormone: radioautography of hippocampus with corticosterone. *Science* 175: 1133–1136.
23. Clinton S, Miller S, Watson SJ, Akil H (2008): Prenatal stress does not alter innate novelty-seeking behavioral traits, but differentially affects individual differences in neuroendocrine stress responsivity. *Psychoneuroendocrinology* 33: 162–177.
24. Kabbaj M, Devine DP, Savage VR, Akil H (2000): Neurobiological correlates of individual differences in novelty-seeking behavior in the rat: differential expression of stress-related molecules. *J Neurosci* 20: 6983–6988.
25. Zhou Z, Blandino P, Yuan Q, Shen P-H, Hodgkinson CA, Virkkunen M, *et al.* (2019): Exploratory locomotion, a predictor of addiction vulnerability, is oligogenic in rats selected for this phenotype. *Proc Natl Acad Sci USA* 116: 13107–13115.
26. Aurbach EL, Inui EG, Turner CA, Hagenauer MH, Prater KE, Li JZ, *et al.* (2015): Fibroblast growth factor 9 is a novel modulator of negative affect. *Proc Natl Acad Sci USA* 112: 11953–11958.
27. Isgor C, Kabbaj M, Akil H, Watson SJ (2004): Delayed effects of chronic variable stress during peripubertal-juvenile period on hippocampal morphology and on cognitive and stress axis functions in rats. *Hippocampus* 14: 636–648.

28. Cohen JL, Glover ME, Pugh PC, Fant AD, Simmons RK, Akil H, *et al.* (2015): Maternal Style Selectively Shapes Amygdalar Development and Social Behavior in Rats Genetically Prone to High Anxiety. *Dev Neurosci* 37: 203–214.
29. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP (2003): Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249–264.
30. Viechtbauer W (2010): Conducting Meta-Analyses in R with The metafor Package. *Journal of Statistical Software* 36. <https://doi.org/10.18637/jss.v036.i03>
31. Pollard KS, Dudoit S, Laan MJ van der (2005): Multiple Testing Procedures: the multtest Package and Applications to Genomics. *Bioinformatics and Computational Biology Solutions Using R and Bioconductor*. Springer, New York, NY, pp 249–271.
32. Shimoyama M, De Pons J, Hayman GT, Laulederkind SJF, Liu W, Nigam R, *et al.* (2015): The Rat Genome Database 2015: genomic, phenotypic and environmental variations and disease. *Nucleic Acids Res* 43: D743-750.
33. De Preter K, Barriot R, Speleman F, Vandesompele J, Moreau Y (2008): Positional gene enrichment analysis of gene sets for high-resolution identification of overrepresented chromosomal regions. *Nucleic Acids Res* 36: e43.
34. Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J, *et al.* (2003): PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 34: 267–273.
35. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, *et al.* (2005): Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA* 102: 15545–15550.

36. Sergushichev A (2016): An algorithm for fast preranked gene set enrichment analysis using cumulative statistic calculation. *bioRxiv*. <https://doi.org/10.1101/060012>
37. Powell JAC (2014): GO2MSIG, an automated GO based multi-species gene set generator for gene set enrichment analysis. *BMC Bioinformatics* 15: 146.
38. Johnson MR, Shkura K, Langley SR, Delahaye-Duriez A, Srivastava P, Hill WD, *et al.* (2016): Systems genetics identifies a convergent gene network for cognition and neurodevelopmental disease. *Nat Neurosci* 19: 223–232.
39. Park CC, Gale GD, de Jong S, Ghazalpour A, Bennett BJ, Farber CR, *et al.* (2011): Gene networks associated with conditional fear in mice identified using a systems genetics approach. *BMC Syst Biol* 5: 43.
40. Cembrowski MS, Wang L, Sugino K, Shields BC, Spruston N (2016): Hipposeq: a comprehensive RNA-seq database of gene expression in hippocampal principal neurons. *Elife* 5: e14997.
41. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, *et al.* (2015): STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 43: D447-452.
42. Hagenauer MH, Schulmann A, Li JZ, Vawter MP, Walsh DM, Thompson RC, *et al.* (2018): Inference of cell type content from human brain transcriptomic datasets illuminates the effects of age, manner of death, dissection, and psychiatric diagnosis. *PLoS ONE* 13: e0200003.
43. Livak KJ, Schmittgen TD (2001): Analysis of relative gene expression data using real-time quantitative PCR and the 2^{(-Delta Delta C(T))} Method. *Methods* 25: 402–408.
44. Yuan JS, Reed A, Chen F, Stewart CN (2006): Statistical analysis of real-time PCR data. *BMC Bioinformatics* 7: 85.
45. Calvo N, Cecchi M, Kabbaj M, Watson SJ, Akil H (2011): Differential effects of social defeat in rats with high and low locomotor response to novelty. *Neuroscience* 183: 81–89.

46. Kabbaj M, Evans S, Watson SJ, Akil H (2004): The search for the neurobiological basis of vulnerability to drug abuse: using microarrays to investigate the role of stress and individual differences. *Neuropharmacology* 47 Suppl 1: 111–122.
47. Albert FW, Carlborg O, Plyusnina I, Besnier F, Hedwig D, Lautenschläger S, *et al.* (2009): Genetic architecture of tameness in a rat model of animal domestication. *Genetics* 182: 541–554.
48. Terenina-Rigaldie E, Moisan M-P, Colas A, Beaugé F, Shah KV, Jones BC, Mormède P (2003): Genetics of behaviour: phenotypic and molecular study of rats derived from high- and low-alcohol consuming lines. *Pharmacogenetics* 13: 543–554.
49. Conti LH, Jirout M, Breen L, Vanella JJ, Schork NJ, Printz MP (2004): Identification of quantitative trait Loci for anxiety and locomotion phenotypes in rat recombinant inbred strains. *Behav Genet* 34: 93–103.
50. Baum AE, Solberg LC, Churchill GA, Ahmadiyeh N, Takahashi JS, Redei EE (2006): Test- and behavior-specific genetic factors affect WKY hypoactivity in tests of emotionality. *Behav Brain Res* 169: 220–230.
51. Johannesson M, Lopez-Aumatell R, Stridh P, Diez M, Tuncel J, Blázquez G, *et al.* (2009): A resource for the simultaneous high-resolution mapping of multiple quantitative trait loci in rats: the NIH heterogeneous stock. *Genome Res* 19: 150–158.
52. Fernández-Teruel A, Escorihuela RM, Gray JA, Aguilar R, Gil L, Giménez-Llort L, *et al.* (2002): A quantitative trait locus influencing anxiety in the laboratory rat. *Genome Res* 12: 618–626.
53. Solberg LC, Baum AE, Ahmadiyeh N, Shimomura K, Li R, Turek FW, *et al.* (2006): Genetic analysis of the stress-responsive adrenocortical axis. *Physiol Genomics* 27: 362–369.
54. Yamazato M, Ohya Y, Nakamoto M, Sakima A, Tagawa T, Harada Y, *et al.* (2006): Sympathetic hyperreactivity to air-jet stress in the chromosome 1 blood pressure quantitative trait locus congenic rats. *Am J Physiol Regul Integr Comp Physiol* 290: R709-714.

55. Llamas B, Contesse V, Guyonnet-Duperat V, Vaudry H, Mormède P, Moisan M-P (2005): QTL mapping for traits associated with stress neuroendocrine reactivity in rats. *Mamm Genome* 16: 505–515.
56. Solberg LC, Baum AE, Ahmadiyeh N, Shimomura K, Li R, Turek FW, *et al.* (2004): Sex- and lineage-specific inheritance of depression-like behavior in the rat. *Mamm Genome* 15: 648–662.
57. Du R-H, Wu F-F, Lu M, Shu X-D, Ding J-H, Wu G, Hu G (2016): Uncoupling protein 2 modulation of the NLRP3 inflammasome in astrocytes and its implications in depression. *Redox Biol* 9: 178–187.
58. Gimsa U, Kanitz E, Otten W, Aheng C, Tuchscherer M, Ricquier D, *et al.* (2011): Alterations in anxiety-like behavior following knockout of the uncoupling protein 2 (*ucp2*) gene in mice. *Life Sci* 89: 677–684.
59. Pekary AE, Sattin A, Lloyd RL (2015): Ketamine modulates TRH and TRH-like peptide turnover in brain and peripheral tissues of male rats. *Peptides* 69: 66–76.
60. Sun X-L, Liu Y, Dai T, Ding J-H, Hu G (2011): Uncoupling protein 2 knockout exacerbates depression-like behaviors in mice via enhancing inflammatory response. *Neuroscience* 192: 507–514.
61. Wang D, Zhai X, Chen P, Yang M, Zhao J, Dong J, Liu H (2014): Hippocampal UCP2 is essential for cognition and resistance to anxiety but not required for the benefits of exercise. *Neuroscience* 277: 36–44.
62. Zeng H, Schimpf BA, Rohde AD, Pavlova MN, Gragerov A, Bergmann JE (2007): Thyrotropin-releasing hormone receptor 1-deficient mice display increased depression and anxiety-like behavior. *Mol Endocrinol* 21: 2795–2804.
63. Xiao B, Harada Y, Kawakami K, Nabika T (2011): A 1.8-Mbp fragment on chromosome 1 affects sympathetic response to stress: evaluation in reciprocal congenic strains between stroke-prone spontaneously hypertensive rat and Wistar-Kyoto rat. *J Hypertens* 29: 257–265.
64. Klimes I, Weston K, Gasperíková D, Kovács P, Kvetnanský R, Jezová D, *et al.* (2005): Mapping of genetic determinants of the sympathoneural response to stress. *Physiol Genomics* 20: 183–187.

65. Potenza MN, Brodtkin ES, Joe B, Luo X, Remmers EF, Wilder RL, *et al.* (2004): Genomic regions controlling corticosterone levels in rats. *Biol Psychiatry* 55: 634–641.
66. Belvindrah R, Rougon G, Chazal G (2002): Increased neurogenesis in adult mCD24-deficient mice. *J Neurosci* 22: 3594–3607.
67. Cerezo A, Guadamillas MC, Goetz JG, Sánchez-Perales S, Klein E, Assoian RK, del Pozo MA (2009): The absence of caveolin-1 increases proliferation and anchorage- independent growth by a Rac-dependent, Erk-independent mechanism. *Mol Cell Biol* 29: 5046–5059.
68. Galach M (2014): Molecular mechanisms involved in epithelial differentiation of human induced pluripotent stem cells [Dissertation]. <https://doi.org/DOI:10.11588/heidok.00017870>
69. Head BP, Peart JN, Panneerselvam M, Yokoyama T, Pearn ML, Niesman IR, *et al.* (2010): Loss of caveolin-1 accelerates neurodegeneration and aging. *PLoS ONE* 5: e15697.
70. Hossain MB, Shifat R, Li J, Luo X, Hess KR, Rivera-Molina Y, *et al.* (2017): TIE2 Associates with Caveolae and Regulates Caveolin-1 To Promote Their Nuclear Translocation. *Mol Cell Biol* 37. <https://doi.org/10.1128/MCB.00142-17>
71. Jo A, Denduluri S, Zhang B, Wang Z, Yin L, Yan Z, *et al.* (2014): The versatile functions of Sox9 in development, stem cells, and human diseases. *Genes Dis* 1: 149–161.
72. Li X-L, Liu L, Li D-D, He Y-P, Guo L-H, Sun L-P, *et al.* (2017): Integrin β 4 promotes cell invasion and epithelial-mesenchymal transition through the modulation of Slug expression in hepatocellular carcinoma. *Sci Rep* 7: 40464.
73. Liu C, Goswami M, Talley J, Chessier-Martinez PL, Lou C-H, Sater AK (2012): TAK1 promotes BMP4/Smad1 signaling via inhibition of erk MAPK: a new link in the FGF/BMP regulatory network. *Differentiation* 83: 210–219.

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

- 649 74. Panaliappan TK, Wittmann W, Jidigam VK, Mercurio S, Bertolini JA, Sghari S, *et al.* (2018): Sox2 is
650 required for olfactory pit formation and olfactory neurogenesis through BMP restriction and
651 Hes5 upregulation. *Development* 145. <https://doi.org/10.1242/dev.153791>
- 652 75. Poché RA, Furuta Y, Chaboissier M-C, Schedl A, Behringer RR (2008): Sox9 is expressed in mouse
653 multipotent retinal progenitor cells and functions in Müller glial cell development. *J Comp*
654 *Neurol* 510: 237–250.
- 655 76. Shakiba N, White CA, Lipsitz YY, Yachie-Kinoshita A, Tonge PD, Hussein SMI, *et al.* (2015): CD24 tracks
656 divergent pluripotent states in mouse and human cells. *Nat Commun* 6: 7329.
- 657 77. Stolt CC, Lommes P, Sock E, Chaboissier M-C, Schedl A, Wegner M (2003): The Sox9 transcription
658 factor determines glial fate choice in the developing spinal cord. *Genes Dev* 17: 1677–1689.
- 659 78. Suzuki Y, Montagne K, Nishihara A, Watabe T, Miyazono K (2008): BMPs promote proliferation and
660 migration of endothelial cells via stimulation of VEGF-A/VEGFR2 and angiopoietin-1/Tie2
661 signalling. *J Biochem* 143: 199–206.
- 662 79. Oldham MC, Konopka G, Iwamoto K, Langfelder P, Kato T, Horvath S, Geschwind DH (2008):
663 Functional organization of the transcriptome in human brain. *Nat Neurosci* 11: 1271–1282.
- 664 80. Fontanet PA, Ríos AS, Alsina FC, Paratcha G, Ledda F (2018): Pea3 Transcription Factors, Etv4 and
665 Etv5, Are Required for Proper Hippocampal Dendrite Development and Plasticity. *Cereb Cortex*
666 28: 236–249.
- 667 81. Jaworski JN, Jones DC (2006): The role of CART in the reward/reinforcing properties of
668 psychostimulants. *Peptides* 27: 1993–2004.
- 669 82. Young LJ, Wang Z (2004): The neurobiology of pair bonding. *Nat Neurosci* 7: 1048–1054.
- 670 83. Higa KK, Young JW, Ji B, Nichols DE, Geyer MA, Zhou X (2017): Striatal dopamine D1 receptor
671 suppression impairs reward-associative learning. *Behav Brain Res* 323: 100–110.

84. Milienne-Petiot M, Groenink L, Minassian A, Young JW (2017): Blockade of dopamine D1-family receptors attenuates the mania-like hyperactive, risk-preferring, and high motivation behavioral profile of mice with low dopamine transporter levels. *J Psychopharmacol (Oxford)* 31: 1334–1346.
85. Spittau B, Rilka J, Steinfath E, Zöller T, Krieglstein K (2015): TGFβ1 increases microglia-mediated engulfment of apoptotic cells via upregulation of the milk fat globule-EGF factor 8. *Glia* 63: 142–153.
86. Kettenmann H, Kirchhoff F, Verkhratsky A (2013): Microglia: new roles for the synaptic stripper. *Neuron* 77: 10–18.
87. Zeisel A, Hochgerner H, Lonnerberg P, Johnsson A, Memic F, Zwan J van der, *et al.* (2018): Molecular architecture of the mouse nervous system. *bioRxiv* 294918.
88. Fischer S, Ehlert U (2018): Hypothalamic-pituitary-thyroid (HPT) axis functioning in anxiety disorders. A systematic review. *Depress Anxiety* 35: 98–110.
89. Maes M, Vandewoude M, Maes L, Schotte C, Cosyns P (1989): A revised interpretation of the TRH test results in female depressed patients. Part I: TSH responses. Effects of severity of illness, thyroid hormones, monoamines, age, sex hormonal, corticosteroid and nutritional state. *J Affect Disord* 16: 203–213.
90. Lanni A, Moreno M, Lombardi A, Goglia F (2003): Thyroid hormone and uncoupling proteins. *FEBS Lett* 543: 5–10.
91. Choi J, Kim J, Kim T-K, Park J-Y, Lee J-E, Kim H, *et al.* (2015): TRH and TRH receptor system in the basolateral amygdala mediate stress-induced depression-like behaviors. *Neuropharmacology* 97: 346–356.

92. Cichon S, Mühleisen TW, Degenhardt FA, Mattheisen M, Miró X, Strohmaier J, *et al.* (2011): Genome-wide association study identifies genetic variation in neurocan as a susceptibility factor for bipolar disorder. *Am J Hum Genet* 88: 372–381.
93. Miró X, Meier S, Dreisow ML, Frank J, Strohmaier J, Breuer R, *et al.* (2012): Studies in humans and mice implicate neurocan in the etiology of mania. *Am J Psychiatry* 169: 982–990.
94. Rauch U, Feng K, Zhou XH (2001): Neurocan: a brain chondroitin sulfate proteoglycan. *Cell Mol Life Sci* 58: 1842–1856.
95. Cole AE, Murray SS, Xiao J (2016): Bone Morphogenetic Protein 4 Signalling in Neural Stem and Progenitor Cells during Development and after Injury. *Stem Cells Int* 2016: 9260592.
96. Harland R (2000): Neural induction. *Curr Opin Genet Dev* 10: 357–362.
97. Bond AM, Bhalala OG, Kessler JA (2012): The dynamic role of bone morphogenetic proteins in neural stem cell fate and maturation. *Dev Neurobiol* 72: 1068–1084.
98. Brooker SM, Gobeske KT, Chen J, Peng C-Y, Kessler JA (2017): Hippocampal bone morphogenetic protein signaling mediates behavioral effects of antidepressant treatment. *Mol Psychiatry* 22: 910–919.
99. Gobeske KT, Das S, Bonaguidi MA, Weiss C, Radulovic J, Disterhoft JF, Kessler JA (2009): BMP signaling mediates effects of exercise on hippocampal neurogenesis and cognition in mice. *PLoS ONE* 4: e7506.
100. Meyers EA, Gobeske KT, Bond AM, Jarrett JC, Peng C-Y, Kessler JA (2016): Increased bone morphogenetic protein signaling contributes to age-related declines in neurogenesis and cognition. *Neurobiol Aging* 38: 164–175.
101. Mira H, Andreu Z, Suh H, Lie DC, Jessberger S, Consiglio A, *et al.* (2010): Signaling through BMPRII regulates quiescence and long-term activity of neural stem cells in the adult hippocampus. *Cell Stem Cell* 7: 78–89.

102. Gomes WA, Mehler MF, Kessler JA (2003): Transgenic overexpression of BMP4 increases astroglial and decreases oligodendroglial lineage commitment. *Dev Biol* 255: 164–177.
103. Caronia G, Wilcoxon J, Feldman P, Grove EA (2010): Bone morphogenetic protein signaling in the developing telencephalon controls formation of the hippocampal dentate gyrus and modifies fear-related behavior. *J Neurosci* 30: 6291–6301.
104. García-Fuster MJ, Perez JA, Clinton SM, Watson SJ, Akil H (2010): Impact of cocaine on adult hippocampal neurogenesis in an animal model of differential propensity to drug abuse. *Eur J Neurosci* 31: 79–89.
105. Duman RS, Monteggia LM (2006): A neurotrophic model for stress-related mood disorders. *Biol Psychiatry* 59: 1116–1127.
106. Bansod S, Kageyama R, Ohtsuka T (2017): Hes5 regulates the transition timing of neurogenesis and gliogenesis in mammalian neocortical development. *Development* 144: 3156–3167.
107. Sakai D, Suzuki T, Osumi N, Wakamatsu Y (2006): Cooperative action of Sox9, Snail2 and PKA signaling in early neural crest development. *Development* 133: 1323–1333.
108. Sun W, Cornwell A, Li J, Peng S, Osorio MJ, Aalling N, *et al.* (2017): SOX9 Is an Astrocyte-Specific Nuclear Marker in the Adult Brain Outside the Neurogenic Regions. *J Neurosci* 37: 4493–4507.
109. Ming G-L, Song H (2011): Adult neurogenesis in the mammalian brain: significant answers and significant questions. *Neuron* 70: 687–702.
110. Fuller ML, DeChant AK, Rothstein B, Caprariello A, Wang R, Hall AK, Miller RH (2007): Bone morphogenetic proteins promote gliosis in demyelinating spinal cord lesions. *Ann Neurol* 62: 288–300.
111. Yan H, Zhu X, Xie J, Zhao Y, Liu X (2016): β -amyloid increases neurocan expression through regulating Sox9 in astrocytes: A potential relationship between Sox9 and chondroitin sulfate proteoglycans in Alzheimer’s disease. *Brain Res* 1646: 377–383.

112. Schafer DP, Stevens B (2015): Microglia Function in Central Nervous System Development and Plasticity. *Cold Spring Harb Perspect Biol* 7: a020545.
113. Wu Y, Dissing-Olesen L, MacVicar BA, Stevens B (2015): Microglia: Dynamic Mediators of Synapse Development and Plasticity. *Trends Immunol* 36: 605–613.
114. Albus E, Sinnigen K, Winzer M, Thiele S, Baschant U, Hannemann A, *et al.* (2016): Milk Fat Globule-Epidermal Growth Factor 8 (MFG-E8) Is a Novel Anti-inflammatory Factor in Rheumatoid Arthritis in Mice and Humans. *J Bone Miner Res* 31: 596–605.
115. Fricker M, Neher JJ, Zhao J-W, Théry C, Tolkovsky AM, Brown GC (2012): MFG-E8 mediates primary phagocytosis of viable neurons during neuroinflammation. *J Neurosci* 32: 2657–2666.
116. Fuller AD, Van Eldik LJ (2008): MFG-E8 regulates microglial phagocytosis of apoptotic neurons. *J Neuroimmune Pharmacol* 3: 246–256.
117. Vogler S, Pahnke J, Rousset S, Ricquier D, Moch H, Miroux B, Ibrahim SM (2006): Uncoupling protein 2 has protective function during experimental autoimmune encephalomyelitis. *Am J Pathol* 168: 1570–1575.
118. Arsenijevic D, Onuma H, Pecqueur C, Raimbault S, Manning BS, Miroux B, *et al.* (2000): Disruption of the uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species production. *Nat Genet* 26: 435–439.
119. Jo WK, Zhang Y, Emrich HM, Dietrich DE (2015): Glia in the cytokine-mediated onset of depression: fine tuning the immune response. *Front Cell Neurosci* 9: 268.
120. Sato K (2015): Effects of Microglia on Neurogenesis. *Glia* 63: 1394–1405.
121. Mastellos DC (2014): Complement emerges as a masterful regulator of CNS homeostasis, neural synaptic plasticity and cognitive function. *Exp Neurol* 261: 469–474.

122. Maras PM, Daugherty J, Hebda-Bauer E, Watson SJ, Akil H (2018): Minocycline treatment reduces anxiety- and depressive-like behaviors in a genetic model of internalizing mood disorders. *Neuroscience Meeting Planner* Program No. 234.03.
123. Hilde KL, Hagenauer MH, Stefanov AV, Birt I, Hebda-Bauer E, Watson SJ, Akil H (2018): Mechanisms of hippocampal development: Molecular and cellular phenotypes in the selectively bred high-responder/low-responder model of affective disorders. *Neuroscience Meeting Planner* Program No. 233.04.
124. O'Connor AM, Pardo T, Birt I, Hagenauer MH, Maras PM, Prater KE, *et al.* (2018): The hippocampal glycome contributes to behavioural phenotype in a novel rodent model of mood disorders. *Neuroscience Meeting Planner* Program No. 233.01.
125. Clinton SM, Stead JDH, Miller S, Watson SJ, Akil H (2011): Developmental underpinnings of differences in rodent novelty-seeking and emotional reactivity. *Eur J Neurosci* 34: 994–1005.

Tables and Table Legends

	MBNI_AffymetrixRae230_F4	MBNI_AffymetrixRgU34A_F6	MBNI_AffymetrixRae230_F15	MBNI_IlluminaRatRef12v1_F15	MBNI_RNASeq_F29	Alabama_Nimblegen_F34	MBNI_RNASeq_F37	MBNI_RNASeq_F43
Generation	4	6	15	15	29	34	37	43
Laboratory	MBNI	MBNI	MBNI	MBNI	MBNI	Alabama	MBNI	MBNI
Lead Scientist	Dr. John Stead	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Peter Blandino	Dr. Cigdem Aydin
Age	Adult	P7, P14, P21	P14	P14	P14, Adult	P7, P14, P21, Adult	Adult	Adult
n per group	6	6	6	6	2	5	6	5
Platform	Affymetrix Rat Expression Set 230 A	Affymetrix Rat Genome (RG) U34A GeneChips	Affymetrix Rat Expression Set 230 A	Illumina RatRef-12v1 Beadchip	RNA-Seq	NimbleGen Rat Gene Expression 12x135	RNA-Seq	RNA-Seq
Tissue Extraction	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Dorsal Hippocampus Tissue Punch	Whole Hippocampus	Whole Hippocampus
Exposure	Basal	Basal	Basal	Basal	Basal	Basal	Basal	Vehicle Injections
Behavioral Testing	Basal	Basal	Basal	Basal	Basal	Basal	Locomotor, Anxiety	Social Interaction After Mild Stress
GEO Data Release	GSE140596	GSE29552	GSE140595	GSE140594	GSE140597	GSE88874	GSE140598	GSE140287
Citation		Clinton <i>et al.</i> 2011				Cohen <i>et al.</i> 2015		

Table 1. An overview of the eight transcriptional profiling studies included in our current meta-analyses of differential gene expression in the bHR and bLR hippocampus at four developmental time points: P7, P14, P21, and adulthood. Citations: (28,125)

Figures and Figure Legends

Figure 1. An overview of the experimental and analytical workflow used to identify top candidate genes for mediating the effects of selective breeding on bHR/bLR phenotype.

Left: Many generations of selective breeding based on exploratory locomotor behavior drove segregation of genetic variants that contribute to internalizing and externalizing behavior within our bHR and bLR rats. The effect of these variants on behavior is mediated by alterations in gene expression and cellular function, which produce local changes in cell type balance and structure within brain regions responsible for affective behavior, such as the hippocampus.

Right: Our concurrent genetic study used exome sequencing to identify genetic variants that segregated bHR/bLR rats in our colony, and then used a sampling of those variants to locate regions of the genome (quantitative trait loci – QTLs) that might contribute to exploratory locomotor activity (25). Our current study used meta-analyses of hippocampal transcriptional profiling studies to identify bHR/bLR differentially-expressed (DE) genes, pathways, cell types, and networks in development and adulthood. In our results, we emphasize DE genes that were 1) consistently DE across multiple developmental stages, 2) central to DE pathways, cell types, and networks, 3) located near genetic variants that segregated bHR/bLR rats in our colony and/or within QTLs for exploratory locomotion. These genes are the top candidates for mediating the effects of selective breeding on bHR/bLR phenotype.

Figure 2. Selectively-bred high responder (bHR) and low responder (bLR) rats model an extreme propensity for internalizing vs. externalizing behavior. A) Over the course of 56 generations of selective breeding (F1-F56), the bHR rats (green) have developed increasingly elevated exploratory activity in a novel field (*y-axis*: average total locomotor score), whereas the bLR rats (red) have become less exploratory. These trends plateaued after F42, when our breeding strategy changed to deaccelerate divergence. Arrows indicate the generations during which hippocampal transcriptomic profiling datasets were collected, along with a name indicating the respective laboratory, platform, and generation for each dataset. **B)** bLR rats have

been highly anxious since the initiation of our breeding colony. The example above is from the behavioral data accompanying the MBNI_RNASeq_F37 transcriptomic dataset showing bLRs spending a smaller percentage of time in the anxiogenic open arms of the elevated plus maze than bHR rats (effect of phenotype: $F(2,15)=6.72$, $p=8.25E-03$, boxes=first quartile, median, and third quartile, whiskers = range). **C)** bLR rats are more reactive to stressors. This example is from the behavioral data accompanying the MBNI_RNASeq_F43 transcriptomic dataset showing bLR rats spending a smaller percentage of time interacting socially following exposure to a single mild stressor ($F(1,8)=5.86$, $p=0.0418$).

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

Adult Meta-Analysis: Top Results (FDR<0.05)											
Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets	Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets
1	Tmem144	3.04E-08	4.95E-04	-3.57	3	38	Trappc6a	6.16E-05	2.61E-02	-2.06	3
2	Asb15	4.05E-07	2.32E-03	2.82	3	39	Cav1	6.47E-05	2.61E-02	1.37	5
3	Kif15	4.27E-07	2.32E-03	-2.21	4	40	Robo3	6.54E-05	2.61E-02	-2.00	3
4	Pkhd1l1	6.13E-07	2.49E-03	-3.17	3	41	C1qa	6.57E-05	2.61E-02	-1.40	5
5	Car9	7.87E-07	2.56E-03	2.78	3	42	Tubg1	6.88E-05	2.65E-02	-1.40	5
6	Klhl34	1.04E-06	2.81E-03	-2.39	4	43	Zfp952	7.17E-05	2.65E-02	2.12	2
7	Mvb12b	2.03E-06	4.32E-03	-2.95	2	44	Slc9a3r1	7.31E-05	2.65E-02	1.32	5
8	Amer2	2.12E-06	4.32E-03	2.10	3	45	Egfem1	7.45E-05	2.65E-02	-2.11	2
9	Rpl17	2.86E-06	5.17E-03	-2.16	4	46	Epb41l4a	7.49E-05	2.65E-02	-2.10	2
10	Nudt4	4.27E-06	6.51E-03	-1.58	5	47	Trmt10a	8.18E-05	2.83E-02	-1.75	3
11	Pkib	4.40E-06	6.51E-03	1.70	5	48	Oard1	8.81E-05	2.98E-02	1.65	3
12	Fxyd7	5.22E-06	6.62E-03	-1.74	4	49	Selenop	1.00E-04	3.33E-02	2.21	2
13	Scube2	5.29E-06	6.62E-03	-2.68	2	50	Ezr	1.17E-04	3.72E-02	1.27	5
14	Ddx20	5.77E-06	6.62E-03	-1.93	4	51	Acox3	1.17E-04	3.72E-02	1.26	5
15	R3hdm4	6.10E-06	6.62E-03	-2.03	3	52	Ints8	1.21E-04	3.74E-02	2.11	2
16	Hmgn5b	7.33E-06	6.94E-03	2.60	2	53	Pld4	1.22E-04	3.74E-02	-1.45	4
17	Slc39a12	7.35E-06	6.94E-03	2.62	3	54	Ucp2	1.27E-04	3.83E-02	1.29	5
18	Acss2	7.68E-06	6.94E-03	-1.87	4	55	Rbm3	1.29E-04	3.83E-02	-1.52	4
19	Rhpn2	8.16E-06	6.98E-03	2.33	3	56	Ankdd1b	1.37E-04	3.93E-02	-2.00	2
20	Exosc7	9.65E-06	7.85E-03	1.80	4	57	Aar2	1.38E-04	3.93E-02	2.02	2
21	Rnls	1.04E-05	8.06E-03	1.92	3	58	Fn3k	1.42E-04	3.97E-02	-1.47	4
22	Slc19a3	1.50E-05	1.11E-02	-2.36	3	59	Slc27a1	1.45E-04	4.01E-02	1.24	5
23	Prss55	1.91E-05	1.35E-02	-2.43	2	60	Afmid	1.52E-04	4.11E-02	1.98	2
24	Apln	2.07E-05	1.41E-02	1.48	5	61	Slc4a11	1.54E-04	4.11E-02	-1.78	3
25	Mfge8	2.70E-05	1.73E-02	1.50	5	62	Zfp110	1.64E-04	4.31E-02	1.24	5
26	C2cd3	2.84E-05	1.73E-02	2.06	3	63	Tek	1.70E-04	4.33E-02	-1.79	3
27	Dnaaf3	2.87E-05	1.73E-02	-2.51	2	64	Fmo5	1.70E-04	4.33E-02	1.42	4
28	Etv4	3.30E-05	1.92E-02	2.02	3	65	Blvra	1.74E-04	4.36E-02	1.52	4
29	Tmem2	3.72E-05	1.99E-02	-2.29	2	66	Prdm5	2.03E-04	4.94E-02	-1.76	3
30	Tdg	3.84E-05	1.99E-02	-1.38	5	67	Samd5	2.09E-04	4.94E-02	1.95	3
31	Mki67	4.03E-05	1.99E-02	-2.01	3	68	Ntn4	2.10E-04	4.94E-02	1.94	2
32	Tnnt1	4.06E-05	1.99E-02	-1.51	5	69	Chd1l	2.11E-04	4.94E-02	1.45	4
33	Zfp90	4.13E-05	1.99E-02	-1.59	4	70	Ccdc137	2.13E-04	4.94E-02	1.97	2
34	LOC363337	4.16E-05	1.99E-02	-2.82	2	71	Trhr	2.16E-04	4.94E-02	1.19	5
35	Uhrf1	4.95E-05	2.29E-02	-1.67	3	72	Zfp821	2.22E-04	4.98E-02	-1.48	3
36	Tuba8	5.06E-05	2.29E-02	-1.44	4	73	Myh6	2.23E-04	4.98E-02	-1.25	5
37	Rltpr	5.58E-05	2.45E-02	-2.28	2	74	Sp3	2.26E-04	4.98E-02	1.93	3

P7: Top Result (FDR<0.05)						P14: Top Result (FDR<0.15)					
Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets	Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets
1	Ncan	1.38E-07	4.86E-04	-4.16	2	1	Bmp4	9.40E-06	1.47E-01	-1.49	5

Figure 3. The top DE genes within the bHR/bLR hippocampal gene expression meta-analyses. P-value=nominal p-value, FDR=false detection rate, Estimate=estimated effect size (i.e., the difference in expression between bHR and bLR rats in units of standard deviation,

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

green/positive=higher expression in bHRs, red/negative= higher expression in bLRs), # of
datasets=number of datasets included in the meta-analysis for that gene.

A. Genetic Variant Near Ncan (<100 kb distant) Segregates bHR and bLR Rats

Nearby SNP (in Nr2c2ap):
16_20887728 (Rnor5)

	GG	GA	AA
bHR	12	0	0
bLR	0	7	5

B. Neurocan is particularly elevated in early development in bLR rats

C. Neurocan is more highly expressed in bLR rats at P7

D. Neurocan is more highly expressed in bLR rats at P14

Figure 4. The extracellular matrix constituent Neurocan (Ncan) has elevated expression in bLR rats at age P7. A) A genetic variant on Chr 16 nearby Ncan (<100 kb) segregated bHR and bLR rats (Fisher's exact test: $p=1.63E-07$). **B)** Boxplots illustrating the effect of age and bHR/bLR phenotype on Ncan expression (log(2) signal) in two microarray studies (boxes=median and interquartile range, whiskers=range, red=bLR, green=bHR). The effect of phenotype is most obvious at an age when Ncan is elevated in development (P7). **C-D)** Forest

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

plots showing that Ncan was more expressed in bLRs than bHRs (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis) **C**) in the P7 meta-analysis ($\beta=-4.16$, $p=1.38E-07$, $FDR=4.86E-04$), **D**) and nominally in the P14 meta-analysis ($\beta=-0.90$, $p=1.01E-02$, $FDR=7.24E-01$).

Figure 5. A regulator of proliferation and differentiation, Bone morphogenetic protein 4 (Bmp4), is more highly expressed in bLR rats than bHR rats at P14 and adulthood. A-B) Two forest plots showing that Bmp4 was consistently elevated in bLR rats (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis) at **A)** P14 ($\beta=-1.49$, $p=9.40E-06$, $FDR=1.47E-01$) and **B)** adulthood (adult: $\beta=-1.04$, $p=1.01E-03$, $FDR=9.38E-02$). This direction of effect mirrors findings in the literature that show that blocking the expression of Bmp4 in mice reduces anxiety

and depressive-like behavior (98). **C)** Using qPCR, we confirmed that bLRs showed greater *Bmp4* expression than bHRs at P14 and adulthood (P90) using hippocampal tissue from later generations (F51, F55). Log(2) fold change in *Bmp4* expression was calculated using the Livak method ($-\Delta\Delta Cq$, (43)), using *Gapdh* as the reference housekeeping gene and bLRs as the reference group (therefore the bLR mean is set to 0 in all panels). $**p < 0.005$ (Data release: doi: 10.6084/m9.figshare.10321658; P14: $Log(2)FC = -3.74$, $T(5.60) = -6.10$, $p = 0.00115$; P90: $T(8.74) = -6.87$, $p = 8.44E-05$). **D)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, we found that *Bmp4* showed a negative relationship with percent time in the open arms ($\beta = -0.034$, $R^2 = 0.28$, $p = 0.024$) and a positive relationship with the number of fecal boli produced on the EPM ($\beta = 0.32$, $R^2 = 0.29$, $p = 0.020$, *data not shown*).

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

A. bHR/bLR Differentially-Expressed Genes Are Often Located Close to Genetic Variants (SNPs) that Segregate bHR and bLR Rats

B. bHR/bLR Differentially-Expressed Genes Often Overlap QTLs for Exploratory Locomotion

C. Differentially-Expressed Genes within QTLs for Exploratory Locomotion:

QTLs from bHR/bLR F2 Intercross: Exploratory Locomotion				Overlapping QTLs from Rat Genome Database		
Chr	QTL range in MB* (LOD>3, +/-1MB)	LOD	Genes w/ FDR<0.1 in Meta-Analysis	Anxiety	Stress Response	Behavioral Despair
1	28-31	3.13			[53]	[56]
1	39-65	4.75	<i>Ezr</i>	[47]	[53]	
1	98-104	3.42	<i>Dbp</i>	[47],[48]	[53]	
1	115-132	7.76	<i>Hmgn5b</i>	[47],[48]	[53],[54]	
1	139-142	3.14	<i>Mfge8</i>	[47],[48]	[54]	
1	156-171	4.32	<i>Olr35, Spcs2, C2cd3, Ucp2</i>	[47],[48]	[54]	
2	43-70	4.26	<i>Pdzd2, Selenop</i>	[49]	[55]	
2	92-95	3.02			[55], [55]	
3	43-65	4.71	<i>Ttc30a1</i>	[49]	[53]	[56]
5	98-102	3.34		[51],[52]		
7	67-85	5.14	<i>Trhr, Pkhd1l1</i>	[49]	[53]	
17	157-177	3.82	<i>Chrm3, Akrl1cl</i>			
18	71-81	4.99		[50]		

D. Top Genomic Regions Enriched with Differentially Expressed Genes:

Genetic Loci Enriched with bHR/bLR Differentially Expressed Genes (PGE)							Overlapping QTLs:			
Chr	Location (Rnor_6.0)	P-value	FDR	Size of Region (kb)	Enrichment Score	Genes w/ FDR<0.1 in Meta-Analysis	Exploratory Locomotion	Anxiety	Stress Response	Behavioral Despair
1	2782900-4653300	7.76E-04	1.02E-02	1,870	0.6	<i>Samd5</i>				[56]
1	164225800-165684000	4.97E-05	4.68E-03	1,458	0.2	<i>Olr35, Spcs2, C2cd3, Ucp2</i>	[25]	[47], [48]	[54]	
1	174385400-174620200	1.77E-04	5.69E-03	235	1	<i>Scube2</i>		[48]	[63]	
3	62800900-63455200	7.76E-04	1.02E-02	654	0.6	<i>Ttc30a1</i>	[25]	[49]	[53]	[56]
3	119361600-119677400	1.22E-04	5.69E-03	316	0.6	<i>Blvra</i>			[53],[55]	[56]
3	151032900-151150600	1.77E-04	5.69E-03	118	1	<i>Acsc2, Myh7b</i>			[55]	
3	176280000-176526000	4.08E-04	7.97E-03	246	0.8					
5	155246400-155264100	1.77E-04	5.69E-03	18	1	<i>C1qa, C1qc</i>		[49]		
7	81919900-83358500	2.02E-04	6.05E-03	1,439	0.5	<i>Trhr</i>	[25]	[49]	[53]	
10	88392200-103730100	6.80E-05	4.68E-03	15,338	0.1	<i>Slc9a3r1, Sox9, Tubg1, Ghdc, Tmem101, Acbd4, Ddx42, Etv4</i>		[50]	[64], [65]	
12	37444100-39302600	4.30E-06	2.96E-03	1,859	0.3	<i>Eif2b1</i>				
16	81035000-81209500	3.35E-05	4.68E-03	175	0.8				[55]	
19	32182900-42110300	2.85E-05	4.68E-03	9,927	0.1	<i>Zfp90</i>				

Figure 6. Many of the top differentially expressed genes are located near genetic variants that segregate bHR/bLR rats within quantitative trait loci (QTLs) for exploratory locomotor activity. **A.** Our concurrent genetic study used exome sequencing to identify variants (SNPs) that segregated bHR/bLR rats in our colony (25). A high percentage of bHR/bLR differentially-expressed genes (adult meta-analysis: $FDR < 0.05$) were found within +/- 100 kb, 250 kb, 500 kb, or 1 MB of these segregating variants, using either traditional (Bonferonni-corrected $p < 0.05$) or more stringent (Bonferonni-corrected $p < 5.00E-5$) criteria to define segregation. Asterisks (*) designate enrichment (Fisher's exact test $p < 0.0001$) in comparison to the non-differentially expressed genes in our meta-analysis ($FDR > 0.05$). **B.** Our concurrent genetic study used a sampling of bHR/bLR segregating variants to identify QTLs for exploratory locomotor activity in a novel field using an bHR/bLR F2 intercross (25). bHR/bLR differentially-expressed genes (adult meta-analysis: $FDR < 0.05$) were 4.5x more likely to overlap (+/- 1MB) QTLs for exploratory locomotion than other genes included in our meta-analysis (Fisher's exact test: $p = 0.0035$). **C.** A table illustrating the top genes from our meta-analyses ($FDR < 0.1$, bold+italic= $FDR < 0.05$) that overlap (+/- 1 MB) significant ($LOD > 4$) and putative ($LOD > 3$) QTLs for exploratory locomotion identified by our concurrent genetic study (25). Also depicted is overlap with QTLs identified in the Rat Genome Database (32) for the following behaviors relevant to the bHR/bLR phenotype: anxiety (47–52), stress-related responses (53–55), and behavioral despair (56). **D.** The top chromosomal loci enriched for bHR/bLR DE genes overlap previously-identified QTLs relevant to externalizing and internalizing behaviors. The top chromosomal loci enriched for bHR/bLR DE genes were identified using Positional Gene Enrichment analysis (PGE): location (full coordinates), p-value=nominal p-value, FDR =false detection rate, enrichment score= ratio of genes with $p < 0.01$ out of all genes in the region. Also depicted is the overlap (+/- 1 MB) of these enriched chromosomal loci with QTLs for exploratory locomotor activity (25), as well as with QTLs identified in the Rat Genome Database (32) for anxiety (47–50), stress-related responses (53–55,63–65), and behavioral despair (56).

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

Figure 7. A region on chromosome 1 implicated in bHR/bLR phenotype contains two genes important for brain function and development, Uncoupling Protein 2 (Ucp2) and C2 Calcium Dependent Domain Containing 3 (C2cd3). A) Genetic variants on Chr 1 within Ucp2 and C2cd3 segregate bHR and bLR rats in our colony (Rnor5 coordinates, Fisher's exact test: SNP 1_171712284: $p=1.66E-09$; SNP 1_171672484: $p=1.27E-13$). **B)** Ucp2 and C2cd3 are located on Chr 1 within a QTL for exploratory locomotor activity. An example of the correlation between genetic variation in this region and behavior is illustrated using the sequencing results

from a nearby SNP and exploratory locomotor activity measured in a bHRxbLR F2 intercross ($n=310$; $adj.R^2=0.049$, $p=2.00E-04$, $FDR=0.0048$). **C)** C2cd3 and Ucp2 were the top DE genes ($FDR<0.05$) within a segment of Chr 1 enriched for DE genes and containing a QTL for exploratory locomotor activity (25). The table illustrates the DE genes within this region: estimate=estimated effect size (green/positive=higher expression in bHRs), bold= $p<0.05$, bold+italic= $FDR<0.05$. **D)** A forest plot showing that Ucp2 had higher expression in bHRs in four out of the five adult datasets included in the adult meta-analysis (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis, effect of phenotype: $\beta=1.29$, $p=1.27E-04$, $FDR=3.83E-02$). This direction of effect mirrors findings in the literature showing that Ucp2 knockout mice have higher anxiety-like behavior and lower locomotor activity, as well as greater sensitivity to stress (57,58,60,61), much like our bLR rats. **E)** A forest plot showing that C2cd3 had higher expression in bHRs in three adult datasets included in the adult meta-analysis (effect of phenotype: $\beta=2.06$, $p=2.84E-05$, $FDR=1.73E-02$). **F)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, C2cd3 (units: log(2) fragments per million (FPM)) showed a positive relationship with exploratory locomotor activity ($\beta=0.000109$, $R^2=0.26$, $p=3.20E-02$). **G)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Ucp2 showed a trend towards a positive relationship with percent of time spent in the anxiogenic open arms of the EPM ($\beta=0.03$, $R^2=0.22$, $p=5.16E-02$).

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

A. Genetic Variant Near Trhr (<200 kb distant) Segregates bHR and bLR Rats

Nearby SNP (in Nudcd1):

7_83331058 (Rnor5)			
	CC	CT	TT
bHR	0	0	12
bLR	12	0	0

B. Trhr is Located within a QTL on Chr 7 for Exploratory Locomotor Activity

Nearby SNP in QTL:
7_83331058 (Rnor5)

C. Trhr is More Highly Expressed in bHR Rats in Adulthood

F. Trhr Correlates with Exploratory Activity

R-squared= 0.22 , P-value= 0.051

D. Trhr is the Top Result in a Section of Chr 7 Enriched for Differentially-Expressed Genes

G. Trhr is the Top Result Within a Hippocampal Co-expression Network Enriched for Differentially-Expressed Genes

E. Trhr is a Top Result in Two Hippocampal-Specific Gene Sets Enriched with bHR/bLR Differential Expression

Top Co-expression Networks in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
From mouse: "paleturquoise"	1.94	-1.08	ADAMTS2,FOXC1,FBLN1,MRC1,OMD,ANGPT2,EMP3,BGN,CD248,HSPB1
From human: "M11"	-1.72	-1.84	FBXO31,NPHP4,PPME1,ARHGAP39,FAM131B,PLXNA1,NCOR2,PPFIA4,DMTN,STRN4
From mouse: "darkred"	-1.68	0.98	MTF2,DUSP11,SNHG11,DDX26B,LRR29,RYR3,NFKBIZ,EML5,TCF25,NRXN3
From human: "M24"	1.54	-0.97	EZR,ITSN1,IDNK,PPAP2B,FAM167A,GFPT2,ITGB4,PSD2,PITPNC1,PHGDH
From mouse: "sienna3"	1.91	-0.64	OTOF,TRHR,SMOC1,ZBTB7C,GLRA2,GRB10,MEIS1,GPR83,ARHGAP6
From mouse: "lightcyan"	-1.27	-1.47	RLTPR,ATMIN,KAT2A,JPH3,ACAD8,REV1,USE1,B9D2,RAB3D,PWWP2B

Top Co-expression Networks at P14 (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
From mouse: "lightyellow"	-1.13	-1.56	ZFP277,ACTA2,DMWD,SEMA7A,UQCRC1,CNNM3,ATM,NPAS2,NCAN,BRD4
From human: "M11"	-1.72	-1.84	DTX3,FBXO31,GFOD1,HYOU1,ADCY9,LRFN4,DYRK1B,PTMS,UBA1,WBSCR17
From mouse: "lightcyan"	-1.27	-1.47	H1FX,ELFN2,ZCRB1,APBA2,SLC2A13,POLK,SNUPN,ATG5,HYOU1,SNAPIN
From human: "M1"	NA	-1.36	EID2,CHST15,RIMBP2,EMX1,NT5C3B,ELFN2,FAM65A,SRSF7,ZDHHC22,APBA2
From mouse: "skyblue"	0.86	-1.84	TSC22D4,MOBP,ADSSL1,SLC35B2,OPALIN,MAG,EFHD1,GJC2,ITGB4
From human: "M7"	-1.45	-1.70	ARHGEF4,STGALNAC6,DHRS11,AGAP3,B3GAT1,ARHGEF25,LRP3,PLSCR3,LMO4,NCS1

Top Regional Markers in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
"dg_v VS dg_d"	2.05	0.88	TRHR,TOX3,PDZRN3,FBLN1,TSHZ1,STK32B,ZFP423,NCAPG,LCORL

Top Regional Markers at P14 (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
"dg_d-dg_v VS ca4-ca3_d-ca3_v-ca2-ca1_d-ca1_v"	-0.88	-1.76	SCN3A,ARHGAP20,MARCKS,SV2C,PDE7B,TRPC6,SLC4A4,BTG2,IGFBP5,RF3

Figure 8. Thyrotropin releasing hormone receptor (Trhr) was the top gene within two hippocampal specific gene sets and within a region of Chromosome 7 implicated in bHR/bLR phenotype. A) A genetic variant on Chr 7 near Trhr (<200 kb distant) segregates bHR and bLR rats in our colony (Fisher's exact test: $p=6.20E-14$). **B)** Trhr is located on Chr 7 within a QTL for exploratory locomotor activity. An example of the correlation between genetic variation in this region and behavior is illustrated using the sequencing results from a SNP nearby Trhr (discussed above) and exploratory locomotor activity measured in a bHRxbLR F2 intercross ($n=315$, $adj.R^2=0.061$, $p=5.79E-05$, $FDR=0.00178$). **C)** A forest plot showing that Trhr expression was consistently elevated in bHR rats since generation F4 (boxes=Cohen's D from each study $\pm$ 95% confidence intervals, "Model"=estimated effect size $\pm$ 95% confidence intervals provided by the meta-analysis, $\beta=1.19$, $p=2.16E-04$, $FDR=4.94E-02$). This direction of effect mirrors findings in the literature that show Trhr-KO mice exhibit greater anxiety and depressive-like behavior (62). **D)** Trhr was the strongest result within a segment of Chromosome 7 enriched for DE genes and overlapping a QTL for exploratory locomotor activity. The table illustrates the DE genes within this region: estimate=estimated effect size (green/positive=greater expression in bHRs), bold= $p<0.05$, bold+italic= $FDR<0.05$. **E)** Trhr was a leading gene in two of the top hippocampal-specific gene sets identified as enriched for bHR/bLR DE genes by GSEA (FDR: False Detection Rate; NES: Normalized Enrichment Score, with positive scores (green) indicating greater expression in bHRs and negative scores (red) indicating greater expression in bLRs, bold: $p<0.05$ in GSEA results, bold+italics: $FDR<0.05$ in GSEA results). The top 10 "Leading Edge" genes for each gene set are shown (bold+italics: $FDR<0.05$ in meta-analysis). These genes have large estimated effect sizes within the meta-analysis and help drive the enrichment of effects within these gene sets. Regional marker gene sets use the following abbreviations: "dg"=dentate gyrus, "v"=ventral, "d"=dorsal, "ca"=Cornu Ammonis subregion. **F)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Trhr (units: log(2) fragments per million (FPM)) showed a trend towards a positive relationship

with exploratory locomotor activity ($\beta=4.48\text{E-}04$, $R^2= 0.22$, $p=5.08\text{E-}02$). **G**) *Trhr* and its ligand, Thyrotropin releasing hormone (Trh), are hub genes within a hippocampal specific co-expression network that is enriched for bHR-upregulated genes. Genes within this network with known protein-protein interactions are illustrated above (STRINGdb: confidence setting=0.15 due to hippocampal co-expression already suggesting potential interaction). Many of these genes have documented associations with reward behavior.

B. An Unbiased PPI-Network Constructed from the Top Results Highlights Genes Involved in Cell Proliferation

Mki-67 Correlates with Exploratory Activity

Figure 9. The top bHR vs. bLR DE results are enriched with genes related to cell proliferation, differentiation, and development, including the canonical Marker of

Proliferation (Mki67). A) A table of the top functional ontology gene sets identified as enriched

for bHR/bLR DE genes by GSEA (FDR: False Detection Rate; NES: Normalized Enrichment Score, with positive scores (green) indicating higher expression in bHRs and negative scores (red) indicating higher expression in bLRs, bold: $p < 0.05$ in GSEA results, bold+italics:

$FDR < 0.05$ in GSEA results). The top 10 “Leading Edge” genes for each gene set are shown (bold+italics: $FDR < 0.05$ in meta-analysis). These genes have large estimated effect sizes within

the meta-analysis and help drive the enrichment of effects within these gene sets. **B)** A PPI

network constructed using the top genes from the adult meta-analysis (192 genes with $FDR < 0.10$, STRINGdb: confidence setting=0.40) had a dominant subnetwork that included

Bmp4 and Mki67 as hub genes. Many of these genes are related to cell proliferation and differentiation within the brain. **C)** Two genetic variants on Chr 1 within Mki-67 fully segregated

bHR and bLR rats in our colony (Rnor5 coordinates, Fisher’s exact test: SNP 1_214939984:

$p = 2.02E-11$, SNP 1_214940284: $p = 2.02E-11$). **D)** A forest plot showing that Mki67 was

consistently elevated in bLR rats in adulthood (boxes=Cohen’s D from each study +/-95% confidence intervals, “Model”=estimated effect size +/-95% confidence intervals provided by the

meta-analysis, adult: $\beta = -2.01$, $p = 4.03E-05$, $FDR = 1.99E-02$). **E)** Within the behavioral data

accompanying the MBNI_RNASeq_F37 dataset, Mki67 (units=log(2) fragments per million

(FPM)) showed a negative relationship with locomotor score ($\beta = -0.000249$, $R^2 = 0.41$, $p = 0.0042$).

Running Head: bHR/bLR Hippocampal Gene Expression Meta-Analysis

bLRs Exhibit Overall Elevated Endothelial and Microglia Specific Gene Expression:

Mfge8, which Promotes Alternative (M2) Activation of Microglia, is a Top bHR/bLR Candidate Gene

E. Genetic Variant in Mfge8 Segregates bHR and bLR rats

SNP within Mfge8:
1, 141829591 (Rnor5)

TT	TA	AA
bHR	12	0
bLR	0	6

F. Mfge8 is More Highly Expressed in bLRs in Adulthood

H. Mfge8 Correlates with Exploratory Activity and Anxiety

C1qa, which Promotes Classical Activation of Microglia, is a Top bHR/bLR Candidate Gene:

I. Genetic Variant Near C1qa (500 kb distant) Segregates bHR and bLR Rats

Nearby SNP (in C1a3b):
5, 159525078 (Rnor5)

CC	CA	AA
bHR	9	2
bLR	0	1

J. C1qa is More Highly Expressed in bLRs in Adulthood

K. C1qa is the Top Result in a Region of Chr 5 Enriched for Differentially-Expressed Genes

L. C1qa Correlates with Exploratory Activity

Figure 10. Microglial-related gene expression differentiates bHR and bLR rats. A) A

heatmap illustrating the effect of bHR/bLR phenotype on cell type specific gene expression, which can reflect overall cell type balance (42) or activation: green=upregulated in bHRs, red=upregulated in bLRs, asterisks(*) indicate $p < 0.05$. Note that only well-characterized non-neuronal cell type categories were included in this analysis. **B-D)** Forest plots (boxes=Cohen's D from each study $\pm$ 95% confidence intervals, "Model"=estimated effect size $\pm$ 95% confidence intervals provided by the meta-analysis) showing an upregulation in bLRs of **(B)** endothelial-specific gene expression at P14 ($\beta = -0.89$, $p = 5.75E-03$) and **(C-D)** microglial-specific gene expression **(C)** at P14 ($\beta = -0.69$, $p = 2.90E-02$) and **(D)** adulthood ($\beta = -0.65$, $p = 4.47E-02$). **E)** A genetic variant on Chr 1 in Milk fat globule-EGF factor 8 (Mfge8) segregates bHR and bLR rats in our colony (Fisher's exact test: $p = 7.53E-08$). Mfge8 promotes alternative (M2) activation of microglia. **F)** A forest plot illustrating that Mfge8 is more highly expressed in bHRs than bLRs in the adult meta-analysis in all five adult datasets ($\beta = 1.50$, $p = 2.70E-05$, $FDR = 1.73E-02$). **G)** Mfge8 is located on Chr 1 within a QTL for exploratory locomotor activity. An example of the correlation between genetic variation in this region and behavior is illustrated using the sequencing results from a nearby SNP (Rnor5 coordinates 1_141117448) and exploratory locomotor activity measured in a bHRxbLR F2 intercross ($n = 317$, $adj.R^2 = 0.061$, $p = 1.81E-05$, $FDR = 0.002$). **H)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Mfge8 (units=log(2) fragments per million (FPM)) showed a positive relationship with total locomotor score ($\beta = 0.000146$, $R^2 = 0.62$, $p = 1.10E-04$) as well as the percent time in the open arms of the EPM ($\beta = 0.00603$, $R^2 = 0.28$, $p = 2.30E-02$). **I)** A genetic variant on Chr 5 near Complement component C1q A Chain (500 kb distant) segregates bHR and bLR rats in our colony (Rnor5 coordinates 5_159525078, Fisher's exact test: $p = 1.09E-07$). C1qa promotes classical activation of microglia via the complement cascade. **J)** A forest plot illustrating that C1qa is more highly expressed in bLRs than bHRs in all five datasets in the adult meta-analysis ($\beta = -1.40$, $p = 6.57E-05$, $FDR = 2.61E-02$). **K)** C1qa is the top DE gene ($FDR < 0.05$) within a

segment of Chr 5 enriched for DE genes. The table illustrates the DE genes within this region:
estimate=estimated effect size (red/negative=higher expression in bLRs), bold= $p < 0.05$,
bold+italic= $FDR < 0.05$. L) Within the behavioral data accompanying the MBNI_RNASeq_F37
dataset, C1qa showed a negative relationship with total locomotor score (C1qa: $\beta = -5.17E-04$,
 $R^2 = 0.29$, $p = 2.09E-02$).