Sleep, regional grey matter volumes, and psychological functioning in adolescents

Adolescent sleep, brain structure, and function

Winok Lapidaire^{1&}, Anna S. Urrila^{1,2,3&}, Eric Artiges^{1,4}, Ruben Miranda¹, Hélène Vulser¹,
Pauline Bézivin-Frere¹, Hervé Lemaître¹, Jani Penttilä^{1,5}, Tobias Banaschewski⁶, Arun L.W.
Bokde⁷, Uli Bromberg⁸, Christian Büchel⁸, Patricia J. Conrod^{9,10}, Sylvane Desrivieres^{9,11},
Vincent Frouin¹², Jürgen Gallinat¹³, Hugh Garavan^{7,14}, Penny Gowland¹⁵, Andreas Heinz¹³,
Bernd Ittermann¹³, Dimitri Papadopoulos-Orfanos¹², Tomáš Paus¹⁶, Michael N. Smolka¹⁷,
Gunter Schumann^{7,11}, Marie-Laure Paillère Martinot^{1,18}, and Jean-Luc Martinot^{1,19*}, for the
IMAGEN consortium⁺

* Corresponding author

Email: jean-luc.martinot@inserm.fr

& These authors contributed equally to this work

Word count main text: 2954

Reference count: 40

Conflict of interest: There is no conflict of interest for any of the authors involved

Author contributions:

Winok Lapidaire: data analysis, writing, editing

23 Anna S. Urrila et al: writing, editing

24 Others: Editing

25 Imagen cohort: data collection

26

27 ¹ National Institute of Health and Medical Research, INSERM Unit 1000 ; “Neuroimaging &
28 Psychiatry”, University Paris Saclay, University Paris Descartes, Paris, France;

29 ² National Institute for Health and Welfare, Department of Health, Unit of Mental Health,
30 Helsinki, Finland;

31 ³ University of Helsinki and Helsinki University Central Hospital, Department of Psychiatry /
32 Adolescent Psychiatry, Helsinki, Finland;

33 ⁴ Department of Psychiatry 91G16, Orsay Hospital, Orsay, France;

34 ⁵ Adolescent Psychiatry Department, Medical school, Tampere University, Tampere, Finland;

35 ⁶ Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University,
36 Mannheim, Germany;

37 ⁷ Institute of Neuroscience, Trinity College Dublin, Dublin, Ireland;

38 ⁸ University Hospital Hamburg Eppendorf, Hamburg, Germany;

39 ⁹ Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, United
40 Kingdom

41 ¹⁰ Department of Psychiatry, University of Montreal, CHU Ste Justine Hospital, Montréal,
42 Canada;

43 ¹¹ MRC Social, Genetic and Developmental Psychiatry (SGDP) Centre, London, United
44 Kingdom;

45 ¹² Neurospin, Commission for Atomic and Alternative Energy, Saclay, France;

¹³ Department of Psychiatry and Psychotherapy, Campus Charité Mitte, Charité – University Medical Centre Berlin, Germany;

¹⁴ Departments of Psychiatry and Psychology, 6436 UHC, University of Vermont., Burlington, VT, USA;

¹⁵ Sir Peter Mansfield Magnetic Resonance Centre, University of Nottingham, Nottingham, United Kingdom;

¹⁶ Bloorview Research Institute, Holland Bloorview Kids Rehabilitation Hospital and Departments of Psychology and Psychiatry, University of Toronto, Canada;

¹⁷ Department of Psychiatry and Psychotherapy, Technical University Dresden, Germany;

¹⁸ APHP, Department of Child and Adolescent Psychiatry, Pitié-Salpêtrière Hospital, Paris, France

¹⁹ CENIR at ICM Institute, Centre of Neuroimaging Research, ICM building, Paris Cedex, France

⁺ Imagen Consortium other collaborators:

M Fauth-Bühler, L Poutska, F Nees, Y Grimmer, M Struve from Central Institute of Mental Health, Mannheim, Germany;

A Ströhle, V Kappel , B M van Noort, from Charité hospital, Berlin, Germany;

N Bordas, Z Bricaud, I Filippi, A Galinowski, F Gollier-Briant, Vincent Ménard, from INSERM, France;

A Cattrell, R Goodman, A Stringaris, C Nymberg, L Reed, from the Institute of Psychiatry, Psychology & Neuroscience, King’s College London, United Kingdom;

- 69 B Ittermann, R Brühl R, from Physikalisch-Technische Bundesanstalt (PTB), Berlin; Germany;
- 70 T Hübner, K Müller, from University of Dresden, Germany;
- 71 U Bromberg, J Gallinat, T Fadai, from University of Hamburg, Germany;
- 72 P Gowland, C Lawrence, from University of Nottingham, United Kingdom;
- 73

Abstract

Changing sleep rhythms in adolescents often lead to sleep deficits and increased variability in sleep schedules. The adolescent brain, and in particular the rapidly developing structures involved in emotional control, are vulnerable to external and internal factors. In our previous study in adolescents at age 14, we observed a strong relationship between weekend sleep schedules and regional medial prefrontal cortex grey matter volumes. Here, we aimed to assess whether this relationship remained in this group of adolescents of the general population at the age of 16 (n=101; mean age 16.8 years; 55% girls). We further examined grey matter volumes in the hippocampi and the amygdalae, calculated with voxel-based morphometry. In addition, we investigated the relationship between regional grey matter volumes with psychological functioning. Sleep was assessed with self-reports and psychological functioning with the Strengths and Difficulties Questionnaire and tests on working memory and impulsivity. Later weekend wake-up times were associated with smaller grey matter volumes in the medial prefrontal cortex and the amygdalae, and greater weekend delays in wake-up time were associated with smaller grey matter volumes in the right hippocampus and amygdala. The medial prefrontal cortex region mediated the correlation between weekend wake up time and both externalising and internalising symptoms. Paying attention to regular sleep habits during adolescence could act as a protective factor against the emergence of psychopathology via enabling favourable brain development.

96 **Abbreviations:** Anatomical Automatic Labeling (AAL); Alcohol Use Disorders Identification
 97 Test (AUDIT); Cambridge Automated Neuropsychological Test Battery (CANTAB); Family
 98 Wise Error (FWE); Magnetisation Prepared Rapid Acquisition Gradient Echo (MPRAGE);
 99 Medial Prefrontal Cortex (mPFC); Montreal Neurologic Institute (MNI); functional Magnetic
 100 Resonance Imaging (fMRI); Region Of Interest (ROI); Statistical Parametric Mapping (SPM);
 101 Strengths and Difficulties Questionnaire (SDQ); Voxel-Based Morphometry (VBM); Weekday
 102 (WD); Weekend (WE);

Introduction

Sleep problems and psychiatric disorders increase sharply hand in hand during adolescence, but our understanding of the potential neurobiological links between them is only emerging (1).

Late sleep, sleep deprivation, and social jet lag (the variability of sleep patterns between weekdays and weekends) have all been associated with a broad range of negative mental health consequences, including increased depressive and anxiety symptoms, increased risk-taking behaviours, as well as deteriorated executive function (2–4). Furthermore, sleep disturbances seem to precede the onset of diverse psychiatric disorders (5).

These studies support the theory that unhealthy sleep habits could affect the developing adolescent brain structure and thereby increase the vulnerability to various kinds of psychopathologies, but few studies on the relationship between adolescents' sleep habits and brain grey matter volumes have been published to date. In a sample of maltreated teenagers, reduced sleep efficiency was recently found to correlate with reduced GMV in hippocampus, inferior frontal gyrus and insula, suggesting that sleep might mediate the negative impact of adverse life events on brain morphology (6). In a mixed sample of children and adolescents, weekday time in bed was found to correlate with regional grey matter volumes of the bilateral hippocampi and the dorsolateral prefrontal cortex (7). In our previous study of 14-year-old adolescents, we found late sleep during the weekend and short sleep during the week to be associated with smaller regional grey matter volumes, particularly in the medial prefrontal cortex (mPFC). In addition, there was a correlation between mPFC GMV and school performance (8). Since sleep characteristics and brain morphology undergo constant changes through adolescence

(9,10), it is important to study their interconnections repeatedly at different points of development.

The mPFC exerts an inhibitory top-down control of subcortical structures (11). Poor sleep and eveningness-prone or irregular sleep rhythms can negatively affect adolescents' emotion regulation, reward-related processing, and impulse inhibition by influencing the mPFC (12–14) as well as the amygdala and the hippocampus (15–19). These structures have also been implicated in the etiology and maintenance of psychiatric disorders (20,21). Studying the effects of sleep especially on the mPFC, the amygdala, and the hippocampus would thus crucially contribute to understanding the development of psychopathology during adolescence. Our general hypothesis is that adolescents' sleep patterns affect brain regional grey matter volumes, which in turn lead to lower psychological functioning or even mild psychopathology.

Understanding the trajectories that lead toward psychiatric disorders as early as possible in development would allow us to develop effective intervention and prevention strategies.

In this follow-up study we aimed to assess whether our previous findings on the correlation between adolescents' sleep habits and brain grey matter volumes remained present at the age of 16, and to extend these findings by examining their relationships with psychological functioning.

Methods

Participants

Participants were recruited from schools near Paris, France, based on their age and absence of any major somatic condition. Written consent was obtained from all subjects in this study. The

study was approved by the regional ethics committee (Comité de Protection des Personnes [CPP] Ile-de-France 7). The adolescents participated in a larger multi-centre study (<http://www.imagen-europe.com/en/the-imagen-study.php>) (22) at age 14 (baseline), and were followed up at age 16. Only the French adolescents were assessed for their sleep habits and were thus eligible for this study. Details of the sample at baseline have been previously reported (8). This study focuses on the sample at age 16, at which time point written informed assent and consent to study participation were obtained from a total of 138 adolescents and their parents, respectively. We excluded participants who did not complete the sleep questionnaires, those whose Magnetic Resonance brain images did not pass the quality control of the raw or the segmented images, participants with brain lesions, and those with marked alcohol consumption (alcohol use disorders identification test (AUDIT) total score >7 (23)). In this study, we present data from the remaining 101 adolescents (mean age =16.83 years, SD=0.61; 56 girls; Table 1). At the time of the study, none of the participants were followed in the psychiatric care system and all participants were attending school regularly.

Table 1. Clinical, behavioral characteristics, and brain volumes in community adolescents at age 16

Variable		Mean or %	SD
Demographic variables	Age (years)	16.83	0.61
	Gender	55 % (n=56) female	
Sleep variables (N=101)	Wake up time WD	7:03	0:43
	Wake up time WE	10:01	1:15
	Bedtime WD	22:42	0:43
	Bedtime WE	00:05	1:10
	Difference wake up time WE-WD	2:59	1:17
	Difference bedtime WE-WD	1:23	1:04
	Time in bed WD	8:18	0:59
	Time in bed WE	9:56	1:00
	Difference time in bed WE-WD	1:40	0:59
Performance scores (N=82)	Delay discounting large amounts	-2.25	0.68
	Delay discounting medium amounts	-2.03	0.73
	Delay discounting geomean	-2.06	0.66
	Spatial working memory	7.59	7.80
Behavioural problems (N=74)	SDQ internalising	4.34	2.40
	SDQ externalising	7.36	2.68
Global brain measures (N=101)	Total grey matter volume	742.02	68.99
	Total white matter volume	477.22	49.28
	Total CSF	395.55	40.89
	Volume scaling factor	1.32	148.21

WD = weekday; WE = weekend; CSF = cerebrospinal fluid.

Sleep assessments

Sleep habits were assessed by asking the adolescents their usual bedtimes and wake up times during weekdays (WD) and weekends (WE). Time in bed was approximated by calculating the number of hours between bedtime and wake up time, separately for WD and WE. WE delay in sleep timing (“social jet lag”) and weekly sleep debt was defined as the difference between weekday and weekend in sleep times and time in bed.

Psychological functioning

Symptom assessment was performed using the Strengths and Difficulties Questionnaire (SDQ) (24), a child and adolescent self-report questionnaire used to identify internalising and externalising problems. It consists of 25 items, five items for each subscale: conduct problems, hyperactivity, emotional problems, peer problems, and prosocial behaviour. The items are scored 0 to 2, reflecting the answers “not true”, “somewhat true” or “certainly true”. The scores are then summed, generating five scale scores ranging from 0 to 10 with higher scores reflecting more problems in the first four scales or more prosocial behaviour in the last scale. In low-risk samples, conduct problems and hyperactivity are best combined into an ‘externalising’ subscale, and emotional problems and peer problems into an ‘internalising’ subscale (25).

Other behavioural measures included the Kirby Delay-Discounting Questionnaire (26), a monetary choice questionnaire assessing cognitive impulsivity through delay discounting by having participants choose between smaller immediately available rewards and larger delayed rewards, and the spatial working memory task, a subtest of the computer-administrated

Cambridge Automated Neuropsychological Test Battery (CANTAB) measuring executive functioning (27). The spatial working memory test has been widely used in typically developing and clinical populations of children and adolescents (28). The error score was used as the outcome variable.

MRI data acquisition and processing

MRI was performed on a 3T scanner (Siemens Trio). High-resolution anatomical MR images were obtained using a standardised 3D T1-weighted magnetisation prepared rapid acquisition gradient echo (MPRAGE) sequence based on the ADNI protocol (<http://adni.loni.usc.edu/methods/mri-analysis/mri-acquisition/>). The parameters were as follows: repetition time=2,300 ms, echo time=2.8 ms, flip angle=8°, 256x256x170 matrix, 1.1x1.1x1.1 mm voxel size.

The images were processed using Statistical Parametric Mapping 8 (SPM 8) using Voxel-Based Morphometry (VBM) (29). The “unified segmentation” algorithm was used to normalise and segment the T1-weighted images into grey matter, white matter and cerebrospinal fluid. Home-made tissue probability maps were used instead of the standard template of SPM based on fully grown and developed brains of adults, who have larger brain volumes. The modulated images were smoothed with a 10-mm full-width at half-maximum isotropic Gaussian kernel. Head size was measured by the volume scaling factor, which is based on the affine transformation performed during spatial normalisation (<https://surfer.nmr.mgh.harvard.edu/fswiki/eTIV>).

Statistical analyses

The outcome measures for the Kirby Delay-Discounting Questionnaire were calculated using an automated calculator (30). A logistic regression that allowed for continuous estimates of k (31), and a logarithmic transformation to normalise the distribution of k -values were applied. The geometric mean of the k -values is bounded by the lowest implied indifference k -value at which subjects chose the larger delayed reward and the highest indifference k -value at which the subject chooses the immediate reward.

Correlations were performed with a Pearson correlation test, applying case-wise deletion for missing values. Group comparisons for socio-demographic and clinical data, and global brain volumes were performed within the framework of the general linear model (GLM) using R software (<http://cran.r-project.org>). Voxel-wise comparisons were carried out within the GLM framework of SPM8. Sleep variables were the main factors while age, gender and volume scaling factor were entered as confounding variables. Alpha was set for 0.05 in all tests.

A priori masks were used to examine the hippocampus, amygdala, and medial prefrontal region with a region of interest (ROI) approach. The hippocampus and amygdala masks were created with the Anatomical Automatic Labeling (AAL) atlas in SPM. The medial prefrontal region was marked with a 10 mm radius from coordinates [-2, 34, -14] taken from our previous results on an analogous sample at age 14, which indicated a strong relationship between sleep habits and GMV in this region (8). Subsequently, it was investigated whether psychometrics measures (spatial working memory, delay discounting, internalising and externalising problems) were associated with grey matter volumes in these regions of interest.

At the voxel-level, statistical significance was set to $p < 0.05$ FWE (Family Wise Error) corrected for multiple comparisons. Brain locations were reported as x, y, and z coordinates in Montreal Neurologic Institute (MNI) space.

In addition, it was examined whether the sleep times that were significant in the ROI VBM analyses were correlated with psychological functioning. Subsequently, causal mediation analyses were performed to determine whether the grey matter clusters could mediate the relation between sleep and psychological functioning variables. These analyses were performed if there was a significant relationship between sleep and psychological functioning variables and GMV of the same ROI.

The mediation analyses were performed with an algorithm using a set of general linear models to derive the mediation and direct effects from the total effect (32). The psychological functioning measures were entered as dependent factor, and sleep variables as independent factor within a linear regression model. For the mPFC ROI analysis, raw volume was extracted from the smoothed, normalized, and modulated images and entered as mediator variable. The hippocampus and amygdala volumes were extracted from the same images using the AAL masks instead of the significant clusters in order to better approximate the complete volumes. Volume scaling factor and age were entered as confounding variables. Gender was added for analyses without a gender interaction between sleep and psychological functioning scores. If there was a gender effect, we performed the mediation analysis separately for boys and girls. This mediation model was performed using 5000 Monte Carlo draws for nonparametric bootstrap. In causal

mediation analysis, a significant mediating effect is defined as a 95% confidence interval that does not include zero.

Results

Participant characteristics

On average, boys went to bed later during the week (boys = 22.53 ± 0.47 , girls = 22.33 ± 0.33 ; $F(99)=2.41$, $p=0.02$) and woke up later during the weekend (mean wake up time: boys = $10.23 \pm 1:17$, girls = $9:44 \pm 1:10$; $F(99)=2.63$, $p=0.01$) as compared to girls. There were no significant gender differences in any of the other sleep variables, nor in delay discounting, spatial working memory or internalising and externalising problems (see Table 1 for descriptives).

Clinically relevant externalising symptoms (SDQ score ≥ 10), were present in 15 out of 81 adolescents (18.3%) and 10 out of 81 (12.2%) showed clinically relevant internalising symptoms (score ≥ 8).

Sleep and regional grey matter volumes

WE wake up time correlated negatively with GMV in the bilateral amygdalae and the mPFC (Table 2, Fig 1). WE delay in wake up time ('social jet lag') correlated negatively with GMV in the right hippocampal region and the right amygdala (Table 2). No other statistically significant correlations between sleep variables and regional grey matter volumes were found.

Table 2. Grey Matter Volume correlations with sleep measures in community adolescents at age 16 using regions-of-interest

Brain region	Cluster		Peak				
	Size	p FWE	T-values	p FWE	coordinates		
					x	y	z
Wake up weekend							
Amygdala							
Amygdala R	237	0.018	3.51	0.011	21	-6	-12
Amygdala L	3	0.047	3.04	0.041	-12	-3	-15
mPFC							
	203	0.029	3.11	0.02	2	42	-11
Variability wake up time							
Amygdala							
Amygdala R	51	0.0350	3.23	0.025	20	-3	-12
Hippocampus							
Hippocampus R	37	0.028	4.23	0.008	20	-25	-23

Size = number of voxels in cluster; MNI = Montreal neurological Institute coordinates in millimeters; R = right; L = left. MNI coordinates are given for the voxel of maximal statistical significance. Statistics significant at the $p < 0.05$ FWE level; analyses are covaried for volume scale factor, age, and gender.

Figure 1. Later wake up time during the weekend was associated with reduced grey matter volumes in the A) left amygdala and B) mPFC

Sleep and psychological functioning

WE wake up time and WE delay in waking up correlated with measures of impulsivity as well as externalizing and internalizing problems (Table 3).

Table 3. Correlations between sleep variables and psychological functioning in community adolescents at age 16

	Wake up time WE	Variability wake up time
Delay discounting large	$r=0.319^{**}$	$r=0.292^{**}$
Delay discounting medium	$r=0.217$	$r=0.194$
Delay discounting geomean	$r=0.265^{*}$	$r=0.247^{*}$
SDQ externalising	$r=0.511^{**}$	$r=0.643^{**}$
SDQ internalising	$r=-0.229$	$r=-0.327^{**}$
Spatial working memory errors	$r=0.093$	$r=0.116$

$^{*}=p<0.05$, $^{**}=p<0.01$, $^{***}=p<0.001$

Grey matter volumes and psychological functioning

Smaller grey matter volumes in the mPFC region were associated with increased delay discounting ($T=3.28$, $p<0.01$) as well as internalising ($T=4.99$, $p<0.001$) and externalising ($T=6.03$, $p<0.001$) problems (Table 4). Additionally, smaller grey matter volumes in the amygdala (most significant cluster $T=10.57$, $p<0.001$), and hippocampal regions ($T=11.01$, $p<0.001$) were associated with internalising problems. There were no other significant relations between grey matter volumes and psychological functioning measures (Table 4).

Table 4. Grey matter volumes correlations with psychological functioning in 138 community adolescents at age 16

Psychological functioning	Amygdala		Hippocampus		mPFC	
	T-value	x,y,z	T-value	x,y,z	T-value	x,y,z
Delay discounting large	-	-	-	-	2.97*	-6,42,-15
Delay discounting medium	-	-	-	-	3.29*	-8,40,-17
Delay discounting geomean	-	-	-	-	3.28**	-8,40,-17
Spatial Working Memory	-	-	-	-	-	-
SDQ externalising	-	-	-	-	6.03***	-2,44,-14
SDQ internalising	10.57***	-30,-3,-29	11.01***	-28,-7,-32	4.99***	4,30,-8
	4.40***	36,0,-24	7.34***	21,-45,-3		
			6.13***	40,-15,-23		

- =no significant results, MNI coordinates and T-values are given for the voxel of maximal statistical significance, *= $p<0.05$, **= $p<0.01$, ***= $p<0.001$

Mediation analyses

Causal mediation analyses showed that variability in mPFC volumes accounted for 30.5% of the total effect between wake up time during the WE and externalising problems, and for 65.2% of the total effect between wake up time during the WE and internalising problems (Table 5).

Table 5. Causal mediation analysis on the relationship between sleep variables and behavioural problems with medial prefrontal cluster volumes as mediator

		Wake up time WE – mPFC - SDQ externalising	Wake up time WE – mPFC -SDQ internalising
Mediation effect	Point estimate	0.393**	-0.354***
	95% CI	0.138-0.720	-0.55
Direct effect	Point estimate	0.9**	-0.176
	95% CI	0.396-1.393	-0.998
Total effect	Point estimate	1.293**	-0.53*
	95% CI	0.776-1.817	-1.004
Proportion total effect via mediation	Point estimate	0.305***	0.652*
	95% CI	0.113-0.579	0.149-3.419

Point estimate, estimate of the size of the effect; 95% CI, 95% confidence interval of the point estimate

= $p < 0.05$, **= $p < 0.01$, *= $p < 0.001$*

Discussion

The present findings confirm and extend our previous findings that sleep patterns in adolescents from the general population correlate with regional brain grey matter volumes and psychological functioning. In particular, later WE wake up times were associated with smaller GMV in the mPFC and the amygdalae. Furthermore, a greater WE delay ('social jetlag') was associated with lower grey matter volume in hippocampus and amygdala regions. Regional grey matter volumes of the mPFC, amygdala, and hippocampus were associated with cognitive impulsivity and

behavioural problems. Causal mediation analyses showed that GMV of the mPFC mediated the relationship between weekend wake up time and externalising as well as internalising problems.

The results corroborate our previous findings in the sample at age 14, which showed an association between grey matter volumes in the mPFC region and weekend wake-up time (8). This could indicate that weekend wake up times affect mPFC grey matter maturation throughout adolescence. The impact of sleep on the brain might be particularly significant and long-lasting during adolescence because the brain is still in development during this phase (33).

The relationship between WE delay ('social jetlag') and grey matter volumes could result from repeated challenges to the circadian regulatory mechanisms (34). This can be compared with having regular *mini-jetlags*. A study on the short term effects of jetlag found reduced resting state activity in the mPFC and the left parahippocampal gyrus as well as other default mode network regions (35). In accordance with our finding of the association between GMV in the right hippocampal region and a difference in wake-up time, a study on chronic functional jetlag reported a relationship between sleep patterns and right temporal lobe atrophy, which included the parahippocampal gyrus (36). An alternative explanation is that later WE wake up times might reflect a late-prone biological rhythm rather than a social jetlag. Chronotype has recently been found to correlate with local GMVs and cortical thickness in a small sample of adult men (37). However, late bedtimes, another characteristic of a late chronotype, were not related to the volumes of the regions of interest in our study.

The results of the current study further show that WE delay and mPFC grey matter volumes are both significantly associated with cognitive impulsivity as well as internalising and externalising problems. Since the mPFC is an important brain area for regulation of the limbic structures, it is plausible that impaired development of this structure due to unhealthy sleep habits can lead to cognitive and emotional control deficits (16,18,38). This is the first study to show a mediating role of grey matter volumes in the mPFC in the relationship between sleep and internalising and externalising symptoms in healthy adolescents.

Internalising symptoms were also associated with smaller GMV in bilateral amygdala and parahippocampal regions. This finding is in accordance with previous research associating abnormalities in the amygdala, the parahippocampus, and the mPFC with emotional difficulties (39). Internalising symptoms are predictors of later educational underachievement, mental disorders, and impaired personal relationships (40). Externalising symptoms and impulsivity are both related to a host of maladaptive behaviours, including drug abuse, gambling, and poor academic and work success (41,42).

Bedtimes and time in bed were not significantly correlated with grey matter volumes at age 16. This is in contrast with our previous study in 14-year-old adolescents and a study by Taki and collaborators (7,8), where weekend bedtimes were associated with smaller mPFC grey matter volumes, but the sample of these studies consisted of younger children, who have not yet developed a delay in circadian rhythms and whose time in bed is still much more regulated by the parents. Therefore, time in bed during the week might be a more important factor in earlier brain developmental phases, while sleep timing and regularity in sleep timing might be more

important later during adolescence. The current results also do not provide evidence of negative effects of sleep debt, as measured by time in bed difference between WD and WE, on the adolescent brain. It must be noted, however, that sleep debt could affect brain regions that were not examined in the current study.

The findings could be confounded by the developmental stage of the participants. However, we used a relatively large sample, drawn from the general population, with a very narrow age range, making the results less likely to be influenced by confounding factors associated with age and sampling bias. A limitation of this study is the use of self-report questionnaires to measure sleep and behavioural problems. Sleep diaries and polysomnographic or actigraphic recordings would have provided a more objective measure of sleep, and including parental reports could have completed the picture of behavioural problems in adolescents. As this is a cross-sectional study, it is difficult to see to which degree sleep habits are a cause or a consequence of reduced grey matter volumes. We theorise that sleep rhythms influence brain development, which in turn causes lower functioning. The results of the causal mediation analyses would also favour this interpretation. However, reduced grey matter volumes might be a pre-existing condition that contributes to cognitive impulsivity and behavioural problems as well as the development of specific sleep patterns. Lastly, the ROI approach does not allow exploration of effects of sleep brain on regions other than the chosen ROIs.

Conclusion

Overall, the present findings are consistent with and extend our previous report in 14 year-old adolescents, suggesting that the negative impact of irregularity of sleep schedules on the

adolescent brain can result in reduced ability to regulate emotions and impulses. This highlights the importance of sleep habits in adolescents and supports the recommendation to keep variability in sleep times to a minimum in order to reduce the risk of psychiatric morbidity.

References

1. Suleiman AB, Dahl RE. Leveraging Neuroscience to Inform Adolescent Health: The Need for an Innovative Transdisciplinary Developmental Science of Adolescence. *J Adolesc Heal* [Internet]. 2017 Mar 1 [cited 2019 Oct 28];60(3):240–8. Available from: <https://www.sciencedirect.com/science/article/pii/S1054139X16309569>
2. Merikanto I, Pesonen A-K, Kuula L, Lahti J, Heinonen K, Kajantie E, et al. Eveningness as a risk for behavioral problems in late adolescence. *Chronobiol Int* [Internet]. 2017 Feb 7 [cited 2019 Oct 28];34(2):225–34. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28055288>
3. Owens J, Adolescent Sleep Working Group, Committee on Adolescence. Insufficient Sleep in Adolescents and Young Adults: An Update on Causes and Consequences. *Pediatrics* [Internet]. 2014 Sep 1 [cited 2019 Oct 28];134(3):e921–32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25157012>
4. Smaldone A, Honig JC, Byrne MW. Sleepless in America: inadequate sleep and relationships to health and well-being of our nation's children. *Pediatrics* [Internet]. 2007 Feb [cited 2019 Oct 28];119 Suppl 1(Supplement 1):S29-37. Available from: <http://pediatrics.aappublications.org/lookup/doi/10.1542/peds.2006-2089F>
5. Tesler N, Gerstenberg M, Huber R. Developmental changes in sleep and their relationships to psychiatric illnesses. *Curr Opin Psychiatry* [Internet]. 2013 Nov [cited 2019 Oct 28];26(6):572–9. Available from: <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00001504-201311000-00010>
6. Teicher MH, Ohashi K, Khan A, Garcia LCH, Klengel T, Anderson CM, et al. Does sleep disruption mediate the effects of childhood maltreatment on brain structure? *Eur J Psychotraumatol* [Internet]. 2017 [cited 2018 Nov 8];8. Available from: <https://doi.org/10.1080/20008198.2018.1450594>
7. Taki Y, Hashizume H, Thyreau B, Sassa Y, Takeuchi H, Wu K, et al. Sleep duration during weekdays affects hippocampal gray matter volume in healthy children. *Neuroimage* [Internet]. 2012 Mar [cited 2019 Oct 28];60(1):471–5. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1053811911013541>

8. Urrila AS, Artiges E, Massicotte J, Miranda R, Vulser H, Bézivin-Frere P, et al. Sleep habits, academic performance, and the adolescent brain structure. *Sci Rep* [Internet]. 2017 Feb 9 [cited 2017 Mar 10];7:41678. Available from: <http://www.nature.com/articles/srep41678>
9. Hagenauer MH, Perryman JI, Lee TM, Carskadon MA. Adolescent changes in the homeostatic and circadian regulation of sleep. *Dev Neurosci* [Internet]. 2009 [cited 2019 May 9];31(4):276–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19546564>
10. Giedd JN, Lalonde FM, Celano MJ, White SL, Wallace GL, Lee NR, et al. Anatomical brain magnetic resonance imaging of typically developing children and adolescents. *J Am Acad Child Adolesc Psychiatry* [Internet]. 2009 May 1 [cited 2019 May 9];48(5):465–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19395901>
11. Cho SS, Pellicchia G, Aminian K, Ray N, Segura B, Obeso I, et al. Morphometric Correlation of Impulsivity in Medial Prefrontal Cortex. *Brain Topogr* [Internet]. 2013 Jul 30 [cited 2019 Oct 28];26(3):479–87. Available from: <http://link.springer.com/10.1007/s10548-012-0270-x>
12. Forbes EE, Ryan ND, Phillips ML, Manuck SB, Worthman CM, Moyles DL, et al. Healthy adolescents' neural response to reward: associations with puberty, positive affect, and depressive symptoms. *J Am Acad Child Adolesc Psychiatry* [Internet]. 2010 Feb [cited 2019 Oct 28];49(2):162-72.e1-5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20215938>
13. Galvan A. Adolescent development of the reward system. *Front Hum Neurosci* [Internet]. 2010 [cited 2019 Oct 28];4:6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20179786>
14. Tashjian SM, Goldenberg D, Galván A. Neural connectivity moderates the association between sleep and impulsivity in adolescents. *Dev Cogn Neurosci* [Internet]. 2017 Oct [cited 2019 Oct 28];27:35–44. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1878929317301111>
15. Kreutzmann JC, Havekes R, Abel T, Meerlo P. Sleep deprivation and hippocampal vulnerability: changes in neuronal plasticity, neurogenesis and cognitive function. *Neuroscience* [Internet]. 2015 Nov 19 [cited 2019 Oct 28];309:173–90. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0306452215003942>
16. Motomura Y, Kitamura S, Nakazaki K, Oba K, Katsunuma R, Terasawa Y, et al. Recovery from Unrecognized Sleep Loss Accumulated in Daily Life Improved Mood Regulation via Prefrontal Suppression of Amygdala Activity. *Front Neurol* [Internet]. 2017 Jun 30 [cited 2019 Oct 28];8:306. Available from: <http://journal.frontiersin.org/article/10.3389/fneur.2017.00306/full>
17. O'Byrne JN, Berman Rosa M, Gouin J-P, Dang-Vu TT. Neuroimaging findings in primary insomnia. *Pathol Biol (Paris)* [Internet]. 2014 Oct [cited 2019 Oct 28];62(5):262–9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S036981141400087X>
18. Yoo S-S, Gujar N, Hu P, Jolesz FA, Walker MP. The human emotional brain without sleep--a prefrontal amygdala disconnect. *Curr Biol* [Internet]. 2007 Oct 23 [cited 2019 Oct 28];17(20):R877-8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960982207017836>
19. Robinson JL, Erath SA, Kana RK, El-Sheikh M. Neurophysiological differences in the adolescent brain following a single night of restricted sleep - A 7T fMRI study. *Dev Cogn Neurosci* [Internet]. 2018 Jun [cited 2019 Oct 28];31:1–10. Available from:

- 481 <https://linkinghub.elsevier.com/retrieve/pii/S1878929317302530>
- 482 20. Hoogman M, Bralten J, Hibar DP, Mennes M, Zwiers MP, Schweren LSJ, et al.
- 483 Subcortical brain volume differences in participants with attention deficit hyperactivity
- 484 disorder in children and adults: a cross-sectional mega-analysis. *The Lancet Psychiatry*
- 485 [Internet]. 2017 Apr [cited 2019 Oct 28];4(4):310–9. Available from:
- 486 <http://www.ncbi.nlm.nih.gov/pubmed/28219628>
- 487 21. Hulvershorn LA, Cullen K, Anand A. Toward dysfunctional connectivity: a review of
- 488 neuroimaging findings in pediatric major depressive disorder. *Brain Imaging Behav*
- 489 [Internet]. 2011 Dec 8 [cited 2019 Oct 28];5(4):307–28. Available from:
- 490 <http://link.springer.com/10.1007/s11682-011-9134-3>
- 491 22. Schumann G, Loth E, Banaschewski T, Barbot A, Barker G, Büchel C, et al. The
- 492 IMAGEN study: reinforcement-related behaviour in normal brain function and
- 493 psychopathology. *Mol Psychiatry* [Internet]. 2010 Dec 19 [cited 2019 Oct
- 494 28];15(12):1128–39. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21102431>
- 495 23. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the
- 496 Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early
- 497 Detection of Persons with Harmful Alcohol Consumption--II. *Addiction* [Internet]. 1993
- 498 Jun [cited 2019 Oct 28];88(6):791–804. Available from:
- 499 <http://doi.wiley.com/10.1111/j.1360-0443.1993.tb02093.x>
- 500 24. Goodman R. The Strengths and Difficulties Questionnaire: A Research Note. *J Child*
- 501 *Psychol Psychiatry* [Internet]. 1997 Jul 1 [cited 2019 Oct 28];38(5):581–6. Available
- 502 from: <http://doi.wiley.com/10.1111/j.1469-7610.1997.tb01545.x>
- 503 25. Goodman A, Lamping DL, Ploubidis GB. When to use broader internalising and
- 504 externalising subscales instead of the hypothesised five subscales on the Strengths and
- 505 Difficulties Questionnaire (SDQ): data from British parents, teachers and children. *J*
- 506 *Abnorm Child Psychol* [Internet]. 2010 Nov 10 [cited 2019 Oct 28];38(8):1179–91.
- 507 Available from: <http://link.springer.com/10.1007/s10802-010-9434-x>
- 508 26. Kirby KN, Maraković NN. Delay-discounting probabilistic rewards: Rates decrease as
- 509 amounts increase. *Psychon Bull Rev* [Internet]. 1996 Mar 1 [cited 2019 Oct 28];3(1):100–
- 510 4. Available from: <http://link.springer.com/10.3758/BF03210748>
- 511 27. Wild K V., Musser ED. The Cambridge Neuropsychological Test Automated Battery in
- 512 the Assessment of Executive Functioning. In: *Handbook of Executive Functioning*
- 513 [Internet]. New York, NY: Springer New York; 2014 [cited 2019 Oct 28]. p. 171–90.
- 514 Available from: http://link.springer.com/10.1007/978-1-4614-8106-5_11
- 515 28. Faridi N, Karama S, Burgaleta M, White MT, Evans AC, Fonov V, et al. Neuroanatomical
- 516 correlates of behavioral rating versus performance measures of working memory in
- 517 typically developing children and adolescents. *Neuropsychology* [Internet]. 2015 Jan
- 518 [cited 2019 Oct 28];29(1):82–91. Available from:
- 519 <http://www.ncbi.nlm.nih.gov/pubmed/25000324>
- 520 29. Ashburner J, Friston KJ. Voxel-Based Morphometry—The Methods. *Neuroimage*.
- 521 2000;821(11):805–21.
- 522 30. Kaplan BA, Lemley SM, Reed DD, Jarmolowicz DP. 21- and 27-Item Monetary Choice
- 523 Questionnaire Automated Scorers. 2014 [cited 2019 Oct 28]; Available from:
- 524 <https://kuscholarworks.ku.edu/handle/1808/15424>
- 525 31. Wileyto EP, Audrain-McGovern J, Epstein LH, Lerman C. Using logistic regression to
- 526 estimate delay-discounting functions. *Behav Res Methods Instrum Comput* [Internet].

- 2004 Feb [cited 2019 Oct 28];36(1):41–51. Available from:
<http://www.ncbi.nlm.nih.gov/pubmed/15190698>
32. Imai K, Keele L, Tingley D, Yamamoto T. Causal Mediation Analysis Using R. In Springer, New York, NY; 2010 [cited 2019 Oct 28]. p. 129–54. Available from:
http://link.springer.com/10.1007/978-1-4419-1764-5_8
33. Casey BJ, Jones RM, Hare TA. The adolescent brain. *Ann N Y Acad Sci* [Internet]. 2008 Mar [cited 2019 Oct 28];1124(1):111–26. Available from:
<http://doi.wiley.com/10.1196/annals.1440.010>
34. Taylor A, Wright HR, Lack LC. Sleeping-in on the weekend delays circadian phase and increases sleepiness the following week. *Sleep Biol Rhythms* [Internet]. 2008 Jul [cited 2019 Oct 28];6(3):172–9. Available from: <http://doi.wiley.com/10.1111/j.1479-8425.2008.00356.x>
35. Coutinho JF, Gonçalves OF, Maia L, Fernandes Vasconcelos C, Perrone-McGovern K, Simon-Dack S, et al. Differential activation of the default mode network in jet lagged individuals. *Chronobiol Int* [Internet]. 2015 Feb 2 [cited 2019 Oct 28];32(1):143–9. Available from: <http://www.tandfonline.com/doi/full/10.3109/07420528.2014.955187>
36. Cho K. Chronic “jet lag” produces temporal lobe atrophy and spatial cognitive deficits. *Nat Neurosci* [Internet]. 2001 Jun [cited 2019 Oct 28];4(6):567–8. Available from: http://www.nature.com/articles/nn0601_567
37. Rosenberg J, Jacobs HIL, Maximov II, Reske M, Shah NJ. Chronotype differences in cortical thickness: grey matter reflects when you go to bed. *Brain Struct Funct* [Internet]. 2018 Sep 15 [cited 2019 Oct 28];223(7):3411–21. Available from: <http://link.springer.com/10.1007/s00429-018-1697-y>
38. Alexander WH, Brown JW. Medial prefrontal cortex as an action–outcome predictor. *Nat Neurosci* [Internet]. 2011 Oct 18 [cited 2019 Oct 28];14(10):1338–44. Available from: <http://www.nature.com/articles/nn.2921>
39. Bootzin RR, Epstein DR. Understanding and Treating Insomnia. *Annu Rev Clin Psychol* [Internet]. 2011 Apr 27 [cited 2019 Oct 28];7(1):435–58. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17716026>
40. Colman I, Wadsworth MEJ, Croudace TJ, Jones PB. Forty-year psychiatric outcomes following assessment for internalizing disorder in adolescence. *Am J Psychiatry* [Internet]. 2007 Jan [cited 2019 Oct 28];164(1):126–33. Available from: <http://psychiatryonline.org/doi/abs/10.1176/ajp.2007.164.1.126>
41. Colman I, Murray J, Abbott RA, Maughan B, Kuh D, Croudace TJ, et al. Outcomes of conduct problems in adolescence: 40 year follow-up of national cohort. *BMJ* [Internet]. 2009 Jan 8 [cited 2019 Oct 28];338(jan08 2):a2981. Available from: <http://www.bmj.com/cgi/doi/10.1136/bmj.a2981>
42. Daugherty JR, Brase GL. Taking time to be healthy: Predicting health behaviors with delay discounting and time perspective. *Pers Individ Dif* [Internet]. 2010 Jan [cited 2019 Oct 28];48(2):202–7. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0191886909004188>

Figure 1. Later wake up time during the weekend was associated with reduced grey matter volumes in the A) left amygdala and B) mPFC