Spatial, climate, and ploidy factors drive genomic diversity and resilience in the widespread grass *Themeda triandra*

Ahrens CW^{1,2}, James EA², Miller AD³, Aitken NC⁴, Borevitz JO⁴, Cantrill DJ², Rymer PD¹

¹ Hawkesbury Institute for the Environment, Western Sydney University, Richmond NSW

² Royal Botanic Gardens Victoria, Melbourne, VIC, Australia

³ Deakin University, Warrnambool, VIC, Australia

⁴ Research School of Biology, Australian National University, Canberra, ACT, Australia

Author Correspondence:

Collin Ahrens

Email: c.ahrens@westernsydney.edu.au

Phone: +61 2 4570 1862

Word Count:

Summary – 195

Introduction – 1393

Methods – 2352

Results – 711

Conclusions – 1896

Total – 6352

Summary

•Fragmented grassland ecosystems, and the species that shape them, are under immense pressure. Restoration and management strategies should include genetic diversity and adaptive capacity to improve success but these data are generally unavailable. Therefore, we use the foundational grass, *Themeda triandra*, to test how spatial, environmental, and ploidy factors shape patterns of genetic variation.

•We used reduced-representation genome sequencing on 487 samples from 52 locations to answer fundamental questions about how the distribution of genomic diversity and ploidy polymorphism supports adaptation to harsher climates. We explicitly quantified isolation-by-distance (IBD), isolation-by-environment (IBE), and predicted population genomic vulnerability in 2070.

36 •We found that a majority (54%) of the genomic variation could be attributed to IBD, while 22% of
 37 the genomic variation could be explained by four climate variables showing IBE. Results indicate
 38 that heterogeneous patterns of vulnerability across populations are due to genetic variation, multiple
 39 climate factors, and ploidy polymorphism, which lessened genomic vulnerability in the most
 40 susceptible populations.

41

42 •These results indicate that restoration and management of *T. triandra* should incorporate knowledge
 43 of genomic diversity and ploidy polymorphisms to increase the likelihood of population persistence
 44 and restoration success in areas that will become hotter and more arid.

45

46 Key words

47 adaptation; genomic diversity; genomic vulnerability; landscape genomics; polyploidy; restoration;
 48 *Themeda triandra* (kangaroo grass)

49

50

51 Introduction

52 Grasses (Poaceae) are one of the most ecologically important vascular plant groups, making up 25%
 53 of the world's vegetation (Shantz, 1954). They provide key ecosystem services that underpin
 54 environmental health (i.e. habitat and food sources for native wildlife, nutrient cycling and carbon
 55 sequestration), and carry significant economic value as they include four of the five major crops in
 56 terms of global production (Raven & Thomas, 2010). Grasses are essential constituents of several
 57 vegetation communities including grasslands, grassy woodlands, and alpine regions. However,
 58 grasslands and grassy woodlands have historically been under immense pressure from rangeland and
 59 agricultural uses (Eldridge *et al.*, 2016; Hopkins & Holz 2006), leading to the fragmentation of natural
 60 populations and reductions in genetic diversity (Harrison *et al.*, 2015). Today, only about 4.6% of the
 61 billions of hectares of grassland ecosystems remain worldwide (IUCN 2016). In Australia, grassland
 62 systems are the most poorly conserved and degraded communities (Hobbs & Yates, 2000), and are
 63 likely to experience major negative long-term effects. Many regions of Australia that support
 64 grasslands are becoming warmer, drier and increasingly fire prone under climate change, highlighting
 65 the importance of preserving genetic diversity and evolutionary potential (Dunlop *et al.*, 2012).
 66 However, most research on genetic diversity in grass species has generally been undertaken on those
 67 of agricultural importance (Buckler *et al.*, 2001) such as wheat, corn, rice, and sorghum, or those that
 68 are being developed for biofuels such as switchgrass (*Panicum* – Casler *et al.*, 2007; Harrison *et al.*,
 69 2015) and sugarcane (*Miscanthus* – Vermerris, 2008). While research on species such as switchgrass
 70 have provided valuable insights into natural patterns of genetic diversity, adaptation across gradients,
 71 and the role ploidy plays between these lines of enquiry (Morris *et al.*, 2011; Lowry *et al.*, 2014,
 72 2019; Grabowski *et al.*, 2014), major gaps in knowledge for other ecologically important grasses
 73 persist and continue to inhibit effective conservation management.

74

75 Genetic diversity is maintained within a species by a combination of selective (such as range shifts
 76 and natural selection) and neutral processes (such as gene flow, mutation, and genetic drift) (Futuyma,
 77 2013). However, grasses often have complex evolutionary histories (Stebbins, 1956) influenced by
 78 factors such as clonality (Fischer & Van Kleunen, 2002), polyploidy (Keeler & Bradshaw, 1998),
 79 intrageneric hybridization, genome size, and different physiologies such as photosynthetic
 80 mechanisms (e.g. C3 versus C4) (Edwards *et al.*, 2010). These complex and often lineage-specific life
 81 histories can complicate our ability to project findings across species, meaning that the species-
 82 specific data needed for practitioners to make informed management decisions is often lacking.
 83 Perhaps the lack of research on ecologically important grass species and their complex life histories
 84 are not mutually exclusive. Regardless, information about how genetic diversity is distributed across
 85 habitats and environmental gradients, often reflecting selection and local adaptation, can help inform
 86 management and restoration strategies (Hoffmann *et al.*, 2015). This is particularly pertinent given
 87 grassland communities are already showing signs of climate stress, and empirical data is urgently

needed to support adaptive management strategies that prepare grasslands for new climate challenges by maximising evolutionary potential. In addition, research that focuses on genetic diversity across species ranges can help identify populations vulnerable to climate stress, allowing practitioners to prioritise management that safeguards populations at risk. For example, genomic signals of selection can be used to predict climate-driven population declines (Bay *et al.*, 2018). Specifically, ‘genomic vulnerability’ of individual populations, defined as the mismatch between current and predicted future genomic variation inferring population susceptibility to the loss of genetic diversity and/or maladaptation, can help identify populations most at risk. As our ability to integrate geospatial and genomic resources continues to grow, so will the ability of researchers to identify genomic vulnerability in ecologically important species, providing practitioners with improved management frameworks for mitigating climate change effects on ecosystems by preserving patterns of endemism and maximising adaptive potential.

Grasses often display ploidy differences among populations across their natural range. Indeed, polyploidy is common among vascular plants with c. 35% of species characterised as having a recent history of polyploidy (Wood *et al.*, 2009). For many species, associations between ploidy and local environmental conditions reflect adaptation, a pattern which has been studied extensively in crop plants (Alix *et al.*, 2017). Further, it has recently been shown that niche differentiation occurs faster in polyploids than diploid relatives (Baniaga *et al.*, 2019). While the causes of polyploidy are poorly understood (Soltis *et al.*, 2010), whole genome duplication events have been shown to coincide with historical climate change events (Cai *et al.*, 2019), and patterns of allopolyploidy have been linked to changes in environment (Wagner *et al.*, 2019). The effects of polyploidy are increasingly evident, with gene expression levels shown to vary from tissue to tissue in polyploids compared to their diploid counterparts (Adams *et al.*, 2003), and polyploid species often having significant fitness advantages (Petit & Thompson, 1997; Bretagnolle & Thompson, 2001; Ramsey, 2011; Hahn *et al.*, 2012; Hoffmann *et al.*, 2015; Wei *et al.*, 2019). Genome duplication may in itself be an advantage because it buffers the organism against deleterious alleles (Voigt-Zielinski *et al.*, 2012; Wagner *et al.*, 2019), and higher rates of heterozygosity reduce risks associated with inbreeding effects (Ronfort, 1999). Despite the potential benefits of polyploidy, there are known disadvantages, including the potential dilution of beneficial mutations (Stebbins, 1971) and disturbance of cellular functions such as epigenetic regulation, mitosis, and meiosis (Comai, 2005). However, ploidy polymorphism may provide an important evolutionary pathway for species to establish in previously unsuitable habitats or adapt *in situ* (Grabowski *et al.*, 2014).

Understanding patterns of genetic diversity and evolutionary mechanisms for adapting to new environments is key to improving the conservation of intact grasslands and the restoration of degraded grassland habitats. Globally, restoration practices largely advocate the use of seed sourced from local provenances, based on the assumption that local genotypes are best matched to stable local

environments and to avoid perceived risks associated with outbreeding (Thornhill, 1993; Edmands, 2006). Yet, in many cases local provenancing can lead to poor restoration outcomes (Broadhurst *et al.*, 2008; Prober *et al.*, 2015). In highly modified landscapes the genetic integrity of many species has been compromised, and local-provenancing can favour the selection of genetically depauperate and maladapted seed (Jones, 2013). Also, local-provenancing gives little consideration to the persistence of plantings under future climates, with growing evidence that genotypes from non-local sources may outperform those sourced locally (Hoffmann *et al.*, 2015; Prober *et al.*, 2015; Breed *et al.*, 2019). In addition, foundation species are especially important during the restoration process because their genetic variation can shape the networks of ecological interaction influencing community assembly, stability, and evolution (Gibson *et al.*, 2012; Lau *et al.*, 2016). Empirically derived restoration strategies are now being widely adopted around the world to support biodiversity, evolutionary potential, and restoration success, and similar approaches should also be employed for ploidy polymorphism.

In this study, we assess patterns of genetic structure, genotype-ploidy-environment associations, and genomic vulnerability in a foundational grassland species. *Themeda triandra*, commonly known as Kangaroo Grass, has a continent wide distribution, is characterised by ploidy polymorphisms (Hayman 1960) and has limited seed dispersal (Everson *et al.* 2009). The species provides critical ecosystem services supporting grassland habitats throughout Australia, and is widely used in grassland restorations, but is suffering major declines, shows signs of climate stress, and is in need of improved restoration guidelines. Notably, several studies suggest that re-establishment of *T. triandra* is an important first step for the restoration of Australia's grasslands (Adair & McDougall 1987; McDonald 2000; Cole & Lunt, 2005), highlighting the importance of research geared toward assessing the resilience of remnant populations, and management approaches that incorporate evolutionary potential. In this context, we assess the likely drivers of genetic structure across a portion of *T. triandra*'s range, predicting both isolation-by-distance (IBD) and isolation-by-environment (IBE) to be key drivers due to the species' limited seed dispersal and broad climatic niche. Based on estimates of gene flow and correlative measures of local adaptation, we test for genomic mismatches between local gene pools and future climates to help identify populations likely to be most vulnerable to new climatic challenges. Lastly, we test for associations between polyploidy and harsh climate zones, to gain insights into the role of polyploidy in historical and future adaptive processes. These results will provide clear pathways on how to incorporate genomic, environmental, and ploidy information into improved guidance for adaptive management plans that aim to protect these dwindling grassland ecosystems.

Materials and Methods

Species and sampling

Themeda triandra is a perennial C4 tussock grass, with ploidy variability, and occurs across three continents (Australia, Asia, and Africa) (Dell'Acqua *et al.*, 2013; Snyman *et al.*, 2013; Linder *et al.*, 2018). It is Australia's most widespread species, being adapted to habitats as diverse as the semi-arid interior and sub-alpine regions (Mitchell & Miller, 1990). In Australia, diploids and tetraploids are the most common ploidy variants, but triploid, pentaploid, hexaploid and aneuploid individuals have also been identified (Hayman, 1960). Past studies suggest that *T. triandra* originally evolved in tropical Asia and migrated through coastal corridors to Australia (Hayman, 1960), with Australian lineages diverging 1.37 mya (0.79 - 3.07 mya) (Dunning *et al.*, 2017). However, dating using secondary calibrations, as in (Dunning *et al.*, 2017) can lead to unreliable and overly young estimates of divergence (Schenk, 2016). *Themeda triandra* is widely considered a foundation species for three reasons: 1) it defines particular ecosystems (Snyman *et al.*, 2013), 2) it controls the distribution and abundance of associated flora and fauna (Morgan, 1998), and 3) it regulates the core ecosystem processes especially through fire (Morgan & Lunt, 1999). The species is also considered to be an indicator of (agro)ecosystem health (Novellie & Kraaij, 2010) and its long-term persistence provides ecosystem stability, ecosystem services, resistance to plant invasions, and facilitates rehabilitation of polluted and degraded habitat (Novellie & Kraaij, 2010; Dell'Acqua *et al.*, 2013). Furthermore, its persistence is critical for the restoration of grasslands in Australia and is reliant on recurring fire to remove old tillers and for seedling establishment (McDougall 1989). The distribution of *T. triandra* is suggestive of a complex evolutionary history with high levels of genetic structuring throughout Australia. Although *T. triandra* itself is not formally listed as an endangered species, it is an important constituent of temperate grassland communities, which have been declared as endangered in the Australian Capital Territory and New South Wales, and threatened in Victoria. The grasslands are under threat due to loss and fragmentation of habitats through inadequate land management practices.

Samples were collected between 2015 and 2017 from 52 populations spanning the heterogeneous climate from its eastern Australian distribution, which deliberately coincides with the densest portion of its distribution. Sampling was structured to ensure different environment combinations were sampled between coastal and inland (west of the Great Dividing Range, see Fig S1) sites. Sites were identified using records on the Atlas of Living Australia public database (ala.org.au) and chosen using the following criteria: herbaria collection or observation was after the year 2000, location data was within 50 m of accuracy, and occurred on land that was publicly accessible. Between 10 and 21 leaf samples were collected per location and placed directly into silica gel to rapidly dessicate leaf samples for DNA preservation. Sampled plants were at least 5 m apart to ensure independence of genotypes by minimising the chance of collecting clonal samples. Our collections comprised a total of 584 individual specimens, which were stored under laboratory conditions until required for genetic analysis.

Using the work of Hayman (1960), we created a predictive map of ploidy levels for populations distributed across our sampling distribution. Hayman measured ploidy levels across Australia, with most of his sites overlapping our sampling distribution. We interpolated his data using nearest neighbor analysis using QGIS v2.14 (Quantum GIS Development team), allowing us to extract predicted ploidy level for each population location to provide us with the number of predicted chromosomes (i.e. diploid = 20; tetraploid = 40; hexaploid = 60). A few individuals were equidistant between two predicted ploidy levels and were assigned ploidy level between 20 and 40. This was interpreted as indicating a mixed ploidy population. Ploidy predictions were verified with population-level heterozygosity, see below for details.

DNA extraction and library preparation

For reduced-representation library preparation and sequencing, genomic DNA from each individual was isolated from approximately 25 mg of silica-dried leaf tissue using the Stratec Invisorb DNA Plant HTS 96 kit (Invitex, Berlin, Germany). Libraries were created similarly to Ahrens et al. (2017). Briefly, extracted DNA was digested with PstI for genome complexity reduction, and ligated with a uniquely barcoded sequencing adapter pair. We then amplified each sample individually by PCR to avoid sample bias. We pooled samples in equimolar ratios and selected amplicons between 350 and 600 bp from an agarose gel. The library pool was sequenced on three Illumina NextSeq400 lanes using a 75bp paired-end protocol on a high output flowcell at the Biomolecular Resources Facility at the Australian National University, generating ~864 million read pairs.

For long-reads via the MinION sequencer (Oxford Nanopore Technologies, UK), we used the open access high molecular weight DNA extraction protocol developed by Jones & Borevitz (2019). Briefly, 30 g of fresh leaf material from a known diploid individual was processed with 150 mL nuclei isolation buffer using a high-powered blender. The homogenate was filtered repeatedly using a funnel, through sequentially 2, 4 and 8 layers of Miracloth. Next, 100% Triton X-100 was added for nuclei isolation and the mixture centrifuged to create a pellet of nuclei. The pellet was washed twice with a pre-chilled nuclei buffer. DNA extraction from the nuclei was initiated by adding fresh lysis buffer with 3% Sodium dodecyl sulfate (SDS) at 50°C. Binding buffer was added to use Sera-Mag beads to remove the lysis buffer from the DNA solution, washing with 70% ethanol 3 times until the beads were clean. The beads were removed by adding 220 uL of ultra-pure H₂O and resuspending the beads with attached DNA. The supernatant was removed and subsequently size selected for fragments longer than 30 kb using a PippinHT (Sage Science, Beverly MA). MinION library preparation and sequencing was performed as per the manufacturer's instructions and specifications, and resulted in 412,906 reads (Fig S2). Median read length was 27,156 bases, and the longest read length was 144,466 bases, with an overall average read-quality of 10 (Fig S2).

SNP calling

We checked the quality of the raw short-read sequencing reads with FastQC (v0.10.1, [Andrews, 2012]). Then, we demultiplexed the raw reads associated with each sample's unique combinatorial barcode using AXE v0.2.6 (Murray & Borevitz, 2018). During this step we were unable to assign 19% of the reads. We trimmed each sequence to 64 basepairs while removing the barcodes and ensured quality of the reads using trimmomatic v 0.38 (Bolger *et al.*, 2014). Quality was assessed using a sliding window of 4 basepairs (the number of bases used to average quality) and a quality score of 15 (the average quality required among the sliding window), and if the average quality dropped below 15, the sequences were cut. Then we indexed the long-reads (Fig S2 for distribution of length and number of reads sequenced) using the BWA software and the *index* argument. We aligned the short-reads to the long-reads for more accurate SNP calling compared to a *de novo* pipeline. Short-reads were aligned using BWA-mem (v0.7.17-r1198, [Li *et al.*, 2013]), as paired reads, with 82.5% of reads successfully mapped. Samtools v 1.9 (Li *et al.*, 2009) was used to transform the SAM files to BAM files for use within STACKS v 2.41 (Catchen *et al.*, 2013). The argument gstacks and populations were used in that order on the BAM files to create a VCF file, minimum thresholds (minor allele frequency = 0.01; one random SNP per read was retained) were set here for further cleaning in R (R core development team 2019). The mean coverage per sample was 15.8× with a standard deviation of 20×, this resulted in many samples being dropped (see below for details). Lastly, VCFtools v 0.1.16 (Danecek *et al.*, 2011) was used to create a 012 file for further cleaning of the snp matrix in R.

The missing data threshold was set to 50% per locus and individual which resulted in an average of 30% missing data from the whole SNP dataframe. Minor allele frequency was set to 0.05 to avoid identifying patterns of population structure that may be due to locally shared alleles (De la Cruz & Raska, 2014). Then we removed SNPs in high linkage disequilibrium (>50% similar). We also removed possible clones in Genodive v 2.0b27 (Meirmans & Van Tienderen, 2004) using the *assign clones* function, removing nine individuals. After conservative SNP filtering, we were left with 487 individuals from 52 populations.

Analysis

Genodive was used to estimate population summary statistics for the total number of alleles observed across loci, total heterozygosity, and the inbreeding coefficient (G_{IS} ; Nei, 1987). We expected that the degree of heterozygosity within populations would reflect ploidy status (i.e. higher heterozygosity would imply polyploids) as described by Soltis & Soltis (2000). Consequently, we validated predicted ploidy level among populations from Hayman's map (see above for details) by comparing those predictions to population-level heterozygosity. G_{IS} is the same as F_{IS} for a single locus with two alleles (Chakraborty & Leimar 1987), and is calculated by the ratio of observed heterozygosity within subpopulations to the expected heterozygosity and ranges from -1 (complete outbreeding) to 1 (complete inbreeding). Genodive was also used for an analysis of molecular variance (AMOVA)

using the Excoffier method (Excoffier et al. 1995). Global F_{ST} with 95% confidence intervals was calculated using the *fstat* argument and the population pairwise F_{ST} was calculated using the *pairwise.fst* argument in the *hierfstat* package in R (Goudet, 2005).

Themeda triandra has a broad geographic distribution spanning a variety of environmental gradients, therefore we wanted to estimate the amount of genetic variation that could be attributed to isolation-by-distance (IBD) and -environment (IBE). First, we downloaded the 19 bioclim variables from worldclim.org (Fick & Hijmans, 2017), and extracted all of the climate variables for each of the sample locations in R using the package *raster* (Hijmans & van Etten 2012). A Principle Components Analysis (PCA) was performed to determine potential correlations between the 19 climate variables and produce an environmental dataset consisting of least correlated variables (Fig S3). We chose to retain variables from six of the loose clusters (temperature mean diurnal range (T_{RANGE}), maximum temperature of the warmest month (T_{MAX}), precipitation seasonality (P_{SEAS}), mean annual temperature (T_{MA}), mean annual precipitation (P_{MA}), and precipitation of the driest month (P_{DM})).

We used sNMF (Frichot *et al.*, 2014) in the LEA package in R (Frichot & François, 2015) to investigate the observed patterns of population structure that include contributions from both geography (IBD) and environment (IBE). sNMF estimates ancestry coefficients based on sparse non-negative matrix factorisation and least-squares optimisation. The sparse non-negative matrix factorisation is robust to departures from traditional population genetic model assumptions, making this algorithm ideal to use with polyploid species such as *T. triandra*. We performed sNMF with the following attributes: $k = 1-10$, 10 replications per k -value (number of ancestral clusters), and 1,000 iterations. Entropy scores for each k -value were compared to choose the optimal number of clusters using the recommendations in the sNMF instruction manual. A consensus for the optimal k -value was created by averaging the results over the 10 replicate runs using CLUMPP v1.1.2 (Jakobsson & Rosenberg, 2007) and drawn using DISTRUCT v1.1 (Rosenberg, 2003).

We used Moran's Eigenvector Maps (MEM) to test if IBD was a major determinant of the species' genetic diversity, as described in previous work (Dray *et al.*, 2006; Legendre & Legendre, 2012) but called PCNM in the first papers. Briefly, MEM calculates a matrix of pairwise Euclidean distances **D** among the sampling sites, then transform the **D** matrix into a similarity matrix to produce the MEM. Eigenvalues are produced corresponding to orthogonal vectors of similarity. To ascertain spatial patterns of genetic diversity we used the R package memgene (Galpern *et al.*, 2014). Memgene identifies spatial neighbourhoods in genetic distance data that adopts a regression framework where the predictors are generated using MEMs, this multivariate technique was developed for spatial ecological analyses but is recommended for genetic applications. Memgene identifies variables (eigenvalues) that represent significant spatial genetic patterns at multiple spatial scales. Each variable

explains a proportion of the total variance explained by spatial patterns. For this study, we show two variables because it explains most of the variation described by IBD.

Using the environmental data layers we employ a generalized dissimilarity model (GDM) to identify the importance of specific climate variables responsible for shaping observed patterns of genetic structure within our dataset. Analyses were performed using the *gdm* package v 1.3.7 in R (Manion *et al.*, 2018) and a pairwise F_{ST} matrix (based on all SNP loci) to estimate allelic turnover through climatic space (deviations in allele frequency associated with environment type). Where GDM holds all variables in the model constant to identify the partial genomic distance associated with the climate factor (Ferrier *et al.*, 2007), whereby accounting for spatial patterns caused by demographic processes (Fitzpatrick & Keller, 2015). After running the GDM analysis, only four of the climates remained (T_{MAX} , P_{SEAS} , T_{MA} , and P_{MA}), as the other two climate factors were removed by a backward elimination procedure. The GDM output includes the deviance explained by the climate and spatial variables, and a spline plot for each climate and spatial variable. Spline plots were predicted across the study area and beyond for every 2.5km grid cell. These predicted grids were mapped using *ggplot* in R (Wickham, 2011) to describe the relative IBE.

We calculated ‘genomic vulnerability’ for the sampling area following Bay *et al.* (2018), which consists of three main components: exposure, sensitivity, and adaptive capacity (Dawson *et al.*, 2011). Genomic vulnerability is the amount of genomic change required to track environmental change over time and is interpreted as expected population decline. To do this, we substituted predictive maps in 2070 using the CCSM4 model with the representative concentration pathway 8.5 (worldclim.org), which is a prediction based on the anthropogenic carbon dioxide output not deviating from its current trajectory. These maps were also downloaded from worldclim and developed in the same way as described above. Lastly, we subtracted the projected genomic differentiation from the current genomic differentiation to get a difference between the two. We estimate genomic vulnerability twice, with and without predicted ploidy levels to understand how ploidy may affect population decline, particularly in the most vulnerable areas.

Results

We estimated patterns of population structure among 487 samples from 52 sample locations for *T. triandra* using a dataset consisting of 3,443 polymorphic SNPs with a minor allele frequency (MAF) of 0.05 and an average of 30% missing data. AMOVA indicated that a significant proportion of the genetic variance (10%) could be attributed to difference among sample sites ($P = 0.001$; $F_{ST} = 0.22$), while the majority of the variance (79.3%) was attributed to differences between individuals ($P < 0.01$; $F_{IT} = 0.31$). Large and significant positive inbreeding coefficients (G_{IS}) were observed for many sites, indicating an excess of homozygotes, while three populations had negative inbreeding coefficients indicating homozygote deficits (Table 1). Levels of genetic diversity (number of alleles

and heterozygosity) was variable among populations, with a mean number of alleles of 1.082 (95% CI 1.078-1.086; range 1.109 - 1.366) and a mean heterozygosity within populations (H_S) of 0.074 (range 0.06 - 0.12; Table 1). Heterozygosity estimates reflect patterns that are consistent with the hypothesis that greater ploidy levels are present in the hotter regions of our sampling distribution (Fig 1). However, this linear model, although significant ($r^2 = 0.086$; $P = 0.035$), explains only a small proportion of the variation. This pattern is likely driven by the three populations in the hottest region. Heterozygosity and predicted chromosome number were in agreement for these three populations, the populations with the highest T_{MAX} (QLD, PR, SWC). Some populations with high heterozygosity were predicted to be diploids (UL, GOR, NAM), but these populations were nearly equidistant to tetraploid and diploid populations and are likely tetraploid populations (Fig 1).

General patterns of population structure show a clear delineation between southern and northern populations (Fig 2) with an optimal k -value of 3 (Fig S4). The third k -value is found in two populations, and partially assigned in two other populations. These populations containing the third ancestral cluster were generally found in the central area of the sampling region. Notably, there are portions of populations, particularly in the south central portion of the sampling region, that have been assigned to the northern ancestral cluster. While there are a few individuals in the north assigned to the southern ancestral cluster.

Isolation-by-distance (IBD) was found to be significant in *T. triandra*. In fact, IBD accounts for 54% of the total genomic variation (Fig 3). Two axes are shown in separate figures, and together they explained 95% of the variation explained by IBD alone. The first axis shows a strong split between the northern and southern sections of the sampling area (Fig 3a), similar to the population structure identified in the sNMF results. A second pattern of IBD occurs in the northern part of the sampling region and is between the inland and coastal populations, while the most westerly population is slightly more similar to the northern sampling region (Fig 3b).

In addition to spatially driven genomic variation, isolation-by-environment (IBE) explains a significant amount of variation. While we chose six independent climate variables to explore IBE, only four were found to be significant (T_{MA} , T_{MAX} , P_{MA} , P_{SEAS} ; maps for climate variables in Fig S5). The GDM analysis was able to identify that 31.3% of the variation was attributable to these climate and spatial variables (Fig 4), and 22.0% of the variation was attributable directly to climate. When performing the same analysis with the inclusion of ploidy level, the variation explained rose by only 0.4%, but under this model, the T_{MAX} variable explained less variation (red lines in Fig 4) while all other variables remained similar. In the current climate, the differences between the two models were negligible (Fig 5a & c). However, when forecasting the differences in 2070, the outputs suggest a heterogeneous population decline by 0 and 25% (Fig 5b) with the highest proportion of change occurring inland of the eastern coast. Critically, the inclusion of ploidy polymorphism showed

genomic vulnerability dropping by 5% in the most vulnerable areas (Fig 5b & d; ploidy map provided in Fig S5), in this output, we find that genomic vulnerability occurs where the land transitions from the alpine region to the inland region. The lowest probability of change (population decline or gene pool turnover) is in the mountainous ecosystems in the southeastern portion of the sampling region.

Discussion

Our study indicates contemporary structuring of genomic diversity in *Themeda triandra* is being driven largely by a combination of spatial and climate factors. These patterns are indicative of a species with limited propagule dispersal and restricted gene flow. The apparent lack of connectivity among remnant populations suggests gene flow is unlikely to help local populations adapt to future climate challenges. Instead, their adaptive potential will rely on trait plasticity and standing genetic variation that allows for adaptation *in situ*. Strong associations between gene pools and climate may reflect patterns of local adaptation, and heterogeneity in climatic conditions at both local and regional scales, suggests that the impacts of climate change on remnant populations are likely to be uneven. This is supported by assessments of mismatches between current and predicted future genomic variation, creating heterogeneous patterns of ‘genomic vulnerability’ across populations. We also demonstrate polyploidy associations with harsh climate zones, suggesting polyploidy is potentially linked to historical adaptation processes and may assist populations in overcoming future climate challenges. This study highlights the need for adaptive management strategies that incorporate evolutionary potential, including seed sourcing and population mixing strategies that can help overcome genomic vulnerability and maladaptation under future climates.

Isolation-by-distance

The majority of genomic variation found in *T. triandra* could be explained by geographic isolation. This is likely to be due to low levels of gene flow and seed dispersal between populations contributing to strong genetic structuring, as found in South African populations (Everson *et al.*, 2009). However, this structure could also be driven by a partially apomictic reproductive system in *T. triandra* (Brown & Emery, 1957; Birari, 1980), with clonal reproduction inflating signals of population-level genetic uniqueness. We found some evidence of clonal *T. triandra* genotypes, but these individuals were removed during the data filtering phase prior to analyses. While our data are unable to confirm the relationship between clonality and polyploidy due to low replication, our data suggests that polyploidy occurs infrequently at milder temperatures, while being dominant among populations occurring in the highest temperature environments. These findings are consistent with Hayman (1960) who argues that the diploid landrace is likely absent in the harsher climates, suggesting the presence of positive selection for polyploid landraces in the hot and dry inland environments.

Perhaps the most germane work of this nature is that of the grass species *Panicum virgatum*. Similar to *T. triandra*, *P. virgatum*’s ploidy level increases with distance from the coast, with higher ploidy

levels found in more arid inland environments (Zhang *et al.*, 2011; Lowry *et al.*, 2014; Grabowski *et al.*, 2014). As demonstrated in *P. virgatum*, we provide evidence for polyploidy evolution through multiple, isolated events rather than the establishment and expansion of polyploids from one duplication event. For example, some populations of predicted polyploids are more closely related to diploid populations rather than other tetraploid populations. This suggests genome doubling can occur spontaneously within populations and is both induced and maintained by selection under certain environmental scenarios. Indeed, it has been shown that polyploids can have an increased fitness advantage under heat- and water- stressed conditions (Rey *et al.*, 2017).

Isolation-by-environment and genomic vulnerability

Along with geography, climate factors describe a large percentage of genomic variation found in *T. triandra*. We found strong associations between gene pools and environments (particularly with T_{MAX} and P_{SEAS}), possibly reflecting adaptation to climate. While quantitative tests are needed to validate these findings (e.g. common garden experiments – Sork, 2017), our results are consistent with the idea that signals of adaptation are ubiquitous throughout genomes (Kern & Hahn, 2018). Maximum temperature of the warmest month or week (T_{MAX}) has been found to be an important driving force of selection in other Australian plants (Steane *et al.*, 2017a,b; Jordan *et al.*, 2017; Ahrens *et al.*, 2019). Interestingly, evidence suggests that climatic factors can have different impacts on patterns of genetic diversity and adaptation in different grass species. For example, *T. triandra* and *Andropogon gerardii* are both dominant C4 grass species, with temperature and precipitation factors being key selective forces driving diversity in *T. triandra*, while lower precipitation suppresses genetic diversity in *A. gerardii* (Avolio *et al.*, 2013). Despite these differences, polyploidy appears dominant in harsher regions in both species indicating there are ploidy based adaptive responses to climate, enabling the expansion of species into habitats unsuitable or less suitable for diploids. The line of adaptation demarcation is stronger for *T. triandra*, where persistence in the semi-arid landscape appears entirely dependent on polyploids, compared to *A. gerardii*, where ploidy mixing occurs in harsher parts of its climate range (Keeler, 1990).

Our analyses of genomic vulnerability across the study area suggest that some populations of *T. triandra* will be more adversely impacted by climate change than others. For example, the most inland populations of our sampling are most vulnerable where we estimate that populations will need to change by over 20%, this region includes both diploid and polyploid populations. The least vulnerable populations are located in the southern and mountainous regions where we would expect populations to change by 0 to 5%. The future mismatch of predicted gene pools in some regions suggests that a change of as much as 25% will be necessary for adaption to the new challenges. Our predictions are based only on correlative analyses, and caution should be taken when interpreting these findings given the uncertainty associated with the genetic mechanisms (i.e. epistatic interactions (Juenger *et al.*, 2005), pleiotropy (Solovieff *et al.*, 2013), chromosomal rearrangements (Juenger *et al.*, 2005;

Yeaman, 2013), and polyploidy (Van de Peer *et al.*, 2017)) and ecological interactions likely to dictate future adaptive responses (Fordyce, 2006). Indeed, our findings further highlight the need for quantitative experiments (i.e. common garden) to validate these findings by testing the physiological limits and safety margins of individual populations.

Not surprisingly, the genomic vulnerability of several populations was buffered by as much as 5% by the presence of polyploids, and this is likely to be an underestimation due to under-predicting which populations are polyploids. Polyploidy is known to provide fitness advantages in many plant species persisting in hot and arid environments, including *T. triandra* populations (Godfree *et al.* 2017). The increased heterozygosity associated with polyploidy may have the effect of slowing the loss of genetic variation and providing more variants for selection to act upon (Comai, 2005). Elevated fitness may also be influenced by duplicated genes and genomes, each set capable of independent selection and evolving new functions (Soltis & Soltis, 2000) by retaining multiple gene copies and acquiring a new function in one copy (Wendel, 2000). Further, increased performance could be due to differential levels of expression between ploidy landraces (e.g. Cromie *et al.*, 2017; Wang *et al.*, 2018; Liqin *et al.*, 2019), and be partially dependent on different epigenetic patterns (Nagymihály *et al.*, 2017). However, quantitative measures are needed to determine how differential expression between diploid and tetraploid landraces may affect their ability to persist in their optimal climates. We argue that these types of processes are likely occurring in *T. triandra* landraces, allowing polyploids to persist and outperform their diploid counterparts in hotter and drier climates.

Management and restoration implications

We are at a critical juncture in history where management and restoration of grassland ecosystems is necessary to preserve these ecosystems and their services. However, the interplay of habitat fragmentation and rapid climate change poses a significant challenge for the conservation and restoration of functionally important plant species. Prioritising investments requires an understanding of species biology and ecology to apply frameworks for identifying the species and populations most at risk. *Themeda triandra*, the most widely distributed species in Australia, is at a critical inflection point due to its use as a food crop (Pascoe, 2018), for native pasture (Fourie *et al.*, 1985), as a foundational species (Snyman *et al.*, 2013), for selective breeding (e.g. *Lolium/Festuca* – Yamada *et al.*, 2005), and in the restoration of degraded lands (Cole & Lunt, 2005; Snyman *et al.*, 2013). Our results provide a critical first step and baseline information to support these new interests, future studies and the development of empirically based management strategies that target grassland and open woodland ecosystems. In Australia, research efforts have mostly focused on *Eucalyptus* species, finding that eucalypt populations are often connected by high levels of gene flow and adapted to local climates (e.g. Steane *et al.*, 2015; Jordan *et al.*, 2017; Supple *et al.*, 2018; Ahrens *et al.*, 2019). In one of the first landscape-scale genomic studies in Australia for an understory species, we show that the iconic grass *T. triandra* has very different patterns of connectivity and adaptation compared with its

Eucalyptus counterparts. Limited dispersal potential and high levels of genetic structuring among remnant populations of *T. triandra* suggests that their adaptability is likely to depend largely on trait plasticity and standing genetic variation that allows for adaptation *in situ*. We provide evidence of genetic and ploidy variation correlated with climate, suggesting that standing genetic variation may be retained within some *T. triandra* populations enabling adaptation to warmer and drier environments emerging under climate change. Indeed, our findings suggest the impacts of climate change may be heterogeneous across the distribution of *T. triandra*. This emphasises the importance of accounting for intraspecific variation, including ploidy, when predicting species responses to new climate challenges. Variability in physiological response to thermal stresses between populations has been established for many plant species (Moran *et al.*, 2016), which may contribute to uneven population responses to thermal stress (Miller *et al.*, 2019). These findings have implications for predicting population responses to climate change, and highlight the importance of interventions (assisted migrations of pre-adapted genotypes) to enhance the resilience of populations showing signs of climate stress given the existence of relatively tolerant populations across the species range.

Conclusion

Successful establishment of *T. triandra* on three continents from its Asian centre-of-origin is likely due to its ability to swiftly meet the challenges of new environmental conditions through mechanisms unique to the species. Genomic analysis of a species can elucidate broad patterns of structure and provide information about how those patterns are distributed across the landscape. While spatial structure was the major component of the species' standing genetic diversity, environmental heterogeneity was also a major component driving patterns of diversity, and patterns of neutral genetic diversity have been shown to be affected by natural selection (Phung *et al.*, 2016). Thus, these findings illustrate that the standing genetic variation can provide a basis for adaptation to changing climates and should be incorporated into restoration projects. We were also able to investigate long standing ploidy questions within a landscape genomics context. Notably, we were able to quantify how ploidy might buffer the species from the most severe climate effects in the future. We found that ploidy, along with standing genetic diversity, could be an important part of the puzzle that increases the probability of grassland ecosystem persistence during a period of dramatic change. Our data suggest that we risk underestimating the adaptive capacity of a species if we do not correct for ploidy polymorphisms and we propose that they should be an integral part of management strategies moving forward. Management of multi-ploidy foundational species should focus on a combination of attributes, including genetic variation, intraspecific ploidy polymorphisms, and trait characteristics to develop populations that are resilient to future climate scenarios ensuring ecosystem health, function, and long-term restoration success.

539

540 Data accessibility

541 Our data will be deposited on Dryad. Including the full SNP data set and population metadata.

542

543 Acknowledgements

544 We would like to thank Dr Ashley Jones for assistance in the lab for DNA prep for long-read
 545 sequences and Dr Robert Godfree for access to a known diploid individual for long-read sequencing.
 546 Royal Botanic Gardens Melbourne Friends supported this work through the Helen McLellan Research
 547 Grant. JOB and NCA were supported by ARC Centre of Excellence in Plant Energy Biology CE
 548 CE140100008.

549

550 Author Contribution

551 Design of the research was by CA and EJ; collection was performed by CA and EJ along with
 552 volunteers; lab work was performed by NA; data analysis was performed by CA; and writing the
 553 manuscript was performed by CA, EJ, and AM and all authors contributed to editing the manuscript.

554

References

- Adair R, McDougall K. 1987.** *Re-establishment of Native Grasses in Lowland Areas Progress Report*. Australian National Parks and Wildlife Service, Canberra.
- Adams KL, Cronn R, Percifield R, Wendel JF. 2003.** Genes duplicated by polyploidy show unequal contributions to the transcriptome and organ-specific reciprocal silencing. *Proceedings of the National Academy of Sciences of the United States of America* **100**: 4649–4654.
- Ahrens CW, Supple MA, Aitken NC, Cantrill DJ, Borevitz JO, James EA. 2017.** Genomic diversity guides conservation strategies among rare terrestrial orchid species when taxonomy remains uncertain. *Annals of botany* **119**, 1267–1277.
- Ahrens CW, Byrne M, Rymer PD. 2019.** Standing genomic variation within coding and regulatory regions contributes to the adaptive capacity to climate in a foundation tree species. *Molecular Ecology* **28**: 2502–2516.
- Alix K, Gérard PR, Schwarzacher T, Heslop-Harrison JSP. 2017.** Polyploidy and interspecific hybridization: partners for adaptation, speciation and evolution in plants. *Annals of Botany* **120**: 183–194.
- Andrews S. 2010.** FastQC: a quality control tool for high throughput sequence data.
- Avolio ML, Beaulieu JM, Smith MD. 2013.** Genetic diversity of a dominant C4 grass is altered with increased precipitation variability. *Oecologia* **171**: 571–581.
- Baniaga AE, Marx HE, Arrigo N, Barker MS. 2019.** Polyploid plants have faster rates of multivariate niche differentiation than their diploid relatives. *Ecology Letters*.
- Bay RA, Harrigan RJ, Le Underwood V, Lisle Gibbs H, Smith TB, Ruegg K. 2018.** Genomic signals of selection predict climate-driven population declines in a migratory bird. *Science* **359**: 83–86.
- Birari SP. 1980.** Apomixis and sexuality in *Themeda* forssk. at different ploidy levels (Gramineae). *Genetica* **54**: 133–139.
- Bolger AM, Lohse M, Usadel B. 2014.** Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.
- Breed MF, Harrison PA, Blyth C, Byrne M, Gaget V, Gellie NJC, Groom SVC, Hodgson R, Mills JG, Prowse TAA, et al. 2019.** The potential of genomics for restoring ecosystems and biodiversity. *Nature Reviews Genetics* **20**: 615–628.

585 **Bretagnolle F, Thompson JD. 2001.** Phenotypic plasticity in sympatric diploid and autotetraploid
586 *Dactylis glomerata*. *International Journal of Plant Sciences* **162**: 309–316.

587 **Broadhurst LM, Lowe A, Coates DJ, Cunningham SA, McDonald M, Vesk PA, Yates C. 2008.**
588 Seed supply for broadscale restoration: maximizing evolutionary potential. *Evolutionary Applications*
589 **1**: 587–597.

590 **Brown WV, Emery WHP. 1957.** Apomixis in the Gramineae, Tribe Andropogoneae: *Themeda*
591 *triandra* and *Bothriochloa ischaemum*. *Botanical Gazette* **118**: 246–253.

592 **Buckler ES 4th, Thornsberry JM, Kresovich S. 2001.** Molecular diversity, structure and
593 domestication of grasses. *Genetical Research* **77**: 213–218.

594 **Cai L, Xi Z, Amorim AM, Sugumaran M, Rest JS, Liu L, Davis CC. 2019.** Widespread ancient
595 whole-genome duplications in *Malpighiales* coincide with Eocene global climatic upheaval. *New*
596 *Phytologist* **221**: 565–576.

597 **Casler MD, Stendal CA, Kapich L, Vogel KP. 2007.** Genetic diversity, plant adaptation regions,
598 and gene pools for switchgrass. *Crop Science* **47**: 2261.

599 **Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. 2013.** Stacks: an analysis tool set
600 for population genomics. *Molecular Ecology* **22**: 3124–3140.

601 **Chakraborty R, Leimar O. 1987.** Genetic variation within a subdivided population. In *Population*
602 *Genetics and Fishery Management*. (Eds N. Ryman and F. Utter.) pp. 89–120.

603 **Cole BI, Lunt ID. 2005.** Restoring Kangaroo Grass (*Themeda triandra*) to grassland and woodland
604 understoreys: a review of establishment requirements and restoration exercises in south-east Australia.
605 *Ecological Management and Restoration* **6**: 28–33.

606 **Comai L. 2005.** The advantages and disadvantages of being polyploid. *Nature Reviews Genetics* **6**:
607 836–846.

608 **Cromie GA, Tan Z, Hays M, Jeffery EW, Dudley AM. 2017.** Dissecting gene expression changes
609 accompanying a ploidy-based phenotypic switch. *G3* **7**: 233–246.

610 **Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,**
611 **Marth GT, Sherry ST, et al. 2011.** The variant call format and VCFtools. *Bioinformatics* **27**: 2156–
612 2158.

613 **Dawson TP, Jackson ST, House JI, Prentice IC, Mace GM. 2011.** Beyond predictions: biodiversity
614 conservation in a changing climate. *Science* **332**: 53–58.

- 615 **De la Cruz O, Raska P. 2014.** Population structure at different minor allele frequency levels. *BMC*
616 *Proceedings* **8**: S55.
- 617 **Dell'Acqua M, Gomasca S, Porro A, Bocchi S. 2013.** A tropical grass resource for pasture
618 improvement and landscape management: *Themeda triandra* Forssk. *Grass and Forage Science* **68**:
619 205–215.
- 620 **Dray S, Legendre P, Peres-Neto PR. 2006.** Spatial modelling: a comprehensive framework for
621 principal coordinate analysis of neighbour matrices (PCNM). *Ecological Modelling* **196**: 483–493.
- 622 **Dunlop M, Hilbert D, Ferrier S, House A, Liedloff A, Prober SM, et al. 2012.** The implications of
623 climate change for biodiversity conservation and the National Reserve System: Final synthesis.: A
624 report prepared for the Department of Sustainability, Environment, Water, Population and
625 Communities, and the Department of Climate Change and Energy Efficiency. CSIRO Climate
626 Adaptation Flagship, Canberra, Australia.
- 627 **Dunning LT, Liabot A-L, Olofsson JK, Smith EK, Vorontsova MS, Besnard G, Simpson KJ,**
628 **Lundgren MR, Addicott E, Gallagher RV, et al. 2017.** The recent and rapid spread of *Themeda*
629 *triandra*. *Botany Letters* **164**: 327–337.
- 630 **Edmands S. 2006.** Between a rock and a hard place: evaluating the relative risks of inbreeding and
631 outbreeding for conservation and management. *Molecular Ecology* **16**: 463–475.
- 632 **Edwards EJ, Osborne CP, Strömberg CAE, Smith SA, C4 Grasses Consortium, Bond WJ,**
633 **Christin P-A, Cousins AB, Duvall MR, Fox DL, et al. 2010.** The origins of C4 grasslands:
634 integrating evolutionary and ecosystem science. *Science* **328**: 587–591.
- 635 **Eldridge DJ, Poore AGB, Ruiz-Colmenero M, Letnic M, Soliveres S. 2016.** Ecosystem structure,
636 function, and composition in rangelands are negatively affected by livestock grazing. *Ecological*
637 *Applications* **26**: 1273–1283.
- 638 **Everson TM, Yeaton RI, Everson CS. 2009.** Seed dynamics of *Themeda triandra* in the montane
639 grasslands of South Africa. *African Journal of Range & Forage Science* **26**: 19–26.
- 640 **Excoffier L. 1995.** AMOVA 1.55 (analysis of molecular variance). *University of Geneva, Geneva*.
- 641 **Ferrier S, Manion G, Elith J, Richardson K. 2007.** Using generalized dissimilarity modelling to
642 analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity and*
643 *Distributions* **13**: 252–264.
- 644 **Fick SE, Hijmans RJ. 2017.** WorldClim 2: new 1-km spatial resolution climate surfaces for global
645 land areas. *International Journal of Climatology* **37**: 4302–4315.

- 646 **Fischer M, Van Kleunen M. 2002.** On the evolution of clonal plant life histories. *Evolutionary*
647 *Ecology* **15**: 565–582.
- 648 **Fitzpatrick MC, Keller SR. 2015.** Ecological genomics meets community-level modelling of
649 biodiversity: mapping the genomic landscape of current and future environmental adaptation. *Ecology*
650 *Letters* **18**: 1–16.
- 651 **Fordyce JA. 2006.** The evolutionary consequences of ecological interactions mediated through
652 phenotypic plasticity. *Journal of Experimental Biology* **209**: 2377–2383.
- 653 **Fourie JH, Opperman DPJ, Roberts BR. 1985.** Evaluation of the grazing potential of grass species
654 in Tarchonanthus veld of the northern cape. *Journal of the Grassland Society of Southern Africa* **2**:
655 13–17.
- 656 **Frichot E, François O. 2015.** LEA: An R package for landscape and ecological association studies.
657 *Methods in Ecology and Evolution* **6**: 925–929.
- 658 **Frichot E, Mathieu F, Trouillon T, Bouchard G, François O. 2014.** Fast and efficient estimation of
659 individual ancestry coefficients. *Genetics* **196**: 973–983.
- 660 **Futuyma DJ. 2013.** *Evolution*. Sinauer Associates Incorporated. Sunderland, MA.
- 661 **Galpern P, Peres-Neto PR, Polfus J, Manseau M. 2014.** MEMGENE: Spatial pattern detection in
662 genetic distance data. *Methods in Ecology and Evolution* **5**: 1116–1120.
- 663 **Gibson DJ, Allstadt AJ, Baer SG, Geisler M. 2012.** Effects of foundation species genotypic
664 diversity on subordinate species richness in an assembling community. *Oikos* **121**: 496–507.
- 665 **Godfree RC, Marshall DJ, Young AG, Miller CH, Mathews S. 2017.** Empirical evidence of fixed
666 and homeostatic patterns of polyploid advantage in a keystone grass exposed to drought and heat
667 stress. *Royal Society Open Science* **4**: 170934.
- 668 **Goudet J. 2005.** hierfstat, a package for r to compute and test hierarchical *F*-statistics. *Molecular*
669 *Ecology Notes* **5**: 184–186.
- 670 **Grabowski PP, Morris GP, Casler MD, Borevitz JO. 2014.** Population genomic variation reveals
671 roles of history, adaptation and ploidy in switchgrass. *Molecular Ecology* **23**: 4059–4073.
- 672 **Hahn MA, van Kleunen M, Müller-Schärer H. 2012.** Increased phenotypic plasticity to climate
673 may have boosted the invasion success of polyploid *Centaurea stoebe*. *PLoS ONE* **7**: e50284.
- 674 **Harrison SP, Gornish ES, Copeland S. 2015.** Climate-driven diversity loss in a grassland
675 community. *Proceedings of the National Academy of Sciences of the United States of America* **112**:

676 8672–8677.

677 **Hayman DL. 1960.** The distribution and cytology of the chromosome races of *Themeda australis* in
678 southern Australia. *Australian Journal of Botany* **8**: 58.

679 **Hijmans RJ, van Etten J. 2012.** raster: Geographic analysis and modeling with raster data. R
680 package version 2.0-12. <http://CRAN.R-project.org/package=raster>

681 **Hobbs RJ, Yates CJ. 2000.** *Temperate Eucalypt Woodlands in Australia: Biology, Conservation,*
682 *Management and Restoration.* Surrey Beatty & Sons Pty. Ltd. Chipping Norton, Australia

683 **Hoffmann A, Griffin P, Dillon S, Catullo R, Rane R, Byrne M, Jordan R, Oakeshott J, Weeks A,**
684 **Joseph L, et al. 2015.** A framework for incorporating evolutionary genomics into biodiversity
685 conservation and management. *Climate Change Responses* **2**: 1.

686 **Hopkins A, Holz B. 2006.** Grassland for agriculture and nature conservation: production, quality and
687 multi-functionality. *Agronomy research* **4**, 3-20.

688 **IUCN. 2016.** <https://www.unep-wcmc.org/resources-and-data/protected-planet-report-2016> accessed:
689 22 October 2019

690 **Jakobsson M, Rosenberg NA. 2007.** CLUMPP: a cluster matching and permutation program for
691 dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**:
692 1801–1806.

693 **Jones TA. 2013.** When local isn't best. *Evolutionary Applications* **6**: 1109-1118.

694 **Jones A, Borevitz JO. 2019.** Nuclear DNA purification from recalcitrant plant species for long-read
695 sequencing <https://www.protocols.io/view/nuclear-dna-purification-from-recalcitrant-plant-s-vmee43e>
696 accessed: 18 July 2019

697 **Jordan R, Hoffmann AA, Dillon SK, Prober SM. 2017.** Evidence of genomic adaptation to climate
698 in *Eucalyptus microcarpa*: Implications for adaptive potential to projected climate change. *Molecular*
699 *Ecology* **26**: 6002–6020.

700 **Juenger TE, Sen S, Stowe KA, Simms EL. 2005.** Epistasis and genotype-environment interaction
701 for quantitative trait loci affecting flowering time in *Arabidopsis thaliana*. *Genetica* **123**: 87–105.

702 **Keeler KH. 1990.** Distribution of polyploid variation in big bluestem (*Andropogon gerardii*,
703 Poaceae) across the tallgrass prairie region. *Genome* **33**: 95–100.

704 **Keeler KH, Bradshaw AD. 1998.** Population biology of intraspecific polyploidy in grasses. In
705 *Population Biology of Grasses.* Cambridge University Press, Cambridge, UK. 183–206.

706 **Kern AD, Hahn MW. 2018.** The neutral theory in light of natural selection. *Molecular Biology and*
707 *Evolution* **35**: 1366–1371.

708 **Lau MK, Keith AR, Borrett SR, Shuster SM, Whitham TG. 2016.** Genotypic variation in
709 foundation species generates network structure that may drive community dynamics and evolution.
710 *Ecology* **97**: 733–742.

711 **Legendre P, Legendre L. 2012.** Numerical Ecology. (3rd. English ed.), Elsevier. Amsterdam,
712 Netherlands.

713 **Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,**
714 **1000 Genome Project Data Processing Subgroup. 2009.** The sequence alignment/map format and
715 SAMtools. *Bioinformatics* **25**: 2078–2079.

716 **Li H. 2013.** Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
717 arXiv:1303.3997

718 **Linder HP, Lehmann CER, Archibald S, Osborne CP, Richardson DM. 2018.** Global grass
719 (Poaceae) success underpinned by traits facilitating colonization, persistence and habitat
720 transformation. *Biological reviews of the Cambridge Philosophical Society* **93**: 1125–1144.

721 **Liqin G, Jianguo Z, Xiaoxia L, Guodong R. 2019.** Polyploidy-related differential gene expression
722 between diploid and synthesized allotriploid and allotetraploid hybrids of *Populus*. *Molecular*
723 *Breeding* **39**: 69.

724 **Lowry DB, Behrman KD, Grabowski P, Morris GP, Kiniry JR, Juenger TE. 2014.** Adaptations
725 between ecotypes and along environmental gradients in *Panicum virgatum*. *The American Naturalist*
726 **183**: 682–692.

727 **Lowry DB, Lovell JT, Zhang L, Bonnette J, Fay PA, Mitchell RB, Lloyd-Reilley J, Boe AR, Wu**
728 **Y, Rouquette FM Jr, et al. 2019.** QTL $\times$ environment interactions underlie adaptive divergence in
729 switchgrass across a large latitudinal gradient. *Proceedings of the National Academy of Sciences of*
730 *the United States of America* **116**: 12933–12941.

731 **Manion G, Lisk M, Ferrier S, Nieto-Lugilde KM, Fitzpatrick MC. 2018.** gdm: Functions for
732 generalized dissimilarity modeling. *R package*.

733 **McDonald T, 2000.** Strategies for the ecological restoration of woodland plant communities:
734 harnessing natural resilience. In: *Temperate Eucalypt Woodlands in Australia: Biological*
735 *Conservation, Management and Restoration* (eds Richard J. Hobbs and Colin J. Yates), pp. 286–297.
736 Surrey Beatty and Sons, Chipping Norton, NSW.

737 **McDougall KL. 1989.** The re-establishment of *Themeda triandra* (kangaroo grass): implications for
738 the restoration of grasslands [Victoria]. *Technical Report Series-Arthur Rylah Institute for*
739 *Environmental Research (Australia).*

740 **Meirmans PG, Van Tienderen PH. 2004.** genotype and genodive: two programs for the analysis of
741 genetic diversity of asexual organisms. *Molecular Ecology Notes* **4**: 792–794.

742 **Miller AD, Hoffmann AA, Tan MH, Young M, Ahrens C, Cocomazzo M, Rattray A,**
743 **Ierodiconou DA, Trembl E, Sherman CDH. 2019.** Local and regional scale habitat heterogeneity
744 contribute to genetic adaptation in a commercially important marine mollusc (*Haliotis rubra*) from
745 southeastern Australia. *Molecular Ecology* **28**: 3053–3072.

746 **Mitchell M, Miller M. 1990.** The identification of some common native grasses in Victoria.
747 *Rutherglen Research Institute, Victoria, Australia.*

748 **Moran EV, Hartig F, Bell DM. 2016.** Intraspecific trait variation across scales: implications for
749 understanding global change responses. *Global Change Biology* **22**: 137–150.

750 **Morgan JW. 1998.** Importance of canopy gaps for recruitment of some forbs in *Themeda triandra*-
751 dominated grasslands in south-eastern Australia. *Australian Journal of Botany* **46**: 609.

752 **Morgan JW, Lunt ID. 1999.** Effects of time-since-fire on the tussock dynamics of a dominant grass
753 (*Themeda triandra*) in a temperate Australian grassland. *Biological Conservation* **88**: 379–386.

754 **Morris GP, Grabowski PP, Borevitz JO. 2011.** Genomic diversity in switchgrass (*Panicum*
755 *virgatum*): from the continental scale to a dune landscape. *Molecular ecology* **20**: 4938–4952.

756 **Murray KD, Borevitz JO. 2018.** Axe: rapid, competitive sequence read demultiplexing using a trie.
757 *Bioinformatics* **34**: 3924–3925.

758 **Nagymihály M, Veluchamy A, Györgypál Z, Ariel F, Jégu T, Benhamed M, Szűcs A, Kereszt A,**
759 **Mergaert P, Kondorosi É. 2017.** Ploidy-dependent changes in the epigenome of symbiotic cells
760 correlate with specific patterns of gene expression. *Proceedings of the National Academy of Sciences*
761 *of the United States of America* **114**: 4543–4548.

762 **Nei M. 1987.** Molecular Evolutionary Genetics.

763 **Novellie P, Kraaij T. 2010.** Evaluation of *Themeda triandra* as an indicator for monitoring the
764 effects of grazing and fire in the Bontebok National Park. *Koedoe* **52**.

765 **Pascoe B. 2018.** Dark Emu: Aboriginal Australia and the Birth of Agriculture, *New Edition*.
766 Magabala Books.

767 **Petit C, Thompson JD. 1997.** Variation in phenotypic response to light availability between diploid
768 and tetraploid populations of the perennial grass *Arrhenatherum elatius* from open and woodland
769 sites. *Journal of Ecology* **85**: 657.

770 **Phung TN, Huber CD, Lohmueller KE. 2016.** Determining the Effect of Natural Selection on
771 Linked Neutral Divergence across Species. *PLoS genetics* **12**: e1006199.

772 **Prober SM, Byrne M, McLean EH, Steane DA, Potts BM, Vaillancourt RE, Stock WD. 2015.**
773 Climate-adjusted provenancing: a strategy for climate-resilient ecological restoration. *Frontiers in*
774 *Ecology and Evolution* **3**.

775 **Ramsey J. 2011.** Polyploidy and ecological adaptation in wild yarrow. *Proceedings of the National*
776 *Academy of Sciences of the United States of America* **108**: 7096–7101.

777 **Raven J, Thomas H. 2010.** Grasses. *Current Biology* **20**: R837–R839.

778 **Rey PJ, Manzaneda AJ, Alcántara JM. 2017.** The interplay between aridity and competition
779 determines colonization ability, exclusion and ecological segregation in the heteroploid
780 *Brachypodium distachyon* species complex. *New Phytologist* **215**: 85–96.

781 **Ronfort J. 1999.** The mutation load under tetrasomic inheritance and its consequences for the
782 evolution of the selfing rate in autotetraploid species. *Genetical Research* **74**: 31–42.

783 **Rosenberg NA. 2003.** distruct: a program for the graphical display of population structure. *Molecular*
784 *Ecology Notes* **4**: 137–138.

785 **Schenk JJ. 2016.** Consequences of Secondary Calibrations on Divergence Time Estimates. *PloS one*
786 **11**: e0148228.

787 **Shantz HL. 1954.** The Place of Grasslands in the Earth's Cover. *Ecology* **35**: 143–145.

788 **Snyman HA, Ingram LJ, Kirkman KP. 2013.** *Themeda triandra*: a keystone grass species. *African*
789 *Journal of Range & Forage Science* **30**: 99–125.

790 **Solovieff N, Cotsapas C, Lee PH, Purcell SM, Smoller JW. 2013.** Pleiotropy in complex traits:
791 challenges and strategies. *Nature Reviews Genetics* **14**: 483–495.

792 **Soltis DE, Buggs RJA, Doyle JJ, Soltis PS. 2010.** What we still don't know about polyploidy.
793 *TAXON* **59**: 1387–1403.

794 **Soltis PS, Soltis DE. 2000.** The role of genetic and genomic attributes in the success of polyploids.
795 *Proceedings of the National Academy of Sciences of the United States of America* **97**: 7051–7057.

- 796 **Sork VL. 2017.** Genomic Studies of Local Adaptation in Natural Plant Populations. *Journal of*
797 *Heredity* **109**: 3–15.
- 798 **Steane DA, Mclean EH, Potts BM, Prober SM, Stock WD, Stylianou VM, Vaillancourt RE,**
799 **Byrne M. 2017a.** Evidence for adaptation and acclimation in a widespread eucalypt of semi-arid
800 Australia. *Biological Journal of the Linnean Society* **121**: 484–500.
- 801 **Steane DA, Potts BM, McLean EH, Collins L, Holland BR, Prober SM, Stock WD, Vaillancourt**
802 **RE, Byrne M. 2017b.** Genomic Scans across Three Eucalypts Suggest that Adaptation to Aridity is a
803 Genome-Wide Phenomenon. *Genome Biology and Evolution* **9**: 253–265.
- 804 **Steane DA, Potts BM, McLean E, Collins L, Prober SM, Stock WD, Vaillancourt RE, Byrne M.**
805 **2015.** Genome-wide scans reveal cryptic population structure in a dry-adapted eucalypt. *Tree*
806 *Genetics & Genomes* **11**.
- 807 **Stebbins GL. 1971.** *Processes of organic evolution*. Prentice Hall.
- 808 **Stebbins GL. 1956.** Cytogenetics and evolution of the grass family. *American Journal of Botany* **43**:
809 890–905.
- 810 **Supple MA, Bragg JG, Broadhurst LM, Nicotra AB, Byrne M, Andrew RL, Widdup A, Aitken**
811 **NC, Borevitz JO. 2018.** Landscape genomic prediction for restoration of a *Eucalyptus* foundation
812 species under climate change. *eLife* **7**: e31835.
- 813 **Thornhill NW. 1993.** *The Natural History of Inbreeding and Outbreeding: Theoretical and*
814 *Empirical Perspectives*. University of Chicago Press. Chicago, IL.
- 815 **Van de Peer Y, Mizrahi E, Marchal K. 2017.** The evolutionary significance of polyploidy. *Nature*
816 *Reviews Genetics* **18**: 411–424.
- 817 **Vermerris W. 2008.** Miscanthus: Genetic Resources and Breeding Potential to Enhance Bioenergy
818 Production. *Genetic Improvement of Bioenergy Crops*: 295–308.
- 819 **Voigt-Zielinski M-L, Piwczyński M, Sharbel TF. 2012.** Differential effects of polyploidy and
820 diploidy on fitness of apomictic *Boechera*. *Sexual plant reproduction* **25**: 97–109.
- 821 **Wagner F, Ott T, Zimmer C, Reichhart V, Vogt R, Oberprieler C. 2019.** At the crossroads
822 towards polyploidy: genomic divergence and extent of homoploid hybridization are drivers for the
823 formation of the ox-eye daisy polyploid complex (*Leucanthemum*, Compositae-Anthemideae). *New*
824 *Phytologist* **223**: 2039–2053.
- 825 **Wang T, Huang D, Chen B, Mao N, Qiao Y, Ji M. 2018.** Differential expression of photosynthesis-

826 related genes in pentaploid interspecific hybrid and its decaploid of *Fragaria* spp. *Genes & genomics*
827 **40**: 321–331.

828 **Wei N, Cronn R, Liston A, Ashman T-L. 2019.** Functional trait divergence and trait plasticity
829 confer polyploid advantage in heterogeneous environments. *New Phytologist* **221**: 2286–2297.

830 **Wendel JF. 2000.** Genome evolution in polyploids. *Plant Molecular Evolution*: 225–249.

831 **Wickham H. 2011.** ggplot2. *Computational Statistics* **3**: 180–185.

832 **Wood TE, Takebayashi N, Barker MS, Mayrose I, Greenspoon PB, Rieseberg LH. 2009.** The
833 frequency of polyploid speciation in vascular plants. *Proceedings of the National Academy of*
834 *Sciences of the United States of America* **106**: 13875–13879.

835 **Yamada T, Forster JW, Humphreys MW, Takamizo T. 2005.** Genetics and molecular breeding in
836 *Lolium/Festuca* grass species complex. *Grassland Science* **51**: 89–106.

837 **Yeaman S. 2013.** Genomic rearrangements and the evolution of clusters of locally adaptive loci.
838 *Proceedings of the National Academy of Sciences of the United States of America* **110**: E1743–51.

839 **Zhang Y, Zalapa JE, Jakubowski AR, Price DL, Acharya A, Wei Y, Brummer EC, Kaeppler**
840 **SM, Casler MD. 2011.** Post-glacial evolution of *Panicum virgatum*: centers of diversity and gene
841 pools revealed by SSR markers and cpDNA sequences. *Genetica* **139**: 933–948.

842
843

Figure Legends

Figure 1. Within population heterozygosity (Hs) versus maximum temperature of the warmest month. Colors indicate diploid (blue), mixed populations (green; equidistant between tetraploid and diploid populations), and tetraploid (red) based on Hayman's (1960) work. Ellipsoid outlines populations that have high heterozygosity and may be tetraploids.

Figure 2. Sparse non-negative matrix factorization (sNMF) for all individuals, points on the map indicate population location, map colors represent T_{MAX} (maximum temperature of the warmest month). Barplot indicates identified genetic ancestral clusters for each individual (bar) with an optimal k -value of three. Inset shows the Australia-wide distribution of *T. triandra* as a heat map and location of the study area.

Figure 3. Identification of the spatial component of genetic variation using Moran's Eigenvector Maps. Two distinct spatial patterns accounted for most of the 54% of genetic variation explained through isolation by distance. The first MEM variable (a) explained a greater proportion of the variation than the second variable (b). Circles of similar size and colour represent individuals with similar scores on this axis.

Figure 4. Generalised dissimilarity modelling (GDM). (a) Non-linear relationship between climate distance and genomic distance, where points are site pairs. (b) Relationship between predicted genomic distance and observed genomic distance, where points are site pairs. (c) The geographic spline showing the relationship between predicted genomic change and geographic distance. (d–g) Predicted splines showing the estimated relationship between genomic distance and individual climate variables: (d) mean annual precipitation (T_{MA}), (e) maximum temperature of the warmest month (T_{MAX}), (f) mean annual precipitation (P_{MA}), and (g) precipitation seasonality (P_{SEAS}); inset is the amount of variation explained by predicted ploidy polymorphisms (red lines are the model that includes ploidy). Variation explained for the climate-only + spatial model is 31.3% (22% attributed to climate), and with climate, ploidy, and spatial is 31.7% (23% attributed to climate).

Figure 5. Predicted spatial variation in genomic composition based on the outputs from the general dissimilarity models (GDM). Maps include the (a) climate-only GDM and (b) the predicted genomic vulnerability based on comparing the current GDM and the predicted GDM for 2070. Whereas, the (c) climate + ploidy GDM, and (d) and the predicted genomic vulnerability are shown for direct comparison to the climate-only model. A 5% reduction in genomic vulnerability is indicated in the most severely affected areas when including ploidy level in the GDM. The greater the difference (dark orange), the more genomic change is needed to adjust to future climate conditions.

882

883 Tables

884 **Table 1.** Locations and genetic diversity indices for sampled populations. T_{MAX} = maximum
885 temperature of the warmest month; P_{SEAS} = precipitation seasonality; H_s = heterozygosity
886 within populations; G_{IS} = inbreeding coefficient; A_N = number of alleles; C_P = predicted
887 chromosome number.

Pop	X	Y	T_{MAX} (°C)	P_{SEAS} (mm)	A_N	H_s	C_P	G_{IS}
MTG	138.754	-34.977	26.7	52	1.357	0.088	40	0.059
BBNP	153.028	-30.420	28.1	43	1.215	0.066	20	0.181
BCR	153.054	-28.646	28.2	45	1.241	0.071	20	0.182
BL	151.737	-29.867	25.6	33	1.233	0.068	20	0.2
BLAPT	152.807	-31.395	26.9	34	1.232	0.079	20	0.317
BLARD	150.444	-35.196	25.3	24	1.21	0.066	20	0.154
BNR	151.997	-29.113	25.2	32	1.169	0.064	20	0.156
BRA	151.996	-32.631	27.2	25	1.212	0.064	20	0.179
BU	151.076	-30.189	29.5	31	1.164	0.064	20	0.207
BYR	153.620	-28.652	28.1	32	1.203	0.064	20	0.143
CB	150.674	-30.885	30.9	34	1.164	0.062	30	0.137
BCG	143.316	-37.612	26.1	23	1.142	0.043	20	0.085
DCD	150.728	-34.013	28.1	29	1.294	0.083	40	0.121
DCR	149.982	-36.356	24.8	24	1.271	0.08	30	0.062
DW	151.997	-29.114	25.2	32	1.203	0.068	20	0.165
RWCK	146.817	-36.583	28.1	32	1.185	0.062	22	0.366
BUR	145.026	-37.834	26.0	17	1.136	0.06	20	0.328
ANG	144.153	-38.335	23.9	22	1.187	0.07	20	0.138
EUN	152.888	-30.811	27.7	39	1.234	0.067	20	0.184
GHK	149.863	-36.979	24.0	18	1.259	0.073	20	0.236
GOR	150.588	-35.009	25.2	23	1.397	0.102	20	-0.04
GRES	151.219	-32.546	30.1	34	1.231	0.068	20	0.171
QLD	149.878	-27.926	33.5	29	1.309	0.12	40	0.034
JG	152.008	-30.514	25.3	38	1.187	0.065	20	0.196
KCK	152.579	-31.795	27.5	37	1.238	0.068	20	0.177
KOZ	148.402	-35.889	22.4	29	1.255	0.075	20	0.28
KUN	152.844	-31.196	27.3	37	1.20	0.069	20	0.205
L	152.292	-28.411	27.9	37	1.195	0.073	20	0.17
LO	149.998	-33.167	26.4	20	1.203	0.082	20	0.318
MGR	149.077	-36.244	25.0	19	1.254	0.07	20	0.051
ML	152.473	-28.380	26.9	40	1.285	0.083	20	0.218
MNP	150.373	-35.457	24.1	13	1.36	0.089	20	0.127
Mong	149.944	-35.426	25.5	16	1.193	0.064	20	0.252
MS	150.881	-29.988	29.8	31	1.161	0.067	40	0.165
MSF	149.055	-34.825	28.1	13	1.293	0.084	20	0.334
NAB	152.370	-32.086	27.7	35	1.217	0.063	20	0.226
NAM	152.976	-30.639	28.0	41	1.109	0.109	20	---
OPC	153.037	-29.820	28.5	40	1.109	0.064	20	0.116
PR	150.186	-31.418	31.7	34	1.366	0.115	40	-0.135
MSCP	140.631	-37.145	28.1	44	1.242	0.072	20	0.257
SIW	153.146	-30.192	27.4	40	1.147	0.062	20	0.17
SOM	151.286	-33.404	26.1	31	1.248	0.065	20	0.187
SPNR	149.747	-37.557	22.6	12	1.247	0.07	20	0.157
STCK	149.314	-35.360	26.7	13	1.257	0.075	20	0.245
SWC	149.707	-31.400	31.3	31	1.18	0.105	41	-0.753
NSW	148.142	-36.542	26.6	17	1.245	0.067	20	0.151
TOO	152.391	-28.453	28.7	39	1.221	0.068	20	0.183
UL	152.073	-30.537	25.2	39	1.234	0.099	20	-0.326
WOL	150.806	-34.434	25.6	31	1.229	0.065	20	0.183
WYE	151.491	-33.175	26.6	30	1.235	0.065	20	0.195
YNGNP	150.693	-33.057	28.2	38	1.223	0.063	26	0.196
YNR	151.076	-30.189	29.5	31	1.213	0.076	20	0.319
Overall					1.842	0.074		0.134

888

889

890 Supplementary information

891 **Table S1.** F_{ST} pairwise table and input for GDM analysis. (tsv file)

892 **Figure S1.** Elevation of the study area.

893 **Figure S2.** Histogram of MinION long-read read-lengths and average read quality.

894 **Figure S3.** Principal components analysis for all 19 bioclim variables.

895 **Figure S4.** Cross entropy plot to determine the k -value for sNMF results.

896 **Figure S5.** Maps for all four climate variables and ploidy distribution.

897

898

climate variables

climate + ploidy prediction