A tradeoff between tolerance and resistance to a major fungal pathogen in elite wheat cultivars

Alexey Mikaberidze, a.mikaberidze@reading.ac.uk

Bruce A. McDonald, bruce.mcdonald@usys.ethz.ch

Plant Pathology, Institute of Integrative Biology, ETH Zurich

Author for correspondence: Alexey Mikaberidze, School of Agriculture, Policy and Development,
University of Reading, Whiteknights, Reading, UK. Telephone: +44 118 378 7649, email:
a.mikaberidze@reading.ac.uk

Summary

- Tolerance and resistance represent two strategies that hosts evolved to protect themselves from pathogens. Tolerance alleviates the reduction in host fitness due to infection without reducing a pathogen's growth, while resistance reduces pathogen growth. We investigated tolerance of wheat to the major fungal pathogen *Zymoseptoria tritici* in 335 elite wheat cultivars.
- We used a novel digital phenotyping approach that included 11,152 infected leaves and counted 2,069,048 pathogen fruiting bodies.
- We discovered a new component of tolerance that is based on the relationship between the green area remaining on a leaf and the number of pathogen fruiting bodies. We found a negative correlation between tolerance and resistance among intolerant cultivars, presenting the first compelling evidence for a tradeoff between tolerance and resistance to plant pathogens. Surprisingly, the tradeoff arises due to limits in the host resources available to the pathogen and not due to metabolic constraints, contrary to what ecological theory suggests.
- The mechanism underlying this tradeoff may be relevant for many plant diseases in which the amount of host resources available to the pathogen can limit the pathogen population. Our analysis indicates that European wheat breeders may have selected for tolerance instead of resistance to an important pathogen.

keywords: *Triticum aestivum*, host-pathogen interaction, host defenses, plant disease, *Zymoseptoria tritici*, digital phenotyping

Introduction

Tolerance and resistance represent two important mechanisms that plants and animals evolved to protect themselves from pathogens (Roy et al., 2000; Baucom & De Roode, 2011). Tolerance to a pathogen is usually defined as the host's ability to alleviate the reduction in its fitness due to infection without reducing the growth of the pathogen (Baucom & De Roode, 2011; Ney et al., 2013). A more tolerant host genotype will suffer a smaller loss in fitness per unit increase of pathogen population present within the host (called the pathogen burden) than a less tolerant host genotype. Hence, tolerance of a host genotype can be quantified as the reduction in host fitness per unit increase in pathogen burden. In contrast, resistance is usually measured as the host's ability to suppress the infection itself and reduce the resulting pathogen burden upon infection. The difference between tolerance and resistance is illustrated in Fig. 1.

Since the concept of tolerance was first coined more than a century ago (Cobb, 1894), numerous studies [reviewed by Pagan & Garcia-Arenal (2018)] investigated tolerance to pathogens in crop plants (Caldwell *et al.* 1958; Schafer, 1971; Newton *et al.*, 1998; Bingham *et al.*, 2009; Ney *et al.*, 2013; Newton, 2016), model plants (Kover & Schaal, 2002; Pagan et al., 2008, Shuckla et al., 2017) and wild plants (Roy et al., 2000; Inglese & Paul, 2006; Carr et al., 2006). Råberg et al. (2007) were the first to demonstrate tolerance to an infectious disease in animals. The therapeutic potential of tolerance in human medicine inspired a surge of further investigations (Medzhitov et al., 2012; Ayres & Schneider, 2012; Råberg, 2014; Soares et al., 2017). Several studies uncovered molecular mechanisms of tolerance in model animal species (Ayres & Schneider, 2008; Shinzawa et al., 2009; Richardson et al., 2010; Maze-Guilmo et al., 2014), while Blanchet et al. (2010); Jackson et al. (2014); Hayward et al. (2014); Maze-Guilmo et al. (2014) characterized tolerance to parasites in wild animal populations. Zeller & Koella (2017) used an experimental evolution approach to determine how tolerance/resistance evolves in mosquito populations exposed to microsporidian parasites.

Figure 1 Differentiation and quantification of tolerance and resistance. The figure illustrates the relationships between host fitness and pathogen load for three hypothetical host genotypes: A (blue), B (red) and C (yellow). For each host genotype, tolerance is quantified as the rate at which the host loses its fitness with an increase in pathogen burden. The difference in resistance between hosts can be measured on the X-axis as the difference in the mean pathogen burden (vertical lines). The position of each genotype on the Y-axis is determined by its fitness in the absence of the pathogen and is not related to tolerance or resistance. Host genotypes A and B have the same resistance because the average pathogen burden that they carry is the same. However, genotype A is more tolerant than genotype B because the fitness of genotype B decreases at a higher rate with increasing pathogen burden. This is reflected in the steeper slope in genotype B compared to genotype A. In contrast, host genotypes A and C have the same tolerance, but genotype A is more resistant than genotype C. Finally, the comparison of host genotype B and host genotype C represents a mixed case: genotype B is more resistant, but less tolerant than genotype C.

86

87 It is generally thought that since both tolerance and resistance are defense strategies that
88 require reallocation of host resources, they should confer fitness costs to the host (Roy & Kirchner,
89 2000; Simms & Triplett, 1994; Brown, 2002). For this reason, a metabolic tradeoff between tolerance
90 and resistance is expected due to a limitation in host resources. A large body of ecological theory has
91 been developed based on this premise (van der Meijden et al., 1988; Herms & Mattson, 1992; Roy &
92 Kirchner, 2000; Fornoni et al., 2004; Restif & Koella, 2004; Miller et al., 2005; Best et al., 2008).
93 However, empirical evidence for a tradeoff between tolerance and resistance remains sparse. A few
94 studies reported a negative relationship between tolerance and resistance to herbivory (Fineblum &
95 Rausher, 1995; Stowe, 1998; Baucom & Mauricio, 2008) and Råberg et al. (2007) presented a similar
96 finding in mice infected with malaria. Other studies reported no correlation between tolerance and
97 resistance in plants subjected to herbivores (Mauricio et al., 1997), in humans infected with HIV
98 (Regoes et al., 2014) or in wild sheep infected with a parasite (Maze-Guilmo et al., 2014).
99 Interestingly, in *Drosophila melanogaster* populations exposed to a bacterial infection, tolerance and
100 resistance exhibited a positive correlation (Howick & Lazzaro, 2014). Likewise, populations of the
101 mosquito *Aedes aegypti* that were infected by the microsporidian parasite *Vavraia culicis* and evolved
102 for 10 generations exhibited a positive relationship between tolerance and resistance (Zeller & Koella,
103 2017). No evidence for a tradeoff between host tolerance and resistance was so far reported in the
104 plant pathology literature.

105 In this study, we investigated tolerance to the fungal pathogen *Zymoseptoria tritici* (formerly
106 *Mycosphaerella graminicola*) in 335 elite European wheat cultivars. *Z. tritici* causes septoria tritici
107 blotch (STB), a disease that is a major constraint on wheat production globally and the most
108 destructive disease of wheat in Europe (Fones & Gurr, 2015). *Z. tritici* spores germinate on wheat
109 leaves and penetrate the leaves through stomata (Kema et al., 1996). After penetration, the fungus
110 grows for several days within leaves without producing visible symptoms. During this asymptomatic
111 period, the pathogen invades the host mesophyll around the position of the initial penetration. After
112 10 to 20 days of asymptomatic growth, the fungus becomes necrotrophic and kills the invaded plant
113 tissue, forming necrotic lesions. Asexual fruiting bodies called pycnidia begin to form in the necrotic
114 lesions soon thereafter. Spores that form in the pycnidia provide inoculum to start the next cycle of
115 pathogen reproduction. The formation of necrotic lesions corresponds to host damage caused by the
116 pathogen that can be quantified as the proportion of leaf area covered by lesions (PLACL). The

number of pycnidia provides a measure of pathogen reproduction that can be quantified by counting the number of pycnidia present on an infected leaf, N_p (Stewart et al., 2016a; Karisto et al., 2018).

Control of STB relies mainly on applications of fungicides and deployment of STB-resistant wheat varieties. However, populations of *Z. tritici* are extremely diverse due to a high degree of sexual reproduction and large effective population sizes. As a result, the pathogen has the capacity to rapidly adapt to both fungicides (Fraaije et al., 2005; Zhan et al., 2006) and host resistances (Cowger et al., 2000; McDonald and Mundt, 2016) as a result of strong directional selection favoring particular pathogen genotypes. In contrast, host tolerance does not impair pathogen reproduction and is not expected to impose strong directional selection. For this reason, tolerance presents a promising alternative to protect wheat yield that is not prone to pathogen adaptation.

Several previous studies investigated tolerance of wheat to STB empirically (Eyal & Ziv, 1974; Zuckerman et al., 1997; Parker et al., 2004; Foulkes et al., 2006; Collin et al., 2018). Van den Berg et al (2017) used mathematical modeling to reveal functional traits in wheat that contribute to tolerance. These studies used wheat yield (measured as tons of grain per hectare or as the thousand kernel weight) to quantify the plant fitness (the Y-axis in Fig. 1) and the PLACL or healthy area duration (HAD, Waggoner & Berger (1987)) to quantify the pathogen burden (the X-axis in Fig. 1). Accordingly, tolerance was quantified as the yield loss associated with each unit increase in PLACL or unit loss in HAD.

PLACL and HAD quantify the damage that the pathogen causes on an infected host plant. However, these quantities do not necessarily accurately reflect the size of the pathogen population present within the infected host plant (Stewart et al., 2016a; Karisto et al., 2018). For this reason, tolerance measured in these traditional ways is considered to be tolerance to the disease, which may not coincide with tolerance to the pathogen (Gaunt, 1981). The goal in this study was to characterize wheat tolerance to its pathogen, *Z. tritici*. With this in mind, we used (i) green leaf area to quantify a component of plant fitness and (ii) the number of pycnidia per leaf to quantify the pathogen burden. Grain yield is usually seen as a more comprehensive measure of fitness in crop plants than green leaf area. However, a number of field experiments have demonstrated that the reduction in the green area of the three upper-most leaf layers in wheat is a major driver of yield loss induced by STB (Eyal & Ziv, 1974; King et al., 1983; Forrer & Zadoks, 1983; Shaw & Royle 1989b, Thomas et al., 1989), thereby justifying our choice (i) (see Discussion for a more detailed justification). The choice (ii) is justified because the number of pycnidia per leaf was shown to be the main factor influencing the

number of pathogen spores produced on an infected leaf (Stewart et al., 2016a). Moreover, the proportion of the leaf area covered by STB lesions was demonstrated to be largely independent from the number of pycnidia produced on a leaf (Karisto et al., 2018). For these reasons, the number of pycnidia per leaf is a better indicator of the pathogen population inhabiting a leaf than the PLACL.

By conducting these measurements on 11,152 individual wheat leaves belonging to 335 different cultivars (counting in total 2,069,048 individual pycnidia), we were able to identify and measure a novel component of wheat tolerance to *Z. tritici* that operates on the scale of individual leaves. We call this “leaf tolerance” as opposed to the “whole-plant tolerance” that was characterized previously. In this study, we focused on leaf tolerance and did not consider whole-plant tolerance. A way to estimate tolerance over a range of pathogen burdens as we describe here is to estimate range tolerance (Baucom & De Roode, 2011). The component of tolerance that we measured represents fecundity tolerance rather than mortality tolerance, because this disease does not kill its host but instead reduces its fecundity.

We used a combination of mathematical modeling and field experimentation to formulate and test several hypotheses connected to leaf tolerance of wheat to *Z. tritici*. First, based on our current understanding of the infection biology of *Z. tritici*, we formulated and tested empirically two alternative hypotheses regarding the relationship between the green leaf area and the number of pycnidia per leaf. Second, we tested the hypothesis that wheat cultivars differ in terms of their leaf tolerance. Finally, we tested the expectation of a tradeoff between leaf tolerance and resistance and found a significant negative relationship between tolerance and resistance. Surprisingly, our analysis indicates that this negative association arises due to the limitation in the leaf area of wheat plants and not as a result of metabolic costs associated with tolerance/resistance as predicted by ecological theory.

Materials and Methods

Here we analyzed a subset of the raw data reported in (Karisto et al., 2018). Below, we describe the main features of the experimental design that are relevant for this analysis. A comprehensive description of the experimental design can be found in (Karisto et al., 2018).

Plant materials and experimental design

In total, 335 elite European winter wheat (*Triticum aestivum*) varieties from the GABI-wheat

panel (Kollers et al., 2013a,b) were evaluated in this experiment. Two biological replicates of the wheat panel were grown during the 2015-16 growing season in two complete blocks separated by approximately 100 m at the Field Phenotyping Platform site of the Eschikon Field Station of the ETH Zurich, Switzerland (coordinates 47.449°N, 8.682°E) (Kirchgessner et al., 2017). The complete blocks were composed of 18 rows and 20 columns consisting of 1.2 x 1.7 m plots, with the genotypes arranged randomly within each block. Best practices recommended for conventional, high-input wheat production were used, including applications of fertilizers and pesticides. Complete details are given in (Karisto et al., 2018).

Septoria tritici blotch inoculum, sampling of infected leaves

All STB infection was natural, with the majority of primary inoculum likely originating from airborne ascospores coming from nearby wheat fields that surround the Eschikon field site. As a result, the infections analyzed in this experiment were caused by thousands of different pathogen strains. For this study we used leaves exhibiting obvious STB lesions that were collected on 4 July 2016 (approximate range of GS 75 [milk development] to GS 85 [dough development]). Up to 16 infected leaves were collected at random for each plot from the leaf layer below the flag leaf (i.e., flag-1 or second leaf). The sampled leaves were placed into paper envelopes, kept on ice in the field, and stored at 4°C for 2 days before mounting onto A4 paper with printed reference marks and sample names, as described by Stewart et al. (2016b). Absorbent paper was placed between each sheet of eight mounted leaves and sheets were pressed with approximately 5 kg at 4°C for 2 to 3 days prior to scanning at 1,200 dpi with a Canon CanoScan LiDE 220 flatbed scanner.

Determination of the green leaf area and the number of pycnidia per leaf

Scanned images were analyzed with the software ImageJ (Schindelin et al., 2015) using the macro described by Karisto et al. (2018). The maximum length of the scanned area for each leaf was 17 cm. When leaves were longer than 17 cm, bases of the leaves were placed within the scanned area, while the leaf tips extended outside the scanned area. For each leaf, the following quantities were automatically recorded from the scanned image: total leaf area (A_{tot}), necrotic and chlorotic leaf area (A_{necr}) and the number of pycnidia (N_p). Necrotic and chlorotic leaf areas were detected based on discoloration of the leaf surface and were not based on the presence of pycnidia. We then calculated the green (healthy) leaf area as $H = A_{\text{tot}} - A_{\text{necr}}$.

Statistical analysis

Statistical analysis was conducted in the Python programming language (version 3.6.2, <https://www.python.org>) using the open-source packages scipy (version 0.19.1), numpy (version 1.11.1) and matplotlib (version 1.5.3; Jones et al., 2001). The Python package rpy2 (version 2.8.6, <https://rpy2.bitbucket.io/>) was used to access statistical routines of R (R Core Team, 2016).

To control for the effect of total leaf area on the number of pycnidia per leaf, we performed the adjustment $N_{p,i} \rightarrow (A_{tot} / A_{tot,i}) N_{p,i}$ prior to the analysis, where $N_{p,i}$ and $A_{tot,i}$ is the number of pycnidia and the total area of an individual leaf i and A_{tot} is the mean total leaf area averaged over the whole dataset. First, we pooled together the data from leaves belonging to different cultivars and fitted the relationship between N_p and H using a linear function $H = H_0(1 - \kappa N_p)$ and an exponential function $H = H_0 \exp(-\kappa N_p)$, where H is the green leaf area and H_0 is the green leaf area in the absence of disease. For both functions, the slope κ can be used to quantify tolerance: small κ -values correspond to high leaf tolerance and large κ -values correspond to low leaf tolerance. This overall fit gave us the baseline to which we then compared the tolerance of individual wheat cultivars. Second, we estimated κ in each of the 335 wheat cultivars by fitting both the linear and the exponential functions to individual leaf data belonging to each of the cultivars. Next, we used multiple one-sided bootstrap t-tests with resampling cases (Davison and Hinkley, 2001) to compare κ -estimates in each cultivar to the baseline κ -estimate, where we used the false discovery rate correction for multiple comparisons. Fits were performed using the nonlinear ordinary least-squares optimization with the Nelder-Mead method in the lmfit package (version 0.9.7) for Python (Newville et al., 2014). We also determined the significance of the effects of the spatial block and the cultivar on tolerance: we used likelihood ratio tests to compare more complex models in which data from each cultivar/spatial block was fitted using separate κ or H_0 -values to simpler models where only single κ or H_0 parameters were fitted to the whole dataset.

To determine whether there is a relationship between tolerance and resistance of wheat to *Z. tritici*, we used the correlation test based on the Spearman's rank correlation coefficient, r_s (routine “scipy.stats.spearmanr” of the scipy package for Python), to analyze correlations between tolerance quantified as the slope κ and resistance quantified as the average number of pycnidia per leaf (Karisto et al., 2018). We chose to use the Spearman's correlation instead of Pearson's correlation, because it is computed in a non-parametric fashion based on relative ranks of the estimates. It does not assume any

specific functional form of the relationship between the two variables and thereby is not influenced by the widths of their distributions. To test whether the correlation is significantly different from zero, the routine uses a t-test that requires a number of assumptions to be fulfilled such as the normality of the probability distribution of the correlation coefficients.

To determine whether these assumptions hold and the conclusions based on this test are valid, we conducted a series of statistical tests based on a more robust bootstrap t-test (Davison et al., 1997). We generated a large number of bootstrap samples ($n_{bs}=10^5$) by resampling with replacement the estimates of leaf tolerance and resistance for each of the 335 cultivars and used them to compute the 95 % confidence interval (CI) of r_s . We then generated the same number of bootstrap samples based on the estimates of tolerance and resistance separately in each of the two groups of cultivars, tolerant and intolerant cultivars. This allowed us to compute the confidence intervals of r_s estimates in each of the two cultivar groups. Finally, we tested whether r_s was significantly different from zero in each of the groups and whether one of the groups had a significantly higher r_s than the other group.

Results

A novel component of tolerance

Recent studies (Karisto et al., 2018; Stewart et al., 2016a) demonstrated that factors responsible for leaf damage during the infection are largely uncoupled from the pathogen's capacity to reproduce on leaves. Based on this knowledge, we devised a simple mathematical model that describes the change in the green leaf area corresponding to an increase in the number of pycnidia on a leaf (for more details see Notes S1).

The following differential equation governs the relationship between the number of pycnidia and the green leaf area remaining on the leaf:

$$\frac{dH}{dN_p} = -a_p, \quad (1)$$

where a_p is the area of the lesion that corresponds on average to a single pycnidium. Equation (1) represents mathematically a rather general statement that the green area remaining on leaves decreases with increasing numbers of pycnidia. If a_p depends neither on H , nor on N_p , then the solution of Eq. (1) is a linear function:

$$H(N_p) = H_0(1 - \kappa N_p), \quad (2)$$

where $\kappa = a_p/H_0$ and H_0 is the green leaf area in the absence of infection. Alternatively, if a_p is

proportional to the green leaf area H , i.e. $a_p = \kappa H$, then the solution of Eq. (1) is an exponential function:

$$H(N_p) = H_0 \exp(-\kappa N_p). \quad (3)$$

In both its linear and exponential versions, the model predicts that the leaf loses its green area as it carries higher numbers of pycnidia. We consider the green leaf area, H , as a quantity representing plant fitness and the number of pycnidia per leaf, N_p , as a proxy for the pathogen population present on the leaf (pathogen burden). Consequently, the slope of the decrease, κ , characterizes the tolerance of wheat to *Z. tritici*: it measures the amount by which the green leaf area decreases when a single pycnidium is added to the leaf. We call κ the intolerance parameter, as the cultivars with higher κ -values will lose their green leaf area at a higher rate than cultivars with lower κ -values when the number of pycnidia is increased.

Using this model we developed a novel way to measure tolerance of wheat to *Z. tritici* that operates on the scale of individual leaves (“leaf tolerance”) as opposed to the “whole-plant tolerance” that was studied previously in this pathosystem (Eyal & Ziv, 1974; Parker et al., 2004; Foulkes et al., 2006; Collin et al., 2018). We demonstrated in Notes S2 that these two components of tolerance contribute to overall tolerance as multiplicative factors (under the assumption that the relationships between the yield and the damaged leaf area and between the yield and the number of pycnidia per leaf are both linear). Based on two sets of biological assumptions, we formulated two hypotheses about the shape of the relationship between the green leaf area and the number of pycnidia on the leaf represented by Eq. (2) and (3). We next tested these hypotheses using the empirical data gathered from wheat leaves naturally infected by *Z. tritici*.

Relationship between green leaf area and yield

To determine whether the green leaf area in our experiment can be considered a good measure of plant fitness, we studied the correlation between the green leaf area measured on infected leaves and yield measured as the weight of grains per unit area of land [tons per hectare (t/ha)] and as the thousand kernel weight (TKW). Figure 2 illustrates the outcomes: green leaf area averaged over leaves belonging to the same cultivar correlates weakly, but significantly, with the yield of the corresponding cultivar ($r_s = 0.13$, $p = 0.017$ for yield measured in t/ha and $r_s = 0.13$, $p = 0.019$ for yield measured as TKW).

We would like to emphasize that in our experiment, the green leaf area was recorded only on

302 infected leaves, while yield was measured from plants sampled without regard to their infection
 303 status, hence the yield measures comprised both healthy and infected plants. If in addition to the
 304 green area of infected leaves, we were able to also measure the STB incidence (that is the proportion
 305 of second leaves that were infected), then the product of the green leaf area on infected leaves times
 306 the STB incidence would give us the average green leaf area on all second leaves. This quantity
 307 would likely explain a much larger percentage of variation in yield. This has been convincingly
 308 demonstrated in a large number of field experiments, in which the reduction in wheat yield was
 309 strongly correlated with the reduction in the green leaf area of second leaves due to STB (e.g., King et
 310 al., 1983; Shaw & Royle, 1989b).
 311

312 **Figure 2** Correlation between the green leaf area (GLA) and wheat yield. (a) Yield in tons per hectare
 313 is plotted against the GLA of infected leaves measured at GS 75-85. Each value on the x-axis

represents the average value over approximately 30 leaves originating from two different plots belonging to the same cultivar. Each value on the y -axis represents the yield averaged over two plots planted with the same cultivar. (b) Yield measured as thousand kernel weight is plotted against the GLA. Each value on the x -axis represents the average value over approximately 15 leaves originating from a single plot belonging to the same cultivar. Each value on the y -axis represents the yield measured in a single plot.

Green leaf area decreases nonlinearly with the number of pycnidia

Figure 3a shows 11,152 individual leaf measurements of the number of pycnidia per leaf, N_p , and the green leaf area, H . Overall, leaves lose more of their green area when they carry a larger number of pycnidia. The exponential function, Eq. (3), provided a better fit (standard error of the estimate $s=472$, coefficient of determination $R^2=0.37$) than the linear function, Eq. (2) ($s=508$, $R^2=0.28$). For this reason, we estimated the overall slope using the exponential function and obtained the best-fit parameter values: for the slope $\kappa=0.00172$ and the intercept $H_0=1816 \text{ mm}^2$. The spatial block had a significant effect on κ (likelihood ratio 6.9, $p=0.008$) and H_0 (likelihood ratio 41, $p=1.4 \times 10^{-10}$), but the cultivar had a much greater effect on κ (likelihood ratio 856, $p=1.3 \times 10^{-47}$) and H_0 (likelihood ratio 2636, $p<10^{-50}$).

There does not appear to be a clear pattern in terms of the goodness of fit neither for cultivars with different levels of tolerance nor for cultivars with different levels of resistance. To illustrate this, we present the goodness of fit metrics for both fit functions for four cultivars representing contrasting levels of tolerance and resistance. In a more tolerant cultivar Intact (blue curve in Fig. 3c), the linear fit yields $s=349$, $R^2=0.26$, while the exponential fit yields $s=345$, $R^2=0.28$. But in a less tolerant cultivar Lynx (red curve in Fig. 3c), the linear fit gives $s=430$, $R^2=0.59$ compared to the exponential fit that gives $s=415$, $R^2=0.62$. In a more resistant cultivar Element the linear fit gives $s=450$, $R^2=0.2$ and the exponential fit gives $s=448$, $R^2=0.21$; in a less resistant Arack the linear fit gives $s=496$, $R^2=0.37$, while the exponential fit gives $s=490$, $R^2=0.4$. The fits for these two cultivars are shown in Fig. S2.

Ranking of cultivars according to their tolerance to *Z. tritici*

We estimated κ for each cultivar by fitting the empirical dependency of the green leaf area on

the number of pycnidia with the exponential function [Eq. (3)], i.e. we obtained tolerance curves for each cultivar (such as the two tolerance curves depicted Fig. 3a in blue and red). The distribution of the κ -estimates is shown in Fig. 3b. Next, we ranked the cultivars according to their tolerance to *Z. tritici* (see Fig. 3c and Table S1). Smaller κ -values corresponded to more tolerant cultivars. We also compared κ -estimates for each cultivar to the baseline value (black vertical line in Fig. 3c). We found that 22 cultivars were significantly more tolerant than the baseline (blue squares in Fig. 3c) and 25 cultivars were significantly less tolerant than the baseline (red squares in Fig. 3c). Thus, the cultivars that we investigated in our field experiment exhibited significant differences with respect to leaf tolerance.

To determine to what extent the ranking of cultivars with respect to their κ -estimates was conserved between the two replicate blocks, we estimated the κ -values for each cultivar separately in each of the blocks. The κ -estimates exhibited a positive and significant correlation between the two replicates ($r_s=0.18$, $p=0.001$). In addition, we obtained similar data for a subset of 38 cultivars in 2015 (Stewart et al., 2016), which allowed us to evaluate the robustness of the outcomes. The κ -estimates exhibited a positive but a non-significant correlation between the two years ($r_s=0.3$, $p=0.07$).

Figure 3 Tolerance of wheat to *Z. tritici* measured on the scale of individual leaves. (a) Green (Healthy) leaf area normalized by the leaf area in the absence of disease, H/H_0 , is plotted versus the

number of pycnidia per leaf N_p . 11152 individual leaf measurements are shown using grey points. Best fit curves based on the exponential function $H/H_0 = \exp[-\kappa N_p]$ are shown for all data (black curve) and for two example cultivars, a tolerant cultivar (Intact, blue curve), and an intolerant cultivar (Lynx, red curve). (b) The distribution of the 335 cultivars with respect to their κ -estimates. The most tolerant cultivars are those with the lowest κ -estimates at the left of the distribution. (c) Ranking of wheat cultivars according to their tolerance. κ -estimates for the 335 cultivars are shown in order of decreasing tolerance, that is increasing slope κ (grey points). Cultivars with tolerance significantly different from the baseline tolerance (black line) are marked with blue points (more tolerant) and red points (less tolerant), according to one-sided bootstrap t-tests with the confidence threshold of 0.05. Cultivars illustrated in panel (a) are marked using larger blue (cultivar Intact) and red (cultivar Lynx) squares.

Relationship between tolerance/resistance and the year of cultivar registration

In a subset of 205 out of 335 cultivars, we had information on cultivar registration years. In those cultivars, tolerance increased with the year of cultivar registration: the correlation between the intolerance parameter κ and the cultivar's registration year was negative and significant ($r_s = -0.17$, $p = 0.02$). In contrast, resistance did not exhibit a significant correlation with the cultivar's registration year ($r_s = -0.06$, $p = 0.36$).

Evidence for a tradeoff between leaf tolerance and resistance

We found that the estimates of tolerance, κ , for each of the 335 cultivars correlated negatively with the mean number of pycnidia per leaf, N_p , the measure of resistance to STB, with $r_s = -0.27$, $p = 5.8 \times 10^{-7}$ (Fig. 4). Interestingly, κ -estimates correlated positively with mean PLACL values in each cultivar ($r_s = 0.31$, $p = 8.6 \times 10^{-9}$).

Why are more tolerant cultivars on average less resistant? In other words, why do more tolerant cultivars carry more pycnidia on their leaves than less tolerant cultivars? A possible explanation is that since the leaf area is limited, this places a constraint on the maximum number of pycnidia that a leaf can carry. This constraint should lead to a negative relationship between N_p and κ in cases where the number of pycnidia on leaves belonging to the same cultivar approach the

maximum allowed by the leaf area. A less tolerant cultivar suffers a larger necrotic area forming, on average, per pycnidium [parameter a_p in Eq. (1)]. Consequently, the maximum number of pycnidia per leaf is lower in a less tolerant cultivar than in a more tolerant cultivar. Therefore, the limitation in the leaf area should affect more strongly pathogen populations infecting less tolerant cultivars, where the green leaf area decreases more steeply with increasing numbers of pycnidia (Fig. 5a). If the limitation in the leaf area is indeed the dominant factor responsible for the negative relationship between tolerance and resistance, then the negative correlation should be present in intolerant cultivars, but absent in tolerant cultivars. This is because only intolerant cultivars have a high proportion of their leaf area covered by lesions already at rather modest numbers of pycnidia (red curves in Fig. 5a).

To test this expectation in a more quantitative fashion, we subdivided all cultivars into two groups according to their tolerance estimates using the baseline tolerance ($\kappa=0.00172$) as the threshold. Next, we conducted the Spearman's correlation test (based on a t-test) in each of the groups separately. We found that intolerant cultivars exhibited a significant correlation between tolerance and resistance ($r_s = -0.34$, $p = 4.3 \times 10^{-6}$), while tolerant cultivars showed no significant correlation between the two traits ($r_s = -0.09$, $p = 0.27$). To test the validity of this outcome, we performed a series of more robust tests based on a bootstrap t-test. We first computed the uncertainty in the estimate $r_s = -0.27$ for all cultivars in the form of the 95 % confidence interval: CI, -0.37 to -0.16. We also computed the uncertainties in r_s -estimates for the two groups of cultivars, tolerant ($r_s = -0.09$, CI, -0.25 to 0.07) and intolerant ($r_s = -0.34$, CI, -0.47 to -0.2). Figure S1 visualizes the bootstrap distributions of r_s and the CIs of r_s -estimates. The bootstrap t-test confirmed the outcome of the conventional t-test: the correlation between tolerance and resistance was significant among intolerant cultivars ($p = 4.0 \times 10^{-6}$) and not significant among tolerant cultivars ($p = 0.28$). Furthermore, we used a more stringent bootstrap t-test to compare the r_s -estimates among tolerant and intolerant cultivars and found that the correlation among intolerant cultivars was significantly more negative than among tolerant cultivars ($p = 0.019$).

Figure 5 illustrates why intolerant cultivars should be more strongly affected by the limitation in the leaf area than tolerant cultivars. Figure 5a shows the tolerance curves for tolerant (blue) and intolerant (red) cultivars. We determined the maximum number of pycnidia that can be reached in each cultivar by computing the number of pycnidia, N_{pm} , at which 95 % of the green leaf area is lost

423 on average (this is given by the intersection of each tolerance curve with the horizontal line $H/$
424 $H_0=0.05$ in Fig. 5a). Next, we computed the ranges in terms of N_{pm} corresponding to tolerant cultivars
425 (blue-shaded area in Fig. 5b) and intolerant cultivars (red-shaded area in Fig. 5b). To determine the
426 extent to which pathogen populations infecting tolerant and intolerant cultivars could be affected by
427 the limitation in the leaf area, we compared these ranges with the overall distribution of leaves with
428 respect to the numbers of pycnidia they carry (cf. the histogram in Fig. 5b and the blue- and red-
429 shaded areas). While the blue-shaded area in Fig. 5b contained 38 leaves that constitute only about
430 0.3 % of the total number of 11,152 infected leaves, the red-shaded area contained a much more
431 substantial proportion (22 %) of the leaves (2,485 leaves out 11,152). Therefore, the intolerant
432 cultivars contained a much greater proportion of the leaves on which pathogen populations were
433 likely to be affected by limitations in the leaf area compared to the tolerant cultivars. Combined with
434 the observation that only intolerant cultivars exhibited a negative relationship between tolerance and
435 resistance, these data support our hypothesis that the negative relationship between tolerance and
436 resistance arises largely due to a limitation in the leaf area.

Figure 4 Correlation between leaf tolerance and resistance of wheat to STB. The measure of resistance, N_p is plotted against the measure of tolerance, κ , for each of 335 wheat cultivars (grey circles). Horizontal line shows the mean resistance and vertical line shows the baseline tolerance; blue/red circles mark the cultivars that are significantly more/less tolerant than the baseline.

Figure 5 Limitation in the leaf area in tolerant and intolerant wheat cultivars. (a) Tolerance curves fitted to individual leaf data (raw data points are shown in Fig. 3a) for 335 wheat cultivars, blue curves correspond to cultivars more tolerant than the baseline (black curve) and red curves correspond to cultivars less tolerant than the baseline. The horizontal line shows the threshold $H/H_0=0.05$. (b) Distribution of 11,152 individual infected leaves with respect to the number of pycnidia per leaf, N_p . Shaded areas illustrate the ranges of N_p -values in which the pathogen population infecting tolerant (blue) and intolerant (red) wheat varieties is affected by the limitation in the leaf area.

Discussion

We discovered a novel component of tolerance in wheat to *Z. tritici* that operates on the scale of individual leaves (leaf tolerance). We devised an approach to quantify leaf tolerance empirically based on automated measurements of the green leaf area and the numbers of pycnidia on individual leaves. We gathered data from 11,152 individual infected leaves and characterized leaf tolerance in 335 elite European wheat cultivars. Cultivars exhibited significant differences in leaf tolerance, suggesting that this trait is at least partially under genetic control. We also found a negative relationship between leaf tolerance and resistance to *Z. tritici*, indicating that there is a tradeoff between tolerance and resistance. Our study presents the first clear evidence for such a tradeoff in the context of plant-pathogen interactions. We discuss the consequences of this possible tradeoff for the selection of tolerance/resistance in agricultural host populations.

Surprisingly, the nature of this tradeoff turned out to be different from what we expected based on ecological theory. Our analysis shows that the tradeoff is only present in cultivars with less than average tolerance (intolerant cultivars) and that the limitation in the leaf area is the dominant factor responsible for its occurrence. This mechanism differs from the host metabolic constraints that are usually implicated in the tradeoff between resistance and tolerance. We expect this novel mechanism underlying a tradeoff between tolerance and resistance could operate across a large class of infectious diseases in plants and animals, in which tolerance to the pathogen can be measured and the amount of host resources available to the pathogen can limit the pathogen population within the host. A key conceptual outcome of our study is that observing a negative relationship between tolerance and resistance is not necessarily indicative of a metabolic tradeoff, whereby tolerance and resistance confer fitness costs. Instead, as we show here, a tradeoff can arise via an entirely different mechanism, namely a limitation in the amount of host tissue or resources available to the pathogen (or more generally a limitation in the degree of fitness a host can lose because of infection). In the two prominent examples of a negative relationship between tolerance and resistance found in the literature (herbivory in plants, Fineblum & Rausher, 1995; malaria in mice, Råberg et al., 2007), the mechanisms underlying this relationship remain unknown.

The limitation in the leaf area is expected to constrain the evolution of pathogen populations towards higher reproductive fitness on intolerant cultivars, but the pathogen may overcome this limitation by evolving lower virulence (Anderson & May, 1982). According to our current understanding in ecological theory, a metabolic tradeoff between tolerance and resistance is expected

that arises due to limitation in resources available to the host. Our data does not exclude the possibility of a metabolic tradeoff, but its detection may require an even more comprehensive dataset than what we have at hand. Evidence for the metabolic tradeoff is more likely to be found in the future by considering a larger number of sufficiently tolerant cultivars, because as we demonstrated here, in more tolerant cultivars the relationship between tolerance and resistance is not dominated by the limitation in the leaf area.

Jackson et al. (2014) reported that mature male voles were more tolerant to macroparasite infection compared to young males. Zeller & Koella (2017) found that the availability of nutrients influenced the magnitude of tolerance to microsporidian parasites in mosquito populations: mosquitos that had a restricted food supply were generally less tolerant to infection. It is plausible that both of these factors influence the tolerance of wheat leaves to *Z. tritici*. First, only the STB-induced damage on the three upper-most wheat leaves correlates strongly with yield loss (e.g., Thomas et al., 1989). Hence leaf tolerance may confer a fitness advantage to plants only during the later developmental stages when these leaf layers have already emerged. As a result, selection may have favored tolerance to manifest only during this late stage of development (similar to the adult plant resistance that is well known for several plant diseases). Second, the severity of STB epidemics is known to increase with increased rates of nitrogen fertilization (Leitch & Jenkins, 1995). This may result from an improved nutritional or physiological status of the leaves or a more disease-conducive physical environment. Hence leaf tolerance and its relationship with resistance may be affected by changing the rate of nitrogen application. Empirical investigation of both of these factors is feasible in the *Z. tritici*-wheat pathosystem and would improve our understanding of the ecological determinants of tolerance.

The dataset we used to characterize tolerance to a plant pathogen is unusually large compared to previous studies. For example, the number of different wheat genotypes used to study tolerance of wheat to STB in earlier studies ranged from 2 to 25 (Eyal & Ziv, 1974; Zuckerman et al., 1997; Parker et al., 2004; Foulkes et al., 2006; Collin et al., 2018). Råberg et al. (2007) investigated tolerance of mice to malaria infection using five mouse strains and three strains of *Plasmodium chabaudi*. Only the study of human tolerance to HIV (Regoes et al., 2014) and the study of tolerance in the wild population of Soay sheep to a gastrointestinal nematode infection (Hayward et al., 2014) had comparably large datasets that included thousands of infected individuals. Remarkably, as we demonstrated here, both tolerance and resistance can be readily quantified from digital images of infected leaves.

Our analyses and interpretations are based on two important assumptions: (i) the reduction in the green area of the second leaf is a major driver of yield loss induced by STB, and; (ii) the number of pycnidia per leaf is a good measure of the size of the pathogen population on a leaf. We justify these assumptions as follows: (i) Compared to infected leaves that have a large fraction of their surface area covered by lesions, leaves with a larger green area intercept a larger fraction of the incoming radiation, which contributes to plant yield. There is overwhelming empirical evidence showing that the reduction in the green leaf area is a major driver of yield loss for many leaf-affecting diseases of wheat (for example, Teng and Gaunt, 1980; Seck et al., 1991; Gaunt 1995; Bhathal et al., 2003), including STB (Eyal & Ziv, 1974; King et al., 1983; Forrer & Zadoks, 1983; Shaw & Royle, 1989b; Thomas et al., 1989). In particular, these studies conclude that the reduction in yield is strongest for the three upper leaves (including the second leaf on which we focused in this study) if the green leaf area is measured during the critical phase of seed development. In our field experiment, we could not determine yield corresponding to each individual infected leaf that we sampled. However, we measured overall yield per plot and found significant correlations between the green leaf area of second leaves (sampled at GS 75-85) and yield, measured both as tons per hectare and as thousand kernel weight ($r_s=0.13$, $p=0.017$ for yield measured in t/ha and $r_s=0.13$, $p=0.019$ for yield measured as TKW, see Results, Fig. 2). Note that the green leaf area was recorded only on infected leaves, while the yield was measured from plants sampled without regard to their infection status, hence the samples used to calculate yield comprised both healthy and infected plants. Therefore, the correlation coefficients we obtained here are likely to considerably underestimate the actual correlations between the green leaf area and yield, consistent with previous studies that found much stronger correlations (Eyal & Ziv, 1974; King et al., 1983; Forrer & Zadoks, 1983; Shaw & Royle 1989b). In particular, Thomas et al., (1989) reported that STB severity on second (flag-1) leaves had a particularly strong effect on yield. Thus, there is compelling empirical evidence in the existing literature and also an indication in the present study showing that the reduction in the green leaf area of second leaves contributes substantially to the yield loss induced by the disease.

To justify (ii) we first note that in this study we investigate tolerance and resistance from an evolutionary perspective. Hence, the measure of pathogen burden should reflect the reproductively active population of the pathogen. Measuring the total number of spores produced per leaf may provide a better way to quantify pathogen burden, but was not possible in our experiment because it could not be automated. However, we believe that the number of pycnidia is a reasonable proxy of

pathogen burden because the number of pycnidia was shown to be the main factor determining the number of pathogen spores produced on an infected leaf (Stewart et al., 2016a). In addition, a recent field experiment showed that the proportion of the leaf area covered by STB lesions was largely independent from the number of pycnidia produced on a leaf (Karisto et al., 2018). Combining these two findings led us to conclude that the number of pycnidia per leaf is a better measure to quantify the pathogen population inhabiting a leaf than the area of a leaf damaged by infection.

According to our statistical analysis, an exponential decrease better fits the empirical dependency of the green leaf area on the number of pycnidia per leaf than a linear decrease, demonstrating that leaf tolerance curves were nonlinear. This deviates from what was reported in earlier analyses of wheat tolerance to *Z. tritici*: tolerance curves were typically fitted using linear functions (Eyal & Ziv, 1974; Parker et al., 2004; Foulkes et al., 2006), with the notable exception of the study by Shaw & Royle (1989b) that used a family of nonlinear curves. It was important to establish the departure from linearity in our study for two reasons. First, it allowed a more accurate comparison of tolerance estimates in different cultivars against the baseline. Second, it provided additional insight into the biology of the infection, because the linear model and the exponential model are based on different biological assumptions.

Our analysis of the model (Notes S1) demonstrates that the linear function Eq. (2) approximates well the relationship between the green leaf area and the number of pycnidia on the leaf when the number of pycnidia on the leaf is sufficiently low. This implies that the number of lesions on the leaf is also likely to be low, with the necrotic area covering only a small proportion of the total leaf area. Under this scenario, lesions develop mostly independently of each other. However, when lesions start to occupy a large proportion of the total leaf area, they become more likely to influence each other's development due to limitations in space and/or resources. Under this scenario, Eq. (2) is no longer a good approximation and the necrotic area $a_p = a_l / n_p$ that corresponds to a single pycnidium may depend on both green leaf area H and the number of pycnidia N_p already present on the leaf (i.e., a density dependence). Above, we considered the simplest case of this dependency when a_p is proportional to H , which resulted in the exponential solution [Eq. (3)]. This dependency may result from the lesion area a_l being proportional to the remaining green leaf area H . Biologically, this means that as more of the green leaf area becomes occupied by lesions, lesions tend to grow to a smaller size due to limitations in available green space and/or resources in the leaf. Alternatively, this dependency may arise due to the number of pycnidia per lesion n_p being inversely proportional to the

577 remaining green leaf area H . This may occur due to an increased activation of plant defenses as more
578 of the leaf area becomes occupied by lesions. Since our analysis shows that Eq. (3) is better supported
579 by the data we collected for *Z. tritici* than Eq. (2), we conclude that density dependence contributes to
580 the relationship between the number of pycnidia and the green leaf area, and is therefore expected to
581 influence epidemiological dynamics on the scale of individual leaves. However, dedicated
582 experiments under controlled conditions will be needed to reveal the mechanism behind the density
583 dependence.

584 We recently identified several chromosomal regions and candidate genes in the wheat genome
585 associated with resistance to STB (quantified as the mean pathogen burden, i.e. the mean number of
586 pycnidia per leaf) using the same phenotypic dataset (Karisto et al., 2018) and a genome-wide
587 association study (GWAS; Yates, et al., 2019). We hypothesize that a GWAS based on the leaf-level
588 tolerance estimates that we report here could also identify significantly associated chromosomal
589 regions. This would indicate that leaf-level tolerance has an underlying genetic basis and is subject to
590 evolutionary processes, and potentially elucidate molecular mechanisms affecting leaf-level
591 tolerance. One possible mechanism could be related to additive actions of toxin sensitivity genes
592 carried by different wheat cultivars that interact with host-specific toxins produced by the pathogen,
593 as demonstrated for *Parastagonospora nodorum* on wheat (Friesen et al., 2008; Oliver et al., 2012).
594 This mechanism would contribute to tolerance if the number of actively interacting toxin - toxin
595 sensitivity gene pairs exceeds a threshold beyond which the removal of a single gene pair does not
596 impair the pathogen reproduction, but nevertheless reduces the host damage, thereby decreasing the
597 average necrotic area per pycnidium and increasing leaf tolerance.

598 Breeding for resistance to STB disease is based on disease assessments that do not quantify
599 pathogen reproduction on the leaves. The amount of disease is typically assessed visually using a
600 categorical scale of severities corresponding to different ranges in terms of the proportion of necrotic
601 area on the leaves (or PLACL). As a result, breeders are likely to select for cultivars with lower
602 PLACL. But as we have shown above, cultivars with lower PLACL are on average more tolerant.
603 Hence, by focusing on PLACL wheat breeders may have inadvertently selected for increased
604 tolerance. Due to the tradeoff between tolerance and resistance, this simultaneously favors lower
605 levels of STB resistance. Some support for this hypothesis is given by our preliminary analysis of the
606 relationship between tolerance/resistance and the year of cultivar registration. In a subset of 205 out
607 of 335 cultivars, we found that estimates of tolerance increased with the year of cultivar registration,

while estimates of resistance did not exhibit a significant change over time κ . This pattern suggests that the selection practices used by plant breeders have led to wheat populations in Europe with higher tolerance but lower resistance to STB over time.

The method to quantify leaf tolerance presented here can potentially be used to measure tolerance to other pathogens that infect plant leaves. The necessary condition is that digital images of infected leaves should enable quantitative measurements of both the damage to the plant induced by the pathogen and the size of the pathogen population on the leaf. This should be possible for many necrotrophic pathogens that form visible fruiting bodies on the leaf surface. Using this approach may facilitate the discovery of similar tradeoffs between tolerance and resistance across a wide array of plant-pathogen systems.

Acknowledgements

We thank P. Karisto, D. Dos Santos Pereira, S. Fouché, and L. Meile for help in collecting and processing leaf samples; H. Zellweger for managing the wheat trial; and A. Hund and J. Anderegg for providing data on wheat yield. A. Mikaberidze would like to thank P. Karisto and R. Regoes for fruitful discussions. A. Mikaberidze gratefully acknowledges financial support from the Swiss National Science Foundation through the Ambizione grant PZ00P3_161453.

Author Contributions

AM and BAM conceived and designed the experiment. AM supervised the field collection and processing of leaf samples. AM analyzed the data in discussions with BAM. AM wrote the manuscript. AM and BAM revised the manuscript.

Data Accessibility Statement

Raw data is available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.171q4>. We confirm that, should the manuscript be accepted, additional data supporting the results will be archived in an appropriate public repository such as Dryad or Figshare, and the data DOI will be provided at the end of the article.

References

- Anderson, R. M. & May, R.M. (1982). Coevolution of hosts and parasites. *Parasitology*, 85, 411-426.
- Ayres, J.S. & Schneider, D.S. (2008). A Signaling Protease Required for Melanization in *Drosophila* Affects Resistance and Tolerance of Infections. *PLoS Biol.*, 6, e305.
- Ayres, J.S. & Schneider, D.S. (2012). Tolerance of Infections. *Annu. Rev. Immunol.*, 30, 271-294.
- Baucom, R.S. & De Roode, J.C. (2011). Ecological immunology and tolerance in plants and animals. *Funct. Ecol.*, 25, 18-28.
- Baucom, R.S. & Mauricio, R. (2008). Constraints on the evolution of tolerance to herbicide in the common morning glory: Resistance and tolerance are mutually exclusive. *Evolution*, 62, 2842-2854.
- van den Berg, F, Paveley ND, Bingham IJ & van den Bosch F. (2017). Physiological Traits Determining Yield Tolerance of Wheat to Foliar Diseases. *Phytopathology*, 107, 1468-1478.
- Best, A., White, A. & Boots, M. (2008). Maintenance of host variation in tolerance to pathogens and parasites. *Proc. Natl. Acad. Sci.*, 105, 20786-20791.
- Bhathal, J.S., Loughman, R. & Speijers, J. (2003). Yield reduction in wheat in relation to leaf disease from yellow (tan) spot and septoria nodorum blotch. *European Journal of Plant Pathology*, 109, 435–443.
- Bingham, I.J., Walters, D.R., Foulkes, M.J. & Paveley, N.D. (2009). Crop traits and the tolerance of wheat and barley to foliar disease. *Ann. Appl. Biol.*, 154, 159-173.
- Blanchet, S., Rey, O. & Loot, G. (2010). Evidence for host variation in parasite tolerance in a wild fish population. *Evol. Ecol.*, 24, 1129-1139.

670

671 Brown, J.K. (2002). Yield penalties of disease resistance in crops. *Curr. Opin. Plant Biol.*, 5, 339-

672 344.

673

674 Caldwell, R.M., Schafer, J.F., Compton, L.E. & Patterson, F.L. (1958). Tolerance to Cereal Leaf

675 Rusts. *Science*, 128, 714-715.

676

677 Carr, D.E., Murphy, J.F. & Eubanks, M.D. (2006). Genetic variation and covariation for resistance

678 and tolerance to Cucumber mosaic virus in *Mimulus guttatus* (Phrymaceae): A test for costs and

679 constraints. *Heredity*, 96, 29-38.

680

681 Cobb, N.A. (1894). Contributions to an economic knowledge of Australian rust (Uredinae) Chap. 10.

682 Agr. Gaz. N. S. W., 5, 239-50.

683

684 Collin, F., Bancal, P., Spink, J., Appelgren, P.K., Smith, J., Paveley, N.D., Bancal, M.O. & Foulkes,

685 M.J. (2018). Wheat lines exhibiting variation in tolerance of *Septoria tritici* blotch differentiated by

686 grain source limitation. *Field Crop. Res.*, 217, 1-10.

687

688 Cowger, C., Hoffer, M.E. & Mundt, C.C. (2000). Specific adaptation by *Mycosphaerella graminicola*

689 to a resistant wheat cultivar. *Plant Pathol.*, 49, 445-451.

690

691 Davison, A. C. & Hinkley, D. V. (1997). Bootstrap methods and their applications. Cambridge

692 University Press, Cambridge, UK.

693

694 Eyal, Z. & Ziv, O. (1974). The relationship between epidemics of *Septoria* leaf blotch and yield

695 losses in spring wheat. *Phytopathology*, 64, 1385-1389.

696

697 Fineblum, W.L. & Rausher, M.D. (1995). Tradeoff between resistance and tolerance to

698 herbivore damage in a morning glory. *Nature*, 377, 517-520.

699

700 Fones, H. & Gurr, S. (2015). The impact of *Septoria tritici* Blotch disease on wheat: An EU

701 perspective. *Fungal Genet. Biol.*, 79, 3-7.

702

703 Fornoni, J., Nunez-Farfan, J., Valverde, P.L. & Rausher, M.D. (2004). Evolution of Mixed Strategies
704 of Plant Defense Allocation against Natural Enemies. *Evolution*, 58, 1685-1695.

705

706 Forrer, H. R. & Zadoks, J. C. (1983). Yield reduction in wheat in relation to leaf necrosis caused by
707 *Septoria tritici*. *Neth. J. Pl. Path.*, 89, 87-98.

708

709 Foulkes, M.J., Paveley, N.D., Worland, A., Welham, S.J., Thomas, J. & Snape, J.W. (2006). Major
710 Genetic Changes in Wheat with Potential to Affect Disease Tolerance. *Phytopathology*, 96, 680-688.

711

712 Friesen, T. L., Faris, J. D., Solomon, P. S., and Oliver, R. P. (2008). Host-specific toxins: Effectors of
713 necrotrophic pathogenicity, *Cell. Microbiol.*, 10, 1421-1428.

714

715 Fraaije, B., Cools, H.J., Fountaine, J., Lovell, D.J., Motteram, J., West, J.S. & Lucas, J. A. (2005).
716 Role of Ascospores in Further Spread of QoI-Resistant Cytochrome b Alleles (G143A) in Field
717 Populations of *Mycosphaerella graminicola*. *Phytopathology*, 95, 933-941.

718

719 Gaunt, R. (1981). Disease Tolerance - an Indicator of Thresholds? *Phytopathology*, 71, 915-916.

720

721 Gaunt, R.E. (1995). The Relationship Between Plant Disease Severity and Yield. *Annual Review of*
722 *Phytopathology*, 33, 119-144.

723

724 Hayward, A.D., Nussey, D.H., Wilson, A.J., Berenos, C., Pilkington, J.G., Watt, K.A., Pemberton,
725 J.M. & Graham, A.L. (2014). Natural Selection on Individual Variation in Tolerance of
726 Gastrointestinal Nematode Infection. *PLoS Biol.*, 12, 1-13.

727

728 Herms, D.A. & Mattson, W.J. (1992). The Dilemma of Plants: To Grow or Defend. *Q. Rev. Biol.*, 67,
729 283-335.

730

731 Howick, V.M. & Lazzaro, B.P. (2014). Genotype and diet shape resistance and tolerance across

732 distinct phases of bacterial infection. *BMC Evol. Biol.*, 14, 1-13.

733

734 Inglese, S.J. & Paul, N.D. (2006). Tolerance of *Senecio vulgaris* to Infection and Disease Caused by

735 Native and Alien Rust Fungi. *Phytopathology*, 96, 718-726.

736

737 Jackson, J., Hall, A., Friberg, I., Ralli, C., Lowe, A., Zawadzka, M., Turner, A.K., Stewart, A.,

738 Birtles, R.J., Paterson, S., Bradley, J.E. & Begon, M. (2014). An immunological marker of tolerance

739 to infection in wild rodents. *PLoS Biol.*, 12, e1001901.

740

741 Jones E., Oliphant E., Peterson P. et al. (2001). SciPy: Open Source Scientific Tools for Python.

742 Available at <http://www.scipy.org/>. Last accessed 30.05.2018.

743

744 Karisto, P., Hund, A., Yu, K., Anderegg, J., Walter, A., Mascher, F., McDonald, B.A. &

745 Mikaberidze, A. (2018). Ranking quantitative resistance to *Septoria tritici* blotch in elite wheat

746 cultivars using automated image analysis. *Phytopathology*, 108, 568-581

747

748 Kema, G.H., Yu, D., Rijckenberg, F.H., Shaw, M.W. & Baayen, R.P. (1996). Histology of the

749 pathogenesis of *mycosphaerella graminicola* in wheat. *Phytopathology*, 86, 777-786.

750

751 King, J.E., Jenkins, J.E. & Morgan, W.A. (1983). The estimation of yield losses in wheat from

752 severity of infection by *Septoria* species. *Plant Pathology*, 32, 239-249.

753

754 Kollers, S., Rodemann, B., Ling, J., Korzun, V., Ebmeyer, E., Argillier, O., Hinze, M., Plieske, J.,

755 Kulosa, D., Ganai, M.W. & Roder, M.S. (2013a). Genetic architecture of resistance to *Septoria*

756 *tritici* blotch (*Mycosphaerella graminicola*) in European winter wheat. *Mol. Breed.*, 32, 411-423.

757

758 Kollers, S., Rodemann, B., Ling, J., Korzun, V., Ebmeyer, E., Argillier, O., Hinze, M., Plieske, J.,

759 Kulosa, D., Ganai, M.W. & Roder, M.S. (2013b). Whole Genome Association Mapping of

760 *Fusarium* Head Blight Resistance in European Winter Wheat (*Triticum aestivum* L.). *PLoS One*, 8.

761

762 Kover, P.X. & Schaal, B.A. (2002). Genetic variation for disease resistance and tolerance among

763 *Arabidopsis thaliana* accessions. *Proc. Natl. Acad. Sci.*, 199, 11270-11274.

764

765 Leitch, M.H. & Jenkins P.D. (1995). Influence of nitrogen on the development of Septoria epidemics

766 in winter wheat. *Journal of Agricultural Science*, 124, 361-368.

767

768 Linde, C.C., Zhan, J. & McDonald, B.A. (2002). Population Structure of *Mycosphaerella*

769 *graminicola*: From Lesions to Continents. *Phytopathology*, 92, 946-55.

770

771 Mauricio, R., Rausher, M.D. & Burdick, D.S. (1997). Variation in the defense strategies of plants:

772 Are resistance and tolerance mutually exclusive? *Ecology*, 78, 1301-1311.

773

774 Maze-Guilmo, E., Loot, G., Paez, D.J., Lefevre, T. & Blanchet, S. (2014). Heritable variation in host

775 tolerance and resistance inferred from a wild host-parasite system. *Proc. R. Soc. B Biol. Sci.*, 281,

776 20132567-20132567.

777

778 Medzhitov, R., Schneider, D.S. & Soares, M.P. (2012). Disease Tolerance as a Defense Strategy.

779 *Science*, 335, 936-942.

780

781 van der Meijden, E., Wijn, M. & Verkaar, H.J. (1988). Defence and Regrowth, Alternative Plant

782 Strategies in the Struggle against Herbivores. *Oikos*, 51, 3550-363.

783

784 Miller, M.R., White, A. & Boots, M. (2005). The evolution of host resistance: Tolerance and control

785 as distinct strategies. *J. Theor. Biol.*, 236, 198-207.

786

787 Newville, M., Stensitzki, T., Allen, D. B., & Ingargiola, A. (2014, September 21). LMFIT: Non-

788 Linear Least-Square Minimization and Curve-Fitting for Python¶. Zenodo.

789 <http://doi.org/10.5281/zenodo.11813>

790

791 Newton, A.C. (2016). Exploitation of Diversity within Crops: the Key to Disease Tolerance? *Front.*

792 *Plant Sci.*, 7, 1-12.

793

794 Newton A.C., Guy D.C., Gaunt R.E., Thomas W.T.B. (2000). The effect of powdery mildew
795 inoculum pressure and fertiliser level on disease tolerance in spring barley. *Journal of Plant Diseases*
796 *and Protection*, 107, 67–73.

797

798 Ney, B., Bancal, M.O., Bancal, P., Bingham, I.J., Foulkes, J., Gouache, D., Paveley, N. & Smith, J.
799 (2013). Crop architecture and crop tolerance to fungal diseases and insect herbivory. Mechanisms to
800 limit crop losses. *Eur. J. Plant Pathol.*, 135, 561-580.

801

802 Oliver, R. P., Friesen, T. L., Faris, J. D., and Solomon, P. S. (2012). *Stagonospora nodorum*: From
803 pathology to genomics and host resistance. *Annu. Rev. Phytopathol.*, 50, 23-43.

804

805 Pagan, I. & Garcia-Arenal, F. (2018). Tolerance to Plant Pathogens: Theory and Experimental
806 Evidence. *Int. J. Mol. Sci.*, 19, 810.

807

808 Pagan, I., Alonso-Blanco, C. & Garcia-Arenal, F. (2008). Host responses in life-history traits and
809 tolerance to virus infection in *Arabidopsis thaliana*. *PLoS Pathog.*, 4.

810

811 Parker, S.R., Shaw, M.W. & Royle, D.J. (1995). The reliability of visual estimates of disease severity
812 on cereal leaves. *Plant Pathol.*, 44, 856-864.

813

814 Parker, S.R., Welham, S., Paveley, N.D., Foulkes, J. & Scott, R.K. (2004). Tolerance of septoria leaf
815 blotch in winter wheat. *Plant Pathol.*, 53, 1-10.

816

817 R Core Team (2016). R: A Language and Environment for Statistical Computing. R Foundation for
818 Statistical Computing, Vienna, Austria.

819

820 Råberg, L. (2014). How to Live with the Enemy: Understanding Tolerance to Parasites. *PLoS Biol.*,
821 12.

822

823 Råberg, L., Sim, D. & Read, A.F. (2007). Disentangling Genetic Variation for Resistance and
824 Tolerance to Infectious Diseases in Animals. *Science*, 318, 812-815.

825

826 Regoes, R.R., McLaren, P.J., Battegay, M., Bernasconi, E., Calmy, A., Gunthard, H.F., Hoffmann,
827 M., Rauch, A., Telenti, A. & Fellay, J. (2014). Disentangling Human Tolerance and Resistance
828 Against HIV. *PLoS Biol.*, 12.

829

830 Restif, O. & Koella, J.C. (2004). Concurrent Evolution of Resistance and Tolerance to Pathogens.
831 *Am. Nat.*, 164, E90-E102.

832

833 Richardson, C.E., Kooistra, T. & Kim, D.H. (2010). An essential role for XBP-1 in hos protection
834 against immune activation in *C. elegans*. *Nature*, 463, 1092-1095.

835

836 Roy, B.A., Kirchner, J.W., (2000). Evolutionary dynamics of pathogen resistance and tolerance.
837 *Evolution*, 54, 51-63.

838

839 Roy, B.A., Kirchner, J.W., Christian C.E. & Rose, L.E. (2000). High disease incidence and apparent
840 disease tolerance in a North American Great Basin plant community. *Evolutionary Ecology*, 14: 421-
841 438

842

843 Schafer, J.F. (1971). Tolerance to Plant Disease. *Annu. Rev. Phytopathol.*, 9, 235-252.

844

845 Schindelin, J., Rueden, C.T., Hiner, M.C. & Eliceiri, K.W. (2015). The ImageJ Ecosystem: An Open
846 Platform for Biomedical Image Analysis. *Mol. Reprod. Dev.*, 529, 518-529.

847

848 Seck, M., Roelfs, A.P. & Teng, P.S. (1991). Influence of leaf position on yield loss caused by wheat
849 leaf rust in single tillers. *Crop Protection*, 10, 222-228.

850

851 Shaw, M.W. & Royle, D. (1989a). Airborne inoculum as a major source of *Septoria tritici*
852 (*Mycosphaerella graminicola*) infections in winter wheat crops in the UK. *Plant Pathology*, 38, 35-
853 43.

854

855 Shaw, M.W. & Royle, D.J. (1989b). Estimation and validation of a function describing the rate at

856 which *Mycosphaerella graminicola* causes yield loss in winter wheat. *Ann. Appl. Biol.*, 115, 425-442.
857

858 Shinzawa, N., Nelson, B., Aonuma, H., Okado, K., Fukumoto, S., Miura, M. & Kanuka, H. (2009).
859 p38 MAPK-Dependent Phagocytic Encapsulation Confers Infection Tolerance in *Drosophila*. *Cell*
860 *Host Microbe*, 6, 244-252.
861

862 Shuckla, A., Pagán, I. & García-Arenal, F. (2017). Effective tolerance based on resource reallocation
863 is a virus-specific defence in *Arabidopsis thaliana*. *Mol. Plant Pathol.*, 19, 1454-1465
864

865 Simms, E. & Triplett, J. (1994). Costs and Benefits of Plant Responses to Disease: Resistance and
866 Tolerance. *Evolution*, 48, 1973-1985.
867

868 Soares, M.P., Teixeira, L. & Moita, L.F. (2017). Disease tolerance and immunity in host protection
869 against infection. *Nat. Rev. Immunol.*, 17, 83-96.
870

871 Stewart, E.L., Croll, D., Lendenmann, M.H., Sanchez-vallet, A., Hartmann, F.E., Palma-guerrero, J.,
872 Ma, X. & McDonald, B.A. (2016a). QTL mapping reveals complex genetic architecture of
873 quantitative virulence in the wheat pathogen *Zymoseptoria tritici*. *Mol. Plant Pathol.*, 19, 201-216.
874

875 Stewart, E.L., Hagerty, C.H., Mikaberidze, A., Mundt, C., Zhong, Z. & McDonald, B.A. (2016b). An
876 improved method for measuring quantitative resistance to the wheat pathogen *Zymoseptoria tritici*
877 using high throughput automated image analysis. *Phytopathology*, 106, 782-788.
878

879 Stowe, K.A. (1998). Experimental Evolution of Resistance in *Brassica rapa*: Correlated Response of
880 Tolerance in Lines Selected for Glucosinolate Content. *Evolution*, 52, 703-712.
881

882 Teng, P.S. & Gaunt, R.E. (1980). Modelling systems of disease and yield loss in cereals. *Agricultural*
883 *Systems*, 6, 131-154.
884

885 Thomas, M.R., Cook, R.J., & King, J.E. (1989). Factors affecting development of *Septoria tritici* in
886 winter wheat and its effect on yield. *Plant Pathology*, 38, 246-257.

887
888 Waggoner, P. & Berger, R. (1987). Defoliation, disease, and growth. *Phytopathology*, 77, 393-398.
889
890 *Yates, S., *Mikaberidze, A., Krattinger, S.G., Abrouk, M., Hund, A., Yu, K., Studer, B., Fouche, S.,
891 Meile, L., Pereira, D., Karisto, P., McDonald, B.A. (2019). Precision phenotyping reveals novel loci
892 for quantitative resistance to septoria tritici blotch, *Plant Phenomics*, 3285904. (*equal contributions)
893
894 Zhan, J., Pettway, R. & McDonald, B. (2003). The global genetic structure of the wheat pathogen
895 *Mycosphaerella graminicola* is characterized by high nuclear diversity, low mitochondrial diversity,
896 regular recombination, and gene flow. *Fungal Genetics and Biology*, 38, 286-297.
897
898 Zhan, J., Stefanato, F. & McDonald, B. (2006). Selection for increased cyproconazole tolerance in
899 *Mycosphaerella graminicola* through local adaptation and in response to host. *Mol. Plant Pathol.*, 7,
900 259-268.
901
902 Zeller, M & Koella, J. (2017). The Role of the Environment in the Evolution of Tolerance and
903 Resistance to a Pathogen. *American Naturalist*, 190, 389-397.
904
905 Zuckerman, E., Eshel, A. & Eyal, Z. (1997). Physiological Aspects Related to Tolerance of Spring
906 Wheat Cultivars to Septoria tritici Blotch. *Phytopathology*, 87, 60-65.

Supporting Information

The following Supporting Information is available for this article:

Notes S1. Simple model of leaf tolerance

Notes S2. Two components of tolerance

Figure S1. Detailed statistical analysis of the correlation between tolerance and resistance among tolerant and intolerant cultivars.

Figure S2. Example fits for cultivars with contrasting levels of resistance.

Table S1. Ranking of wheat cultivars according to their tolerance to *Zymoseptoria tritici*.

Notes S1. Simple model of leaf tolerance

Here we derive a simple model of infection of wheat leaves by *Z. tritici*. We assume that each lesion has an area a_l and contains n_p pycnidia. How much will the green leaf area H diminish under a small increase in the number of pycnidia ΔN_p ? To find out, we express a small decrease in the green leaf area ΔH in terms of a small increase in the number of lesions: $\Delta H = -a_l \Delta N_l$. At the same time, the increment in the number of pycnidia ΔN_p is related to the increment in the number of lesions: $\Delta N_p = n_p \Delta N_l$. A simple rearrangement yields the relationship between the increment in the number of pycnidia and the decrease in the green leaf area: $\Delta H = -a_l/n_p \Delta N_p$. By taking the limit $\Delta N_p \rightarrow 0$ and $\Delta H \rightarrow 0$ we obtain the differential equation:

$$\frac{dH}{dN_p} = -a_p, \quad (4)$$

where $a_p = a_l/n_p$ represents the area of the lesion that corresponds to a single pycnidium.

Assumption 1.

We first assume that a_p depends neither on H , nor on N_p . In this case, the solution of Eq. (4) is a linear function that can be written as:

$$H(N_p) = H_0(1 - \kappa N_p), \quad (5)$$

where $\kappa = a_p H_0$ and H_0 is the green leaf area in the absence of infection.

Assumption 2.

Here, we relax Assumption 1 and consider the case when the area a_p is proportional to the green leaf area H , i. e.

$$a_p = \kappa H. \quad (6)$$

In this case, the solution of Eq. (4) is an exponential function that can be expressed as:

$$H(N_p) = H_0 \exp(-\kappa N_p). \quad (7)$$

Note, that when κN_p is small, we can expand the function $H(N_p)$ Eq. (7) in the Taylor's series with respect to its argument κN_p . Retaining the zeroth and the first terms of the series yields the linear dependency Eq. (5).

Notes S2. Two components of tolerance

Here we demonstrate mathematically that the novel component of tolerance of wheat to STB that we measured in this experiment and the tolerance components that were measured previously contribute to overall tolerance as multiplicative factors. We use the critical point model (King et al., 1983; Shaw & Royle, 1989a) to relate the grain yield to the loss in the green leaf area in the three upper-most leaves:

$$Y = Y_0 - \sum_{i=1}^3 \kappa_{0i} I_i, \quad (8)$$

where Y is the yield in the presence of disease (measured for example in tons per hectare), Y_0 is the yield in the absence of disease and I_i is the area of the i th leaf ($i=1$ for flag leaf, $i=2$ for flag-1 leaf, $i=3$ for flag-2 leaf) that became chlorotic or necrotic as a result of infection, measured during the critical stage of grain development (around GS 75). The slope κ_{0i} represents the intolerance parameter that may have different values in different leaf layers. We call this component of tolerance "whole-plant tolerance", because it includes various mechanisms of compensation for the leaf damage at the level of the whole plant.

In this study, we argue that the number of pycnidia per leaf, N_{pi} , corresponding to the i th

leaf layer allows for a more accurate quantification of the pathogen population than the area of the leaf I_i damaged due to disease. Similarly to Eq. (8), this leads to the following relationship between the yield and the number of pycnidia per leaf:

$$Y = Y_0 - \sum_{i=1}^3 \kappa_{toti} N_{pi}, \quad (9)$$

where κ_{toti} is the total intolerance parameter and as in Eq. (8), we perform a summation over the three upper-most leaf layers. Further, we assume for simplicity that the linear model Eq. (2) describes well the relationship between the green leaf area, H_i , and the number of pycnidia per leaf N_{pi} . The damaged leaf area, I_i , can then be determined from Eq. (2), but for each individual leaf layer i , according to

$$I_i = H_{0i} - H_i = a_{pi} N_{pi}, \quad (10)$$

where a_{pi} is the damaged leaf area that corresponds on average to a single pycnidium, which quantifies the degree of tolerance on the scale of individual leaves that we measured in this study for the second leaf (i.e., flag-1 leaf). Here we used the linear approximation of the dependency of the green leaf area on the number of pycnidia, which works well when the number of pycnidia per leaf is sufficiently low. In this case, a_{pi} is related to the intolerance parameter, κ_i , that we measured here through H_{0i} , the green leaf area in the absence of disease: $a_{pi} = H_{0i} \kappa_i$. To establish the relationship between the overall tolerance quantified by κ_{tot} , whole-plant tolerance quantified by κ_0 and leaf tolerance quantified by κ , we substitute I from Eq. (10) in Eq. (8)

$$Y = Y_0 - \sum_{i=1}^3 \kappa_{0i} a_{pi} N_{pi}, \quad (11)$$

Comparison of Eq. (11) and Eq. (9) reveals that for each leaf layer i

$$\kappa_{toti} = \kappa_{0i} a_{pi}, \quad (12)$$

the total intolerance parameter is a product of the whole-plant and leaf intolerance parameters. If the relationships between the yield, Y , and the damaged leaf area, I_i , and between the yield, Y , and the number of pycnidia per leaf, N_{pi} , are not linear, then the yield would still be a decreasing, but a nonlinear function of the number of pycnidia per leaf, and the multiplicative relation between the total tolerance and the two components of intolerance in Eq. (12) will not be retained.

989

991 **Figure S1** Detailed statistical analysis of the correlation between tolerance and resistance among
 992 tolerant and intolerant cultivars. The analysis is based on creating a large number of bootstrap
 993 samples ($n_{bs}=1'000'000$) on the basis of the estimates of tolerance and resistance in 335 wheat
 994 cultivars shown in Fig. 3 of the main text. The histogram shows the distributions of the values of the
 995 Spearman's correlation coefficient, r_s , among tolerant (blue) and intolerant (red) cultivars. These
 996 distributions approximate the underlying probability distributions. Solid vertical lines show the
 997 point-estimates of r_s among tolerant (blue) and intolerant (red) cultivars. Dotted vertical lines show
 998 the 95 % confidence intervals of the corresponding point estimates.

Figure S2 Example fits for cultivars with contrasting levels of resistance. Cultivar Element (estimate of resistance $N_p = 66$ estimate of tolerance $\kappa = 0.0033$,) is more resistant than cultivar Arack (estimate of resistance $N_p = 412$, estimate of tolerance $\kappa = 0.0014$). However, in terms of tolerance the two cultivars are not significantly different from the baseline (black curve).