Global drivers of diversification in a marine species complex

Catarina N.S. Silva¹, Nicholas P. Murphy², James J. Bell³, Bridget S. Green⁴, Guy Duhamel⁵, Andrew C. Cockcroft⁶, Cristián E. Hernández⁷, Jan M. Strugnell^{1, 2}

¹ Centre of Sustainable Tropical Fisheries and Aquaculture, James Cook University, Townsville, Qld 4810, Australia

² Department of Ecology, Environment & Evolution, La Trobe University, Melbourne, Vic 3086, Australia

³ School of Biological Sciences, Victoria University of Wellington, Wellington, 6140, New Zealand

⁴ Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS 7001, Australia

⁵ Département Adaptations du Vivant, BOREA, MNHN, Paris, 75005, France

⁶ Department of Agriculture, Forestry and Fisheries, Cape Town, 8012, South Africa

⁷ Departamento de Zoología, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Casilla 160C, Concepción, Chile

Abstract

Investigating historical gene flow in species complexes can indicate how environmental and reproductive barriers shape genome divergence before speciation. The processes influencing species diversification under environmental change remain one of the central focal points of evolutionary biology, particularly for marine organisms with high dispersal potential. We investigated genome-wide divergence, introgression patterns and inferred demographic history between species pairs of all extant rock lobster species (*Jasus* spp.), a complex with long larval duration, that has populated continental shelf and seamount habitats around the globe at approximately 40°S. Genetic differentiation patterns revealed the effects of the environment and geographic isolation. Species associated with the same habitat structure (either continental shelf or seamount/island) shared a common ancestry, even though the habitats were not adjacent. Differences in benthic temperature explained a significant proportion (41.3%) of the genetic differentiation. The Eastern Pacific species pair of *J. caveorum* and *J. frontalis* retained a signal of strict isolation following ancient migration, whereas species pairs from Australia and Africa and seamounts in the Indian and Atlantic oceans included events of introgression after secondary contact. Parameters estimated for time in isolation and gene flow were congruent with genetic differentiation metrics suggesting that the observed differentiation patterns are the product of migration and genetic drift. Our results reveal important effects of habitat and demographic processes on the divergence of species within the genus *Jasus* providing the first empirical study of genome-wide drivers of diversification that incorporates all extant species in a marine genus with long pelagic larval duration.

Introduction

The discrete categorization of speciation modes as sympatric, allopatric or parapatric is now considered to be overly simplistic (Butlin *et al.* 2008). Several events (or modes of speciation) can influence the biogeographic states of populations at different time periods during divergence, and as a result, the speciation process is now generally considered to be gradual and reticulated (Smadja & Butlin 2011; Feder *et al.* 2012). However, the processes responsible for influencing species diversification are still poorly understood and remain one of the central focal points of ecology and evolutionary biology (Arendt *et al.* 2016).

Reconstructing the diversification history for species complexes can be challenging for non-model marine organisms (e.g. Palero *et al.* 2009; Momigliano *et al.* 2017) as many have large population sizes and the potential for long distance dispersal, which frequently result in weak genetic differentiation (Ovenden 2013). For these species, it is often difficult to determine whether weak genetic differentiation is actually present or masked by the large population sizes (Lowe & Allendorf 2010). In addition, marine species with long distance dispersal can quickly fill available niches, leaving fewer opportunities for *in situ* cladogenesis (Pinheiro *et al.* 2017). As a result, only a few studies have reconstructed the history of diversification in marine species (e.g. Crow *et al.* 2010; Le Moan *et al.* 2016; Momigliano *et al.* 2017; Souissi *et al.* 2018; Titus *et al.* 2019).

Changes in the distribution of marine species resulting from historical climatic variation have been an important driver of diversification across taxa (Davis *et al.* 2016). Climatic fluctuations during the late Pleistocene, in particular, resulted in periods of isolation intercalated by contact and gene flow between lineages (Hewitt 2000). These events dramatically transformed available habitat causing major shifts in species distribution ranges and shaping the genetic structure of many marine species worldwide (Benardine Jeffrey *et al.* 2007; Kenchington *et al.* 2009; Van Oppen *et al.* 2011; Strugnell *et al.* 2012). Sequential glacial and interglacial periods have then further shaped the divergence history of species as a result of periods of isolation intercalated by gene flow (Weigelt *et al.* 2016). A better understanding of the species-specific historical context of divergence is therefore needed to estimate the actual timing and role of gene flow during speciation. Understanding how historical climatic fluctuations shaped species divergence provides clues on how species might respond to future environmental changes, which is vital for effective conservation and management plans (Olivieri *et al.* 2016).

Advances in next-generation sequencing (NGS) now provide the opportunity to investigate genome-wide patterns of differentiation along the speciation continuum, allowing the better detection of changes as two lineages diverge from one another on the path to reproductive isolation (Feder *et al.* 2012). In particular, these methods provide effective tools solutions for species with no reference genomes (Catchen *et al.* 2017), which is the case for many marine species including rock lobsters. This technology has also allowed the

integration of genomic and environmental data which can be used for testing the hypothesis that selection is more efficient than drift in opposing the homogenizing effects of migration (Manel & Holderegger 2013). In addition, this approach can also detect candidate markers underlying adaptation to local environments for species with moderate to long distance dispersal potential (e.g. Benestan *et al.* 2016; Sandoval-Castillo *et al.* 2018). This robust approach is particularly useful in the marine environment where isolation and speciation is increasingly found to be associated with selection/local adaptation (Rocha *et al.* 2005; Momigliano *et al.* 2017). Improvements in methodology have further enabled the use of genome-wide polymorphism data to infer complex demographic histories and the relative influence of gene flow and historical processes on the genomic landscape. For example, in the marine environment this approach has been used in the European anchovy *Engraulis encrasicolus* (Le Moan *et al.* 2016), the Atlantic Salmon *Salmo salar* (Rougemont & Bernatchez 2018) and the corkscrew sea anemone *Bartholomea annulate* (Titus *et al.* 2019). One increasingly popular approach is demographic inference based on the computation of a joint allele frequency spectrum (JAFS) from genetic polymorphism data (Gutenkunst *et al.* 2009; Excoffier *et al.* 2013). This approach allows an estimation of several demographic parameters such as population sizes, migration rates and time intervals since specific events using a composite likelihood. Therefore, the role of historical events in the diversification and speciation of marine species can now be more accurately determined.

Rock lobsters (*Jasus* spp.) are a useful model to study the role of historical climatic variations and gene flow on divergence. The six extant *Jasus* lobster species (*J. caveorum*, *J. edwardsii*, *J. frontalis*, *J. lalandii*, *J. paulensis* and *J. tristani*) are distributed in a narrow latitudinal band (~25° to 47°; Fig. 1) in the Southern Hemisphere (Booth 2006) up to 200 m (Holthuis 1991) and possibly up to 600 m depth (Duhamel personal communication). These animals have a long pelagic larval duration (PLD; up to two years for *J. edwardsii*), with the potential for extensive dispersal (Bradford *et al.* 2015). Despite such a long PLD, all species have a restricted latitudinal distribution; for example, *J. caveorum* is only known from a single seamount in the eastern South Pacific Ocean (Webber & Booth 1995). Phylogenetic relationships between *Jasus* species have been investigated with a limited number of mtDNA markers (Brasher *et al.* 1992; Ovenden *et al.* 1997). Ovenden *et al.* (1997) identified a clade containing *J. edwardsii*, *J. lalandii* and *J. frontalis*, however, the relative branching order was not resolved by analysis of sequence variation in the cytochrome c oxidase subunit I (COI) and the 16S ribosomal RNA sequences. In addition, the species *J. tristani* and *J. paulensis*, which occur in islands and seamounts off the southern Atlantic and Indian Oceans, respectively, were hypothesized to have come into secondary contact during past glacial periods, resulting in low levels of mtDNA differentiation (Ovenden *et al.* 1997; Groeneveld *et al.* 2012). At the species level, population genetic studies have demonstrated a general pattern of low, yet often significant, differentiation (Ovenden *et al.* 1992; Matthee *et al.* 2007; Porobić *et al.* 2013; Thomas & Bell 2013; Villacorta-Rath *et al.* 2016). Post-settlement selection and chaotic genetic patchiness, also described as a shifting, ephemeral genetic pattern, has also been observed in *J. edwardsii*,

highlighting the uncertainties in predicting connectivity between populations of highly dispersive marine organisms (Villacorta-Rath *et al.* 2018).

Although a few studies suggest a recent divergence between *Jasus* lineages (Pollock 1990; Ovenden *et al.* 1997), relatively little attention has focused on investigating diversification processes in *Jasus* lobsters. Here we investigate speciation processes in all the extant lobster species of the genus *Jasus*. This study aims to test for evidence of admixture/introgression between species, investigating the genetic patterns associated with habitat structure (continental shelf or seamount/island) and inferring the demographic history of *Jasus* spp. using genome-wide single nucleotide polymorphisms (SNP).

Methods

Sampling, DNA extractions and sequencing

Tissue samples of *Jasus* spp. were collected between 1995 and 2017 from 17 locations throughout the entire genus' range (Fig. 1). A total of 375 samples were collected (11 *Jasus caveorum*, 53 *J. frontalis*, 41 *J. tristani*, 129 *J. lalandii*, 49 *J. paulensis* and 92 *J. edwardsii*). Tissue was stored in 70% ethanol before processing. Total genomic DNA of *J. caveorum* museum samples was extracted using the QIAamp DNA Micro Kit (Qiagen) according to the manufacturer's instruction. The remaining tissue samples were extracted using NucleoMag® Tissue (Macherey-Nagel) following to the manufacturer's instructions.

Library preparation and sequencing was conducted by Diversity Arrays Technology, Canberra, Australia and followed standard protocols of DArTseq™ genotyping technology (Kilian *et al.* 2012). Briefly, approximately 100 ng (2 µL) of each sample was digested with the restriction enzymes PstI and SphI, and unique barcode sequences simultaneously ligated onto the ends of each resulting fragment as per Kilian et al. 2012. The PstI-compatible adapter included an Illumina flow-cell attachment sequence, a primer sequence and unique barcode, with the reverse SphI-compatible adaptor contained in the flow-cell attachment region. A minimum of 15% random technical replicates were included for downstream quality control. Each sample with fragments containing both PstI and SphI cut sites was amplified in PCR reactions using the following conditions: 94°C for 1 min then 30 cycles of 94 °C for 20 s, 58 °C for 30 s, 72 °C for 45 s, and 72 °C for 7 min. Samples were checked visually on an agarose gel to ensure complete digestion and uniform range of fragment sizes. Using approximately 10 µL of each sample, samples were sequenced on a single flow-cell lane on the Illumina HiSeq2500 for 77 cycles.

De novo assembly and variant calling

Libraries were demultiplexed and reads were filtered for overall quality (`-c`, `-q` and `-r` options) using *process_radtags* in STACKS v.2.0b9 (Catchen *et al.* 2013). The Stacks pipeline *denovo_map.pl* was executed to run each of the Stacks modules individually (*ustacks*, *cstacks*, *sstacks* and *populations*). The formation of loci was allowed with a maximum of two nucleotides between stacks ($M = 2$) and a minimum stack depth of three ($m = 3$) among reads for accurate calling (*ustacks* module). Reads were aligned *de novo* with each other, and a catalogue of putative RAD tags was created (*cstacks* module). Putative loci were searched against the catalog (*sstacks* module) and further filtering was then conducted in the *populations* module.

Retained reads were present in at least 70% of samples within each species, were detected in all species, had a rare allele frequency of at least 2% and had no more than 2 alleles detected. Potential paralogs were excluded by removing markers with heterozygosity > 0.50 within samples and analyses were restricted to one random SNP per locus (using `--write_random_snp`). These filtering steps aimed to exclude as many SNPs as was possible with genotyping errors and missing data.

Genetic diversity and population structure

Allelic richness, pairwise F_{ST} values and respective p-values were estimated using *hierfstat* package in R (Goudet 2005). The R package *adeigenet* was used to estimate observed and expected heterozygosity, inbreeding coefficients and for discriminant analyses of principal components (DAPC) and membership probability plots (Jombart 2008). Outlier analyses were conducted in BayeScan to look for signatures of selection. Prior odds were set to 100 to minimize chances of false positives with 5,000 pilot runs, followed by 100,000 iterations (5,000 samples, a thinning interval of 10, and a burn-in of 50,000).

Environmental data collection and analyses

Initially, 13 environmental variables were obtained from Bio-Oracle (Assis *et al.* 2018; Table S1). Only uncorrelated variables ($r < 0.6$) were retained resulting in seven layers (surface and benthic temperature mean, surface salinity, surface and benthic current velocity, benthic iron and surface phytoplankton). Multiple regression of distance matrices (MRDM; Legendre *et al.*, 2014) was used to examine the association of geographic distance (estimated as the shortest path distance in the ocean) with patterns of genetic differentiation (measured as pairwise F_{ST} values), using the R package *ecodist* (Goslee & Urban 2007). Redundancy analysis (RDA; Forester *et al.* 2018) was used to investigate genotype-environment associations

using the R package *vegan* (Oksanen *et al.* 2019). Significance was assessed using a permutation test (999 permutations) for redundancy analysis using the function *anova.cca()*.

Relationships among lineages

The program TREEMIX v1.12 (Pickrell & Pritchard 2012) was used to further investigate historical relationships among lineages. A maximum-likelihood (ML) phylogeny was first inferred and then single migration events between branches were sequentially added to determine whether migration/admixture events improve the likelihood fit. To formally test for admixture between *Jasus* spp., the three-population test (Reich *et al.* 2009) included with TREEMIX was used. In this test, the $f_3(X; A, B)$ statistic is negative when a population X does not form a simple tree with populations A and B, but rather may be a mixture of A and B.

Demographic modelling

Previous analysis suggests evidence of admixture between species pairs, and so we tested several hypothesis of divergence modes, aiming to identify speciation events through time, for each closely related pair of species: *J. caveorum* - *J. frontalis*, *J. edwardsii* - *J. lalandii* and *J. tristani* - *J. paulensis*. The species pairs were selected based on their genetic and morphological relationships (Holthuis & Sivertsen 1967; George & Kensler 1970; Brasher *et al.* 1992; Ovenden *et al.* 1997; Groeneveld *et al.* 2012; this study). For each pair, six models were built representing alternative modes of divergence considering possible scenarios: (SI) Strict Isolation, where the environment (e.g. sea level change and ocean currents) promoted allopatry; (IM) Isolation-with-Migration, with continuous gene flow through the speciation process; (AM) Ancient Migration, with an ancient gene flow event but recent isolation; (SC) Secondary Contact, with a recent gene flow event after past isolation; (PAM) Ancient Migration with two periods of gene flow, with two ancient gene flow events but recent isolation; and (PSC) Secondary Contact with two periods of contact, two recent gene flow events after past isolation. Six additional models were built to include expansion/contraction events in the initial models (suffix 'ex'). All models were implemented allowing for asymmetric migration rates (m12, m21).

Demographic inference was performed using the diffusion approximation method implemented in the software *daði* (Gutenkunst *et al.* 2009). The function *vcf2dadi* in the R package *radiator* (Gosselin 2017) was used to create *daði* SNP input files. We used the folded joint site frequency spectrum (JSFS) for model selection because the closest out-group (*Sagmariasus verreauxi*) was too distant (diverged around 11 Mya; Ovenden *et al.* 1997), which resulted in a highly reduced number of orientable polymorphisms.

In total, 12 models were tested per species pair, fitted with the observed joint site frequency spectrum (SFS) using 20 replicate runs per model and the best model was retained (Fig. S4). The Akaike information criterion (AIC) was used to perform comparisons among models (Sakamoto *et al.* 1986).

To compare among nested models of increasing complexity and address over-parametrization issues we used the comparative framework of Tine *et al.* (2014) by penalizing models which contain more parameters. For each species pair, a score was estimated for each model using:

$$\text{Score} = \frac{(\Delta_{\max} - \Delta \text{AIC}_i)}{\Delta_{\max}} \quad (1)$$

where, $\Delta_{\max}$ corresponds to the difference in AIC between the worst and the best performing model ($\Delta_{\max} = \text{AIC}_{\max} - \text{AIC}_{\min}$) and $\Delta \text{AIC}_i = \text{AIC}_i - \text{AIC}_{\min}$. Therefore, the worst model has a score of 0 and the best model has a score of 1. To evaluate the relative probabilities of the different models within each species pair, Akaike weights (W_{AIC}) were also calculated following:

$$W_{\text{AIC}} = \frac{e^{\frac{-(\Delta \text{AIC}_i)}{2}}}{\sum_{i=1}^R e^{\frac{-(\Delta \text{AIC}_i)}{2}}} \quad (2)$$

where R corresponds to the total number of models considered (R=12).

Demographic parameters were converted into indicative biologically units, given the missing crucial information about mutation rate per generation in *Jasus* spp. The ancestral effective population size (N_{ref}) before split for each species pair was estimated following:

$$N_{\text{ref}} = \frac{\theta}{4L\mu} \quad (3)$$

with θ being the optimal multiplicative scaling factor, μ the mutation rate (fixed at 8×10^{-8} mutations per site per generation; Obbard *et al.* 2012) and L the effective length of the genome explored:

$$L = \frac{zy73}{x} \quad (4)$$

where x is the number of SNPs originally detected from y RAD-tags of 73 bp present in the initial data set, and z the number of SNPs retained, following Rougeux *et al.* (2017). Estimated units in $2N_{\text{ref}}$ were converted to years assuming a generation time of 10 years (Pech *et al.* 2009). Estimated migration rates were divided by $2N_{\text{ref}}$ to obtain the proportion of migrants in every generation.

Results

Genetic diversity and population structure

Sequencing yielded a total of 1,501,921,855 quality-trimmed sequencing reads, providing an average depth of coverage per individual over all SNPs of 58.9x. After applying the different filtering steps, 2,596 SNPs common to all species were retained for subsequent analyses. The lowest values of observed heterozygosity, expected heterozygosity, and allelic richness were observed for *J. caveorum*. While *J. frontalis* had the highest inbreeding coefficients (0.502), all species had very similar values. The highest values of allelic richness were observed for *J. lalandii* (Table 1). The highest pairwise F_{ST} values were observed for *J. tristani* – *J. caveorum* and *J. paulensis* – *J. caveorum* ($F_{ST} = 0.463$ and $F_{ST} = 0.436$, respectively, $p < 0.05$), while the lowest values were observed for *J. tristani* – *J. paulensis* ($F_{ST} = 0.022$, $p < 0.01$; Table 2).

Table 1 Summary statistics of genetic diversity per species using 2,596 SNPs. H_O : observed heterozygosity, H_E : expected heterozygosity, F_{IS} : inbreeding coefficient, A_R : allelic richness

Species	Sample size	H_O	H_E	F_{IS}	A_R
<i>J. caveorum</i>	11	0.012	0.012	0.499	1.04
<i>J. frontalis</i>	53	0.064	0.065	0.502	1.32
<i>J. tristani</i>	41	0.092	0.104	0.500	1.31
<i>J. lalandii</i>	129	0.086	0.104	0.499	1.60
<i>J. paulensis</i>	49	0.087	0.103	0.498	1.31
<i>J. edwardsii</i>	92	0.084	0.100	0.501	1.34
	375				

Table 2 Pairwise F_{ST} values (below diagonal) and corresponding p-values (above diagonal) estimated using *hierfstat* package in R.

	<i>J. caveorum</i>	<i>J. frontalis</i>	<i>J. tristani</i>	<i>J. lalandii</i>	<i>J. paulensis</i>	<i>J. edwardsii</i>
<i>J. caveorum</i>	0	0.010	0.013	0.010	0.011	0.012
<i>J. frontalis</i>	0.081	0	0.007	0.005	0.006	0.007
<i>J. tristani</i>	0.463	0.206	0	0.007	0.008	0.009
<i>J. lalandii</i>	0.305	0.137	0.387	0	0.006	0.007
<i>J. paulensis</i>	0.436	0.229	0.022	0.408	0	0.007
<i>J. edwardsii</i>	0.413	0.106	0.441	0.230	0.452	0

No signatures of selection were detected by the outlier detection analyses (Fig. S1). Lobster species were grouped into three main clusters by discriminant analyses of principal components when using 2 PCs (52.3% variation) (Fig. 2). There was evidence of admixture, in particular between *J. paulensis* - *J. tristani* in the membership probability plot and the DAPC results, and pairwise F_{ST} values (Fig. 1 and 2). The first DAPC axis (LD1) explained 29.9% of the variation and highlighted the divergence between habitat structure (i.e. *J. edwardsii* and *J. lalandii* vs. remaining species; Fig. S2a), while the second DAPC axis (LD2), which explained 22.4% of the variation, showed three main clusters and highlighted the differences between *J. paulensis* and *J. tristani* and the remaining species (Fig. S2b).

Fig. 1. Sample locations, approximate distribution range of *Jasus* spp. (adapted from Booth, 2006) and membership probability plot using 2 principal components.

Fig. 2. Discriminant analyses of principal components (DAPC) of *Jasus* spp. using 2 principal components (explaining 52.3% variation).

Genotype-environment associations

Geographic distance explained 23.3% of the total genetic variation (F_{ST}) between rock lobster species ($p < 0.01$), using multiple regression of distance matrices (MRDM). The model with all seven environmental variables explained 52.9% of the total genetic variation between lobster species ($p < 0.001$), while the model with the four most important environmental variables explained 51.4% of the total genetic variation ($p < 0.001$). Benthic mean temperature was the single environmental variable that explained most of the genetic differentiation while controlling for the effects of geographic distance (41.3%; $p < 0.001$; Table 3).

All seven environmental variables explained 18% of the variation in rock lobster species ($p < 0.001$) when using the constrained ordination in RDA analyses. All values of the variance inflation factors were below five, indicating that multicollinearity among the predictor variables is not inflating the model. The first five constrained axes were significant in explaining the genetic variation between species (each explaining 53.3%, 25.3%, 12%, 5.2% and 2.2%, respectively; $p < 0.001$; Fig. 3). Genetic variation of *J. caveorum* and *J. frontalis* was associated with higher surface temperature, while *J. paulensis* and *J. tristani* were associated with lower surface temperature. *J. edwardsii* and *J. lalandii* were associated with higher benthic temperature, benthic current velocity and benthic iron. Finally, *J. lalandii* was associated with higher surface phytoplankton, while *J. edwardsii* was associated with higher surface current velocity (Fig. 3).

Table 3 Summary of the models for the multiple regression of distance matrices (MRDM) analyses using F_{ST} as a measure of genetic differentiation. GeoDist: geographic distance (km); SurfTemp and BenTemp: Surface and benthic temperature ($^{\circ}\text{C}$); SurfSal: Surface salinity (PSS); SurfCurren and BenCurren: Surface and benthic current velocity (m.s^{-1}); BenIron: Benthic dissolved iron (mmol.m^{-3}); SurfPhyto: surface phytoplankton (mmol.m^{-3}).

Factors	r^2	p-value
GeoDist	0.233	<0.01
GeoDist + SurfTemp	0.235	<0.001
GeoDist + BenTemp	0.413	<0.001
GeoDist + SurfSal	0.233	<0.001
GeoDist + SurfCurren	0.238	<0.01
GeoDist + BenCurren	0.298	<0.001
GeoDist + BenIron	0.237	<0.001
GeoDist + SurfPhyto	0.290	<0.001
GeoDist + SurfTemp + BenTemp + SurfSal + SurfCurren + BenCurren + BenIron + SurfPhyto	0.529	<0.001
GeoDist + SurfTemp + BenTemp + BenCurren + SurfPhyto	0.514	<0.001

Fig. 3. Ordination plot of redundancy analysis (RDA) of *Jasus* spp. The vectors are the environmental predictors (see Table 3 for a detailed description).

Relationships among lineages

Results from TREEMIX identified three ancestral events of admixture (Fig. S3). However, from the three-population test of admixture, only two f_3 values were negative with associated Z-scores < -0.6 , indicating evidence that *J. tristani* does not form a simple tree with *J. paulensis*, *J. lalandii* and *J. edwardsii*, but rather may be a mixture of these (Table S2, Supporting information). Therefore, the three-population test supported the ancestral event of admixture detected by TREEMIX from the most recent common ancestor (MRCA) of *J. lalandii* and *J. edwardsii* to *J. tristani*. The genetic relationships among species inferred by TREEMIX revealed similar patterns to the genetic differentiation analyses, clearly separating species pairs *J. lalandii*–*J. edwardsii*, *J. paulensis* – *J. tristani* and *J. caveorum* – *J. frontalis* (Fig. S3).

Demographic modelling

In general, SC and PSC models provided better fits to the data with good predictions of the joint site frequency spectrum (JSFS) asymmetry for the *J. paulensis* – *J. tristani* and *J. edwardsii* – *J. lalandii* pairs while AM, PAM and SI had better support for the *J. caveorum* – *J. frontalis* pair (Table S3, Fig. S4). Incorporating population expansion events (suffix 'ex') improved the fit of PSC models for all species pairs but there was not a clear pattern for PAM models. In contrast, the strict isolation (SI) and ancient migration (AM, PAM) models were weakly supported for the *J. paulensis* – *J. tristani* and *J. edwardsii* – *J. lalandii* pairs while the secondary contact models (SC, PSC) were weakly supported for the *J. caveorum* – *J. frontalis* pair (Table S3, Fig. S4).

Asymmetries in gene flow with ratios of m_{21}/m_{12} indicated a stronger migration from population two to population one in all species pairs, and the lower proportion of migrants was observed for the *J. edwardsii* – *J. lalandii* pair (Fig. 4, Table 4). Detailed results for demographic inferences are provided in Table S3, Fig. 4, Table 4, Fig. S4 and Fig. S5 (supporting information).

The best supported model for the *J. paulensis* – *J. tristani* pair was PSCex (Table S3). Within this model, total divergence time between species was approximately $25,826 \pm 6,286$ years ago (Table 4). The period without contact was approximately 5.4 times longer than the period with secondary contact. The best supported model for the *J. edwardsii* – *J. lalandii* pair was PSCex (Table S3). Total divergence time between *J. edwardsii* and *J. lalandii* was approximately $38,460 \pm 12,242$ years ago (Table 4). The period without contact was approximately 38.8 times longer than the period with secondary contact. Finally, the best supported model for the *J. caveorum* – *J. frontalis* pair was PAMex (Table S3). Within this model, total divergence time between species was approximately $28,709 \pm 12,674$ years (Table 4) and the period without contact was approximately 8.1 times longer than the period with ancient migration. Therefore, divergence times with errors overlap across the three species pairs and was estimated to be between 19,540 and 32,112 years for *J. paulensis* – *J. tristani*, 26,218 and 50,702 years for *J. edwardsii* – *J. lalandii* and 16,035 and 41,383 years for *J. caveorum* – *J. frontalis*.

Fig. 4. Representation of the best demographic model for each species pair; *J. caveorum* – *J. frontalis*: ancient migration with two periods of ancient gene flow and recent population contraction (PAMex); *J. lalandii* – *J. edwardsii* and *J. tristani* – *J. paulensis*: secondary contact with two periods of contact and recent population expansion (PSCex). Asymmetric migration rates (m_{21} and m_{12}) are represented by the arrows with higher rates of migration from population two to population one for all species pairs (thicker lines in arrows). Width of the boxes represent sizes of the ancestral population (N_{ref}), population sizes before expansion/contraction (N_1 , N_2) and population sizes after expansion/contraction (N_{1e} , N_{2e}). T_s is the time of divergence in strict isolation, $T_{SC/AM}$ the time of secondary contact or ancient migration and T_e the time of expansion.

Table 4. Parameters estimates for the best model of each species pair with standard deviation. *J. paulensis* – *J. tristani*: secondary contact with two periods of contact and recent population expansion (PSCex); *J. edwardsii* – *J. lalandii*: secondary contact and recent population expansion (SCex); *J. caveorum* – *J. frontalis*: ancient migration with two periods of ancient gene flow and recent population contraction (PAMex).

Species group	1: <i>J. paulensis</i> , 2: <i>J. tristani</i>	1: <i>J. edwardsii</i> , 2: <i>J. lalandii</i>	1: <i>J. caveorum</i> , 2: <i>J. frontalis</i>
Best Model	PSCex	PSCex	PAMex
K	9	9	9
N_{ref}	43.96	85.64	64.19
N₁	3.50 ± 0.84	1.50 ± 0.25	1.30 ± 0.35
N₂	49.67 ± 19.38	41.01 ± 15.61	19.19 ± 9.09
N_{1e}	0.67 ± 0.16	2.44 ± 0.41	0.05 ± 0.24
N_{2e}	2.52 ± 0.53	1.28 ± 0.34	0.20 ± 0.23
m₁₂	11.00 ± 3.77	1.77 ± 0.23	7.71 ± 2.33
m₂₁	2.59 ± 2.22	0.09 ± 0.16	1.99 ± 0.36
T_S	11.94 ± 2.67	10.66 ± 3.37	9.74 ± 4.48
T_{SC/AM}	2.20 ± 0.70	0.27 ± 0.10	1.20 ± 0.19
T_e	0.51 ± 0.20	0.29 ± 0.11	0.24 ± 0.27
T_{total}	29.29 ± 7.15	22.46 ± 7.15	22.36 ± 9.87
*m₁₂	0.12 ± 0.04	0.01 ± 0.001	0.06 ± 0.02
*m₂₁	0.03 ± 0.03	0.00 ± 0.00	0.01 ± 0.00
*T_S	10,524 ± 2,346	18,258 ± 5,769	12,507 ± 5,746
*T_{SC/AM}	1,941 ± 618	470 ± 167	1,541 ± 239
*T_e	447 ± 179	503 ± 185	306 ± 352
*T_{total}	25,826 ± 6,286	38,460 ± 12,242	28,709 ± 12,674

K: The number of free parameters in the model

N_{ref}: The effective size of the ancestral population before the split

N₁: The effective size of population 1 before expansion

N₂: The effective size of population 2 before expansion

N_{1e}: The effective size of population 1 after expansion

N_{2e}: The effective size of population 2 after expansion

m₁₂: The neutral movement of genes from population 2 to population 1 in units of 2N_{ref} generations

m₂₁: The neutral movement of genes from population 1 to population 2 in units of 2N_{ref} generations

T_S: The time of divergence in strict isolation in units of 2N_{ref} generations

T_{SC/AM}: The time of secondary contact/ancient migration in units of 2N_{ref} generations

T_e: The time of expansion in units of 2N_{ref} generations

T_{total}: The total time since the split in units of 2N_{ref} generations

*m₁₂: The proportion of migrants per generation from population 2 to population 1

*m₂₁: The proportion of migrants per generation from population 1 to population 1

*T_S: The time of divergence in strict isolation in units of numbers of years

*T_{SC/AM}: The time of secondary contact/ancient migration in units of numbers of years

*T_e: The time of expansion in units of numbers of years

*T_{total}: The total time since the split in units of numbers of years

Discussion

Here we investigated genome-wide divergence and introgression patterns in all extant species of rock lobsters (*Jasus* spp.) for the first time. Genetic differentiation patterns revealed the effects of the environment and geographical isolation. Species that were associated with the same habitat structure (continental shelf or seamount/island) were more closely related to each other than with species from a different habitat structure. Benthic temperature was the single environmental variable that explained most of the genetic differentiation (F_{ST}) while controlling for the effects of geographic distance (41.3%), and *J. edwardsii* and *J. lalandii* were associated with higher mean benthic temperatures. We also detected multiple introgression events (gene flow) since the first divergence in all species pairs.

Divergence during Pleistocene

Divergence times between species pairs estimated by demographic modelling overlap across all species comparisons and suggest that global/widespread processes might have driven initial divergence across all species. During the divergence period estimated for *Jasus* spp. (from 38,460 to 25,826 years ago) global temperature and sea levels were decreasing (Clark *et al.* 2009). Sea level was 65 to 125 metres lower than today possibly creating more shallow benthic habitat for lobsters in the open ocean in contrast with a reduction in the available continental shelf habitat (Schaaf 1996). A significant increase in the Southern Ocean barotropic stream function occurred between 49,000 and 28,500 years ago (Fogwill *et al.* 2015), which may have increased the dispersal potential of larvae. Also, a large increase in subantarctic productivity occurred 35–65,000 years ago during Heinrich events H4–H6 (Sachs & Anderson 2005), which may have enabled planktotrophic larvae to survive for longer periods of time. These processes may have facilitated worldwide dispersal and colonization of circumpolar habitats by the *Jasus* spp. ancestor during this period (Fig. 5).

During the last glacial maximum (LGM) about 19,000–22,000 years ago, temperature and sea levels reached minimum values (Clark *et al.* 2009). New ecological zones could have appeared, favouring species associated with seamount and island habitats, which could explain the ancient secondary contact events for *J. caveorum* and *J. frontalis* (Schaaf 1996). On the other hand, there was a shrinkage of the shallow continental margins habitat and species therein (Schaaf 1996; Holland 2012). Species associated with continental shelf habitat could also have shifted northwards while tracking their thermal optima. These changes in habitat may have resulted in longer periods of isolation between the continental shelf associated species *J. lalandii* and *J. edwardsii*.

Fig. 5. Changes in global sea level (adapted from Huybrechts 2002) and proportion of available habitat in the photic zone (Schaaf 1996) relative to present day. 1) Antarctic sea ice expands around 28,500 years ago (Fogwill *et al.* 2015); 2) Ocean currents reduction (around 22,000 years ago; Alloway *et al.* 2007); 3) West Antarctic Ice Sheet deglaciation (around 14,500 years ago; Clark *et al.* 2009).

Transition from the LGM to the Holocene precipitated further changes in the available shallow benthic habitat for lobsters with the increase in sea levels and temperature (Clark *et al.* 2009). There were two major expansions of available shallow benthic habitat at 14,000 and 11,500 years (Schaaf 1996) which, for example, could have increased the suitable habitat for *J. frontalis*, having a positive effect on the population size and the effective population size (Porobić *et al.* 2013). However, the variation of the photic sea-bottom area was not linear nor directly correlated with the sea-level oscillations, but reflected topography patterns (Schaaf 1996). These fluctuations in available habitat resulted in alternating periods of isolation and gene flow that have shaped the present genetic signatures of *Jasus* lobsters (Fig. 5).

Are species genetically isolated?

The present study suggests that global processes might have driven initial divergence across all *Jasus* species. However, current genetic signatures highlight the complexities exclusive to each species evolution. For example, *J. edwardsii* and *J. lalandii* were the most genetically differentiated species pair ($F_{ST}=0.230$), followed by *J. caveorum* – *J. frontalis* ($F_{ST}=0.081$) and *J. tristani* – *J. paulensis* ($F_{ST}=0.022$). This was in agreement with the parameters estimated from demographic modelling, in particular the period of gene flow estimated for each species pair (approximately 470 ± 167 years, $1,541 \pm 239$ years and $1,941 \pm 618$ years, respectively).

Our study provides genome-wide evidence of admixture between *J. paulensis* - *J. tristani* that also showed great genetic similarity. Although George & Kensler (1970) have noted that *J. tristani* and *J. paulensis* possess a significant difference in the abdominal sculpture index, genetic evidence suggests that these species can be synonymized as *J. paulensis*, which was also proposed by Groeneveld *et al.* (2012) using the mitochondrial cytochrome oxidase I gene. As a comparison, since initial divergence, *J. tristani* and *J. paulensis* spent 4.1 times longer in secondary contact than *J. edwardsii* – *J. lalandii* and 1.2 times longer than *J. caveorum* and *J. frontalis*. The Tristan da Cunha and Gough Islands (current distribution of *J. tristani*) and the Amsterdam and St. Paul Islands (current distribution of *J. paulensis*) have been grouped in the same zoogeographic province (called the West Wind Drift Islands Province) based on endemic fish fauna distribution (Collette & Parin 1991). Therefore, the distribution of marine species with a pelagic larvae stage may be influenced by the currents of the West Wind Drift and the long periods of gene flow may explain the close relationship between *J. tristani* and *J. paulensis*.

Species associated with continental shelf habitat *J. edwardsii* and *J. lalandii* were genetically more closely related to each other than to *J. tristani* and *J. paulensis* (island/seamount habitat) despite their geographic locations (i.e. *J. lalandii* is geographically closer to *J. tristani* and *J. paulensis* than to *J. edwardsii*; Fig. 1). Although connectivity would be possible between these species groups given the dispersal potential (indeed, *J. lalandii* larvae have been found in the southwest Indian Ocean as far east as Amsterdam Island, adjacent to the *J. paulensis* habitat (Booth & Ovenden 2000)), species appear to be adapted to local environmental conditions. Our results show that benthic temperature might be a limiting factor affecting gene flow between species from island/seamount and continental shelf habitat. Temperature is important for regulating the rate of embryological development in lobsters (Phillips 2013) and could limit reproduction of species adapted to local benthic temperatures.

It has been shown that during the post-larval or puerulus stage, *J. edwardsii* are able to recognize environmental cues such as chemical, acoustic and substrate cues and increase settlement success in suitable habitats (Stanley *et al.* 2015; Hinojosa *et al.* 2016, 2018). Magnetism is also an important cue for adult Western Rock Lobster, *Panulirus cygnus* (Lestang 2014) and adults and postlarvae of spiny lobster *Panulirus argus* (Boles & Lohmann 2003; Kough *et al.* 2014; Ernst & Lohmann 2016). Our results show that genetic diversity of *J. lalandii* and *J. edwardsii* is associated with benthic iron. Therefore, it is possible that *Jasus* larvae are able to use environmental cues such as magnetism for orientation and different species have adapted to recognize local ecotypes and settle on habitats with contrasting structure. Ovenden *et al.* (1997) also suggested that the common ancestor of *J. tristani* and *J. paulensis* may have been able to recognize environmental cues from island and seamount habitats that allowed them to colonize mid-ocean habitats during glacial periods.

In highly dispersive marine taxa, interglacial recolonization of high-latitude habitats can occur rapidly (Hewitt 2000). Such patterns have been established for a range of Northern Hemisphere marine species (e.g. Marko 2004; Ledoux *et al.* 2018), but relatively little is known about the genetic effects of recent glaciations in the Southern Hemisphere (but see e.g. Fraser *et al.* 2009; Strugnell *et al.* 2012; Porobić *et al.* 2013). This study revealed genome-wide patterns of divergence and introgression in all extant species of a highly dispersive marine taxa for the first time. While results point to the important role of demographic and neutral processes of differentiation between species pairs, it also suggests a possible effect of selection in promoting genetic divergence between habitats. Future studies should address the role of adaptive processes to elucidate their relative contribution in shaping genome divergence and speciation of *Jasus* lobsters and to better understand how future environmental change will impact species distribution.

Data availability

Raw demultiplexed sequencing data will be available at Dryad. Pipelines for *de novo* assembly, genetic structure, environmental association and demographic inference analyses will be available at github after publication.

Acknowledgments

Funding for this research was provided by an Australian Research Council Discovery Project grant (Project No. DP150101491) awarded to J.M.S., J.J.B., B.S.G. and N.P.M. We would like to thank Gary Carlos (University of Tasmania), Colin Fry (University of Tasmania), Daniel Ierodiaconou (Deakin University), Kent Way, Andrew Kent, Geoff Liggins, Marcus Miller, Giles Ballinger, Darrel Sykes (DAFF), Rick Webber (Te Papa Museum), Jason How (Department of Fisheries, Western Australia) and Sadie Mills (NIWA) for field assistance and sample collection; T.A.A.F. (Terres Australes et Antarctiques Françaises), for their French fisheries observer service "COPEC", the fishery observer Sophie Cascade on board the F.V. "AUSTRAL" and Charlotte Chazeau to have made available biological scientific samples and data of *Jasus paulensis* from catches in the Saint-Paul/Amsterdam French EEZ; the help of crew has also been appreciated. C.E.H was supported by FONDECYT grants #1170815.

Author contributions

All authors contributed insights about data analysis, interpretation of results and edited the final drafts of the manuscript. C.N.S.S. analysed the data. C.N.S.S., J.M.S and N.P.M. conceived the study.

References

- Alloway B V, Lowe DJ, Barrell DJA, Newnham RM, Almond PC, Al E (2007) Towards a climate event stratigraphy for New Zealand over the past 30 000 years (NZ-INTIMATE project). *Journal of Quaternary Science*, **22**, 9–35.
- Arendt D, King N, Carroll SB, Wittopp P, Koonin E, Kruglyak L (2016) Big Questions in Evolution. *Cell*, **166**, 528–529.
- Assis J, Tyberghein L, Bosch S, Verbruggen H, Serrão EA, De Clerck O (2018) Bio-ORACLE v2.0: Extending marine data layers for bioclimatic modelling. *Global Ecology and Biogeography*, **27**, 277–284.
- Benardine Jeffrey I, Hale P, Degnan BM, Degnan SM (2007) Pleistocene isolation and recent gene flow in *Haliotis asinina*, an Indo-Pacific vetigastropod with limited dispersal capacity. *Molecular Ecology*, **16**, 289–304.
- Benestan L, Quinn BK, Maaroufi H, Laporte M, Clark FK, Greenwood SJ, Rochette R, Bernatchez L (2016) Seascape genomics provides evidence for thermal adaptation and current-mediated population structure in American lobster (*Homarus americanus*). *Molecular Ecology*, **25**, 5073–5092.
- Boles LC, Lohmann KJ (2003) True navigation and magnetic maps in spiny lobsters. *Nature*, **421**, 60–63.
- Booth JD (2006) *Jasus species*. In ‘*Lobsters: Biology, Management, Aquaculture and Fisheries*’. (Ed. B. F. Phillips.). Blackwell Scientific Publications: Oxford.
- Booth JD, Ovenden JR (2000) Distribution of *Jasus* spp. (Decapoda: Palinuridae) phyllosomas in southern waters: Implications for larval recruitment. *Marine Ecology Progress Series*, **200**, 241–255.
- Bradford RW, Griffin D, Bruce BD (2015) Estimating the duration of the pelagic phyllosoma phase of the southern rock lobster, *Jasus edwardsii* (Hutton). *Marine and Freshwater Research*, **66**, 213–219.
- Brasher D, Ovenden J, White R (1992) Mitochondrial DNA variation and phylogenetic relationships of *Jasus* spp. (Decapoda: Palinuridae). *Journal of Zoology*, **227**, 1–16.
- Butlin RK, Galindo J, Grahame JW (2008) Sympatric, parapatric or allopatric: The most important way to classify speciation? *Philosophical Transactions of the Royal Society B: Biological Sciences*, **363**, 2997–3007.
- Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: An analysis tool set for population genomics. *Molecular Ecology*, **22**, 3124–3140.
- Catchen JM, Hohenlohe PA, Bernatchez L, Funk WC, Andrews KR, Allendorf FW (2017) Unbroken: RADseq remains a powerful tool for understanding the genetics of adaptation in natural populations. *Molecular Ecology Resources*, **17**, 362–365.
- Clark PU, Dyke AS, Shakun JD, Carlson AE, Clark J, Wohlfarth B, Mitrovica JX, Hostetler SW, McCabe AM (2009) The Last Glacial Maximum. *Science*, **325**, 710–714.
- Collette BB, Parin N V (1991) Shallow-Water Fishes of Walters Shoals, Madagascar Ridge. *Bulletin of Marine*

516 *Science*, **48**, 1–22.

517 Crow KD, Munehara H, Bernardi G (2010) Sympatric speciation in a genus of marine reef fishes. *Molecular*

518 *Ecology*, **19**, 2089–2105.

519 Davis KE, Hill J, Astrop TI, Wills MA (2016) Global cooling as a driver of diversification in a major marine

520 clade. *Nature Communications*, **7**, 1–8.

521 Ernst DA, Lohmann KJ (2016) Effect of magnetic pulses on Caribbean spiny lobsters: Implications for

522 magnetoreception. *Journal of Experimental Biology*, **219**, 1827–1832.

523 Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M (2013) Robust Demographic Inference from

524 Genomic and SNP Data. *PLoS Genetics*, **9**.

525 Feder JL, Egan SP, Nosil P (2012) The genomics of speciation-with- gene-flow. *Trends in Genetics*, **28**, 342–

526 350.

527 Fogwill CJ, Turney CSM, Hutchinson DK, Taschetto AS, England MH (2015) Obliquity Control On Southern

528 Hemisphere Climate During The Last Glacial. *Scientific Reports*, **5**, 1–10.

529 Forester BR, Lasky JR, Wagner HH, Urban DL (2018) Comparing methods for detecting multilocus adaptation

530 with multivariate genotype-environment associations. *Molecular Ecology*, 2215–2233.

531 Fraser CI, Nikula R, Spencer HG, Waters JM (2009) Kelp genes reveal effects of subantarctic sea ice during

532 the Last Glacial Maximum. *Proceedings of the National Academy of Sciences of the United States of*

533 *America*, **106**, 3249–3253.

534 George RW, Kensler CB (1970) Recognition of marine spiny lobsters of the *Jasus lalandii* group (crustacea:

535 Decapoda: Palinuridae). *New Zealand Journal of Marine and Freshwater Research*, **4**, 292–311.

536 Goslee SC, Urban DL (2007) The ecodist package for dissimilarity-based analysis of ecological data. *Journal Of*

537 *Statistical Software*, **22**, 1–19.

538 Gosselin T (2017) radiator: RADseq Data Exploration, Manipulation and Visualization using R. R package

539 version 0.0.5.

540 Goudet J (2005) HIERFSTAT, a package for r to compute and test hierarchical F-statistics. *Molecular Ecology*

541 *Notes*, **5**, 184–186.

542 Groeneveld JC, Von der Heyden S, Matthee CA (2012) High connectivity and lack of mtDNA differentiation

543 among two previously recognized spiny lobster species in the southern Atlantic and Indian Oceans.

544 *Marine Biology Research*, **8**, 764–770.

545 Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD (2009) Inferring the joint demographic

546 history of multiple populations from multidimensional SNP frequency data. *PLoS Genetics*, **5**.

547 Hewitt G (2000) The genetic legacy of the Quaternary ice ages. *Nature*, **405**, 907–913.

548 Hinojosa I, Gardner C, Green B, Jeffs A (2018) Coastal chemical cues for settlement of the southern rock

549 lobster, *Jasus edwardsii*. *Bulletin of Marine Science*, 1–16.

Hinojosa IA, Green BS, Gardner C, Hesse J, Stanley JA, Jeffs AG (2016) Reef sound as an orientation cue for shoreward migration by pueruli of the rock lobster, *Jasus edwardsii*. *PLoS ONE*, **11**, 1–15.

Holland SM (2012) Sea level change and the area of shallow-marine habitat: implications for marine biodiversity. *Paleobiology*, **38**, 205–217.

Holthuis LB (1991) *Marine lobsters of the world. An annotated and illustrated catalogue of species of interest to fisheries known to date*. FAO Fisheries Synopsis. No. 125, Vol. 13. Rome, FAO.

Holthuis LB, Sivertsen E (1967) *The Crustacea Decapoda, Mysidacea and Cirripedia of the Tristan da Cunha Archipelago, with a revision of the “frontalis” subgroup of the genus Jasus*. Res. Norv. Scient. Exp. *Tristan da Cunha 1937-1938*, **52**, 1-55.

Huybrechts P (2002) Sea-level changes at the LGM from ice-dynamic reconstructions of the Greenland and Antarctic ice sheets during the glacial cycles. *Quaternary Science Reviews*, **21**, 203–231.

Jombart T (2008) Adegnet: A R package for the multivariate analysis of genetic markers. *Bioinformatics*, **24**, 1403–1405.

Kenchington EL, Harding GC, Jones MW, Prodöhl P a (2009) Pleistocene glaciation events shape genetic structure across the range of the American lobster, *Homarus americanus*. *Molecular Ecology*, **18**, 1654–67.

Kilian A, Wenzl P, Huttner E, Carling J, Xia L, Blois H, Caig V, Heller-Uszynska K, Jaccoud D, Hopper C, Aschenbrenner-Kilian M, Evers M, Peng K, Cayla C, Hok P, Uszynski G (2012) Diversity Arrays Technology: A Generic Genome Profiling Technology on Open Platforms. In: *Data Production and Analysis in Population Genomics. Methods in Molecular Biology*, vol 888. (eds Pompanon F, Bonin A), pp. 67–89. Humana Press, Totowa.

Kough AS, Paris CB, Staaterman E (2014) In situ swimming and orientation behavior of spiny lobster (*Panulirus argus*) postlarvae. *Marine Ecology Progress Series*, **504**, 207–219.

Ledoux JB, Frleta-Valić M, Kipson S, Antunes A, Cebrian E, Linares C, Sánchez P, Leblois R, Garrabou J (2018) Postglacial range expansion shaped the spatial genetic structure in a marine habitat-forming species: Implications for conservation plans in the Eastern Adriatic Sea. *Journal of Biogeography*, **45**, 2645–2657.

Legendre P, Lapointe F-J, Casgrain P (1994) Modeling Brain Evolution from Behavior: A Permutational Regression Approach. *Evolution*, **48**, 1487–1499.

Lestang S (2014) The orientation and migratory dynamics of the western rock lobster, *Panulirus cygnus*, in Western Australia. *ICES Journal of Marine Science*, **71**, 1052–1063.

Lowe WH, Allendorf FW (2010) What can genetics tell us about population connectivity? *Molecular ecology*, **19**, 3038–51.

Manel S, Holderegger R (2013) Ten years of landscape genetics. *Trends in Ecology and Evolution*, **28**, 614–

584 621.

585 Marko PB (2004) “What’s larvae got to do with it?” Disparate patterns of post-glacial population structure in
586 two benthic marine gastropods with identical dispersal potential. *Molecular Ecology*, **13**, 597–611.

587 Matthee CA, Cockcroft AC, Gopal K, Von Der Heyden S (2007) Mitochondrial DNA variation of the west-coast
588 rock lobster, *Jasus lalandii*: Marked genetic diversity differences among sampling sites. *Marine and*
589 *Freshwater Research*, **58**, 1130–1135.

590 Le Moan A, Gagnaire PA, Bonhomme F (2016) Parallel genetic divergence among coastal-marine ecotype
591 pairs of European anchovy explained by differential introgression after secondary contact. *Molecular*
592 *Ecology*, **25**, 3187–3202.

593 Momigliano P, Jokinen H, Fraimout A, Florin A-B, Norkko A, Merilä J (2017) Extraordinarily rapid speciation in
594 a marine fish. *Proceedings of the National Academy of Sciences*, 201615109.

595 Obbard DJ, MacLennan J, Kim KW, Rambaut A, O’Grady PM, Jiggins FM (2012) Estimating divergence dates
596 and substitution rates in the drosophila phylogeny. *Molecular Biology and Evolution*, **29**, 3459–3473.

597 Oksanen J, Blanchet F, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O’Hara RB, Simpson GL,
598 Solymos P, Stevens MHH, Szoecs E, Wagner H (2019) vegan: Community Ecology Package. R package
599 version 2.5-6.

600 Olivieri I, Tonnabel J, Ronce O, Mignot A (2016) Why evolution matters for species conservation:
601 Perspectives from three case studies of plant metapopulations. *Evolutionary Applications*, **9**, 196–211.

602 Van Oppen MJH, Peplow LM, Kininmonth S, Berkelmans R (2011) Historical and contemporary factors shape
603 the population genetic structure of the broadcast spawning coral, *Acropora millepora*, on the Great
604 Barrier Reef. *Molecular Ecology*, no-no.

605 Ovenden JR (2013) Crinkles in connectivity: combining genetics and other types of biological data to
606 estimate movement and interbreeding between populations. *Marine and Freshwater Research*, **64**,
607 201.

608 Ovenden JR, Booth JD, Smolenski AJ (1997) Mitochondrial DNA phylogeny of red and green rock lobsters
609 (genus *Jasus*). *Marine and Freshwater Research*, **48**, 1131–1136.

610 Ovenden JR, Brasher DJ, White RWG (1992) Mitochondrial DNA analyses of the red rock lobster *Jasus*
611 *edwardsii* supports an apparent absence of population subdivision throughout Australasia. *Marine*
612 *Biology*, **112**, 319–326.

613 Palero F, Lopes J, Abelló P, Macpherson E, Pascual M, Beaumont MA (2009) Rapid radiation in spiny lobsters
614 (*Palinurus* spp) as revealed by classic and ABC methods using mtDNA and microsatellite data. *BMC*
615 *Evolutionary Biology*, **9**, 263.

616 Pecl G, Frusher S, Gardner C, Haward M, Hobday A, Jennings S, Nursey-Bray M, Punt A, Revill H, van Putten I
617 (2009) *The east coast Tasmanian rock lobster fishery – vulnerability to climate change impacts and*

618 *adaptation response options. Report to the Department of Climate Change, Australia.*

619 Phillips BF (2013) *Lobsters: Biology, Management, Aquaculture and Fisheries* (BF Phillips, Ed.). Department of
620 Environment & Agriculture, Curtin University, Western Australia.

621 Pickrell JK, Pritchard JK (2012) Inference of Population Splits and Mixtures from Genome-Wide Allele
622 Frequency Data. *PLoS Genetics*, **8**, e1002967.

623 Pinheiro HT, Bernardi G, Thorrold, Simon R, Joyeux J-C, Macieira RM, Gasparini JL, Rocha C, Rocha LA (2017)
624 Island biogeography of marine organisms. *Nature*, **549**, 82–85.

625 Pollock DE (1990) Palaeoceanography and Speciation in the Spiny Lobster Genus *Jasus*. *Bulletin of Marine*
626 *Science*, **46**, 387–405.

627 Porobić J, Canales-Aguirre CB, Ernst B, Galleguillos R, Hernández CE (2013) Biogeography and historical
628 demography of the Juan Fernández rock lobster, *Jasus frontalis* (Milne Edwards, 1837). *Journal of*
629 *Heredity*, **104**, 223–233.

630 Reich D, Thangaraj K, Patterson N, Price AL, Singh L (2009) Reconstructing Indian population history. *Nature*,
631 **461**, 489–494.

632 Rocha LA, Robertson DR, Roman J, Bowen BW (2005) Ecological speciation in tropical reef fishes. *Proceedings*
633 *of the Royal Society B: Biological Sciences*, **272**, 573–579.

634 Rougemont Q, Bernatchez L (2018) The demographic history of Atlantic salmon (*Salmo salar*) across its
635 distribution range reconstructed from approximate Bayesian computations*. *Evolution*, **72**, 1261–1277.

636 Rougeux C, Bernatchez L, Gagnaire PA (2017) Modeling the multiple facets of speciation-with-gene-flow
637 toward inferring the divergence history of lake whitefish species pairs (*Coregonus clupeaformis*).
638 *Genome Biology and Evolution*, **9**, 2057–2074.

639 Sachs JP, Anderson RF (2005) Increased productivity in the subantarctic ocean during Heinrich events.
640 *Nature*, **434**.

641 Sakamoto Y, Ishiguro M, Kitagawa G (1986) *Akaike Information Criterion Statistics*. D. Reidel.

642 Sandoval-Castillo J, Robinson NA, Hart AM, Strain LWS, Beheregaray LB (2018) Seascape genomics reveals
643 adaptive divergence in a connected and commercially important mollusc, the greenlip abalone (*Haliotis*
644 *laevigata*), along a longitudinal environmental gradient. *Molecular Ecology*, 1603–1620.

645 Schaaf A (1996) Sea level changes, continental shelf morphology, and global paleoecological constraints in
646 the shallow benthic realm: a theoretical approach. *Palaeogeography, Palaeoclimatology,*
647 *Palaeoecology*, **121**, 259–271.

648 Smadja CM, Butlin RK (2011) A framework for comparing processes of speciation in the presence of gene
649 flow. *Molecular Ecology*, **20**, 5123–5140.

650 Souissi A, Bonhomme F, Manchado M, Bahri-Sfar L, Gagnaire P-A (2018) Genomic and geographic footprints
651 of differential introgression between two divergent fish species (*Solea* spp.). *Heredity*, **121**, 579–593.

652 Stanley JA, Hesse J, Hinojosa IA, Jeffs AG (2015) Inducers of settlement and moulting in post-larval spiny
653 lobster. *Oecologia*, **178**, 685–697.

654 Strugnell JM, Watts PC, Smith PJ, Allcock AL (2012) Persistent genetic signatures of historic climatic events in
655 an Antarctic octopus. *Molecular Ecology*, **21**, 2775–2787.

656 Thomas L, Bell JJ (2013) Testing the consistency of connectivity patterns for a widely dispersing marine
657 species. *Heredity*, **111**, 345–54.

658 Tine M, Kuhl H, Gagnaire P, Louro B, Desmarais E, Martins RST, Hecht J, Knaust F, Belkhir K, Klages S,
659 Dieterich R, Stueber K, Piferrer F, Guinand B, Bierne N, Volckaert FAM, Bargelloni L, Power DM,
660 Bonhomme F *et al.* (2014) European sea bass genome and its variation provide insights into adaptation
661 to euryhalinity and speciation. *Nature Communications*, **5**, 5770.

662 Titus BM, Blischak PD, Daly M (2019) Genomic signatures of sympatric speciation with historical and
663 contemporary gene flow in a tropical anthozoan (Hexacorallia: Actiniaria). *Molecular Ecology*, 3572–
664 3586.

665 Villacorta-Rath C, Ilyushkina I, Strugnell JM, Green BS, Murphy NP, Doyle SR, Hall NE, Robinson AJ, Bell JJ
666 (2016) Outlier SNPs enable food traceability of the southern rock lobster, *Jasus edwardsii*. *Marine*
667 *Biology*, **163**, 1–11.

668 Villacorta-Rath C, Souza CA, Murphy NP, Green BS, Gardner C, Strugnell JM (2018) Temporal genetic patterns
669 of diversity and structure evidence chaotic genetic patchiness in a spiny lobster. *Molecular Ecology*, **27**,
670 54–65.

671 Webber WR, Booth JD (1995) A new species of *Jasus* (Crustacea: Decapoda: Palinuridae) from the eastern
672 South Pacific Ocean. *New Zealand Journal of Marine and Freshwater Research*, **29**, 613–622.

673 Weigelt P, Steinbauer MJ, Cabral JS, Kreft H (2016) Late Quaternary climate change shapes island
674 biodiversity. *Nature*, **532**, 99–102.

675