Proteomics Analysis of Differential Expression of Lung Proteins in Response to Highly Pathogenic Avian Influenza Virus Infection in Chicken

Periyasamy Vijayakumar^{1, #}, Ashwin Ashok Raut¹, Santhalembi Chingtham¹, Harshad V Murugkar², Diwakar D. Kulkarni², Vijendra Pal Singh², Anamika Mishra^{1, *}

¹Pathogenomics Laboratory, ICAR-National Institute of High Security Animal Diseases, OIE Reference lab for Avian Influenza, Bhopal - 462021, Madhya Pradesh, India

² ICAR -National Institute of High Security Animal Diseases, OIE Reference lab for Avian Influenza, Bhopal - 462021, Madhya Pradesh, India

[#] Present affiliation of author: Veterinary College and Research Institute, Tamil Nadu Veterinary and Animal Sciences University, Orathanadu- 614625, Tamil Nadu, India

* corresponding author: reach2anamika@yahoo.com (A.M)

Abstract

Elucidation of molecular pathogenesis underlying virus-host interaction is important for the development of new diagnostic and therapeutic strategies against highly pathogenic avian influenza (HPAI) infection in chicken. However, chicken HPAI viral pathogenesis is not completely understood. To elucidate the intracellular signaling pathways and critical host proteins associated with influenza pathogenesis, we characterized the lung proteome of chicken infected with HPAI H5N1 virus (A/duck/India/02CA10/2011/Agartala). The chicken mass spectra data sets comprised 1,47,451 MS scans and 19,917 MS/MS scans. At local FDR 5% level, we identified total 3313 chicken proteins with presence of at least one unique peptide. At 12 hrs, 247 proteins are downregulated while 1754 proteins are downregulated at 48 hrs indicating that the host has succumbed to infection. There is expression of proteins of the predominant signaling pathways, such as TLR, RLR, NLR and JAK-STAT signaling. Activation of these pathways is associated with cytokine storm effect and thus may be the cause of severity of HPAI H5N1 infection in chicken. Further we identified proteins like MyD88, IKKB, IRAK4, RELA, and MAVS involved in the critical signaling pathways and

some other novel proteins(HNF4A, ELAVL1, FN1, COPS5, CUL1, BRCA1 and FYN) as main hub proteins that might play important roles in influenza pathogenesis in chicken. Taken together, we characterized the signaling pathways and the proteomic determinants responsible for disease pathogenesis in chicken infected with HPAI H5N1 virus.

Key words: Avian influenza virus; proteome; molecular pathways; proteomic determinants; chicken; lung tissues

Introduction

The control measures to any emerging and re-emerging infection, requires an adequate understanding of virus-host interaction (Fauci, 2006). The complete understanding of virus-host interaction will provide important clues for the development of new diagnostic and therapeutic strategies against emerging diseases. The high-throughput functional genomics approach provides a deeper understanding of emerging disease by encompassing both the pathogen and the host response. Virus-host interactions are multidimensional in nature, and this interaction alters various host components such as transcriptome, proteome, miRNA, metabolome, and lipidome (Mishra et al., 2017). In past years, most of the high-throughput studies on H5N1-host interactions have focused at the transcriptome level.

However, transcriptomic studies do not reveal any posttranscriptional regulation, posttranslational modifications, and protein–protein interactions (Josset et al., 2013). The virus-host interaction involves the alternation of hundreds to thousands of host proteins. Hence, applying high throughput proteomics approach for the analysis of virus-induced innate immune responses in conjunction with the transcriptomic analyses promises better understanding of molecular mechanism of influenza pathogenesis (Zak et al., 2014). Proteomics refers to the large-scale study of protein expression, protein-protein interactions or posttranslational modifications based on high-resolution mass spectrometry (Gingras et al., 2007; Altelaar et al., 2013).

Proteomics studies of influenza virus infection in macaques (Brown et al., 2010), mice (Kumar et al., 2014), continuous cell lines (Vester et al., 2009; Coombs et al., 2010; Kummer et al., 2014), chicken (Zou et al., 2010; Li et al., 2017), dogs (Su et al., 2015) and primary human cells (Liu et al., 2008; Lietzen et al., 2011; Kroeker et al., 2012; Liu et al., 2012) have provided information on alteration of host proteome at cellular and organism levels. Recent studies of chicken proteome response against H5N1 infection revealed several alterations of cytoskeleton, metabolic process, cellular component, and transcription regulation proteins (Zou et al., 2010; Li et al., 2017). However, detailed knowledge of signalling pathways and proteomic determinants responsible for the HPAI H5N1 viral pathogenesis in avian species is not known. In this study, the intracellular signaling pathways and critical host proteins responsible for influenza pathogenesis in chicken lungs infected with HPAI H5N1 virus are characterized at the proteome level.

Materials and Methods

Experimental infection of chicken

Six weeks old healthy domestic chicken, sero-negative for Avian Influenza virus (AIV) were used for this study. The animal experiments were approved by the Institutional Animal Ethics Committee of ICAR-NIHSAD (Approval no. 68/IAEC/HSADL/12 dated 11.05.2012) and all the experiments were conducted in biosafety level 3 containment facility of ICAR-National Institute of High Security Animal Diseases, Bhopal, India. The chicken were separated into four groups (n=5 birds/group). Among the four groups, three groups were intranasally inoculated with 10^6 EID₅₀ of H5N1 virus (A/duck/India/02CA10/2011/Agartala) and one group (control) was inoculated with PBS. The birds were observed daily for clinical signs. Lung tissues were collected from five birds from each infected group at 12hr, 24 hr, and 48hr post-infection. Lung tissues were collected from the control group at 12hr post-inoculation. The tissues were snap chilled in Liquid Nitrogen and stored at -80°C until protein

extraction. Avian influenza virus infection of lung tissues was confirmed by virus isolation upon inoculation in embryonated chicken eggs (ECs) and RT-PCR.

Protein extraction

150mg lung tissue from each sample was washed in 50mM NH_4HCO_3 washing buffer. The lung tissue was cut into small pieces and 650 μl of SDS protein extraction lysis buffer [0.1% SDS (Invitrogen); 50mM NH_4HCO_3 (Sigma); 1X cOmplete™ Protease Inhibitor Cocktail (Roche-11836145001)] was added. Tissue samples were homogenized in LZ-Lyser homogenizer at 30 HZ for 2 min. After complete homogenization, the tissue lysate was incubated on ice for 90 mins for complete protein lysis. The lysate was centrifuged at 20,000g for 60 mins at 4°C and the supernatant was collected. The supernatants were immediately snap heat treated at 56°C for 30 min in a dry bath for inactivation of HPAIV H5N1 in the protein extracts. All the heat treated samples were stored at -80°C for mass spectrometry analysis.

Sample preparation for LC-MS analysis

Lung protein quality was checked by 8% SDS-PAGE. A pool for each time point was prepared by pooling 50 μg protein each from 3 best samples at that time point. Protein samples were reduced for 1 hr at 95°C in 100mM dithiothreitol solution followed by alkylation for 45min by 55mM iodoacetamide in the dark at RT. Trypsin was added to all protein samples at a 1:20 (wt/wt) trypsin-to-protein ratio for overnight at 37°C. After trypsin digestion, the sample quality was again checked by SDS PAGE. Digested peptide samples were concentrated to 50 μl total volume using a centrifugal vacuum. The non-infected and infected peptide samples were labeled with iTRAQ 4-Plex (P/N: 4352135) reagents. The labels used for sample pools of different groups were as follows: iTRAQ label 114- control; iTRAQ label 115-12hr; iTRAQ label 116-24hr; iTRAQ label 117-48hr. These iTRAQ labeled samples were pooled and then purified using a strong cation exchange (SCX) chromatography. The fractions from SCX chromatography were collected and pooled

according to RT values. The pooled fractions were vacuum dried and dissolved in 10µL of 0.1% formic acid. The sample was injected (1µL) onto C18 Nano-LC column for separation of peptides followed by mass spectrometry analysis on the WATERS Q-TOF instrument.

Bioinformatics analysis

The WATRES specific raw data set files were converted to proteomics standard format mzXML format by proteowizard tool MSconvert with default parameters option (Chambers et al., 2012). The database search and other downstream bioinformatic analysis were done in Trans-Proteomic Pipeline (TPP) (Deutsch et al., 2010). TPP includes modules for validation of database search results, quantitation of isotopically labeled samples, and validation of protein identifications. MS/MS ion spectra were searched against chicken UniProt reference database using comet search engine (Eng et al., 2013). All 17719 proteins that comprise the chicken UniProt reference database and 11 protein sequences of HPAIV H5N1 virus (A/duck/India/02CA10/2011/Agartala from NCBI protein database ([http://www.ncbi.nlm.nih.gov/protein/?term=txid1004759\[Organism:noexp\]](http://www.ncbi.nlm.nih.gov/protein/?term=txid1004759[Organism:noexp])) were downloaded. The following parameters were used for database search, precursor/peptide mass tolerance- 1.8Da, fragment tolerance -1.6Da, fixed modification – carboxamidomethylation for cys (C) and iTRAQ N terminal, Variable modification- oxidation (M) and phosphorylation of (STY), and number of missed cleavage 2. The Peptide-Spectrum assignments of comet search engine were validated with PeptideProphet and iProphet tool of TPP (Keller et al., 2002). Protein identifications were validated with ProteinProphet tool of TPP based on PeptideProphet or iProphet results (Nesvizhskii and Aebersold, 2004). Proteins identified from ProteinProphet result were filtered based on protein probability above 0.95 (local FDR 5% level) and contained at least 1 unique peptide. This filtered protein list was used for further downstream functional analysis. Up and downregulation of particular protein was calculated as infected sample intensity divided by

control sample intensity (i.e. 115-12hr/114-control).Likewise for all proteins and specific post-infection time interval, fold change values were calculated. Functional classification of the proteins was performed for gene ontology (GO) in Database for Annotation, Visualization and Integrated Discovery (DAVID) (Huang et al., 2009) and pathway analysis in Kyoto Encyclopedia of Genes and Genomes (KEGG) (www.genome.jp/kegg/) (Kanehisa et al., 2017). Heatmap generated using ‘Clustvis’ web tool (Metsalu et al., 2015). We used online web server NetworkAnalyst for construction of protein-protein interaction (PPI) networks (Xia et al., 2014). The main driving or hub proteins were identified on the basis of two topological measures, degree centrality and betweenness centrality.

Meta-analysis of transcriptome datasets

For meta-analysis of chicken transcriptome datasets, we selected the microarray datasets of 2 independent studies. Hu et al. (2015) studied immune response of primary chicken lung cells infected with two HPAI H5N1 viruses using microarray technology (Hu et al., 2015). Ranaware et al. (2016) studied global immune response of chicken infected with HPAI H5N1 (A/duck/India/02CA10/2011) virus infection (Ranaware et al., 2016). The microarray datasets were analysed in GEO2R online tool of the NCBI (<http://www.ncbi.nlm.nih.gov/geo/geo2r/>).The original submitter-supplied processed microarray data tables were identified using the GEO query. Then, we identified control and test group samples and then samples belonging to each group were assigned. Further, the logFC, p-value and adj. p-value were calculated by limma R package. Adjusted p-values below 0.05 were used as threshold to find differentially expressed genes.

Bayesian networks were reconstructed for TLR, RLR, IL1R, NLR and JAK-STAT signaling pathways using ‘bnlearn’ package (Scutari, 2017). The networks were constructed based on intensity values of microarray dataset of our pervious published work (Ranaware et al., 2016). Bayesian network structure was learned from transcriptomic dataset with prior

knowledge using Hill Climbing (HC) algorithm. HC is a score-based structure learning algorithm. After learning the network structure, the conditional probability tables (CPTs) at each node were found by running bn.fit function. Bn.fit runs the EM algorithm to learn CPT for different nodes in the learned graph.

Results and discussion

Virus-host interactions are multidimensional, they include the alterations in transcriptome, proteome, metabolome and lipidome of the host. In recent years, most of the high-throughput studies have focused majorly on the host transcriptomic responses and other host components have received less attention. This study presents a comprehensive characterization of the lung proteome, critical signaling pathways and proteomic determinants responsible for disease pathogenesis at the proteome level in chicken lung tissues infected with HPAI H5N1 virus over different time points post-infection.

Clinical signs

Chicken in the control group did not show any clinical signs during the experimental period. In the test group, the birds were normal up to 12 hr post-infection. Mild clinical symptoms like depression, decreased feed as well as water consumption and ruffled feathers were observed at 24 hr post-infection. The clinical symptoms like dullness, lacrimation, cyanotic combs and wattles, edema and red discoloration of the shanks and feet were seen in the birds at 48 hr post-infection.

Raw mass spectra dataset analysis

Total 15 fractions of WATERS QTOP raw data sets were generated, with data size 48 GB, including 1, 47,451 MS scans and 19,917 MS/MS scans. Total 19,917 MS/MS spectra were searched against chicken protein database by comet search engine. The iProphet algorithm identified 17,273 unique peptides and 8,516 unique proteins at minimum probability threshold of above 0.05 and minimum above 7 amino acids. In order to increase

the accuracy of validation of peptides and proteins, we applied local FDR 5% level (probability cut off above 0.95) in the ProteinProphet output. At higher probability threshold cut off, we identified total 3313 proteins with the presence of at least one unique peptide. ProteinProphet predicted sensitivity and error rate information is shown in Figure S1. This is the highest protein identification reported till date in chicken. Our proteomic approach identified H5N1 viral peptides such as NA, NP and PB1 in the chicken lung proteome. NA is a sialidase responsible for releasing sialic acid from glycoprotein and glycolipid sialoconjugates of bound influenza virus to assist virus release (Wagner et al., 2001). NP is an important viral protein responsible for the packaging of the viral RNA and also shown to be involved in many aspects of influenza viral replication (Portela and Digard, 2002; Wasilenko et al., 2008). PB1 viral protein has been shown to be associated with the high pathogenicity of H5N1 viruses infection in ducks (Hulse-Post et al., 2007). Influenza viral infection status of lung tissues was confirmed by virus isolation in embryonated egg inoculation and RT-PCR method as well as by identification of viral peptides in lung proteome dataset.

Differential protein expression analysis

At 12 hr post-infection, 820 proteins were upregulated and 2493 proteins were downregulated in chicken lung tissues. A total of 827 proteins were upregulated and 2441 proteins were downregulated at 24 hr post-infection. At 48 hr post-infection, 693 proteins were upregulated and 2620 proteins were downregulated in chicken lung proteome (Table 1). A total of 470 and 2235 proteins were found to be commonly upregulated and downregulated in all time intervals post-infection in chicken lungs infected with H5N1 virus, respectively (Figure 1). The protein profile showed that 70, 70 and 101 proteins were exclusively upregulated in chicken lung tissue at 12hr, 24hr and 48hr post-infection, respectively (Figure 1). Downregulation of 87, 35 and 245 proteins were observed exclusively at 12 hr, 24hr and 48hr post-infection, respectively (Figure 1). The fold change value of the upregulated proteins ranged from 42 to 1. Interestingly, at 48 hr time point (fold change value below 1.5), higher

number of proteins were downregulated (n=1754) as compare to 12 hr time point (n=247) post-infection condition (Table 1, Figure 2). This result indicates that most of the host proteins were downregulated at the later stage of infection.

Gene ontology analysis of the commonly upregulated and downregulated lung tissue proteins of H5N1 infected chicken lung enriched following GO terms such as cytoskeleton, regulation of cell cycle, regulation of protein kinase activity etc. (Table 2). The upregulation (KRT6A, MCPH1, MYH7, MICAL1 and MICAL3) and downregulation (ACTL9, CTNNB1, DNM1, FILIP1L, and MYO1B) of cytoskeletal proteins were observed in the lung tissue of chicken. Cytoskeletal proteins have been reported to interact with viral proteins to regulate viral replication and assembly as well as the transport of viral components in the cell (Avalos et al., 1997; Radtke et al., 2006). Similar associations of cytoskeletal proteins with influenza infection condition are also reported previously (Coombs et al., 2010; Sui et al., 2014; Su et al., 2015). The cyclin-dependent kinases (CDKs) such as CDK13, DDB1, DRD3, FOXG1, and TCF3 that are involved in the regulation of cell cycle were all upregulated in chicken lung proteome. Söderholm *et al.* (2016) reported that the cyclin-dependent kinase activity is required for efficient viral replication and for activation of the host antiviral responses (Soderholm et al., 2016). Protein kinase activity associated proteins, namely, JAK3, MARK3, TBK1, EIF2AK3, PRKCA, and TRPM6 were downregulated in the chicken lung tissues. Differential expression of proteins including signal transduction molecules, kinases and other biochemical metabolism related enzymes that are associated with the repair of damaged lung tissues are reported in dogs infected with influenza virus infection (Su et al., 2015). In addition, some apoptosis and tumor-associated proteins (BCL6, FAF1, TBX5, AKAP13, TNFSF10 and TGFB1) were also identified in the chicken lung proteome. Further we did GO term analysis of proteins that were exclusively expressed in particular post-infection condition in chicken (Table 3). This analysis result provides information how the disease

progress from onset of disease to outcome of disease in chicken. At 12 hr interval, most of cellular homeostasis were affected and at 48 hr post-infection condition the host activated critical pathways such as Influenza A, chemokine signaling pathway, Jak-STAT signaling pathway, apoptosis, MAPK signaling pathway (Table 3). These results indicate that the virus initially inhibit the cellular homeostasis and activates the critical pathways at the later stage of infection.

Molecular pathogenesis of H5N1 infection in chicken

Previous transcriptomics study of our group found that highly pathogenic H5N1 virus induced excessive expression of type I IFNs, cytokine, chemokines and ISGs in the lung tissues (Ranaware et al., 2016). This atypical expression of immune genes (cytokine storm) might be the cause for the high mortality in chickens. However, the information on the pathways activated, constituents of cytokine storm and therapeutic strategies against cytokine storm is lacking in avian species. Intensive molecular studies in humans and human animal models system have identified, (1) Activation of TLR3 and 7, as well as endosome (TLR3 and 7) and cytosolic (RIG-I) pathways; (2) activation of IL-1R signaling pathway; and (3) activation of MAVS/MyD88/TRIF signaling as essential pathways involved in the cytokine storm (Teijaro et al., 2014).

In order to check whether these pathways are activated and if activated what are the levels of expression of the immune genes in chicken lung tissues, we applied meta-analysis of lung transcriptome datasets. We utilised the previously published microarray datasets of chicken infected with HPAIVs, because transcriptomic data can capture the complete gene expression dynamics at a particular condition. We mapped the differentially expressed genes retrieved from meta-analysis results of chicken into influenza reference pathways in KEGG database. Activation of TLR signaling pathway, RIG I signaling pathway, NOD like receptor signaling pathway and JAK-STAT signaling pathway were observed in chicken lung transcriptome (Figure S2). Similarly, activation of these pathways was evident in the chicken

proteome datasets (Figure 3). Further Bayesian networks (BN) were constructed with prior knowledge using chicken meta-analysis transcriptome datasets (Figure 4, Figure S3). The combined *in silico* analysis of transcriptome and proteome datasets confirmed the activation of TLRs, RLRs, NLR and Jak-STAT signaling pathways in lung tissues infected with HPAIVs in chickens. The fact that most of the influenza pathogenesis possesses abnormalities in all of these core pathways suggests they play a central role in the cytokine storm.

Next we examined the cytokine storm responsive genes (i.e expression level of cytokines, chemokines and ISGs) as a result of activation of the above mentioned pathways. Cytokine storm responsive genes lists were compiled for chickens based on information in literature (Mishra et al., 2017). The expression levels (fold change) of these genes were retrieved from meta-analysis transcriptome datasets of chicken. Cytokines, chemokines and ISGs genes were found to be upregulated in chicken lung tissues and these may be the basis of the increase in the severity of HPAI H5N1 infection in chickens (Figure 5). In summary, we characterized the immune pathways involved in the cytokine storm and identified cytokine storm responsive genes in chicken lung tissues infected with HPAIVs.

Identification of proteomic determinants responsible for disease pathogenesis in chicken

To know the main driver or hub proteins responsible for disease pathogenesis, we constructed protein-protein interaction (PPI) network based on chicken lung proteome dataset (Figure 6). The proteins such as MyD88, IKBKB, IRAK4, RELA, and MAVS involved in the TLRs, RLRs, IL1R and NLR signaling pathways were identified with high degree centrality and betweenness centrality measures (Table 4).

MYD88 gene encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the IL1R and TLR signaling pathways. These pathways in turn regulate the activation of numerous proinflammatory genes (Deguine and Barton, 2014). The IKBKB protein

phosphorylates the inhibitor in the inhibitor/NF-kappa-B complex, causing dissociation of the inhibitor and activation of NF-kB signaling pathway (Rothwarf and Karin, 1997). MAVS acts downstream of DDX58/RIG-I and IFIH1/MDA5 genes, as an essential signal transducer in the beta interferon signaling pathways and that contribute to antiviral immunity (Seth et al., 2005). RELA/NF-kB is a ubiquitous transcription factor involved in several biological processes. This transcription factor is activated through degradation of its specific inhibitor in the cytoplasm; NF-kB moves to the nucleus and activates transcription of specific genes. The NF-kBp65-p65 complex appears to be involved in invasin-mediated activation of IL-8 expression (Kunsch and Rosen, 1993). Teijaro et al. (2014) reported that MyD88 and MAVS as the predominant signaling molecule required for innate immune cell recruitment and for the majority of cytokine amplification (i.e. cytokine storm) in mice infected with influenza virus (Teijaro et al. 2014). Further they suggested that therapeutic control of cytokine storm is possible through a common pathway inhibition downstream of multiple innate pathogen-sensing molecules of cytokine amplification. In our study the identified hub proteins (MyD88, IKBKB, IRAK4, RELA, and MAVS) were all involved in different components of MyD88 and MVAS signaling pathways. Based on this literature, we suggest that successful therapeutic intervention for cytokine storm in chicken should target these proteins as drug targets to blunt the cytokine amplification. Further the S1P1R agonist therapy may suppress global cytokine amplification in chicken as in mice (Teijaro et al. 2014). However biological validation of this hypothesis by *in vivo* experiment is needed.

In Jak-STAT signalling pathway, we found STAT1, STAT2, STAT3 and SOCS3 proteins as the main driving proteins in the PPI network (Table 4). However, these proteins had protein probability range from 0.63 to 0.90, hence were not evident in our main proteome dataset of chicken. The STAT1, STAT2 and STAT-3 proteins are a key constituents of JAK-STAT signalling pathway, play critical roles in the IFN signalling pathway and are required for a robust IFN-induced antiviral response (Yang et al., 1998; Ho and Ivashkiv, 2006).

SOCS1 and SOCS3 are reported to be critical regulators of IFN responses through the inhibition of STAT phosphorylation and induction of ISGs through a RIG-I/MAVS/IFNAR1-dependent pathway (Pauli et al., 2008; Pothlichet et al., 2008).

Further we identified some novel main driver/hub proteins with a very high degree of centrality and betweenness of centrality measures which are not reported previously to be associated with influenza infection condition in human, human models or avian species. These novel hub proteins are HNF4A, ELAVL1, FN1, COPS5, CUL1, CAND1, BRCA1, CTNNB1 and FYN (Table 4, Figure 6). The HNF4A protein is a transcriptionally controlled transcription factor and required for the transcription of alpha 1-antitrypsin, apolipoprotein CIII, transthyretin genes and HNF1-alpha genes (Ryffel, 2001). ELAVL1 is a member of the ELAVL family of RNA-binding proteins and selectively bind AU-rich elements (AREs) found in the 3' untranslated regions of mRNAs. The ELAVL family of proteins play a role in stabilizing ARE-containing mRNAs (Brennan and Steitz, 2001). This gene has been implicated in a variety of biological processes and has been linked to a number of diseases, including cancer (Wang et al., 2013).

FN1 bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin. FN1 is involved in cell adhesion and migration processes during embryogenesis, wound healing, blood coagulation, host defense, and metastasis (Liao et al., 2018). COPS5 is one of the eight subunits of COP9 signalosome and functions as an important regulator in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8 signalling (Bech-Otschir et al., 2002). CUL1 is a core component of multiple cullin-RING-based SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of proteins involved in cell cycle progression, signal transduction and transcription (Kipreos et al., 1996).

BRCA1 encodes a nuclear phosphoprotein that has a role in maintaining genomic stability, and acts as a tumour suppressor. This protein is involved in transcription, DNA

repair of double-stranded breaks, and recombination (Foulkes and Shuen, 2013). FYN gene encodes a membrane-associated tyrosine kinase that has been implicated in the control of cell growth (Zheng et al., 2017). In summary, many proteins involved in the TLRs, RLRs, NLR and Jak-STAT signaling pathways and other novel proteins were identified as main protein determinant and these proteins might be linked with disease pathogenesis in chicken H5N1 infection. However critical functional role of these proteins in avian influenza pathogenesis in chicken requires further biological validation by *in vivo* and *in vitro* experiments.

In conclusion, we characterized the comprehensive proteome profile of chicken lung tissues infected with HPAI H5N1 virus at different time points post-infection. There are differences in the protein profile at different time points post-infection indicated by the remarkable variations in the levels of expression of proteins at these time points. Combined analysis of transcriptome and proteome datasets revealed activation of TLR, RLR, NLR, and JAK-STAT signaling pathways associated with a cytokine storm effect in chicken infected with HPAI H5N1 virus. Further, we identified many of the important hub proteins linked with influenza pathogenesis in chicken.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD010358.

Ethics statement

The experiments were approved by the Institutional Animal Ethics Committee of ICAR-NIHSAD (Approval no. 68/IAEC/HSADL/12 dated 11.05.2012), and performed under the guidance of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Ministry of Environment and Forests, Govt. of India.

Acknowledgements

This work was funded by the Department of Biotechnology (grant number: BT/IN/Indo-UK/FADH/48/AM/2013). We thank Director, ICAR-National Institute of High

Security Animal Diseases, Director, ICAR-Indian Veterinary Research Institute, and Indian Council of Agricultural Research, India for providing necessary facilities to carry out this work.

Author Contributions

Conceived and designed the experiments: A.M., A.A.R. and P.V. Performed the experiments: A.M., S.C., A.A.R and P.V. Analyzed the data: P.V. and A.M. Contributed reagents/materials/analysis tools: H.V.M., D.D.K. and V.P.S Wrote the paper: P.V. and A.M.

References

1. Altelaar, A.F., Munoz, J. and Heck, A.J. (2013). Next-generation proteomics: towards an integrative view of proteome dynamics. *Nat. Rev. Genet.* 14, 35–48.
2. Avalos, R.T., Yu, Z. and Nayak, D.P. (1997). Association of influenza virus NP and M1 proteins with cellular cytoskeletal elements in influenza virus-infected cells. *J. Virol.* 71, 2947–58.
3. Bech-Otschir, D., Seeger, M. and Dubiel, W. (2002). The COP9 signalosome: at the interface between signal transduction and ubiquitin-dependent proteolysis. *J. Cell Sci.* 115, 467-73.
4. Brennan, C.M. and Steitz, J.A. (2001). HuR and mRNA stability. *Cell Mol. Life Sci.* 58, 266-77.
5. Brown, J.N. et al. (2010). Macaque proteome response to highly pathogenic avian influenza and 1918 reassortant influenza virus infections. *J. Virol.* 84, 12058-68.
6. Chambers, M.C. et al. (2012). A cross-platform toolkit for mass spectrometry and proteomics. *Nat. Biotechnol.* 30, 918-20.
7. Coombs, K.M. et al. (2010). Quantitative proteomic analyses of influenza virus-infected cultured human lung cells. *J. Virol.* 84, 10888-906.
8. Deguine, J. and Barton, G.M. (2014). MyD88: a central player in innate immune signaling. *F1000Prime Rep.* 4, 6:97.

9. Deutsch, E.W. et al. (2010). A guided tour of the Trans-Proteomic Pipeline. *Proteomics*, 10, 1150-9.
10. Eng, J.K., Jahan, T.A. and Hoopmann, M.R. (2013). Comet: an open-source MS/MS sequence database search tool. *Proteomics*, 13, 22-4.
11. Fauci, A.S (2006). Emerging and re-emerging infectious diseases: influenza as a prototype of the host-pathogen balancing act. *Cell*, 124, 665-70.
12. Foulkes, W.D. and Shuen, A.Y. (2013). In brief: BRCA1 and BRCA2. *J. Pathol.* 230, 347-9.
13. Gingras, A.C., Gstaiger, M., Raught, B. and Aebersold, R. (2007). Analysis of protein complexes using mass spectrometry. *Nature Rev. Mol. Cell Biol.* 8, 645–654.
14. Ho, H.H. and Ivashkiv, L.B. (2006). Role of STAT3 in type I interferon responses. Negative regulation of STAT1-dependent inflammatory gene activation. *J. Biol. Chem.* 281, 14111-14118.
15. Hu, J. et al. (2015). PA-X decreases the pathogenicity of highly pathogenic H5N1 influenza A virus in avian species by inhibiting virus replication and host response. *J. Virol.* 89, 4126-42.
16. Huang, D.W., Sherman, B.T. and Lempicki, R. A. (2009). Systematic and integrative analysis of large gene lists using DAVID Bioinformatics Resources. *Nature Protoc.* 4, 44-57.
17. Hulse-Post, D.J. et al. (2007). Molecular changes in the polymerase genes (PA and PB1) associated with high pathogenicity of H5N1 influenza virus in mallard ducks. *J. Virol.* 81, 8515-24.
18. Josset, L., Tisoncik-Go, J. and Katze, M.G. (2013). Moving H5N1 studies into the era of systems biology. *Virus Res.* 178, 151-67.

19. Kanehisa, M. et al. (2017). KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* 45, D353-D361.
20. Keller, A., Nesvizhskii, A.I., Kolker, E. and Aebersold, R. (2002). Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search. *Anal. Chem.* 74, 5383-92.
21. Kipreos, E.T. et al. (1996). cul-1 is required for cell cycle exit in *C. elegans* and identifies a novel gene family. *Cell*, 85, 829-39.
22. Kroeker, A.L., Ezzati, P., Halayko, A.J. and Coombs, K.M. (2012). Response of primary human airway epithelial cells to influenza infection: a quantitative proteomic study. *J. Proteome Res.* 11, 4132–4146.
23. Kumar, Y. et al. (2014). Molecular analysis of serum and bronchoalveolar lavage in a mouse model of influenza reveals markers of disease severity that can be clinically useful in humans. *PLoS One* 9, e86912.
24. Kummer, S. et al. (2014). Alteration of protein levels during influenza virus H1N1 infection in host cells: a proteomic survey of host and virus reveals differential dynamics. *PLoS One*, 9, e94257.
25. Kunsch, C. and Rosen, C.A. (1993). NF-kappa B subunit-specific regulation of the interleukin-8 promoter. *Mol. Cell Biol.* 13, 6137-46.
26. Li, Y. et al. (2017). Proteome Response of Chicken Embryo Fibroblast Cells to Recombinant H5N1 Avian Influenza Viruses with Different Neuraminidase Stalk Lengths. *Sci. Rep.* 7, 40698.
27. Liao, Y.X., Zhang, Z.P., Zhao, J. and Liu, J.P. (2018). Effects of Fibronectin 1 on Cell Proliferation, Senescence and Apoptosis of Human Glioma Cells Through the PI3K/AKT Signaling Pathway. *Cell Physiol. Biochem.* 48, 1382-1396.
28. Lietzen, N. et al. (2011). Quantitative subcellular proteome and secretome profiling of influenza A virus-infected human primary macrophages. *PLoS Pathog.* 7, e1001340.

29. Liu, L., Zhou, J., Wang, Y., Mason, R.J., Funk, C.J. and Du, Y. (2012). Proteome alterations in primary human alveolar macrophages in response to influenza A virus infection. *J. Proteome Res.* 11, 4091–4101.
30. Liu, N. et al. (2008). Proteomics analysis of differential expression of cellular proteins in response to avian H9N2 virus infection in human cells. *Proteomics*, 8, 1851-8.
31. Metsalu, T. and Vilo, J. (2015). ClustVis: a web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap. *Nucleic Acids Res.* 43, W566-70.
32. Mishra, A., Vijayakumar, P. and Raut, A. A. (2017). Emerging avian influenza infections: Current understanding of innate immune response and molecular pathogenesis. *Int. Rev. Immunol.* 36, 89-107.
33. Nesvizhskii, A.I. and Aebersold, R. (2004). Analysis, statistical validation and dissemination of large-scale proteomics datasets generated by tandem MS. *Drug Discov. Today.* 9, 173–181.
34. Pauli, E.K. et al. (2008). Influenza A virus inhibits type I IFN signaling via NF- κ B dependent induction of SOCS-3 expression. *PLoS Pathog.* 4, e1000196.
35. Portela, A. and Digard, P. (2002). The influenza virus nucleoprotein: a multifunctional RNA-binding protein pivotal to virus replication. *J. Gen. Virol.* 83, 723-34.
36. Pothlichet, J. Chignard, M. and Si-Tahar M. (2008). Cutting edge: innate immune response triggered by influenza A virus is negatively regulated by SOCS1 and SOCS3 through a RIG-I/IFNAR1-dependent pathway. *J. Immunol.* 180, 2034–2038.
37. Radtke, K., Dohner, K. and Sodeik, B. (2006). Viral interactions with the cytoskeleton: a hitchhiker's guide to the cell. *Cell Microbiol.* 8, 387–400.
38. Ranaware, B. et al. (2016). Genome Wide Host Gene Expression Analysis in Chicken Lungs Infected with Avian Influenza Viruses. *PLoS One*, 11, e0153671.

39. Rothwarf, D.M. and Karin, M. (1999). The NF-kappa B activation pathway: a paradigm in information transfer from membrane to nucleus. *Sci. STKE*. 1999, RE1.
40. Ryffel, G.U. (2001). Mutations in the human genes encoding the transcription factors of the hepatocyte nuclear factor (HNF) 1 and HNF4 families: functional and pathological consequences. *J. Mol. Endocrinol.* 27,11-29.
41. Scutari, M. (2017). Bayesian Network Constraint-Based Structure Learning Algorithms: Parallel and Optimized Implementations in the bnlearn R Package. *Journal of Statistical Software*, 77, 1-20.
42. Seth, R.B., Sun, L., Ea, C.K. and Chen, Z.J. (2005). Identification and characterization of MAVS, a mitochondrial antiviral signaling protein that activates NF-kappaB and IRF 3. *Cell*, 122, 669-82.
43. Soderholm, S. et al. (2016). Phosphoproteomics to characterize host response during influenza A virus infection of human macrophages. *Mol. Cell Proteomics*. 15, 3203-3219.
44. Su, S. et al. (2015). Global and quantitative proteomic analysis of dogs infected by avian-like H3N2 canine influenza virus. *Front. Microbiol.* 6, 228.
45. Sui, Z., Wen, B., Gao, Z. and Chen, Q. (2014). Fusion-Related Host Proteins Are Actively Regulated by NA during Influenza Infection as Revealed by Quantitative Proteomics Analysis. *PLoS ONE*, 9, e105947.
46. Teijaro, J.R., Walsh, K.B., Rice, S., Rosen, H. and Oldstone, M.B. (2014). Mapping the innate signaling cascade essential for cytokine storm during influenza virus infection. *Proc. Natl. Acad. Sci. U S A*. 111, 3799-804.
47. Vester, D., Rapp, E., Gade, D., Genzel, Y. and Reichl, U. (2009). Quantitative analysis of cellular proteome alterations in human influenza A virus-infected mammalian cell lines. *Proteomics*, 9, 3316-27.

48. Wagner, R., Matrosovich, M. and Klenk, H.D. (2002). Functional balance between haemagglutinin and neuraminidase in influenza virus infections. *Rev. Med. Virol.* 12, 159-66.
49. Wang, H. et al. (2013). The structure of the ARE-binding domains of Hu antigen R (HuR) undergoes conformational changes during RNA binding. *Acta. Crystallogr. D. Biol. Crystallogr.* 69, 373-80.
50. Wasilenko, J.L. et al. (2008). NP, PB1, and PB2 viral genes contribute to altered replication of H5N1 avian influenza viruses in chickens. *J. Virol.* 82, 4544-53.
51. Xia, J., Benner, M.J. and Hancock, R.E. (2014). NetworkAnalyst--integrative approaches for protein-protein interaction network analysis and visual exploration. *Nucleic Acids Res.* 42:W167-74.
52. Yang, C.H., Murti, A. and Pfeffer, L.M. (1998). STAT3 complements defects in an interferon-resistant cell line: evidence for an essential role for STAT3 in interferon signaling and biological activities. *Proc. Natl. Acad. Sci. U S A.* 95, 5568-5572.
53. Zak, D.E., Tam, V.C. and Aderem, A. (2014). Systems-level analysis of innate immunity. *Annu. Rev. Immunol.* 32, 547-77.
54. Zheng, J., Li, H., Xu, D. and Zhu, H. (2017). Upregulation of Tyrosine Kinase FYN in Human Thyroid Carcinoma: Role in Modulating Tumor Cell Proliferation, Invasion, and Migration. *Cancer BiotherRadiopharm.* 32, 320-326.
55. Zou, W. et al. (2010). Proteomics analysis of differential expression of chicken brain tissue proteins in response to the neurovirulent H5N1 avian influenza virus infection. *J. Proteome Res.* 9, 3789-98.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

512 **Table 1.** Differential protein expression analysis in chicken infected with HPAI H5N1 virus.

Condition	Upregulated proteins	Downregulated proteins	Upregulated proteins (>1.5 fold)	Downregulated Proteins (<1.5 fold)
12hr interval	820	2493	138	247
24hr interval	872	2441	157	222
48hr interval	693	2620	173	1754

513

514 **Table 2.** Gene Ontology term analysis of commonly upregulated and downregulated lung
515 tissue proteins in chicken infected with HPAI H5N1 virus.

516

GO terms	Count	P-Value
Upregulated proteins		
Golgi apparatus	32	7.70E-04
Cellular homeostasis	21	1.10E-03
Protein heterodimerization activity	12	2.20E-03
Ion homeostasis	18	3.70E-03
Protein domain specific binding	15	4.40E-03
Acid-amino acid ligase activity	11	5.20E-03
Intracellular organelle lumen	50	6.40E-03
Fatty acid transport	4	1.80E-02
Regulation of cell cycle	13	3.50E-02
Vesicle-mediated transport	19	4.20E-02
Positive regulation of protein kinase cascade	8	5.10E-02
Negative regulation of gene expression	16	8.20E-02
Cytoskeleton	35	8.60E-02
Adenyl nucleotide binding	47	1.90E-03

ATP binding	43	4.90E-03
Downregulated proteins		
Adenyl nucleotide binding	236	1.20E-13
ATP binding	222	5.40E-13
GTPase regulator activity	74	4.50E-08
Cytoskeleton	176	1.20E-06
Protein kinase activity	82	9.20E-04
Regulation of small GTPase mediated signal transduction	47	8.10E-06
Regulation of Rho protein signal transduction	25	1.10E-05
Proteinaceous extracellular matrix	51	8.80E-05
Regulation of Ras protein signal transduction	38	1.30E-04
Protein amino acid phosphorylation	89	6.20E-04
Cell death	93	1.30E-03
Enzyme linked receptor protein signalling pathway	49	3.10E-03
Endosome	44	4.10E-03
Intracellular signaling cascade	146	4.30E-03
Positive regulation of gene expression	70	2.50E-02
Calcium ion binding	105	3.20E-02
Negative regulation of translation	8	3.60E-02
Regulation of phosphorylation	56	4.60E-02
JUN kinase binding	3	4.70E-02
Proteolysis	114	7.20E-02
Positive regulation of MAP kinase activity	15	9.60E-02

517

518

519

520

521 **Table 3.** Functional annotation analysis of proteins exclusively expressed in chicken at

522 different time points post infection with HPAI H5N1 virus.

Pathway activation	No. of proteins	P Value
12 hr post infection		
Cell cycle	31	2.88E-26
mRNA surveillance pathway	21	4.95E-18
RNA transport	24	2.62E-17
p53 signaling pathway	11	1.22E-07
Pathways in cancer	13	0.0128
24 hr post infection		
Cell cycle	12	1.17E-08
Gap junction	8	7.31E-06
RNA degradation	6	6.13E-05
T cell receptor signaling pathway	7	0.000127
Fc epsilon RI signaling pathway	6	0.000215
ErbB signaling pathway	6	0.000485
Phagosome	4	0.00326

Fc gamma R-mediated phagocytosis	5	0.00529
Chemokine signaling pathway	7	0.00615
B cell receptor signaling pathway	4	0.0112
Neurotrophin signaling pathway	5	0.0141
Natural killer cell mediated cytotoxicity	5	0.0221
Focal adhesion	6	0.0287
48 hr post infection		
RNA transport	47	1.77E-18
T cell receptor signaling pathway	37	6.08E-15
Regulation of actin cytoskeleton	52	1.15E-14
Focal adhesion	54	4.67E-14
Pathways in cancer	70	1.13E-13
ErbB signaling pathway	31	6.41E-12
Adipocytokine signaling pathway	26	6.92E-12
B cell receptor signaling pathway	28	1.91E-11
Neurotrophin signaling pathway	35	3.79E-10
mRNA surveillance pathway	26	6.14E-09
Chemokine signaling pathway	41	7.57E-08

Influenza A	28	1.68E-07
Cell cycle	30	3.95E-07
Jak-STAT signaling pathway	25	1.62E-06
Fc epsilon RI signaling pathway	21	1.86E-06
mTOR signaling pathway	14	2.72E-05
Toll-like receptor signaling pathway	20	0.000377
Apoptosis	18	0.000385
Natural killer cell mediated cytotoxicity	24	0.00139
MAPK signaling pathway	38	0.00297
Leukocyte transendothelial migration	19	0.00374
RIG-I-like receptor signaling pathway	11	0.0039

523

524

525

526 **Table 4.** Hub proteins identified in chicken PPI networks based on degree centrality and
527 betweenness centrality measures.

Proteins	Degree centrality	Betweenness centrality
HNF4A	254	254342.21
ELAVL1	254	253172.78
FN1	117	79642.94
COPS5	117	73208.97

CUL1	105	55049.71
CAND1	105	45936.7
CTNNB1	80	75126.54
BRCA1	72	41422.24
FYN	65	35789.72
MYD88	10	2614.04
IKBKB	42	16555.52
RELA	48	23497.55
MAVS	6	1721.53
STAT1	81	46280.33
STAT2	14	1579.16
STAT3	90	66853.77
SOCS3	23	11524.2
IRAK4	8	311.94

528

529

530

531

532

533

534

535

536

537

538

Fig 1. Comparative analysis of upregulated (a) and downregulated (b) proteins expression changes at different time intervals in chicken lung tissues.

Fig 2. Heatmap of differentially expressed proteins of HPAIV infected lung tissues. The expression levels are visualized using a gradient colour scheme.

Fig 3. KEGG pathway analysis of commonly upregulated and downregulated proteins in chicken lung tissues infected with HPAI H5N1 virus.

Fig 4. Bayesian network was constructed based on meta-analysis of transcriptome dataset of chicken. BN constructed with prior knowledge of TLR, RLR, IL1R and NLR signaling pathways in chicken.

Fig 5. Expression level of cytokine storm responsive genes in chicken lung tissues infected with HPAIVs.

611 Fig 6. The protein-protein interaction network of chicken lung tissues infected with HPAI
 612 H5N1 virus. The important hub genes involved in influenza molecular pathogenesis were
 613 highlighted with blue colour in the PPI network.

614