

1 **Sex chromosome and sex locus characterization in the goldfish,**

2 ***Carassius auratus*.**

3
4 Ming Wen^{1,2}, Romain Feron^{2,3,4}, Qiaowei Pan^{2,3}, Justine Guguin², Elodie Jouanno², Amaury
5 Herpin², Christophe Klopp^{5,6}, Cedric Cabau⁶, Margot Zahm⁶, Hugues Parrinello⁷, Laurent
6 Journot⁷, Shawn M. Burgess⁸, Yoshihiro Omori^{9,10}, John H. Postlethwait¹¹, Manfred Schartl¹²,
7 Yann Guiguen^{2*}

8

9 * Correspondance: Yann Guiguen: yann.guiguen@inra.fr

10

11 **AFFILIATIONS:**

12

13 ¹ State Key Laboratory of Developmental Biology of Freshwater Fish, College of Life Science,
14 Hunan Normal University, Changsha, China

15 ² INRAE, UR 1037 Fish Physiology and Genomics, F-35000 Rennes, France.

16 ³ Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland.

17 ⁴ Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland.

18 ⁵ Plate-forme bio-informatique Genotoul, Mathématiques et Informatique Appliquées de
19 Toulouse, INRAE, Castanet Tolosan, France.

20 ⁶ SIGENAE, GenPhySE, Université de Toulouse, INRAE, ENVT, Castanet Tolosan, France.

21 ⁷ Montpellier GenomiX (MGX), c/o Institut de Génomique Fonctionnelle, 141 rue de la
22 Cardonille, 34094, Montpellier Cedex 05, France.

23 ⁸ Translational and Functional Genomics Branch, National Human Genome Research Institute,
24 Bethesda, MD, USA.

25 ⁹ Laboratory of Functional Genomics, Graduate School of Bioscience, Nagahama Institute of
26 Bioscience and Technology, Nagahama, Shiga, Japan.

27 ¹⁰ Laboratory for Molecular and Developmental Biology, Institute for Protein Research, Osaka
28 University, Suita, Osaka, Japan.

29 ¹¹ Institute of Neuroscience, University of Oregon, Eugene, Oregon, USA.

30 ¹² Developmental Biochemistry, Biozentrum, University of Würzburg, Würzburg, Germany and
31 The Xiphophorus Genetic Stock Center, Department of Chemistry and Biochemistry, Texas
32 State University, San Marcos, Texas, USA.

33

34

35

36

37

38 **Abstract**

39 **Background:** Goldfish is an important model for various areas of research, including neural
40 development and behavior and a species of significant importance in aquaculture, especially as
41 an ornamental species. It has a male heterogametic (XX/XY) sex determination system that
42 relies on both genetic and environmental factors, with high temperatures being able to produce
43 female-to-male sex reversal. Little, however, is currently known on the molecular basis of
44 genetic sex determination in this important cyprinid model. We used sequencing approaches to
45 better characterize sex determination and sex-chromosomes in goldfish.

46 **Results:** Our results confirmed that sex determination in goldfish is a mix of environmental
47 and genetic factors and that its sex determination system is male heterogametic (XX/XY).
48 Using reduced representation (RAD-seq) and whole genome (pool-seq) approaches, we
49 characterized sex-linked polymorphisms and developed male specific genetic markers. These
50 male specific markers were used to distinguish sex-reversed XX neomales from XY males and
51 to demonstrate that XX female-to-male sex reversal could even occur at a relatively low rearing
52 temperature (18°C), for which sex reversal has been previously shown to be close to zero. We
53 also characterized a relatively large non-recombining region (~11.7 Mb) on goldfish linkage
54 group 22 (LG22) that contained a high-density of male-biased genetic polymorphisms. This
55 large LG22 region harbors 373 genes, including a single candidate as a potential master sex
56 gene, i.e., the anti-Mullerian hormone gene (*amh*). However, no sex-linked polymorphisms
57 were detected in the goldfish *amh* gene or its 5 kb proximal promoter sequence.

58 **Conclusions:** These results show that goldfish have a relatively large sex locus on LG22, which
59 is likely the goldfish Y chromosome. The presence of a few XX males even at low temperature
60 also suggests that other environmental factors in addition to temperature could trigger female-
61 to-male sex reversal. Finally, we also developed sex-linked genetic markers in goldfish, which
62 will be important for future research on sex determination and aquaculture applications in this
63 species.

64 **Key words:** Goldfish, RADseq, Poolseq, Sex determination, Sex markers, Male genome
65 assembly

66

67 **BACKGROUND**

68 Goldfish, *Carassius auratus*, is a domesticated fish species originating from central Asia and
69 China that has been introduced throughout the world. Goldfish belongs to the Cyprinidae family
70 and is considered as an important fish model for research in endocrinology [1, 2],
71 developmental biology [3, 4] or fish pathology [5]. Thanks to the recent availability of a whole
72 genome sequence assembly [6], goldfish is also now becoming a key model species for studies
73 on genomics and cyprinid genome evolution. It is also a species of high aquaculture importance
74 especially as an ornamental species, with many beautiful and sometimes bizarre phenotypes [7].
75 Unlike birds and mammals, sex determination in teleost is highly dynamic, with frequent
76 turnovers of both sex determination (SD) systems [8] and master sex determining genes (MSD)
77 [9, 10]. Currently about half a dozen different master sex determining genes have been
78 identified in teleosts, including *dmrt1* in the Japanese medaka, *Oryzias latipes* [11], *sdY* in
79 rainbow trout [12], *amh* in Northern pike, Nile tilapia and pejerrey [13-15], *amhr2* in yellow
80 perch and the Takifugu pufferfish [16, 17], *gsdf* in sablefish and Luzon medaka, *O. luzonensis*
81 [18, 19], *gsdf6a* in the turquoise killifish [20] and *sox3* in the Indian ricefish *O. dancena* [21].
82 MSD turnover can be evolutionarily rapid as has been shown for instance in various ricefish
83 species [22]. In addition to genetic determinants, environmental factors -- especially
84 temperature -- have also been shown to play a pivotal role in teleost sex determination [23].
85 Since the late 1960s, the goldfish sex determination system has been characterized as male
86 heterogametic (XX/XY) [24]. More recently, a strong temperature influence on sex-ratios has
87 also been characterized in goldfish, with high rearing temperature treatments inducing complete
88 masculinization of chromosomally all-female genotypes (XX neomales) when applied during
89 early 3 months development [25]. The molecular mechanisms of genetic sex determination,

90 however, are still unknown not only in goldfish, but also in any member of the Cyprinidae
91 family.

92 Because of new high throughput sequencing technologies and the availability of a whole
93 genome sequence assembly for goldfish [26], we implemented both reduced representation (i.e.,
94 Restriction-site associated DNA sequencing (RAD-seq) [27, 28], and whole genome (i.e., Pool
95 sequencing (Pool-seq) [29, 30]) approaches to identify sex-linked genetic polymorphisms in
96 goldfish. We verified that identified sex-linked markers strictly segregated with the Y
97 chromosome, and we characterized the extent of Y chromosome differentiation. Although our
98 experiments did not identify a strong candidate sex-determining gene, these results lay a solid
99 foundation for further molecular exploration of goldfish sex determination.

100

101 **RESULTS**

102 **Characterization of goldfish sex-linked Y chromosome markers**

103 Because goldfish sex determination is highly sensitive to temperature [21], with high
104 temperature leading to the masculinization of some XX females producing XX neomales, we
105 first searched for sex-linked markers using a RAD-seq approach that kept track of phenotypes
106 and genotypes, potentially enabling the discrimination of XX neomales from XY genetic males.

107 From our RAD-seq data, we identified 32 polymorphic/specific RAD-tags that were present in
108 12-15 males among the 30 phenotypic males used in this experiment, and completely absent in
109 all the 30 phenotypic females (Fig. 1A, Supplementary excel file1). These results suggest a
110 male heterogametic genetic sex determination system (XX/XY) as previously shown in goldfish
111 [24], but with a rather high occurrence of XX neomales (around 50 %) in this population of

112 two-year old animals raised outdoor and obtained from different batches of animals with
113 different spawning times i.e., from May-June to late September.

114 To validate the hypothesis that these markers were linked to the heterogametic sex (XY) and
115 the Y chromosome, we first sequenced using Illumina reads and assembled a draft genome
116 sequence of a male goldfish identified as a putative XY male based on the polymorphic/specific
117 RAD-tags (see Material & Methods) and blasted these 32 marker sequences against this genome
118 assembly. This analysis returned 20 contigs with highly significant matches (Supplementary
119 excel file2) spanning a total of 0.24 Mb. By anchoring these sex-linked RAD sequences on our
120 genome assembly, we were able to design three putative Y-allele specific primer pairs that were
121 used to genotype the same individual animals that were used for the RAD-seq analysis. PCR
122 genotyping using these three primer pairs accurately discriminated putative XY genetic males
123 from putative XX neomales and females (Fig. 1B), validating that these primers accurately
124 identified the two types of males found in our RAD-seq analysis. We then genotyped male
125 breeders from our experimental stock with these primers and selected one putative XX neomale
126 (breeder 1, negative PCR amplifications) and one putative XY male (breeder 2, positive PCR
127 amplifications); and both individuals were crossed to the same XX female to generate two
128 separate batches of fish. If our Y-allele specific primers correctly identify the Y chromosome,
129 then our putative XX neomale should give only female offspring and the putative XY male
130 should give both male and female offspring. These two experimental populations were then
131 reared at low temperature (18°C) during the first three months after fertilization to minimize
132 high temperature masculinization [25], and were subsequently maintained at 24°C for nine
133 additional months before the identification of the phenotypic sex. Results from the histological

134 examination of the offspring gonads of the putative XX neomale identified 7 fish with testes,
135 83 fish with ovaries, and 41 fish with undifferentiated gonads. Disregarding fish with
136 undifferentiated gonads suggests a sex ratio of 7.8% males and 92.2% females for the offspring
137 of the XX neomale (Table 1). Gonadal histology of the offspring of the putative XY revealed
138 48 animals with testes, 65 with ovaries, and 14 with undifferentiated gonads, which gives a sex
139 ratio of 42.5% male and 57.5% female, ignoring the offspring with undifferentiated gonads
140 (Table 1). These sex ratio differences (Table 1), strongly support the hypothesis that male
141 breeder 1 is an XX neomale with an offspring sex ratio not significantly different from an
142 expected all-female population with a slight percentage of female-to-male sex-reversal, and that
143 breeder 2 is a genetic XY male with an offspring sex ratio not significantly different from an
144 expected 50:50 sex ratio. In agreement with these results, none of the XX neomale offspring
145 produced a positive PCR amplification for our three Y-allele specific primer pairs (Figure S3,
146 Table 2), and all 48 phenotypic males but only one of 65 phenotypic females offspring from the
147 XY phenotypic male produced positive amplifications (Figure S4, Table 2). This result also
148 indicates that no neomales were detected in offspring from the XY genetic male if we do not
149 consider the undetermined individuals compared to the 7.8% of neomales in the XX neomale
150 offspring population.

151

152 **Characterization of the goldfish sex chromosome and sex-determining region (SDR)**

153 Using the three Y-allele specific primer pairs described above, we genotyped goldfish
154 individuals and selected 30 phenotypic and genotypic males that were used along with 30
155 phenotypic females to contrast whole genome sex differences by pool-sequencing analysis [29].

156 Pool-sequencing reads from the respective XY male and phenotypic female pools were mapped
157 to the high contiguity goldfish female genome assembly [6] to characterize genomic regions
158 enriched for sex-biased signals, i.e., sex coverage differences or sex-biased Single Nucleotide
159 Polymorphism (SNP) differences. Whole genome analysis of SNP distribution (Figure 2)
160 revealed a strong sex-linked signal in males on linkage group 22 (LG22) and two unplaced
161 scaffolds (NW_020523543.1 and NW_020523609.1) with a high density of observed SNPs
162 being heterozygous in the male pool and homozygous in the female pool (Y-specific allele).
163 Interestingly, of the 32 markers found using the RAD-Seq approach, 7 tags were enriched in
164 the unplaced scaffold NW_020523543.1 (Fig. 3C), confirming by a second approach that this
165 scaffold is part of the SD locus in goldfish. These regions with a high density of male-specific
166 SNPs (Figure 3) are potential sex-determining regions that could contain the goldfish master
167 sex determining gene. LG22, being the only linkage group with a large sex determining region
168 (SDR, highlighted by a black box on Fig. 3A, C, D) containing a high-density of male-specific
169 SNPs (~11.7 Mb), likely corresponds to the goldfish Y sex chromosome.

170 We also observed, however, some smaller signals with less dense sex-linked SNPs in other
171 linkage groups (Figure 2A) like for instance on LG47 (Fig. S1) with both male and female sex-
172 linked signals. Interestingly, LG47 is paralogous to LG22 stemming from the Cyprinidae whole
173 genome duplication [6]. Indeed, due to this recent common ancestry, these two chromosomes
174 share large homologous and syntenic regions (Fig. S2) that could have resulted in some false
175 remapping of the pool-sequencing reads leading to some of these secondary minor signals.

176

177 **Identification of candidate SD genes**

178 Searches for annotated genes by BLAST within the 20 contigs found in our male goldfish draft
179 genome assembly based on the RAD-Seq approach did not return any matches for a candidate
180 SD gene, but mostly transposable elements (Supplementary excel file 3). All genes within the
181 SDR (N= 373) were extracted because they are potential candidates for being SD gene(s)
182 (Supplemental excel file 4). Interestingly, among these genes the anti-Mullerian hormone gene
183 (*amh*) was found at the beginning of the SDR on LG22 (Fig. 3B). This gene has been reported
184 to be a sex-determining gene in other fish species [14, 15]. However, we did not identify any
185 male-specific SNPs in the coding sequence of goldfish *amh*. In addition, other male specific
186 alleles within the 5kb promoter region did not show any sex-linkage.

187

188 **DISCUSSION**

189 Though goldfish is an important economic ornamental fish and a useful model for studying
190 development, evolution, neuroscience, and human disease [3], characterization of goldfish sex-
191 specific sequences and potential sex chromosomes have not been reported. In this study, we
192 explored goldfish sex determination using two complementary whole-genome approaches and
193 found that this species has a XX/XY sex determination system as previously described [24] and
194 a large, non-recombining sex determination region on LG22. Although RAD-sequencing or
195 pool-sequencing have been often used separately to explore sex determination in vertebrates
196 [16, 30, 31], we choose to combine these two approaches in goldfish because of the significant
197 female-to-male sex reversal induced by temperature [25] that would have prevented a clear
198 identification of the sex determining region using only a pooled strategy, which mixes genetic
199 XY males and XX males resulting from the sex reversal of genetic females. Because RAD-

200 sequencing keeps track of each individual, we were able to identify sex-reversed individuals in
201 goldfish that might have masked sex-linked markers in Pool-seq.

202

203 Sex markers identification is an important step to characterize SD systems [32-38]. Using two
204 complementary whole-genome approaches, we characterized genomic regions containing sex-
205 linked markers. In goldfish, these sex-linked markers are genomic DNA variations including
206 gaps, indels and SNPs that present heterozygote polymorphisms in all males and complete
207 homozygosity in all females. This male-specific heterozygosity pattern agrees with a male
208 heterogametic XX/XY system as previously reported using progeny testing of hormonally sex-
209 reversed breeders [24]. We found, however, a strong environmental influence leading to a
210 relatively high proportion (around 50%) of female-to-male sex-reversal in the first experimental
211 population that we used for the RAD-Sequencing approach. These animals were actually two-
212 year old goldfish raised in an outdoor experimental facility and obtained at different spawning
213 times i.e., from May-June to late September. Some of these animals experienced early
214 development during summer time at potentially higher temperature and others had their early
215 developmental period at lower temperatures. Considering the known effects of high temperature
216 on female-to-male sex reversal in goldfish [25], the fact that some of these fish were exposed
217 to a high summer temperature could explain this relatively high percentage of female-to-male
218 sex-reversed animals. This high percentage was not found in our other experiments in which
219 fish were raised in indoor recirculating system facilities with a tightly controlled low
220 temperature (18°C) maintained throughout the whole early development phase (3 months). This
221 situation indeed confirms earlier findings showing that temperature is probably a major trigger

222 of neomasculinization in goldfish, but we also found that even at this low temperature there
223 was still a small percentage of female-to-male sex-reversal (7.8%), suggesting that other
224 environmental factors, potentially social factors as demonstrated in other species [8, 39], could
225 also play a role on goldfish sex determination. Apart from goldfish, sex determination in other
226 teleost fish, including Tilapia [40], medaka [41] and tongue sole [42] is also regulated by
227 temperature, which overrides the genetic sex determination mechanisms and leads to female-
228 to-male sex reversal. By developing genetic sexing tools in goldfish that allows the
229 identification of Y-allele carrying animals, we also brought additional evidence that some of
230 these phenotypic males were indeed sex-reversed XX genetic females. These genetic sexing
231 tools are indeed important for better deciphering genetic and environmental sex determination
232 in goldfish. But these PCR primers could be also used to facilitate the industrial production of
233 commercial goldfish-related hybrid fish in China [43, 44], by helping to identify neomales i.e.,
234 XX female-to-male sex reversed animals.

235

236 Sex determination in vertebrates is highly variable with the major exceptions of Eutherian
237 mammals and birds in which XX/XY and ZZ/ZW monofactorial sex determination systems
238 have been conserved over a long evolutionary period [45, 46]. In contrast, fish exhibit much
239 more diverse and dynamic sex determination [9, 10, 47], with monofactorial and polyfactorial
240 [48, 49] genetic systems and frequent switches and turnovers of master sex-determining genes
241 [12, 14, 15, 17, 21, 50]. In goldfish, we identified male-specific markers and obvious male-
242 specific SNPs strongly enriched on LG22. This result confirms that goldfish has an XX-XY
243 system [24] and also indicated that LG22 is the sex chromosome in that species. Evidence is

244 accumulating for the hypothesis that sex chromosomes, in most cases, evolve from autosomes
245 with *de novo* initial evolution of a new sex determination mechanism that subsequently
246 becomes fixed and extended by the suppression of recombination on the sex chromosome in
247 the vicinity of the initial sex locus, which may increase the size of this non recombining sex
248 determination locus [51]. In goldfish, ~11.7 Mb of LG22 contains numerous male-specific
249 SNPs. A similar large size of the non-recombining region on the sex chromosomes was also
250 found in tilapia including 17.9 Mb in *Sarotherodon melanotheron* and 10.7 Mb in *Oreochromis*
251 *niloticus* [30, 31]. The large non-recombining region on LG22 contains 373 gene models based
252 on the goldfish genome annotation and also a large number of transposable elements (TEs) that
253 were found to be strongly enriched in the male specific contigs identified by our RAD-Sex and
254 our draft genome analysis. Enrichment of TEs around sex loci has been found in other vertebrate
255 species [52] and may play a crucial role for suppression of recombination leading to an
256 expansion of sex chromosome divergence.

257

258 With LG22 being the potential sex chromosome in goldfish, it is reasonable to believe that the
259 non-recombining region that we characterize on LG22 contains the goldfish master sex
260 determining gene. But the only “usual suspect” master sex determining candidate found in this
261 region and the additional non-assembled scaffolds containing sex-linked markers is the anti-
262 Mullerian hormone gene (*amh*) that is located at the beginning of the LG22 non-recombining
263 region. Duplications of *amh* have been characterized as the master sex determining gene in
264 different fish species [14, 15], making *Amh* and members of the TGF-beta pathway [17, 19, 20]
265 likely candidates for this sex-determining function. But we have not been able to characterize

266 sex-linked variation neither in the *amh* coding DNA sequence nor in its 5 kb proximal promoter
267 sequence. Even if we cannot rule out the hypothesis that *amh* regulation could be affected by
268 sex-specific cis-regulatory elements located very far upstream from *amh*, our results do not
269 provide any clear and direct evidence that this gene is the goldfish master sex determining gene.
270 Indeed, not all master sex determining genes are classical “usual suspects” known to be
271 involved in the sex-differentiation pathway like TGF-beta members [17, 19, 53], Sox3 [21], or
272 Dmrt1[50, 54]. For instance, the rainbow trout master sex determining gene arose from the
273 duplication / transposition / evolution of an immune-related gene [12]. This finding suggests
274 that goldfish could also have an unusual master sex determining gene, preventing an easy and
275 direct identification just with simple genome-wide analyses and candidate gene approaches.

276

277 The goldfish genome, like the genomes of the common carp and other species of the cyprinid
278 subfamily cyprininae is characterized by a relatively recent whole genome duplication (WGD)
279 that occurred approximately 14 million years ago [6]. This WGD adds an extra complexity to
280 our search for sex-linked regions and sex determining candidate genes because some of these
281 duplicated regions may still retain large blocks of high sequence similarity. The cyprininae
282 genome duplication probably explains why we found an additional sex-biased signal on LG47
283 that stems from the duplication of the same ancestral chromosome that LG22. In addition to the
284 cyprininae WGD, the current goldfish reference genome sequence [6] was assembled from the
285 sequences of an XX gynogenetic animal, meaning that the LG22 sex chromosome sequence is
286 an X chromosome sequence in which potential Y specific regions may be not present. We
287 indeed produced a first draft genome sequence of an XY male but a higher contiguity male

288 genome including long-read technology would be needed to better explore sex-chromosome
289 differences and characterize potential sex-determining candidates.

290

291 **CONCLUSIONS**

292 Our results confirm that sex determination in goldfish is a complex mix of environmental and
293 genetic factors, and that its genetic sex determination system is male heterogametic (XX/XY).
294 We also characterized a relatively large non-recombining region (~11.7 Mb) on LG22 that is
295 likely to be the goldfish Y chromosome. This large non-recombining region on LG22 contains
296 a single obvious candidate as a potential master sex gene, namely the anti-Mullerian hormone
297 gene (*amh*). No sex-linked polymorphism, however, was detected in the goldfish *amh* gene and
298 its 5 kb proximal promoter sequence. Our work provides the foundation required for additional
299 studies that are now required to better characterize sex determination in goldfish and to
300 characterize its master sex-determining gene.

301

302 **MATERIALS AND METHODS**

303 **Experiment fish**

304 Fish used for RAD-seq and Pool-seq were reared outdoors and obtained from different
305 spawning times i.e., between May-June and late September. Putative XY and XX males were
306 selected using Y-allele specific primers and these two males were crossed with the same female
307 to produce two goldfish populations that were incubated and reared indoor at 18°C during three
308 months after fertilization to minimize the chance of sex reversal induced by temperature
309 according to previous research [25]. After these 3 months at 18°C, the rearing temperature was
310 gradually increased to 24°C over a period of 7 days to avoid suddenly dramatic temperature
311 variation. One-year old fish were euthanized with Tricaine before dissection. Gonads of
312 goldfish were fixed in Bouin's fixative solution for 24 hours and then embedded gonads were

313 cut serially into 7 μm sections and stained with Hematoxylin to characterize ovarian or testicular
314 features. Fin clips were stored in 90% alcohol for DNA extraction and genotyping. Statistics
315 were applied to test for significant sex ratio differences and genotype/phenotype sex-linkage
316 with a Chi-squared test ($p < 0.05$).

317

318 **DNA extraction and genotyping**

319 For genotyping, fin clips were lysed with 5% Chelex and 20 mg Proteinase K at 55°C for 2
320 hours, and subsequently denatured by Proteinase K at 99°C for 2 min. Supernatant containing
321 genomic DNA (gDNA) was collected to a new tube after a brief centrifugation. Finally, DNA
322 was diluted to half and stored at -20°C. For genome sequencing, gDNA was extracted with
323 NucleoSpin Kits for Tissue (Macherey-Nagel, Duren, Germany) following the manufacturer's
324 instructions. gDNA concentration and quality were measured with a NanoDrop ND2000
325 spectrophotometer (Thermo Scientific, Wilmington, DE) and a Qubit3 fluorometer (Invitrogen,
326 Carlsbad, CA).

327 Primers were designed from the sequences of male-biased contigs for sex genotyping and a
328 positive control (Table S1) based on contig flattened_line_0 from our Illumina male genome
329 assembly (Accession number: WSJC00000000) using Primer3 version 0.4.0
330 (<http://primer3.ut.ee>). PCRs were performed with 0.1 μM of each primer, 50 ng of gDNA
331 adjusted at 50 ng/ μl , 100 μM dNTP mixture, and 1 μl of 10 \times PCR Buffer (Sigma Aldrich) with
332 0.25 units of JumpStart Taq DNA Polymerase (Sigma Aldrich) in a total volume of 25 μl . The
333 PCR thermal cycle procedures were: 94°C for 30s for denaturing, 58°C for 30s for annealing
334 and 72°C for 30s for extending for 35 cycles. Finally, PCR products were electrophoresed on

335 1.5% agarose gels.

336

337 **Restriction-site association sequencing (RAD-seq) and male-marker discovery**

338 Genomic DNA was extracted from 30 males and 30 females and digest with the restriction
339 enzyme *SbfI* for constructing a RAD-seq library according to standard protocols [55]. Briefly,
340 for each sample, 1µg of DNA was digested using *SbfI*. Digested DNA was purified using
341 AMPure PX magnetic beads (Beckman Coulters) and ligated to indexed P1 adapters (one index
342 per sample) using concentrated T4 DNA ligase (NEB). Ligated DNA was purified using
343 AMPure XP magnetic beads. Each sample was quantified using microfluorimetry (Qubit
344 dsDNA HS assay kit, Thermofisher) and all samples were pooled in equal amount. The pool
345 was fragmented on a Biorputor (Diagenode) and purified using a Minelute column (Qiagen).
346 Sonicated DNA was size selected on an 1,5 % agarose cassette aiming for an insert size of 300
347 bp to 500 bp. Size selected DNA was extracted from the gel using the Qiaquick gel extraction
348 kit (Qiagen), repaired using the End-It DNA-end repair kit (Tebu Bio) and adenylated on its 3'
349 ends using Klenow (exo-) (Tebu-Bio). P2 adapter was ligated using concentrated T4 DNA
350 ligase (NEB) and 50 ng of the ligated product was engaged in a 12 cycles PCR. After AMPure
351 XP beads purification, the resulting library was checked on a Bioanalyzer (Agilent) using the
352 DNA 1000 kit and quantified by qPCR using the KAPA Library quantification kit (Roche, ref.
353 KK4824). The library was sequenced on one lane of Hiseq2500 in single read 100nt mode using
354 the clustering and SBS v3 kit following the manufacturer's instructions.

355 Raw reads were demultiplexed with the program *process_radtags.pl* of Stacks with default
356 settings. 135,019,110 (79.1%) reads were kept after this procedure. Demultiplexed reads were

357 subsequently processed by the RADSex software version 2.0.0
358 (<http://github.com/RomainFeron/RadSex>). The distribution of sequences between male and
359 female were calculated with function *distrib* with all settings to default. This distribution of
360 sequences was visualized with *plot_sex_distribution* function of *radsex-vis*
361 (<http://github.com/RomainFeron/RADSex-vis>) (Fig 1.A). Sequences significantly associated
362 with sex were extracted using the function *signif*, which identifies sex-bias tags.
363 Male-biased tags were compared to the male *de novo* assembly with *ncbi-blast+* (version: 2.6.0)
364 setting the e-value cutoff to 1^{e-20} to identify long, homologous male-biased contigs. Male
365 specific PCR primers were designed from these contigs sequences (see Table S1) using Primer3
366 version 0.4.0 (<http://primer3.ut.ee>).

367

368 **Pooled genome sequencing (Pool-seq) and sex differentiated region identification**

369 Genomic DNA extracted from the fin clips of 13 phenotypic females and 13 genotypic males
370 selected from the animals used for the RAD-Seq experiment, were used for the Pool-Seq
371 analysis. The 13 genotypic males were genotyped using the three Y-allele PCR primers
372 described above. Genomic DNA were pooled in equimolar ratio according to sex and Pool-
373 seq libraries were generated using the Truseq nano DNA sample prep kit (Illumina, ref. FC-
374 121-4001) following the manufacturer's instructions. Briefly, each pool was sonicated using a
375 Bioruptor (Diagenode). The sonicated pools were repaired, size selected on magnetic beads
376 aiming for a 550 pb insert size and adenylated on their 3' ends. Adenylated DNA was ligated to
377 Illumina's specific adapters and, after purification on magnetic beads, was amplified in an 8
378 cycles PCR. Libraries were purified using magnetic beads, checked on a Fragment Analyzer

379 (Agilent) using the HS NGS Fragment kit (DNF-474-33) and quantified by qPCR using the
380 KAPA Library quantification kit (Roche, ref. KK4824). Each library was sequenced on half a
381 lane of a rapid v2 flow cell (Illumina) in paired end 2x250nt mode.

382 Reads from the male and female pools were remapped to a genome sequence coming from a
383 gynogenesis-derived female [QPKE00000000] using BWA mem version 0.7.17 with default
384 parameters. Then, BAM files were sorted and merged with Picard tools version 2.18.2 with
385 default parameters. After that, PCR duplicates were removed with Picard tools. Reads with
386 mapping quality less than 20 and that did not map uniquely were also removed with Samtools
387 version 1.8. Subsequently, the two sex BAM files were used to generate a pileup file using
388 samtools mpileup with per-base alignment quality disabled (-B). A sync file was created using
389 popoolation mpileup2sync version 1.201 (parameters: --min-qual 20), which contains the
390 nucleotide composition of each sex for each position in the reference. Finally, with this sync
391 file, SNPs and coverage between the two sexes of all reference positions were overall calculated
392 with PSASS (version 2.0.0, doi:10.5281/zenodo.2615936). We used a 100kb sliding window
393 with an output point every 500bp to identify sex-specific SNPs enriched regions with PSASS.
394 The PSASS parameters were as follows: minimum depth set to 10 (--min-depth 10), range of
395 heterozygous SNP frequency for a sex-linked locus 0.5 ± 0.2 (--freq-het 0.5, --range-het 0.2),
396 homologous SNP frequency for a sex-linked locus >0.98 (--freq-hom 1, --range-hom 0.02),
397 overlapped sliding window (--window-size 100000, --output-resolution 500). Data
398 visualization was implemented with an R package ([http://github.com/RomainFeron/PSASS-](http://github.com/RomainFeron/PSASS-vis)
399 [vis](http://github.com/RomainFeron/PSASS-vis)).

400

401 **Sequencing and *de novo* assembly of a goldfish male genome**

402 One genetic male was selected for *de novo* assembly using the Y-specific primers described
403 above. Library was generated using the Truseq nano DNA sample prep kit (Illumina, ref. FC-
404 121-4001) following the manufacturer's instructions. Briefly, DNA from a single male
405 individual was sonicated using a Bioruptor (Diagenode). The sonicated DNA was repaired, size
406 selected on magnetic beads aiming for a 550 pb insert size and adenylated on its 3' ends.
407 Adenylated DNA was ligated to Illumina's specific adapters and, after purification on magnetic
408 beads, was amplified in an 8 cycles PCR. Library was purified using magnetic beads, checked
409 on a Fragment Analyzer (Agilent) using the HS NGS Fragment kit (DNF-474-33) and
410 quantified by qPCR using the KAPA Library quantification kit (Roche, ref. KK4824). The
411 library was sequenced on one lane of a rapid v2 flow cell (Illumina) in paired end 2*250nt
412 mode. Illumina paired-end reads were assembled using DiscoverDeNovo (reference
413 <https://software.broadinstitute.org/software/discover/blog/>) with standard parameters.

414

415 **ABBREVIATIONS:** RAD-seq: Restriction site-associated DNA sequencing; SNP: Single
416 nucleotide polymorphism; SD: Sex determination; SDR: Sex differentiated region, MSD:
417 master sex determining genes.

418

419 **DECLARATIONS**

420 **Ethics approval:** Research involving animal experimentation conformed to the principles for
421 the use and care of laboratory animals, in compliance with French ("National Council for
422 Animal Experimentation" of the French Ministry of Higher Education and Research and the

423 Ministry of Food, Agriculture, and Forest) and European (European Communities Council
424 Directive 2010/63/UE) guidelines on animal welfare.

425

426 **Consent for publication:** Not applicable

427

428 **Availability of data and material:** This Whole Genome Shotgun project has been deposited
429 at DDBJ/ENA/GenBank under the accession WSJC00000000. The version described in this
430 paper is version WSJC01000000. Genome sequencing reads of the male genome, the male and
431 female pool-sequencing reads and the RAD-seq demultiplexed sequences have been deposited
432 in the Sequence Read Archive (SRA), under BioProject PRJNA592334.

433

434 **Competing interests:** The authors declare that they have no competing interests.

435

436 **Funding:** This project was supported by funds from the “Agence Nationale de la Recherche”
437 and the “Deutsche Forschungsgemeinschaft” (ANR/DFG, PhyloSex project, 2014-2016) to MS
438 and YG. Montpellier Genomics (MGX) facility was supported by France Génomique National
439 infrastructure, funded as part of “Investissement d’avenir” program managed by Agence
440 Nationale pour la Recherche (contract ANR-10-INBS-09). This work was also supported by
441 Grant-in-Aid for Scientific Research (19K22426) to YO, and grants R01OD011116 and
442 5R01GM085318 from the USA National Institutes of Health to JHP.

443

444 **Authors' contributions:**

445 Conceived and designed the experiments: YG, MW

446 Funding acquisition: YG, MS, JP, LJ

447 Investigation: MW, MP, JG, EJ, AH, CR, HP, SB, YO

448 Bioinformatics analysis: RF, CK, CC, MZ

449 Visualization: MW

450 Wrote the paper: MW, YG

451

452 **Acknowledgements:** We are grateful to the Genotoul bioinformatics platform Toulouse Midi-

453 Pyrenees (Bioinfo Genotoul) for providing computing and/or storage resources and to the

454 INRA-LPGP experimental facilities for taking care of goldfish experiments.

455

456 **Supplementary information**

457 **Supplementary excel file 1:** Sequences of putative Y-allele RAD-tags (N= 32) found in some

458 males but absent from all females.

459 **Supplementary excel file 2:** Contigs from a goldfish Illumina male genome assembly with

460 homologies with the putative Y-allele RAD-tags.

461 **Supplementary excel file 3:** Annotation of potential Y chromosome contigs by sequence

462 comparisons to NCBI Non-redundant protein sequence database using blastx.

463 **Supplementary excel file 4:** Detailed information of annotated genes in the goldfish sex

464 determination regions extracted from the NCBI genome annotation file (accession number

465 QPKE00000000).

REFERENCES

1. Blanco AM, Sundarrajan L, Bertucci JI, Unniappan S: **Why goldfish? Merits and challenges in employing goldfish as a model organism in comparative endocrinology research.** *General and comparative endocrinology* 2018, **257**:13-28.
2. Popesku JT, Martyniuk CJ, Mennigen J, Xiong H, Zhang D, Xia X, Cossins AR, Trudeau VL: **The goldfish (*Carassius auratus*) as a model for neuroendocrine signaling.** *Molecular and cellular endocrinology* 2008, **293**(1-2):43-56.
3. Omori Y, Kon T: **Goldfish: an old and new model system to study vertebrate development, evolution and human disease.** *J Biochem* 2019, **165**(3):209-218.
4. Ota KG, Abe G: **Goldfish morphology as a model for evolutionary developmental biology.** *Wiley Interdisciplinary Reviews: Developmental Biology* 2016, **5**(3):272-295.
5. Choe Y, Yu JE, Park J, Park D, Oh J-I, Kim S, Moon KH, Kang HY: **Goldfish, *Carassius auratus*, as an infection model for studying the pathogenesis of *Edwardsiella piscicida*.** *Veterinary research communications* 2017, **41**(4):289-297.
6. Chen Z, Omori Y, Koren S, Shirokiya T, Kuroda T, Miyamoto A, Wada H, Fujiyama A, Toyoda A, Zhang S: **De novo assembly of the goldfish (*Carassius auratus*) genome and the evolution of genes after whole-genome duplication.** *Science Advances* 2019, **5**(6):eaav0547.
7. Mohammad T, Moulick S, Mukherjee CK: **Economic feasibility of goldfish (*Carassius auratus* Linn.) recirculating aquaculture system.** *Aquaculture Research* 2018, **49**(9):2945-2953.
8. Devlin RH, Nagahama Y: **Sex determination and sex differentiation in fish: an overview of genetic, physiological, and environmental influences.** *Aquaculture* 2002, **208**(3-4):191-364.
9. Pan Q, Anderson J, Bertho S, Herpin A, Wilson C, Postlethwait JH, Schartl M, Guiguen Y: **Vertebrate sex-determining genes play musical chairs.** *C R Biol* 2016, **339**(7-8):258-262.
10. Herpin A, Schartl M: **Plasticity of gene-regulatory networks controlling sex determination: of masters, slaves, usual suspects, newcomers, and usurpaters.** *EMBO reports* 2015, **16**(10):1260-1274.
11. Matsuda M, Nagahama Y, Shinomiya A, Sato T, Matsuda C, Kobayashi T, Morrey CE, Shibata N, Asakawa S, Shimizu N: **DMY is a Y-specific DM-domain gene required for male development in the medaka fish.** *Nature* 2002, **417**(6888):559.
12. Yano A, Guyomard R, Nicol B, Jouanno E, Quillet E, Klopp C, Cabau C, Bouchez O, Fostier A, Guiguen Y: **An immune-related gene evolved into the master sex-determining gene in rainbow trout, *Oncorhynchus mykiss*.** *Curr Biol* 2012, **22**(15):1423-1428.
13. Hattori RS, Murai Y, Oura M, Masuda S, Majhi SK, Sakamoto T, Fernandino JI, Somoza GM, Yokota M, Strüssmann CA: **A Y-linked anti-Müllerian hormone duplication takes over a critical role in sex determination.** *Proceedings of the National Academy of Sciences* 2012, **109**(8):2955-2959.
14. Li M, Sun Y, Zhao J, Shi H, Zeng S, Ye K, Jiang D, Zhou L, Sun L, Tao W *et al*: **A Tandem Duplicate of Anti-Müllerian Hormone with a Missense SNP on the Y**

-
- Chromosome Is Essential for Male Sex Determination in Nile Tilapia, *Oreochromis niloticus*.** *PLOS Genetics* 2015, **11**(11):e1005678.
15. Pan Q, Feron R, Yano A, Guyomard R, Jouanno E, Vigouroux E, Wen M, Busnel J-M, Bobe J, Concordet J-P: **Identification of the master sex determining gene in Northern pike (*Esox lucius*) reveals restricted sex chromosome differentiation.** *BioRxiv* 2019:549527.
 16. Feron R, Zahm M, Cabau C, Klopp C, Roques C, Bouchez O, Eche C, Valiere S, Donnadiou C, Haffray P: **Characterization of a Y-specific duplication/insertion of the anti-Mullerian hormone type II receptor gene based on a chromosome-scale genome assembly of yellow perch, *Perca flavescens*.** *BioRxiv* 2019:717397.
 17. Kamiya T, Kai W, Tasumi S, Oka A, Matsunaga T, Mizuno N, Fujita M, Suetake H, Suzuki S, Hosoya S *et al*: **A trans-species missense SNP in *Amhr2* is associated with sex determination in the tiger pufferfish, *Takifugu rubripes* (fugu).** *PLoS Genet* 2012, **8**(7):e1002798.
 18. Rondeau EB, Messmer AM, Sanderson DS, Jantzen SG, von Schalburg KR, Minkley DR, Leong JS, Macdonald GM, Davidsen AE, Parker WA: **Genomics of sablefish (*Anoplopoma fimbria*): expressed genes, mitochondrial phylogeny, linkage map and identification of a putative sex gene.** *BMC genomics* 2013, **14**(1):452.
 19. Myosho T, Otake H, Masuyama H, Matsuda M, Kuroki Y, Fujiyama A, Naruse K, Hamaguchi S, Sakaizumi M: **Tracing the emergence of a novel sex-determining gene in medaka, *Oryzias luzonensis*.** *Genetics* 2012, **191**(1):163-170.
 20. Reichwald K, Petzold A, Koch P, Downie Bryan R, Hartmann N, Pietsch S, Baumgart M, Chalopin D, Felder M, Bens M *et al*: **Insights into Sex Chromosome Evolution and Aging from the Genome of a Short-Lived Fish.** *Cell* 2015, **163**(6):1527-1538.
 21. Takehana Y, Matsuda M, Myosho T, Suster ML, Kawakami K, Shin IT, Kohara Y, Kuroki Y, Toyoda A, Fujiyama A *et al*: **Co-option of *Sox3* as the male-determining factor on the Y chromosome in the fish *Oryzias dancena*.** *Nat Commun* 2014, **5**:4157.
 22. Matsuda M, Sakaizumi M: **Evolution of the sex-determining gene in the teleostean genus *Oryzias*.** *General and comparative endocrinology* 2016, **239**:80-88.
 23. Ospina-Alvarez N, Piferrer F: **Temperature-dependent sex determination in fish revisited: prevalence, a single sex ratio response pattern, and possible effects of climate change.** *PloS one* 2008, **3**(7):e2837.
 24. Yamamoto TO, Kajishima T: **Sex hormone induction of sex reversal in the goldfish and evidence for male heterogamity(XX/XY).** *Journal of Experimental Zoology* 1968, **168**(2):215-221.
 25. Goto-Kazeto R, Abe Y, Masai K, Yamaha E, Adachi S, Yamauchi K: **Temperature-dependent sex differentiation in goldfish: Establishing the temperature-sensitive period and effect of constant and fluctuating water temperatures.** *Aquaculture* 2006, **254**(1-4):617-624.
 26. Chen Z, Omori Y, Koren S, Shirokiya T, Kuroda T, Miyamoto A, Wada H, Fujiyama A, Toyoda A, Zhang S *et al*: **De Novo assembly of the goldfish (*Carassius auratus*) genome and the evolution of genes after whole genome duplication.** 2018.
 27. Anderson JL, Mari AR, Braasch I, Amores A, Hohenlohe P, Batzel P, Postlethwait JH: **Multiple sex-associated regions and a putative sex chromosome in zebrafish**

-
- revealed by RAD mapping and population genomics. *PloS one* 2012, **7**(7):e40701.
28. Gamble T: **Using RAD-seq to recognize sex-specific markers and sex chromosome systems.** *Molecular ecology* 2016, **25**(10):2114-2116.
29. Schlötterer C, Tobler R, Kofler R, Nolte V: **Sequencing pools of individuals—mining genome-wide polymorphism data without big funding.** *Nature Reviews Genetics* 2014, **15**(11):749-763.
30. Gammerdinger WJ, Conte MA, Baroiller J-F, D’cotta H, Kocher TD: **Comparative analysis of a sex chromosome from the blackchin tilapia, *Sarotherodon melanotheron*.** *BMC genomics* 2016, **17**(1):808.
31. Gammerdinger WJ, Conte MA, Acquah EA, Roberts RB, Kocher TD: **Structure and decay of a proto-Y region in Tilapia, *Oreochromis niloticus*.** *Bmc Genomics* 2014, **15**(1):975.
32. Kincaid-Smith J, Boissier J, Allienne JF, Oleaga A, Djuikwo-Teukeng F, Toulza E: **A Genome Wide Comparison to Identify Markers to Differentiate the Sex of Larval Stages of *Schistosoma haematobium*, *Schistosoma bovis* and their Respective Hybrids.** *PLoS Negl Trop Dis* 2016, **10**(11):e0005138.
33. Charlesworth D, Mank JE: **The birds and the bees and the flowers and the trees: lessons from genetic mapping of sex determination in plants and animals.** *Genetics* 2010, **186**(1):9-31.
34. Pan ZJ, Li XY, Zhou FJ, Qiang XG, Gui JF: **Identification of Sex-Specific Markers Reveals Male Heterogametic Sex Determination in *Pseudobagrus ussuriensis*.** *Mar Biotechnol (NY)* 2015, **17**(4):441-451.
35. Carmichael SN, Bekaert M, Taggart JB, Christie HR, Bassett DI, Bron JE, Skuce PJ, Gharbi K, Skern-Mauritzen R, Sturm A: **Identification of a sex-linked SNP marker in the salmon louse (*Lepeophtheirus salmonis*) using RAD sequencing.** *PLoS One* 2013, **8**(10):e77832.
36. Kafkas S, Khodaeiaminjan M, Guney M, Kafkas E: **Identification of sex-linked SNP markers using RAD sequencing suggests ZW/ZZ sex determination in *Pistacia vera* L.** *BMC Genomics* 2015, **16**:98.
37. Gamble T, Zarkower D: **Identification of sex-specific molecular markers using restriction site-associated DNA sequencing.** *Mol Ecol Resour* 2014, **14**(5):902-913.
38. Fowler BL, Buonaccorsi VP: **Genomic characterization of sex-identification markers in *Sebastes carnatus* and *Sebastes chrysomelas* rockfishes.** *Mol Ecol* 2016, **25**(10):2165-2175.
39. Kobayashi Y, Nagahama Y, Nakamura M: **Diversity and plasticity of sex determination and differentiation in fishes.** *Sex Dev* 2013, **7**(1-3):115-125.
40. Wessels S, Hörstgen-Schwark G: **Temperature dependent sex ratios in selected lines and crosses with a YY-male in Nile tilapia (*Oreochromis niloticus*).** *Aquaculture* 2011, **318**(1-2):79-84.
41. Hattori R, Gould R, Fujioka T, Saito T, Kurita J, Strüssmann C, Yokota M, Watanabe S: **Temperature-dependent sex determination in Hd-rR medaka *Oryzias latipes*: gender sensitivity, thermal threshold, critical period, and DMRT1 expression profile.** *Sexual Development* 2007, **1**(2):138-146.
42. Shao C, Li Q, Chen S, Zhang P, Lian J, Hu Q, Sun B, Jin L, Liu S, Wang Z: **Epigenetic**

-
- modification and inheritance in sexual reversal of fish.** *Genome research* 2014, **24**(4):604-615.
43. Wen M, Peng L, Hu X, Zhao Y, Liu S, Hong Y: **Transcriptional quiescence of paternal mtDNA in cyprinid fish embryos.** *Scientific reports* 2016, **6**:28571.
44. Xu K, Wen M, Duan W, Ren L, Hu F, Xiao J, Wang J, Tao M, Zhang C, Wang J: **Comparative analysis of testis transcriptomes from triploid and fertile diploid cyprinid fish.** *Biology of reproduction* 2015, **92**(4):95, 91-12.
45. Wallis M, Waters P, Graves J: **Sex determination in mammals—before and after the evolution of SRY.** *Cellular and molecular life sciences* 2008, **65**(20):3182.
46. Ellegren H: **Evolution of the avian sex chromosomes and their role in sex determination.** *Trends in Ecology & Evolution* 2000, **15**(5):188-192.
47. Mank JE, Avise JC: **Evolutionary Diversity and Turn-Over of Sex Determination in Teleost Fishes.** *Sexual Development* 2009, **3**(2-3):60-67.
48. Roberts NB, Juntti SA, Coyle KP, Dumont BL, Stanley MK, Ryan AQ, Fernald RD, Roberts RB: **Polygenic sex determination in the cichlid fish *Astatotilapia burtoni*.** *BMC Genomics* 2016, **17**:835.
49. Liew WC, Bartfai R, Lim Z, Sreenivasan R, Siegfried KR, Orban L: **Polygenic Sex Determination System in Zebrafish.** *PLOS ONE* 2012, **7**(4):e34397.
50. Cui Z, Liu Y, Wang W, Wang Q, Zhang N, Lin F, Wang N, Shao C, Dong Z, Li Y: **Genome editing reveals *dmrt1* as an essential male sex-determining gene in Chinese tongue sole (*Cynoglossus semilaevis*).** *Scientific reports* 2017, **7**:42213.
51. Wright AE, Dean R, Zimmer F, Mank JE: **How to make a sex chromosome.** *Nat Commun* 2016, **7**:12087.
52. Chalopin D, Volff JN, Galiana D, Anderson JL, Scharl M: **Transposable elements and early evolution of sex chromosomes in fish.** *Chromosome Res* 2015, **23**(3):545-560.
53. Rondeau EB, Laurie CV, Johnson SC, Koop BF: **A PCR assay detects a male-specific duplicated copy of Anti-Müllerian hormone (*amh*) in the lingcod (*Ophiodon elongatus*).** *BMC research notes* 2016, **9**(1):230.
54. Nanda I, Kondo M, Hornung U, Asakawa S, Winkler C, Shimizu A, Shan Z, Haaf T, Shimizu N, Shima A *et al*: **A duplicated copy of *DMRT1* in the sex-determining region of the Y chromosome of the medaka, *Oryzias latipes*.** *Proc Natl Acad Sci U S A* 2002, **99**(18):11778-11783.
55. Baird NA, Etter PD, Atwood TS, Currey MC, Shiver AL, Lewis ZA, Selker EU, Cresko WA, Johnson EA: **Rapid SNP discovery and genetic mapping using sequenced RAD markers.** *PLoS One* 2008, **3**(10):e3376.

TABLES:

Table 1. Statistics of phenotypic sex in two populations

Population	N. of male	N. of female	N. of fish with undetermined sex	Sex ratio: %♂/♀	Significance ($\neq 0\%♂/100\%♀$)	Significance ($\neq 50\%♂/50\%♀$)
P(XX)	7	83	41	8.4	*	***
P(XY)	48	65	14	73.8	***	NS

P(XX): putative neomale (XX) offspring population; P(XY): putative genetic male (XY) offspring population; NS: Non-significant.

Chi-squared test was applied for statistics in R

Table 2. Goldfish Y-allele sex-linkage

Population	Male [#]	Female [#]	Undetermined sex [#]	Sex linkage
P(XX)	0 / 7	0 / 83	0 / 41	NS
P(XY)	48 / 48	1 / 65	10 / 14	***

[#] Y-allele positive genotyping / total number of samples. P(XX): putative neomale (XX) offspring population; P(XY): putative genetic male (XY) offspring population; NS: Non-significant. Fisher's exact test was applied for statistics in R

Table S1. Sequences of the primers used for Y-allele genotyping in goldfish.

Primers		PCR product (bp)	Genome location	
names	Sequence(5' - 3')		Male assembly	NCBI_genome
Marker 1	Forward: AATACAACATTCCCAGGGAGTGCA Reverse: CATCAAGGGCTATCTGACCAAGA	1169	Flattened_line_39456 0:620-1788	NW_020523543.1
Marker 2	Forward: GTGCTCAATAGACGACGGATTCTC Reverse: GTCTGTCTGTTAGCCTGTTCTCCA	1189	Flattened_line_27079 8:2006-3194	NW_020525535.1
Marker 3	Forward: GATGAAGGTCTCGGTCTGTTGTTA Reverse: CCCTGTTATGTTTGTATTGGCTAC	2548	Flattened_line_35862: 4409-6956	NC_039250.1 (LG8)
Positive control	Forward: AAGAGCGCCTCCTAGTGTTT Reverse: GAGACGGAGGAGTGGTATCG	994	Flattened_line_0:6858 -7842	NC_039245.1 (LG3)

Three Y-allele primer pairs (marker 1 to 3) and one autosomal primer pair (positive control) were designed on our XY male genome assembly (male assembly). Name of the contig and nucleotide position (3'-5') are given in the genome location column.

FIGURES:

Figure 1




Figure 1. RAD-sex tags and male-specific markers in goldfish. (A) Haplotypes heatmap in phenotypic males and females' goldfish. Each cell in the heatmap represents the number of haplotypes presented in x phenotypic males and y phenotypic females (x: cumulative number of males, y: cumulative number of females). Haplotypes present in more than 12 males and absent in all females were identified as male-specific haplotypes (highlighted by red box). **(B)** Genotyping of goldfish males and females with three Y-allele primer pairs and one autosomal primer used as a positive control. Goldfish are categorized into three groups i.e., putative genetic males (XY), putative XX neomales, and genetic females by combining the results of both Y-allele genotyping and sex phenotyping.

Figure 2




Figure 2. Sex determining regions identified by remapping the Pool-seq male and female reads onto the female genome assembly. SNPs were counted using 100kb sliding window with an output point every 500bp. **(A)** Circular plot showing the genome wide metrics of the Pool-seq analysis. All the 50 goldfish linkage groups (LGs) are labelled with their LG number and all unplaced scaffolds are fused together. Outer to inner tracks show respectively: the male-specific SNPs, the female-specific SNPs, and the reads depth ratio between males and females. **(B)** Manhattan plot of the male- and female-specific SNPs showing a strong enrichment of male-specific SNPs on LG22.

Figure 3




Figure 3. Distribution of male-specific SNPs on LG22 and unplaced scaffolds NW_020523543.1 and NW_020523609.1. SNPs were counted using 100kb sliding window with an output point every 500bp and female- and male-specific SNPs were respectively indicated by red and blue color. (A) A large sex-determination region was identified on LG22, which is highlighted with a black box. The candidate sex-determining gene *amh* is located on this LG22, but not in the high density, male-specific SNP region. The region from 8Mb to 10Mb containing *amh* is zoomed in panel (B). (C) The NW_020523543.1 unplaced scaffold exhibits a region around 0.1Mb harboring a small region (200 kb) with a high-density of male-specific SNPs. Meanwhile, sequence comparisons demonstrate that 7 male-biased RAD-tags (colored circles) on a total of 32 map with a high identity onto this scaffold. In contrast, few female-specific SNPs were enriched on this scaffold (red area). (D) The unplaced NW_020523609.1 scaffold is enriched in male-specific SNPs.

SUPPLEMENTARY FIGURES

Figure S1




Figure S1: Distribution of sex-biased SNPs on LG47. SNPs were counted using 100kb sliding window with an output point every 500bp. The top panel displays the profile of male-specific SNPs (blue area), while the bottom panel displays the profile of female-specific SNPs (red area).

Figure S2




Figure S2: Dot plot comparison of LG22 and LG47 showing conserved synteny between these two linkage groups.

Figure S3



Figure S3: Sex genotyping with Y-allele primers of the offspring of a putative XX neomale with a normal XX female. Genotyping was conducted with three Y-allele primers and one autosomal primer used as a gDNA quality control. Phenotypic sex was determined by gonadal histology and males and females are shown using red and yellow color respectively. Female-to-male sex-reversed animals (N= 7) are highlighted by red boxes. Hashes indicate animals with unknown phenotypic sex with undifferentiated gonads based on histology.

Figure S4




Figure S4. Sex genotyping with Y-allele primers of the offspring of a putative XY male with a normal XX female. Genotyping was conducted with three Y-allele primers and one autosomal primer used as a gDNA quality control. Phenotypic sex was determined by gonadal histology and males and females are shown using red and yellow color respectively. The female-to-male sex-reversed animal (N= 1) is highlighted by a red box. Hashes indicate animals with unknown phenotypic sex with undifferentiated gonads based on histology.