1 ***Staphylococcus aureus* staphyloxanthin expression is not controlled by Hfq**

2

3 Wenfeng Liu, Pierre Boudry, Chantal Bohn and Philippe Boulloc*

4 Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Université Paris-Sud,

5 Université Paris-Saclay, 91198, Gif-sur-Yvette cedex, France

6 * Corresponding author: philippe.boulloc@cnrs.fr

7

8 **ABSTRACT**

9 **Objective:** The golden color of *Staphylococcus aureus* is due to the synthesis of carotenoid
10 pigments. In Gram-negative bacteria, Hfq is a global posttranscriptional regulator, but its
11 function in *S. aureus* remains obscure. The absence of Hfq in *S. aureus* was reported to
12 correlate with production of carotenoid pigment leading to the conclusion that Hfq was a
13 negative regulator of the yellow color. However, we reported the construction of *hfq* mutants
14 in several *S. aureus* strains and never noticed any color change; we therefore revisited the
15 question of Hfq implication in *S. aureus* pigmentation.

16 **Results:** The absence or accumulation of Hfq does not affect *S. aureus* pigmentation.

17 **Keywords (6):**

18 *Staphylococcus aureus*, Hfq, pigmentation, staphyloxanthin, regulation.

19

20

21 INTRODUCTION.

22 *Staphylococcus aureus* is a major pathogen responsible for numerous diseases from minor
 23 skin infection to septicemia, affecting humans and other animals. Its name “*aureus*” comes
 24 from the golden color of strains that express carotenoid pigments [1]. These pigments
 25 contribute to oxidative stress and neutrophil resistance, and virulence [2]. The carotenoid
 26 biosynthetic operon (*crtMNOPQ*) leading to the synthesis of staphyloxanthin regulated by σ^B
 27 [3, 4], an alternative σ factor that also controls a large number of general stress genes. σ^B
 28 activity depends on RsbU, its positive regulator [5, 6]. Numerous strains, including the *S.*
 29 *aureus* model NCTC8325, have *rsbU* mutations that prevent σ^B activity and *crt* operon
 30 expression, such that colonies are white. In addition, mutations in 37 genes were show to
 31 result in the loss of a yellow pigmentation [5, 7].

32 Hfq is an RNA chaperone needed for activity of numerous regulatory RNAs in Gram-negative
 33 bacteria [8]. However, its role in Gram-positive bacteria, with the exception of *Clostridium*
 34 *difficile* [9], remains enigmatic [10]. Hfq functionality from different species is often tested by
 35 interspecies complementation tests. However, expression of *hfq* genes from Gram-positive
 36 bacteria *S. aureus* and *Bacillus subtilis* in *Salmonella* could not compensate the absence of
 37 endogenous *hfq*, indicating a functional difference between Gram positive and negative Hfq
 38 [11, 12].

39 We previously compared phenotypes of *S. aureus hfq* mutants with their isogenic parental
 40 strains and observed no detectable difference associated with the absence of Hfq in the
 41 tested conditions [13]. However, our results were partly challenged by a publication reporting
 42 that carotenoid pigment production was increased in *hfq*-negative strains [14]. Here we use
 43 nine different *S. aureus* strains to show that Hfq absence or overexpression has no effect on
 44 pigment expression.

45

46 MAIN TEXT

47 Methods

48 Bacterial strains, plasmids and growth conditions

49 Bacterial strains, plasmids and primers used in this study are listed in Table 1. Allelic
50 replacements of *hfq*⁺ by $\Delta hfq::cat$ were either performed by λ 11-phage mediated
51 transduction using RN4220 *hfq::cat* as a donor strain or by homologous recombination using
52 pMAD $\Delta hfq::cat$ [13, 15]. The $\Delta hfq::cat$ deletion in SAPHB5 was verified by Southern blot and
53 subsequent $\Delta hfq::cat$ transductants were verified by PCR as described [13].

54 Engineered plasmids were constructed as described [16]. Conditional *hfq* expression was
55 obtained by cloning *hfq* under the xyl/tetO promoter in pRMC2 [17] and pRMC2FLAG (Table
56 1). pRMC2Hfq allowing *hfq* conditional expression was obtained as follows: pRMC2 and
57 PCR-amplified *hfq* (using primers 39/49 on HG003 DNA) were KpnI-EcoRI digested and
58 ligated together. pRMC2FLAG was engineered for conditional expression of 3xFLAG-tagged
59 proteins as followed: pRMC2 and pSUB11 [18] were PCR-amplified using primers 856/918
60 and 858/919, respectively. The two resulting products, *i.e.* pRMC2 and *3xflag* coding
61 sequence, were assembled using the Gibson method [19]. pRMC2HfqFLAG, allowing
62 conditional expression of Hfq::3xFLAG, was obtained as follows: pRMC2FLAG and *hfq*
63 HG003 were PCR-amplified using primers 918/865 and 939/940, respectively. The two
64 resulting products were assembled using the Gibson method.

65 Bacteria were grown in BHI medium at 37°C. For strains containing pRMC2 and derivatives,
66 chloramphenicol 5 $\mu\text{g} \cdot \text{ml}^{-1}$ was added to media. Expression from pRMC2 and derivatives was
67 achieved by anhydrotetracycline (aTc) 250 $\text{ng} \cdot \text{ml}^{-1}$ addition to growth media.

68 Protein extraction and Western blotting

69 Overnight cultures were diluted 1000 times in fresh medium. After 3h, aTc was added. 10
70 min and 30min later, cells were harvested by centrifugation (16.000 g for 2 min),
71 resuspended in 400 μl Tris HCl buffer (50 mM, pH 6.8) and lysed using a FastPrep (3 cycles
72 of 45 sec at 6.5 $\text{m} \cdot \text{s}^{-1}$). Cell debris was removed by centrifugation (16.000 g for 10 min).
73 Protein concentration was determined by Bradford assays [20]. For each sample, 3 μg of

74 protein extract was separated on a polyacrylamide gel (Blot™ 4-12% Bis-Tris Plus,
75 Invitrogen). After electrophoresis, proteins were transferred to a polyvinylidene fluoride
76 membrane (iBlot 2 PVDF Mini Stacks, Invitrogen). For blotting and washing, an iBind™ Flex
77 Western System was used according to supplier's instructions. Membranes were probed with
78 the primary polyclonal ANTI-FLAG antibody produced in rabbit (Sigma) at a 1/15,000 dilution.
79 A rabbit secondary antibody conjugated to horseradish peroxidase (Advansta) was used at a
80 1/25,000 dilution. Bioluminescent signal was detected with the WesternBright™ ECL-spray
81 (Advansta) using a digital camera (ImageQuant™ 350, GE Healthcare).

82

83 **Results**

84 **The absence of Hfq does not alter *S. aureus* pigmentation**

85 In 2010, Liu *et al.* reported that “deletion of *hfq* gene in *S. aureus* 8325-4 can increase the
86 surface carotenoid pigments” [14]. Their work was performed using an allele called Δhfq -
87 8325 in which the *hfq* coding sequence was replaced by a kanamycin cassette. The *hfq*
88 chromosomal deletion was constructed in strain RN4220 and then transduced into
89 NCTC8325-4, RN6390, COL and ATCC25923 by phage ϕ 11. We constructed a similar *hfq*
90 deletion in RN4220, except that the *hfq* coding sequence was replaced by a chloramphenicol
91 resistant gene ($\Delta hfq::cat$); this allele was transduced into RN6390, COL and Newman by
92 ϕ 11-phage mediated transduction [13]. Note that RN4220, RN6390 and COL strains were
93 used for both studies. As we did not notice a change of color when the $\Delta hfq::cat$ allele was
94 introduced into these strains, this information was not reported [13]. In view of the previous
95 report, we focused this work on the possibility that Hfq could affect *S. aureus* pigment
96 expression.

97 NCTC8325 isolated in 1960 from a sepsis patient is the progenitor of numerous strains
98 including NCTC8325-4 (cured of three prophages) which itself gave RN6390 and RN4220
99 [21]. As these descendants were mutagenized, they carry several mutations that may affect
100 their phenotypes. NCTC8325 has a deletion of 11 bp in *rsbU* and a point mutation in *tcaR*.
101 The derivatives HG001 (*rsbU* restored), HG002 (*tcaR* restored), HG003 (*rsbU* and *tcaR*

restored) were constructed to perform physiological studies in a non-mutagenized background [21]. All these strains derived from NCTC8325, except HG001 and HG003 (which have a functional σ^B factor), give rise to white colonies (Figure 1). In addition to those reported [13], we constructed $\Delta hfq::cat$ derivatives in NCTC8325, NCTC8325-4, HG001, HG002 and HG003 (Table 1). In contrast to results reported in Liu *et al.*, deletion of the *hfq* gene in all tested strain backgrounds had no effect on pigmentation (Figure 1A). Note that COL, Newman are not NCTC8325 derivatives.

Spectral profiles highlighting *S. aureus* carotenoid production were determined as described [1] for three strains and their *hfq* derivatives after growth for 24h in BHI. HG003 and HG003 $\Delta hfq::cat$ gave equivalent profiles with three pics characteristic of carotenoid production. In contrast, NCTC8325-4 and RN1 had spectra characteristic of no or very little carotenoid production. As expected from our visual observation (Figure 1A), the spectra of Δhfq derivatives did not differ from those of their respective parental strains (Figure 1B).

Hfq overexpression does not alter *S. aureus* pigmentation

In the above-described strains, *hfq* is possibly poorly expressed, in which case *hfq* deletions would not lead to detectable phenotypes. We therefore tested the effects of an inducible Hfq expression system on pigment production. If the absence of Hfq leads to yellow colonies as proposed [14], the presence of Hfq could lower pigment production and lead to white colonies. To address this point, *hfq* was cloned under the control of the $P_{xyl/tetO}$ promoter in multi-copy plasmid pRMC2 [17] leading to pRMC2Hfq. *hfq* expression in strains harboring pRMC2Hfq was induced upon aTc addition to media. To confirm that $P_{xyl/tetO}$ was effectively driving *hfq* expression, a pRMC2Hfq derivative was engineered harboring a *3xflag* sequence inserted in frame at the end of the *hfq* open reading frame. The resulting plasmid, pRMC2HfqFLAG is a proxy for expression from pRMC2Hfq. HG003 was transformed with pRMC2, pRMC2Hfq and pRMC2HfqFLAG. The protein Hfq::3xFLAG was detected upon aTc induction by western blotting using FLAG antibodies (Figure 2A). We inferred from this result that addition of aTc to strains harboring pRMC2Hfq lead to Hfq synthesis. The RN4220 white

and HG003 yellow colors were not affected by the presence of either pRMC2, pRMC2Hfq or pRMC2HfqFLAG and remained identical upon aTc addition to growth medium (Figure 2B).

Conclusion

Our results show that neither the absence, nor the accumulation of Hfq affects pigmentation of *S. aureus*: Hfq does not appear to regulate staphyloxanthin synthesis. Our conclusions are supported by E.J. Tarrant PhD dissertation showing an NCTC8325 *hfq* mutant that remained unpigmented [22]. Of note, *Pseudomonas aeruginosa* reportedly induces pigment production of *S. aureus*, however, this effect was independent of *hfq* transcription [23]. In addition, color variation in USA300 strain was screened in a genome-wide transposon mutant library, and the *hfq* inactivation was not reported to affect *S. aureus* pigmentation [7].

While the *hfq* gene is absent in some Firmicutes (e.g. Lactobacillales), it is conserved in all *S. aureus*, suggesting that it plays a crucial function, however not related to pigment expression. The quest to find the Staphylococcal Hfq function remains open.

LIMITATIONS

Our conclusion is in contradiction with Liu *et al.* results concerning the effect of Hfq on *S. aureus* pigmentation [14]. We cannot rule out that our observation is limited to specific *S. aureus* strains. However, we used an NCTC8325-4 *hfq* derivative similar the one used in the previous study. Furthermore, the present results are strengthened by the construction of *hfq* mutants in numerous *S. aureus* backgrounds. The discrepancy between our and Liu *et al.* 2010 [14] results, is a possible inadvertent selection of mutants with altered color patterns (as shown in [22]) in the former study.

List of abbreviations

PCR: polymerase chain reaction

BHI: brain heart infusion

aTc: anhydrotetracycline

DECLARATIONS

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during this study are included in this published article. Strains and plasmids are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by the Agence Nationale pour la Recherche (ANR) (grant # ANR-15-CE12-0003-01 “sRNA-Fit”) and by the Fondation pour la Recherche Médicale (FRM) (grant # DBF20160635724 “Bactéries et champignons face aux antibiotiques et antifongiques”). WL was the recipient of fellowships from the Chinese scholarship council. PiB was the recipient of fellowships from the FRM.

Authors' contributions

PhB designed the experiments and wrote the manuscript. WL, PiB and CB performed the experiments, analyzed data and revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Sandy Gruss, Annick Jacq and Yan Jing for critical reading of the manuscript, helpful discussions and warm support.

Authors' information

Affiliations

Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Université Paris-Sud, Université Paris-Saclay, 91198, Gif-sur-Yvette cedex, France

ORCID numbers

WL: <https://orcid.org/0000-0003-0747-1852>

CB: <https://orcid.org/0000-0003-1569-9072>

PhB: <https://orcid.org/0000-0003-4601-3387>

Corresponding author

Correspondence to Philippe Bouloc (philippe.bouloc@cnrs.fr).

REFERENCES.

1. Marshall JH, Wilmoth GJ: Pigments of *Staphylococcus aureus*, a series of triterpenoid carotenoids. *J Bacteriol* 1981, 147(3):900-913.
2. Liu GY, Essex A, Buchanan JT, Datta V, Hoffman HM, Bastian JF, Fierer J, Nizet V: *Staphylococcus aureus* golden pigment impairs neutrophil killing and promotes virulence through its antioxidant activity. *J Exp Med* 2005, 202(2):209-215.
3. Bischoff M, Dunman P, Kormanec J, Macapagal D, Murphy E, Mounts W, Berger-Bachi B, Projan S: Microarray-based analysis of the *Staphylococcus aureus* sigmaB regulon. *J Bacteriol* 2004, 186(13):4085-4099.
4. Pelz A, Wieland KP, Putzbach K, Hentschel P, Albert K, Gotz F: Structure and biosynthesis of staphyloxanthin from *Staphylococcus aureus*. *J Biol Chem* 2005, 280(37):32493-32498.
5. Palma M, Cheung AL: sigma(Beta) activity in *Staphylococcus aureus* is controlled by RsbU and an additional factor(s) during bacterial growth. *Infect Immun* 2001, 69(12):7858-7865.
6. Giachino P, Engelmann S, Bischoff M: Sigma(B) activity depends on RsbU in *Staphylococcus aureus*. *J Bacteriol* 2001, 183(6):1843-1852.
7. Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW: A Genetic Resource for Rapid and Comprehensive Phenotype Screening of Nonessential *Staphylococcus aureus* Genes. *MBio* 2013, 4(1).
8. Vogel J, Luisi BF: Hfq and its constellation of RNA. *Nat Rev Microbiol* 2011, 9(8):578-589.
9. Boudry P, Gracia C, Monot M, Caillet J, Saujet L, Hajnsdorf E, Dupuy B, Martin-Verstraete I, Soutourina O: Pleiotropic role of the RNA chaperone protein Hfq in the human pathogen *Clostridium difficile*. *J Bacteriol* 2014, 196(18):3234-3248.
10. Bouloc P, Repoila F: Fresh layers of RNA-mediated regulation in Gram-positive bacteria. *Curr Opin Microbiol* 2016, 30:30-35.
11. Rochat T, Bouloc P, Yang Q, Bossi L, Figueroa-Bossi N: Lack of interchangeability of Hfq-like proteins. *Biochimie* 2012, 94(7):1554-1559.
12. Rochat T, Delumeau O, Figueroa-Bossi N, Noirot P, Bossi L, Dervyn E, Bouloc P: Tracking the Elusive Function of *Bacillus subtilis* Hfq. *PLoS One* 2015, 10(4).
13. Bohn C, Rigoulay C, Bouloc P: No detectable effect of RNA-binding protein Hfq absence in *Staphylococcus aureus*. *BMC Microbiol* 2007, 7:10.
14. Liu Y, Wu N, Dong J, Gao Y, Zhang X, Mu C, Shao N, Yang G: Hfq is a global regulator that controls the pathogenicity of *Staphylococcus aureus*. *PLoS One* 2010, 5(9).
15. Le Lam TN, Morvan C, Liu W, Bohn C, Jaszczyszyn Y, Bouloc P: Finding sRNA-associated phenotypes by competition assays: An example with *Staphylococcus aureus*. *Methods* 2017, 117:21-27.

16. Rochat T, Bohn C, Morvan C, Le Lam TN, Razvi F, Pain A, Toffano-Nioche C, Ponien P, Jacq A, Jacquet E *et al*: The conserved regulatory RNA RsaE down-regulates the arginine degradation pathway in *Staphylococcus aureus*. *Nucleic Acids Res* 2018.
17. Corrigan RM, Foster TJ: An improved tetracycline-inducible expression vector for *Staphylococcus aureus*. *Plasmid* 2009, 61(2):126-129.
18. Uzzau S, Figueroa-Bossi N, Rubino S, Bossi L: Epitope tagging of chromosomal genes in *Salmonella*. *Proc Natl Acad Sci U S A* 2001, 98(26):15264-15269.
19. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO: Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* 2009, 6(5):343-345.
20. Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976, 72:248-254.
21. Herbert S, Ziebandt AK, Ohlsen K, Schafer T, Hecker M, Albrecht D, Novick R, Gotz F: Repair of global regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical isolates. *Infect Immun* 2010, 78(6):2877-2889.
22. Tarrant EJ: The role of Hfq in *S. aureus* gene regulation. *PhD*. University of Leicester; 2013.
23. Antonic V, Stojadinovic A, Zhang B, Izadjoo MJ, Alavi M: *Pseudomonas aeruginosa* induces pigment production and enhances virulence in a white phenotypic variant of *Staphylococcus aureus*. *Infect Drug Resist* 2013, 6:175-186.
24. Nair D, Memmi G, Hernandez D, Bard J, Beaume M, Gill S, Francois P, Cheung AL: Whole-genome sequencing of *Staphylococcus aureus* strain RN4220, a key laboratory strain used in virulence research, identifies mutations that affect not only virulence factors but also the fitness of the strain. *J Bacteriol* 2011, 193(9):2332-2335.
25. Novick RP, Richmond MH: Nature and interactions of the genetic elements governing penicillinase synthesis in *Staphylococcus aureus*. *J Bacteriol* 1965, 90:467-480.
26. Novick R: Properties of a cryptic high-frequency transducing phage in *Staphylococcus aureus*. *Virology* 1967, 33(1):155-166.
27. Novick RP: Molecular biology of the staphylococci. New York, N.Y.: VCH Publishers; 1990.
28. Gill SR, Fouts DE, Archer GL, Mongodin EF, Deboy RT, Ravel J, Paulsen IT, Kolonay JF, Brinkac L, Beanan M *et al*: Insights on evolution of virulence and resistance from the complete genome analysis of an early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing methicillin-resistant *Staphylococcus epidermidis* strain. *J Bacteriol* 2005, 187(7):2426-2438.
29. Baba T, Bae T, Schneewind O, Takeuchi F, Hiramatsu K: Genome sequence of *Staphylococcus aureus* strain Newman and comparative analysis of staphylococcal genomes: polymorphism and evolution of two major pathogenicity islands. *J Bacteriol* 2008, 190(1):300-310.

264 Table 1: *Staphylococcus aureus* strains, plasmids and primer used for this study.

Strain name	Key features	Reference or construction
RN4220	Transformable by DNA from <i>E. coli</i>	[24]
SAPhB5	RN4220 $\Delta hfq::cat$	[13]
NCTC8325	Clinical isolate	[25]
SAPhB224	NCTC8325 $\Delta hfq::cat$	NCTC8325 + $\square 11$ (SAPhB5)
NCTC8325-4	NCTC8325 $\Delta \square 11 \Delta \square 12 \Delta \square 13$	[26]
SAPhB197	NCTC8325-4 $\Delta hfq::cat$	NCTC8325-4 + $\square 11$ (SAPhB5)
RN6390	NCTC 8325-4 $\square 6390$	[27]
SAPhB22	RN6390 $\Delta hfq::cat$	[13]
HG001	NCTC8325 <i>rsbU</i> repaired	[21]
SAPhB199	HG001 $\Delta hfq::cat$	HG001 + $\square 11$ (SAPhB5)
HG002	NCTC8325 <i>tcaR</i> repaired	[21]
SAPhB201	HG002 $\Delta hfq::cat$	HG002 + $\square 11$ (SAPhB5)
HG003	NCTC8325 <i>rsbU</i> and <i>tcaR</i> repaired	[21]
SAPhB203	HG003 $\Delta hfq::cat$	HG003 + $\square 11$ (SAPhB5)
COL	Methicillin resistant clinical isolate	[28]
SAPhB16	COL $\Delta hfq::cat$	[13]
Newman	Clinical isolate	[29]
SAPhB17	Newman $\Delta hfq::cat$	[13]
SAPhB142	RN4220 pRMC2	RN4220 + pRMC2
SAPhB248	RN4220 pRMC2Hfq	RN4220 + pRMC2Hfq
SAPhB251	RN4220 pRMC2HfqFLAG	RN4220 + pRMC2HfqFLAG
SAPhB233	HG003 pRMC2	HG003 +pRMC2
SAPhB249	HG003 pRMC2Hfq	HG003 +pRMC2Hfq
SAPhB257	HG003 pRMC2HfqFLAG	HG003 +pRMC2HfqFLAG
Plasmid name	Key features	Reference/construction*
pRMC2	Anhydrotetracycline (aTc) inducible promoter $P_{xyl/tetO}$	[17]
pRMC2Hfq	<i>hfq</i> inducible expression	See Methods
pRMC2FLAG	pRMC2 derivative for translational gene fusions with <i>3xflag</i> coding sequence	See Methods
pRMC2HfqFLAG	<i>hfq::3xflag</i> inducible expression	See Methods

Primer name	Sequence	purpose
39	GGGGTACCATGATTGCAAACGAAAAC	<i>hfq</i> amplification (with a KpnI site)
49	GGGGAATTCTTATTCTTCACTTTCAGTAGAT GC	<i>hfq</i> amplification (with an EcoRI site)
856	GGTACCGTTAACAGATCTGAG	pRMC2 amplification
918	GCTTATTTTAATTATACTCTATCAATGATAG AG	pRMC2 and pRMC2FLAG amplifications
858	TCAGATCTGTTAACGGTACCGGAATTAGCTT GCATGGAA	<i>3xflag</i> amplification
919	GATAGAGTATAATTAAAAATAAGCGAGCTCGA CTACAAAGACCA	<i>3xflag</i> amplification
865	GACTACAAAGACCATGACGG	pRMC2FLAG amplification
939	GATAGAGTATAATTAAAAATAAGCGTAAAAGG AGTCCGACAGATGA	<i>hfq</i> amplification for cloning in pRMC2FLAG
940	CCGTCATGGTCTTTGTAGTCTTCTTCACTTT CAGTAGATGCTTG	<i>hfq</i> amplification for cloning in pRMC2FLAG

265

Figure legends

Fig 1: Absence of Hfq does not affect *S. aureus* pigmentation. The indicated strains were grown overnight in BHI and then A) streaked on BHI agar or B) assayed for spectral profiles as described [1].

Fig 2: Accumulation of Hfq does not affect *S. aureus* pigmentation. The indicated strains were grown in BHI supplemented with chloramphenicol. They were then A) assayed by Western blotting for *hfq::3xflag* expression (see Methods) and B) streaked on BHI agar supplemented with aTc when indicated.

A

B