Relationship between neural phase-locked responses to speech and perception of speech in noise in young and older adults

Guangting Mai and Peter Howell*

*Corresponding author

Email address of corresponding author: p.howell@ucl.ac.uk

Full postal address: Department of Experimental Psychology, University College London, Gower St., London WC1E 6BT, UK

Running title: Phase-locked responses to speech sounds across age groups

Conflict of Interest: The authors declare no competing conflicts of interest.

Abstract

Auditory phase-locked responses are affected by aging and it has been proposed that this increases the challenges experienced during speech perception in noise (SPiN). However, this proposal lacks direct support. This issue was addressed by measuring speech-evoked phase-locked responses at subcortical (frequency-following responses, FFRs) and cortical (theta-band phase-locking, θ -PLV) levels, and studying the relationship between these phase-locked responses and SPiN (word report accuracies of sentences listened to in noise) in adults across a wide age-range (19-75 years old). It was found that: (1) FFR magnitudes declined with age after hearing loss was controlled for; (2) θ -PLV increased with age, suggesting cortical hyperexcitability in audition; (3) SPiN correlated positively with FFR magnitudes obtained in quiet and with θ -PLV obtained in noise, suggesting that impacts of aging (smaller FFR magnitudes and greater θ -PLV) on SPiN differ at subcortical and cortical levels. Thus, the current study provided evidence for different mechanisms at subcortical and cortical levels through which age affects speech-evoked phase-locked activities and SPiN.

Keywords: Speech perception in noise, frequency following responses, theta-band phase-locking values, age-related hearing loss

1 Introduction

Older adults frequently experience increased difficulty with speech perception in noise (SPiN) (Hume and Dubno, 2010). The factors governing this change with age have been investigated extensively. Declines in cognitive functions such as working memory and attention (Lin et al., 2013) as well as hearing loss, degrade SPiN. Older adults, for example, are poorer than young adults at attending to auditory targets when they have to ignore distractor sounds (Andres et al., 2006). This reduction in attentional ability could lead to poor processing of target speech sounds in noisy environments (Tun et al., 2002, 2009). In addition to degraded cognitive functions, older adults have reduced spectro-temporal acuity that could affect SPiN. This is evident in reduced frequency selectivity (Sommers and Gehr, 1998), impaired temporal gap detection (Schneider and Hamstra, 1999; Humes et al., 2009; Harris et al., 2010) and poorer temporal fine structure (TFS) sensitivity (Hopkins and Moore, 2011; Fullgrabe et al., 2015).

The present study focused on the recent claim that impaired SPiN in older adults is due to degraded temporal encoding of speech sounds (Anderson et al., 2011, 2012; Presacco et al., 2016a). The speech-evoked frequency following responses (FFRs) that originate primarily from the auditory brainstem (Chandrasekaran and Kraus, 2010; Bidelman 2018) require precise temporal processing. FFRs reflect the neural phase-locked activity in response to speech envelope modulations at the fundamental frequency (F_0) (FFR_{ENV_F0}) and TFS for higher harmonics in speech (FFR_{TFS}) (Aiken and Picton, 2008; Skoe and Kraus, 2010). Previous studies have shown that FFRs are smaller in magnitude in older than in young adults (Anderson et al., 2012; Presacco et al., 2016a) which may indicate that reduced FFR magnitudes lead to impaired SPiN in older adults (Anderson et al., 2012). Furthermore, FFRs are associated with SPiN in older adults (Anderson et al., 2011; Fujihira and Shiraishi, 2015; Mai et al., 2018). Specifically, greater FFR_{ENV_F0} magnitude was associated with better SPiN with speech-shaped noise (Anderson et al., 2011). Greater magnitude of FFR_{TFS} in the resolved harmonics region has also been associated with better SPiN when there is reverberation (Fujihira and Shiraishi, 2015) and multi-talker babble noise (Mai et al., 2018).

Cortical phase-locked activity during speech processing also requires neuro-temporal precision. Auditory phase-locking at low frequencies, especially in the theta range (4 ~ 8 Hz), tracks slowly-fluctuating speech envelopes at the cortical level (Luo and Poeppel, 2007; Howard and Poeppel, 2010; Peelle et al., 2013). Theta phase-locking (θ -PLV) in response to amplitude-modulated tones, increases with age (Tlumač et al., 2015; Goossens et al., 2016). This is consistent with prior findings that showed older adults had increased auditory-evoked responses (Alain et al., 2014; Herrmann et al., 2013, 2016) and that cortical tracking of speech envelopes was enhanced (Presacco et al., 2016a). Previous studies showed that theta phase-locking to auditory stimuli with envelope modulations at similar frequencies can predict greater neural firing (Ng et al., 2013) and haemodynamic responses in the auditory cortex (Oya et al., 2018). This indicates that increases in cortical phase-locking could be due to hyperexcitability of the auditory cortex in older adults (Casparly et al., 2008). The increased cortical excitability may also alter the balance between inhibitory and excitatory neural processes in older adults that changes network connectivity and over-represents speech envelopes relative to other speech features (Presacco et al., 2016a).

Despite the reported age effects on subcortical and cortical auditory phase-locked responses, it is unclear at present whether they are related to poor SPiN due to aging. Although subcortical speech-evoked FFRs are associated with SPiN in older adults (Anderson et al., 2011; Fujihira and Shiraishi,

2015; Mai et al., 2018), there is no definitive evidence that shows how age effects on FFRs are related to impaired SPiN. For instance, two recent studies (Presacco et al. 2016a; Schoof and Rosen, 2016) tested the relationship between FFR magnitudes and SPiN in adults using young (< 30 years old) and older (> 60 years old) age groups. Although declines in both FFRs and SPiN were found in older adults compared to young adults, neither study found significant FFR-SPiN correlations. The failure to find neural-behavioral correlations at subcortical levels in Schoof and Rosen (2016) and Presacco et al. (2016a) could be because they were conducted in older adults with relatively normal audiometric hearing (thresholds < 30 dB HL at frequencies $\leq$ 4 kHz). Hence, there were no participants with significant hearing decline due to normal aging, meaning that the older test group did not represent the complete range of hearing present in typical aging populations (Gopinath et al., 2009; Humes et al., 2010).

At the cortical level, Presacco et al. (2016a) argued that increased phase-locking to speech in older adults may reflect a loss of excitation-inhibition balance. It is not clear whether such loss is a mechanism for impaired SPiN during aging and Presacco et al. (2016a) failed to find any correlation between cortical neural activity and SPiN. This absence of correlation may be because different types of background noises were used when neural data and SPiN were measured (single-talker background during neural recording and four-talker babble noise in SPiN tasks; see Presacco et al. (2016a)). On the other hand, studies have shown that greater cortical phase-locking to speech is associated with higher intelligibility of degraded speech stimuli in normal-hearing young adults (Ahissar et al., 2001; Peelle et al., 2013; Doelling et al., 2014) as well as better SPiN in older adults (Mai et al., 2018). These findings are consistent with other studies which showed that greater magnitudes of cortical auditory evoked potentials (CAEPs) can predict better SPiN in both young and older adults (Billings et al., 2013, 2015). Therefore, the relation between age effects on cortical processing and SPiN has not been clarified so far, i.e., whether increased cortical phase-locking according to aging relates to impaired SPiN (due to dampened excitation-inhibition balance) or better SPiN (due to greater sensitivity to speech).

The present study addressed whether age effects on subcortical/cortical phase-locked encoding of speech were associated with impaired SPiN. Behavioral and neural assessments were conducted in healthy adults across a wide age-range (19-75 years). Older adults in the present study had audiometric thresholds at frequencies between 2 and 4 kHz indicative of normal hearing to mild/moderate hearing loss. Therefore individual variability associated with peripheral hearing that

101 occur during normal aging was present in the sample ([Gopinath et al., 2009](#); [Humes et al., 2010](#)).
 102 For the behavioral assessment, participants completed SPiN tasks under two types of background
 103 noise: steady-state speech-shaped noise (SpN) and 16-talker babble noise (BbN). For the neural
 104 assessments, participants listened to a repeated syllable under the same types of noise as in the
 105 behavioral assessment, whilst speech-evoked phase-locked activity was recorded at both subcortical
 106 (FFRs) and cortical (θ -PLV) levels using scalp-electroencephalography (EEG). SPiN and the neural
 107 signatures were compared across the two age groups and multiple linear regressions were conducted
 108 to investigate whether the age-related neural signatures were associated statistically with SPiN.

109 Based on past evidence, it was predicted that older, relative to young, adults would have: (1) smaller
 110 subcortical (FFRs) magnitudes ([Anderson et al., 2012](#); [Presacco et al., 2016a](#)); (2) greater cortical
 111 (θ -PLV) phase-locked responses to speech ([Presacco et al. 2016a](#)); and (3) decreased SPiN ([Hume](#)
 112 [and Dubno, 2010](#)). The predictions for testing the hypotheses that age effects on neural measures
 113 relate to behavioral performance are that decreased SPiN with age should be statistically associated
 114 with: (1) reduced FFR magnitudes; and (2) greater θ -PLV. At the same time, this study explores
 115 which neural (subcortical and/or cortical) signatures optimally model SPiN which is an issue that is
 116 not clear to date.

117 2 Methods

118 The present study followed the same procedure and used parts of the older adults' data from [Mai et](#)
 119 [al. \(2018\)](#)¹.

120 2.1 Participants

121 Participants comprised 23 young (19-42 years; Mean $\pm$ SD = 26.3 $\pm$ 5.5 years; 15 males) and 18
 122 older adults (53-75 years; Mean $\pm$ SD = 67.0 $\pm$ 5.6 years; 7 males). All were native UK English
 123 speakers with no reports of neurological diseases, language-related or psychiatric problems. **Figure**
 124 **1** (left panel) shows the pure-tone audiometric thresholds (PTAs) for frequencies 0.25 ~ 8 kHz
 125 (averaged across both ears) measured using an MA41 Audiometer (MAICO Diagnostics, Germany).
 126 All young participants had normal hearing (PTA $\leq$ 25 dB HL). In older participants, inter-individual

¹ The present study used the data of non-hearing-aid older adults in Group 2 of [Mai et al. \(2018\)](#), where participants listened to the same acoustic stimuli as in the present study whilst neural recordings were made. Data from hearing aid users in [Mai et al. \(2018\)](#) were not included here, since PTAs could not be measured precisely in these participants and hearing aids may introduce additional effects.

variability was high particularly at frequencies of 2 kHz and above. The older adults showed significantly higher low-frequency PTAs (PTA_{Low} ; averaged across 0.25 ~ 1 kHz) and high-frequency PTAs (PTA_{High} ; averaged across 2 ~ 4 kHz)² compared to the young group (both $ps < 10^{-6}$). The boxplot (**Figure 1**, right panel) indicated that PTA_{High} in the older group ranged from normal hearing (≤ 25 dB HL) to mild/moderate (25 ~ 50 dB HL) hearing loss, comparable with the distribution pattern reported in other older samples (Gopinath et al., 2009; Humes et al., 2010). A two-way mixed-design ANOVA was conducted for PTA with factors of Frequency (PTA_{High} vs. PTA_{Low}) and Age Group (young vs. older). A significant [Frequency $\times$ Age Group] interaction occurred ($p = 0.001$), indicating that older adults had significantly greater declines in hearing at the high compared to the low frequencies. Since PTAs differed across frequencies and age groups, PTA_{Low} and PTA_{High} were used as well as PTA averaged across the wider frequency range (0.25 ~ 4 kHz; PTA_{Wide}) as separate covariates and predictors during statistical analyses (see 2.4).

139

Figure 1 Boxplots of pure-tone audiometric thresholds (PTAs) for the young (red) and older (blue) adults averaged across ears. The left panel shows the audiogram for the range from 0.25 to 8 kHz³. Data for 6 and 8 kHz (dashed lines) were not used in the subsequent statistics because spectral distribution of the speech stimuli used in the present study only extended to 4 kHz. The right panel shows the PTAs at the low (0.25 ~ 1 kHz) and high frequency (2 ~ 4 kHz) ranges.

145 2.2 Behavioral experiment

² Since the spectral distribution in the speech stimulus used in both behavioral and neural assessments extended to ≤ 4 kHz, PTAs at 6 and 8 kHz were not included in the statistical analyses in the present study.

³ Three older participants had PTAs that were higher than the measurable limit of the audiometer (85 dB) at 8 kHz in either the left or the right ear and thresholds for them were set at 85 dB when calculating averages over both ears.

SPiN tasks involved participants listening to BKB sentences (Bench et al., 1979) under two types of background noise: steady-state speech-shaped (SpN) and 16-talker babble (BbN) noise. All sentences were pre-recorded utterances spoken by a male British English speaker whose F_0 ranged from 80 to 200 Hz. Each sentence included three key (content) words, e.g., “The clown has a funny face” with key words “clown”, “funny” and “face”. BbN was a mixture of 16 different utterances spoken by 16 male British English speakers with similar voice quality to the target speaker. SpN was formed by randomizing the phases of the long-term spectrum of BbN and transforming the spectrum back to the time domain. As a result, SpN has the same long-term power spectrum as BbN and stable time-domain properties (Rosen et al., 2013).

Participants were seated comfortably in a sound-treated booth facing a Fostex 6301B loudspeaker (Canford Group Ltd.) at zero-degree azimuth. Distance between the loudspeaker and participants’ ears was constant at 1 meter. After eight trials of practice, participants listened to two different sets of 30 sentences (for the backgrounds of SpN and BbN, respectively) at an intensity at this distance of 70dB SPL. Participants repeated as many words as they could from each sentence. Sentences were presented via Matlab 2010a (Mathwork, USA) and SNR varied adaptively to track for the speech reception threshold (SRT, Plomp and Mimpen, 1979) at which 50% of words were correct. For each background type, the first sentence was played at a relatively high SNR (8 and 10 dB for SpN and BbN, respectively). SNR was decreased by 4 dB for subsequent sentences until < 50% words correct (i.e., < 2 words) were reported. SNR was then increased/decreased by 2 dB when word correctness was less/more than 50% in each of the following sentences. SRT was calculated by linear interpolation using the two SNRs which had > 50% and < 50% correct across the minimal step distance (i.e., 2 dB).

2.3 EEG experiment

2.3.1 Acoustic stimuli

Participants listened to a repeatedly-presented, 120-ms-long /i/ syllable produced by a male speaker (Figure 2A). The F_0 contour of the syllable fell from ~ 160 to ~110 Hz (Figure 2B). The F_0 contour covered a similar frequency range and direction of change as those in the F_0 s of the target speaker in the BKB sentences used in the SPiN tasks (BKB sentences are narratives that generally have a falling F_0 contour). The three formants in the syllable were at ~ 280 Hz (F_1), ~ 2400 Hz (F_2) and ~

175 3100 Hz (F3). The amplitude envelope profile was stable except that 5-ms-long rising and falling
176 cosine windows were applied at the onset and offset to avoid transients.

177

178 **Figure 2** The syllable /i/ used during EEG recording. **(A)** The temporal waveform (top) and
179 spectrogram (bottom) of the syllable. F1, F2 and F3 frequencies are around 280, 2400 and 3100 Hz,
180 respectively. **(B)** The falling F_0 contour ranging from around 160 to 110 Hz obtained by
181 autocorrelation. The waveform, spectrogram and F_0 were generated via PRAAT ([Boersma and](#)
182 [Weenink, 2013](#)).

183 The syllable was presented repeatedly at both original (positive) and inverted (negative) polarities
184 in random order with inter-stimulus intervals (ISIs) that varied randomly between 60 and 120 ms
185 (syllable repetition rate was approximately 5 syllables per second). The stimuli were presented in
186 quiet, SpN and 16-talker BbN backgrounds (the last two were the same backgrounds that were used
187 in the SPiN tasks). The SNRs were set at -1 dB, which led to neural responses that correlated
188 significantly with SPiN in older adults ([Mai et al., 2018](#)). There were 6400 sweeps under each
189 background type (3200 sweeps for each polarity). Recordings at each background type were split
190 into 16 segments of equal duration giving 48 segments in total with 400 sweeps per segment. The
191 segments were played in succession in an intermixed order.

192 2.3.2 EEG data acquisition

193 Scalp-EEGs were recorded on an ActiveTwo system (Biosemi, The Netherlands) at a sampling rate
194 of 16384 Hz. Three active electrodes were placed at Cz (vertex), C3 and C4 according to the 10/20
195 configuration. Cz was used to obtain FFRs ([Skoe and Kraus, 2010](#)). Cortical responses were
196 measured on C3 and C4 that reflects activity in the auditory cortex ([Carpenter and Shahin, 2013](#);

Noguchi et al., 2015) and allows reliable cortical phase-locked activity that is significantly associated with SPiN to be recorded (Mai et al., 2018). Bilateral earlobes were used as the reference. Ground electrodes were CMS/DRL. Electrode impedance was kept below 35 mV. The experiment was conducted in an electromagnetic-shielded and sound-treated booth. The stimuli were played via a Rogers LS3/5A loudspeaker (Falcon Acoustics, UK) at zero-degree horizontal azimuth relative to participants' heads when they were reclined (the chair was adjustable). The stimulus level (measured across time including ISIs) at the distance between the loudspeaker and participants' ears (constant at 1 meter) was calibrated at 74.5 dB before background noise was added. The stimulus level was at 79.5 dB after either SpN or BbN was added.

Participants were instructed to relax, close their eyes and keep still in order to avoid movement artefacts. They did not have to make any response to the stimuli (passive listening) and they were not stopped from falling asleep. A webcam monitored the participants throughout the test and no significant changes in head or body position were observed. Participants were not stopped from falling asleep because another purpose of the current experiment was to study the effects of arousal on speech-evoked neural processing across ages (Mai et al., 2019). This investigation was separate from the present paper. As Mai et al. (2019) found that arousal significantly affected the phase-locked responses, only EEG data from periods with high arousal were used here (see 2.4.3 for details).

2.4 Signal processing for EEG data

The signal processing procedure used Matlab 2014a (Mathwork, USA).

2.4.1 Frequency following responses (FFRs)

EEGs at Cz were re-referenced to the average of bilateral earlobes and bandpass filtered between 70 and 4000 Hz using a zero-phase 2nd-order Butterworth filter. Baseline was adjusted using the pre-stimulus period of 50 ms. Sweeps exceeding $\pm 25 \mu\text{V}$ were rejected to exclude movement artefacts. FFRs with positive (FFR_{pos}) and negative (FFR_{neg}) polarities were obtained by averaging across sweeps with their respective polarities. FFRs that represent envelope modulations (FFR_{ENV}) and TFS (FFR_{TFS}) respectively were obtained by addition and subtraction of FFR_{pos} and FFR_{neg} that were then divided by 2 (Aiken and Picton, 2008). **Figure 3** shows an example of FFRs obtained in the present study (FFRs of a single participant recorded in BbN).

226 Three FFR magnitudes were measured: (1) $\text{FFR}_{\text{ENV}_F0}$ that represents neural encoding of envelope
 227 modulations at F_0 , quantified as the magnitude along the F_0 trajectory using FFR_{ENV} (**Figure 3A**);
 228 (2) $\text{FFR}_{\text{TFS}_H2}$ that represents neural encoding of TFS at the resolved harmonics region (2nd
 229 harmonics H2 at 220 ~ 330 Hz in the neighborhood of F1), quantified as the magnitudes along the
 230 H2 trajectory using FFR_{TFS} (**Figure 3B**); and (3) $\text{FFR}_{\text{TFS}_{F2F3}}$ that represents neural encoding of TFS
 231 in the unresolved harmonics region (frequency range around F2 and F3), quantified as the
 232 magnitudes along the F2 and F3 trajectories using FFR_{TFS} (**Figure 3B**). The procedures for spectral
 233 magnitude calculations followed [Mai et al. \(2018\)](#).

234 First, ENV- F_0 (F_0 based on the acoustic envelope), H2, F2 and F3 trajectories of the /i/ syllable
 235 were calculated. To obtain the ENV- F_0 trajectory, a set of 40-ms sliding windows (1-ms per step)
 236 was applied to the syllable's Hilbert envelope. Each 40-ms segment was Hanning-windowed, zero-
 237 padded to 1 second (to achieve 1 Hz frequency resolution) and Fourier-transformed. The frequency
 238 with the highest Fourier magnitude between 110 and 160 Hz (the F_0 range) was chosen as the F_0
 239 value at each step. H2, F2 and F3 trajectories were obtained in the same way, except that: 1) sliding
 240 windows were applied to the syllable rather than the Hilbert envelope; 2) H2 values were selected
 241 within the H2 range (220 ~ 320 Hz); 3) instead of choosing values based on the Fourier spectrum
 242 after zero-padding, F2 and F3 values were chosen based on the spectral profile via cepstral
 243 smoothing ([Proakis and Manolakis, 2007](#)) in the F2 and F3 ranges respectively (2200 ~ 2600 Hz
 244 and 2800 ~ 3500 Hz). Second, to calculate the $\text{FFR}_{\text{ENV}_F0}$ magnitude, the same set of 40-ms sliding
 245 windows was applied to FFR_{ENV} with Hanning-windowing, zero-padding and Fourier-transforms.
 246 The mean log-magnitude was measured across a 20 Hz bandwidth centered at the frequency of the
 247 ENV- F_0 trajectory at that step. The magnitudes were then averaged across all steps along the ENV-
 248 F_0 trajectory. $\text{FFR}_{\text{TFS}_H2}$ and $\text{FFR}_{\text{TFS}_{F2F3}}$ magnitudes were obtained in the same way, except that: 1)
 249 the procedure was applied on FFR_{TFS} along the H2 (for $\text{FFR}_{\text{TFS}_H2}$) and the F2 and F3 (for
 250 $\text{FFR}_{\text{TFS}_{F2F3}}$) trajectories; 2) instead of obtaining magnitudes based on the Fourier spectrum after
 251 zero-padding, $\text{FFR}_{\text{TFS}_{F2F3}}$ magnitude at each step was the summed magnitude of the spectral profile
 252 (via cepstral smoothing) across a 150 Hz and 300 Hz bandwidth respectively centered at F2 and F3
 253 of the syllable at that step.

254

255 **Figure 3** FFR waveforms (top) and spectrograms (bottom) of a single participant recorded in BbN.
 256 **(A)** FFR_{ENV} (bandpass filtered at 70 ~ 2000 Hz); **(B)** FFR_{TFS} (bandpass filtered at 70 ~ 4000 Hz).
 257 The waveforms were based on sweeps with normalized numbers (1450 ~ 1550) during the high
 258 arousal state (see 2.4.3 and 2.4.4). FFR_{ENV_F0} (at F₀ range between 160 and 110 Hz), FFR_{TFS_H2} (at
 259 H2 range between 220 and 320 Hz) and FFR_{TFS_F2F3} (at F2-F3 range between 2000 and 4000 Hz)
 260 are indicated by the boxes surrounded by dashes and their labels. ‘0’ corresponds to the syllable
 261 onset.

262 In addition, neural transmission from the cochlea to the auditory brainstem for FFR_{ENV} takes
 263 between 5 and 10 ms (Chandrasekaran and Kraus, 2010; Skoe and Kraus, 2010), while FFR_{TFS}
 264 occurs at earlier stages in the auditory periphery (Aiken and Picton, 2008). Hence, the maximum
 265 magnitude for time lags in the range 8 to 13 ms and 3 to 8 ms (at 1-ms steps; including an additional
 266 3 ms of air transmission from the loudspeaker to the cochlea) were used as the final FFR_{ENV_F0} and
 267 FFR_{TFS} (FFR_{TFS_H2} and FFR_{TFS_F2F3}) magnitudes, respectively.

268 As well as FFR magnitudes, inter-trial phase-locking values (PLV) at the F₀ (FFR_{PLV_F0}) were also
 269 calculated. This was because FFR_{PLV_F0} reflects pure phase-locking that excludes the influence of
 270 single-trial spectral magnitudes and has a better signal-to-noise ratio than does FFR magnitudes
 271 (Zhu et al., 2013). FFR_{PLV_F0} was calculated in a similar way to FFR_{ENV_F0} along the ENV-F₀
 272 trajectory, except that, after zero-padding in each step (without Hanning-windowing), PLV was
 273 calculated (Morillon et al., 2012) instead of spectral magnitudes:

$$PLV = \frac{1}{n} \left| \sum_{i=1}^n e^{j\phi_i} \right|$$

where n denotes the total number of sweeps, ϕ_i denotes the Fourier phase value at the frequency of the ENV-F₀ trajectory for the i th sweep at that step, and j is $\sqrt{-1}$. As PLV is restricted to values between 0 and 1, FFR_{PLV_F0} at each step was quantified by logit-transforming PLV to $[-\infty, +\infty]$, making it appropriate for linear regression analysis (Waschke et al., 2017):

$$FFR_{PLV_F0} = \ln \frac{PLV}{1-PLV}$$

FFR_{PLV_F0} values were then averaged across all steps along the ENV-F₀ trajectory. The final FFR_{PLV_F0} value was taken as the maximal value for the time lags between 8 and 13 ms as in measurement of the FFR_{ENV_F0} magnitude.

2.4.2 Cortical responses

Cortical responses were measured as theta-band (4 ~ 6 Hz, to correspond to the stimulus repetition rate of 5 syllables per second) phase-locking values (θ -PLV) at C3 and C4. EEGs were decimated to 1024 Hz, re-referenced to the average of the bilateral earlobes and bandpass filtered (4 ~ 6 Hz) using a 2nd-order zero-phase Butterworth filter. Sweeps exceeding ± 15 μ V on either electrode were rejected (Mai et al., 2018)⁴. θ -PLV time series ($PLV(t)$) were calculated and then logit-transformed:

$$PLV(t) = \frac{1}{n} \left| \sum_{i=1}^n e^{j\phi_i(t)} \right|$$

$$Logit-\theta-PLV(t) = \ln \frac{PLV(t)}{1-PLV(t)}$$

where n denotes the total number of sweeps, $\phi_i(t)$ denotes the Hilbert phase series of the filtered EEG of the i th sweep time-locked to the syllable onset and j is $\sqrt{-1}$. Hilbert phase was used as it reflects phase-locking to stimuli even when EEG amplitude variation occurs due to relaxation and eye closure (Thatcher, 2012). Logit- θ -PLV(t) values were then averaged across the stimulus period (120 ms). Neural transmission from cochlea to auditory cortex takes 10 to 30 ms in primates

⁴ Lower rejection thresholds were used than with FFRs (± 25 μ V) because the θ -band signal usually does not have excessively high amplitude since it occupies a relatively narrow frequency range. More than 80% of the sweeps were retained in all participants after artefact rejection.

(Lakatos et al., 2007). Hence Logit- θ -PLV was taken as the maximum value for time lags between 13 and 33 ms (at 1-ms steps) with the 3 ms for air transmission included. Finally, the Logit- θ -PLV(t) was averaged across the two electrodes.

2.4.3 Classification of arousal states

Participants were not required to remain awake during the EEG recording because a separate purpose of the present experiment was to investigate the effect of arousal on speech-evoked responses (Mai et al., 2019). Mai et al. (2019) reported that both subcortical and cortical responses showed significant suppression in low arousal/nREM states compared to high arousal states. This accords with earlier functional imaging work that showed that neural responses to speech in subcortical (Portas et al., 2000) and cortical (Czisch et al., 2004; Wilf et al., 2016) auditory regions reduce during sleep compared to wakefulness. Also, significant correlations between behavioral measures and EEG parameters (FFRs and θ -PLV) were only found in the high arousal state (Mai et al., 2018). Hence, only EEG data from periods with high arousal were used to avoid any influences that arousal has on the neural-behavioral relationship.

Sleep spindles were used to determine arousal state (Martin et al., 2013). Sections of the Cz EEG recordings were categorized into three types of epochs (all epochs were 21-second long) based on the occurrence of sleep spindles: (1) epochs in high arousal states (wakefulness or nREM Stage 1); (2) epochs in low arousal states (nREM Stage 2); and (3) epochs in transition between (1) and (2). After the experiment, participants gave a subjective ranking concerning how much they had slept. There was a significant correlation between the sleep ranking and the percentage of epochs classified as ‘low arousal’ ($p = 0.002$), which validated the spindle-based method. Further methodological details are available in Mai et al (2019).

2.4.4 Normalization of sweep numbers

Robust FFRs require around 1500 artefact-free sweeps (c.f., Dajani et al., 2005; Wong et al., 2007). Hence, participants’ data for a particular background type were not included in subsequent analyses if there were $< 1,450$ artefact-free sweeps in high arousal epochs for that background type. This resulted in 17%-30% of participants being rejected from further analyses (depending on the types of analyses conducted; see 3.2 and 3.3 for details). Moreover, as magnitudes of phase-locked activity are sensitive to the number of sweeps (Aviyente et al., 2011), problems can arise during statistical

analyses if the number of sweeps differ significantly across participants. Therefore, the number of sweeps was normalized to around 1,500 for both FFRs and θ -PLV for each participant in each background type. Normalization was achieved by selecting high arousal epochs at random that included 1,450 to 1,550 sweeps and EEG signatures (magnitudes of FFR_{ENV_F0} , FFR_{TFS_H2} and FFR_{TFS_F2F3} , and Logit- θ -PLV) were obtained from the selected epochs. The random selection procedure was repeated 100 times, giving 100 estimates for each EEG signature. Averages over the 100 estimates were used in the final statistical analyses. Therefore, this process ensured EEG signatures were based on around 1500 sweeps regardless of artefact rejection rates or different number of epochs of the three arousal states across participants (Mai et al., 2018).

2.4.5 Confirming FFR robustness

FFR magnitudes are small and their robustness was tested by statistically comparing the FFR magnitudes with the EEG noise floors. The noise floors were quantified as the EEG magnitudes at the corresponding frequency range (110 ~ 160 Hz (F_0) for FFR_{ENV_F0} ; 220 ~ 320 Hz ($H2$) for FFR_{TFS_H2} ; 150-Hz bandwidth centred at 2400 Hz ($F2$) and 300-Hz bandwidth centred at 3100 Hz ($F3$) for FFR_{TFS_F2F3}) at the 50-ms FFR pre-stimulus period (Mai et al., 2018). The quantification procedure was similar to that used in calculating FFR magnitudes, in which a set of 40-ms sliding windows was applied on FFR_{ENV} (for FFR_{ENV_F0}) or FFR_{TFS} (for FFR_{TFS_H2} and FFR_{TFS_F2F3}) which used 1-ms steps over the pre-stimulus period. Magnitudes of noise floors were measured as the spectral magnitudes (summed magnitude of the cepstrally-smoothed profile for FFR_{TFS_F2F3}) across the corresponding frequency ranges averaged across all steps. These were all conducted along with the calculations of FFR magnitudes during the processes for normalization of sweep numbers.

2.5 Statistical analyses

Statistical analyses were conducted using SPSS 13.0 (SPSS Inc., USA).

2.5.1 Analyses of variance (ANOVAs) and Analyses of covariance (ANCOVAs)

Mixed-design ANOVAs were conducted with the behavioral (SRT in the SPiN tasks) and EEG signatures (FFR_{ENV_F0} , FFR_{PLV_F0} , FFR_{TFS_H2} , FFR_{TFS_F2F3} and Logit- θ -PLV) as the dependent variables, and with the within-subject factor of Noise Type (SpN and BbN for SRT; Quiet, SpN and BbN for EEG) and the between-subject factor of Age Group (young vs. older) as independent

variables. Post-hoc t-tests were conducted if a significant [Noise Type $\times$ Age Group] interaction occurred. The ANOVAs were conducted for two reasons: (1) main effects of Age Group were tested to look into the effects of age on SPiN and neural responses to speech; (2) since older adults experience more difficulties in SPiN under BbN compared to other types of noise (Helfer and Freyman, 2008; Schoof and Rosen, 2014), testing the [Noise Type $\times$ Age Group] interaction should help reveal what neural mechanisms underlie this.

The EEG signatures were further used in mixed-design ANCOVAs that included PTAs (PTA_{Low} , PTA_{High} and PTA_{Wide} , averaged across 0.25 ~ 1 kHz, 2 ~ 4 kHz and 0.25 ~ 4 kHz, respectively; all mean-centred) as covariates, with the same within-subject and between-subject factor as in ANOVAs. These tested for age effects after the variability in peripheral hearing loss was controlled for. PTA_{Low} and PTA_{High} reflect hearing loss at low and high frequencies, respectively, while PTA_{Wide} reflects the combined effect of both. The three PTA variables were used as covariates in separate analyses to avoid the risk of collinearity. A concern about the ANCOVAs is that PTAs and Age are correlated (effects of PTA and Age can overlap), hence including PTAs as covariates may carry risks of partially diminishing the Age effects despite controlling for hearing loss. For this reason, ANOVAs and ANCOVAs were both conducted separately. Alpha values for testing significance in ANOVAs/ANCOVAs were not adjusted.

2.5.2 Multiple linear regressions

Multiple linear regressions were then conducted to test for any behavioral-neural relationship using data from both the young and older groups. EEG signatures (FFR_{ENV_F0} , FFR_{PLV_F0} , FFR_{TFS_H2} , and FFR_{TFS_F2F3} , and Logit-0-PLV) were used as predictors of SPiN (SRTs; dependent variables) for the corresponding noise types. Specifically, EEG signatures obtained in SpN were used to predict SRT in SpN, while EEG signatures obtained in BbN were used to predict SRT in BbN. This avoided problems that could arise due to behavioral and neural recordings being made under different types of noise (Presacco et al., 2016a). Additionally, since FFRs in quiet have been suggested to be associated with SPiN (Anderson et al., 2011), we also used EEG signatures in quiet to predict SRTs. Age was not included as a predictor, since the regressions investigated the contributions of age-related factors (which were identified by significant main effects of Age in the ANOVAs), rather than age itself, to SPiN.

380 The Best-Subset Regression approach was used that selected predictors of EEG signatures that
 381 generated the lowest BIC value. This approach provided the optimal model with best goodness of fit
 382 and least chance of overfitting (Burnham and Anderson, 2003). PTAs (PTA_{Low} , PTA_{High} and
 383 PTA_{Wide}) were also included as predictors to generate the Best-Subsets which take into account the
 384 effects of peripheral hearing loss. To avoid any spurious regression results caused by
 385 multicollinearity, subsets with variance inflation factors (VIFs) > 1.5 excluded (c.f., Stine, 1995). It
 386 is noteworthy that Best-Subset Regression is exploratory in nature and is often used when there is
 387 lack of a priori theory for a given topic. Here, it is not clear which neural parameter(s) can optimally
 388 model SPiN, and such an approach was used to answer this question and aid identification of the
 389 neural substrates that underlie SPiN.

390 After regression analyses using data from both young and older adults, the analyses were further
 391 conducted separately for the young and the older group, to evaluate whether they employ different
 392 neural mechanisms for SPiN.

393 **3 Results**

394 **3.1 Behavioral results**

395 ANOVA was conducted for SRT with Noise Type (SpN vs. BbN) and Age Group (young vs. older)
 396 as factors. The SRTs are plotted as boxplots⁵ in **Figure 4** for SpN and BbN of the 23 young and 18
 397 older participants and the statistics are summarized in **Table 1**. The significant main effect of Noise
 398 Type ($F_{(1, 39)} = 382.850$, $p < 10^{-21}$; $SRT_{SpN} < SRT_{BbN}$) is consistent with previous finding that speech
 399 is better recognized in SpN than in BbN (Rosen et al., 2013). The significant main effect of Age
 400 Group ($F_{(1, 39)} = 5.527$, $p = 0.024$; $SRT_{Young} < SRT_{Older}$) showed that young adults had better
 401 performance than older adults. A significant [Noise Type $\times$ Age Group] interaction occurred ($F_{(1, 39)}$
 402 $= 10.010$, $p = 0.003$) and post-hoc t-tests showed that young adults had significantly better
 403 performance than older adults in BbN ($t_{(26.667)} = -3.399$, $p = 0.002$, Cohen's $d = 1.132$; equal
 404 variances not assumed), but not in SpN ($t_{(39)} = -0.135$, $p = 0.893$, Cohen's $d = 0.043$; equal
 405 variances assumed).

⁵ N.B., circles in boxplots represent outliers which are defined as data points that fall outside the 1.5 times the interquartile range i.e. are above/below the upper/lower quartile. Outliers were *not* excluded during the statistical analyses.

Figure 4. Boxplots for SRT as a function of Noise Type (SpN vs. BbN) and Age Group (young vs. older). Low SRTs represent good SPiN.

Table 1. Statistical results of ANOVAs for SRT with factors of Noise Type (SpN vs. BbN) and Age Group (young vs. older). DV, df, F, and *p* refer to the dependent variable, degrees of freedom, F values, *p* values, respectively. Significant *p* values (< 0.05) are indicated in bold. * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

DV	Factors	<i>df1</i>	<i>df2</i>	<i>F</i>	<i>P</i>
SRT	Noise Type	1	39	382.850	$< 10^{-21}$***
	Age Group	1	39	5.527	0.024*
	Noise Type $\times$ Age Group	1	39	10.010	0.003**

3.2 Neural results

Robustness of FFR was first confirmed by testing whether FFR magnitudes were statistically greater than their corresponding EEG noise floors (see 2.4.5). It was shown that spectral magnitudes of FFRs were all significantly greater than noise floors ($\text{FFR}_{\text{ENV_F0_Quiet}}$: $p < 10^{-8}$; $\text{FFR}_{\text{ENV_F0_SpN}}$ and $\text{FFR}_{\text{ENV_F0_BbN}}$: $p < 0.001$; $\text{FFR}_{\text{TFS_H2_Quiet}}$: $p < 10^{-11}$; $\text{FFR}_{\text{TFS_H2_SpN}}$ and $\text{FFR}_{\text{TFS_H2_BbN}}$: $p < 10^{-10}$; $\text{FFR}_{\text{TFS_F2F3_Quiet}}$, $\text{FFR}_{\text{TFS_F2F3_SpN}}$ and $\text{FFR}_{\text{TFS_F2F3_BbN}}$: $p < 10^{-6}$). Furthermore, the response signal-to-noise ratio (SNR, i.e., difference in magnitudes between FFRs and the corresponding noise floors) did not differ between age groups (SNR for $\text{FFR}_{\text{ENV_F0_Quiet}}$: $p > 0.07$; SNRs for other FFR signatures: $p > 0.2$), indicating that SNR would not be a good index for measuring age

differences. Additional simulations were also conducted in the present study showing that FFR magnitudes can more reliably quantify the FFR fidelity compared to response SNRs (see *Appendix 2*).

ANOVAs were then conducted for EEG signatures (FFR_{ENV_F0} , FFR_{PLV_F0} , FFR_{TFS_H2} , FFR_{TFS_F2F3} and Logit- θ -PLV) including Noise Type (Quiet, SpN and BbN) and Age Group (young vs. older) as factors. Participants' data were not included if artefact-free sweeps in the high arousal periods (wakefulness and nREM stage 1) were < 1450 in Quiet, SpN or BbN to ensure good EEG signal quality (see 2.4.4). 29 (15) participants were retained (i.e., 12 (8) were rejected; rejection rate 29% where the numbers in brackets represent the numbers of young adults). Time series of the EEG signatures are given in **Figure 5** and the boxplots are shown in **Figure 6**.

Statistics on the ANOVAs are summarized in **Table 2**. Greenhouse-Geisser corrections were conducted to correct for violations of sphericity. For FFRs, significant main effects of Noise Type and [Noise Type $\times$ Age Group] interactions were found for both FFR_{ENV_F0} and FFR_{PLV_F0} . Post-hoc comparisons following the main effects showed that FFR_{ENV_F0} and FFR_{PLV_F0} were significantly greater in Quiet than in noise ($FFR_{ENV_F0_Quiet} > FFR_{ENV_F0_SpN}$, $p < 10^{-7}$, Cohen's $d = 1.297$; $FFR_{ENV_F0_Quiet} > FFR_{ENV_F0_BbN}$, $p < 10^{-8}$, Cohen's $d = 1.651$; $FFR_{PLV_F0_Quiet} > FFR_{PLV_F0_SpN}$, $p < 10^{-6}$, Cohen's $d = 1.201$; $FFR_{PLV_F0_Quiet} > FFR_{PLV_F0_BbN}$, $p < 10^{-7}$, Cohen's $d = 1.346$), but they did not differ between SpN and BbN (both $p > 0.5$). Post-hoc t-tests following the interaction effects found significant greater FFR_{PLV_F0} for young than older adults in Quiet ($t_{(27)} = -2.105$, $p = 0.045$, Cohen's $d = 0.782$). No main effects nor interactions were found for FFR_{TFS_H2} or FFR_{TFS_F2F3} . For Logit- θ -PLV, there were significant main effects of Noise Type and Age Group, but no significant [Noise Type $\times$ Age Group] interaction. Post-hoc comparisons found that Logit- θ -PLV was greater in Quiet than in noise, and greater in SpN than in BbN ($Logit-\theta-PLV_{Quiet} > Logit-\theta-PLV_{SpN}$, $p = 0.002$, Cohen's $d = 0.654$; $Logit-\theta-PLV_{Quiet} > Logit-\theta-PLV_{BbN}$, $p < 10^{-6}$, Cohen's $d = 1.334$; $Logit-\theta-PLV_{SpN} > Logit-\theta-PLV_{BbN}$, $p < 10^{-4}$, Cohen's $d = 0.961$); Logit- θ -PLV was greater for older than young adults ($p < 0.001$, Cohen's $d = 1.504$).

ANCOVAs were conducted next that included PTAs (PTA_{Low} , PTA_{High} and PTA_{Wide}) as covariates. For FFR_{ENV_F0} , an additional significant main effect of Age Group was found ($F_{(1, 26)} = 8.842$, $p = 0.006$, $FFR_{ENV_F0_Young} > FFR_{ENV_F0_Older}$) compared to the previous ANOVA when PTA_{Low} was the covariate. The main effect of PTA_{Low} was also significant ($F_{(1, 26)} = 12.943$, $p = 0.001$), where higher

PTA_{Low} correlated with greater FFR_{ENV_F0} magnitude (see **Table 3**). No significant main effects of Age or PTA were found when PTA_{High} or PTA_{Wide} were used as covariates (all $p > 0.2$) (see *Appendix Tables A2-A3*). For Logit- θ -PLV, the significant main effect of Age Group was maintained ($F_{(1, 26)} = 4.608$, $p = 0.041$ and $F_{(1, 26)} = 5.937$, $p = 0.022$) when PTA_{Low} and PTA_{High} were used as the covariates, respectively, however, this became non-significant ($F_{(1, 26)} = 3.516$, $p = 0.072$) when PTA_{Wide} was used as the covariate. No significant main effects of PTA_{Low}, PTA_{High}, or PTA_{Wide} occurred (all $p > 0.2$) (see *Appendix Tables A1-A3*). For FFR_{PLV_F0}, FFR_{TFS_H2} and FFR_{TFS_F2F3}, no significant main effects occurred (all $p > 0.1$) (see *Appendix Tables A1-A3*). The overall ANCOVA results showed: (1) FFR_{ENV_F0} declined with age when PTA_{Low} was controlled for, where higher PTA_{Low} was related to greater FFR_{ENV_F0}; (2) the age effect for Logit- θ -PLV was maintained when PTAs were controlled for (though dropped below significance when PTA_{Wide} was used as the covariate) and the increased Logit- θ -PLV cannot be statistically explained by increased PTA.

Figure 5. Time series of FFR_{ENV_F0}, FFR_{TFS_H2}, FFR_{TFS_F2F3} and Logit- θ -PLV in Quiet (upper row), SpN (mid row) and BbN (lower row). The series were based on sweeps with normalized numbers (1,450 ~ 1,550) during the high arousal periods (see 2.4.3 and 2.4.4) averaged across young (red) and older (blue) adults. Series of FFR_{ENV_F0} are shown as FFR_{ENV} bandpass filtered at 90 ~ 180 Hz (corresponding to the F₀ range). Series of FFR_{TFS_H2} and FFR_{TFS_F2F3} are shown as FFR_{TFS} bandpass filtered at 200 ~ 340 Hz (corresponding to the H2 range), and at 2000 ~ 4000 Hz (corresponding to the F2 and F3 range), respectively. '0' represents the syllable onset.

474

475 **Figure 6.** Boxplots for the five EEG signatures as a function of Noise Type (Quiet, SpN and BbN)
 476 and Age Group (young vs. older). FFR_{ENV_F0} and FFR_{TFS_H2} were measured as log-power (dB);
 477 FFR_{TFS_F2F3} was measured as power of cepstral spectrum (dB) at F2 and F3 range; FFR_{PLV_F0} and
 478 Logit- θ -PLV were measured as logit-transformed phase-locking values (a.u., arbitrary units). Young
 479 and older adults are plotted in red and blue respectively.

480 **Table 2.** Statistical results for ANOVAs (with Greenhouse-Geisser correction) for the five EEG
 481 signatures with factors of Noise Type (Quiet, SpN and BbN) and Age Group (young vs. older). DV,
 482 df, F, p refer to the dependent variable, degrees of freedom, F values, and p values (uncorrected),
 483 respectively. Significant p values (< 0.05) are indicated in bold. ** = $p < 0.01$; *** = $p < 0.001$.

DV	Factors	<i>df1</i>	<i>df2</i>	<i>F</i>	<i>p</i>
FFR_{ENV_F0}	Noise Type	1.606	43.359	49.536	$< 10^{-10}$***
	Age Group	1	27	0.326	0.573
	Noise Type $\times$ Age Group	1.606	43.359	5.608	0.011*
FFR_{PLV_F0}	Noise Type	1.559	42.082	42.827	$< 10^{-9}$***
	Age Group	1	27	2.225	0.147
	Noise Type $\times$ Age Group	1.559	42.082	3.522	0.049*
FFR_{TFS_H2}	Noise Type	1.898	51.244	0.324	0.713
	Age Group	1	27	0.005	0.946
	Noise Type $\times$ Age Group	1.898	51.244	2.061	0.140
FFR_{TFS_F2F3}	Noise Type	1.599	43.181	1.592	0.217
	Age Group	1	27	2.479	0.127
	Noise Type $\times$ Age Group	1.599	43.181	0.051	0.918

Logit- θ -PLV	Noise Type	1.728	46.654	31.068	< 10⁻⁷***
	Age Group	1	27	16.386	< 0.001***
	Noise Type $\times$ Age Group	1.728	46.654	0.972	0.375

484

485 **Table 3.** Statistical result of ANCOVA (with Greenhouse-Geisser correction) for FFR_{ENV_F0} with
486 PTA_{Low} as the covariate. DV, df, F, and *p* refer to the dependent variable, degrees of freedom, F
487 values, and *p* values (uncorrected), respectively. Significant *p* values (< 0.05) are indicated in bold.
488 * = *p* < 0.05; ** = *p* < 0.01.

DV	Factors	<i>df1</i>	<i>df2</i>	<i>F</i>	<i>p</i>
FFR _{ENV_F0}	Noise Type	1.506	39.152	50.277	< 10⁻⁹***
	Age Group	1	26	8.842	0.006**
	PTA _{Low}	1	26	12.943	0.001**
	Noise Type $\times$ Age Group	1.506	39.152	3.118	0.069
	Noise Type $\times$ PTA _{Low}	1.506	39.152	1.445	0.246

489

490 3.3 Behavioral-neural relationship

491 Behavioral-neural relationships were assessed by linear regressions in which SRT in SpN was
492 predicted by EEG signatures obtained in SpN, whilst SRT in BbN was predicted by EEG signatures
493 obtained in BbN. SRTs were further predicted by EEG signatures obtained in Quiet. PTAs (PTA_{Low},
494 PTA_{High} or PTA_{Wide}) were also included as predictors provided that including them improved the
495 statistical capacity of EEG signatures to predict SRTs in the Best-Subsets. Similar to the procedure
496 in 3.2, participants' data obtained under a particular noise type (Quiet, SpN or BbN) were not
497 included if artefact-free sweeps in the high arousal periods were < 1450 for that noise type (see
498 2.4.4). This resulted in 31 (17) participants retained (i.e., 10 (6) were excluded; rejection rate 24%)
499 for analyses in SpN, 34 (18) participants retained (i.e., 7 (5) were excluded; rejection rate 17%) for
500 analyses in BbN, and 32 (18) participants retained (9 (5) were excluded; rejection rate 22%) for
501 analyses in Quiet (the numbers in brackets represent the numbers of young adults).

3.3.1 Regression results including data of both young and older adults

Statistics for the Best-Subset Regressions are shown as in **Tables 4 and 5**. Results that included data of both young and older adults were analyzed first. When SRTs were predicted by EEG signatures obtained in the respective noise types, SRTs were significantly correlated with Logit- θ -PLV (SpN, $t_{(28)} = -3.104$, $p = 0.004$; BbN, $t_{(31)} = -2.508$, $p = 0.018$; greater Logit- θ -PLV correlated with better SPiN) after PTA was controlled for (PTA_{High} for SpN and PTA_{Wide} for BbN) (**Table 4**). When SRTs were predicted by EEG signatures obtained in Quiet, a significant correlation was found between FFR_{ENV_F0} magnitude and SRT in BbN ($t_{(29)} = -2.698$, $p = 0.012$; greater FFR_{ENV_F0} magnitude correlated with better SPiN) after PTA_{Low} was controlled for (**Table 5**).

Table 4. Results for the Best-Subset Regressions in which SRTs were predicted by EEG signatures obtained in the corresponding noise types (i.e., SRT_{SpN} was predicted by EEGs obtained in SpN; SRT_{BbN} was predicted by EEGs obtained in BbN). DV refers to the dependent variables; β , CI, T , p , VIF refer to standardized β -coefficient, 95% confidence interval for standardized β , t values, p values (uncorrected) and variance inflation factors, respectively. N denotes the numbers of participants. F denotes the F values of the models (with corresponding p values in the brackets). Significant p values (< 0.05) are in bold. * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

DV	Participants	N	Predictors	β	CI	T	p	VIF	F
SRT _{SpN}	Young + older	31	Logit- θ -PLV _{SpN}	-0.499	[-0.828, -0.170]	-3.104	0.004**	1.040	6.104
			PTA _{High}	0.351	[0.022, 0.680]	2.183	0.038*	1.040	(0.006**)
	Young	17	PTA _{High}	0.445	[-0.048, 0.938]	1.922	0.074	1.000	3.695 (0.074)
	Older	14	FFR _{ENV_F0_SpN}	-0.475	[-0.903, -0.048]	-2.480	0.033*	1.042	6.120
			Logit- θ -PLV _{SpN}	-0.475	[-0.903, -0.047]	-2.475	0.033*	1.045	(0.012*)
			PTA _{Wide}	0.363	[-0.064, 0.791]	1.892	0.088	1.045	
SRT _{BbN}	Young + older	34	Logit- θ -PLV _{BbN}	-0.377	[-0.684, -0.070]	-2.508	0.018*	1.365	14.738
			PTA _{Wide}	0.814	[0.506, 1.122]	5.418	< 10⁻⁵***	1.365	(< 10⁻⁴***)
	Young	18	PTA _{Wide}	0.318	[-0.184, 0.820]	1.343	0.198	1.000	1.804 (0.198)
	Older	16	FFR _{TFS_H2_BbN}	-0.351	[-0.578, -0.124]	-3.368	0.006**	1.047	28.054
			Logit- θ -PLV _{BbN}	-0.769	[-1.005, -0.533]	-7.103	< 10⁻⁴***	1.128	(< 10⁻⁴***)
			PTA _{Wide}	0.696	[0.456, 0.936]	6.350	< 10⁻⁴***	1.157	

Table 5. Results for the Best-Subset Regression in which SRTs (both in SpN and BbN) were predicted by EEG signatures obtained in Quiet. DV refers to the dependent variables; β , CI, T , p , VIF refers to standardized β -coefficient, 95% confidence interval for standardized β , t values, p values (uncorrected) and variance inflation factors, respectively. N denotes the numbers of participants. F denotes the F values of the models (with corresponding p values in the brackets). Significant p values (< 0.05) are in bold. * = $p < 0.05$; ** = $p < 0.01$.

DV	Participants	N	Predictors	β	CI	T	p	VIF	F
SRT _{SpN}	Young + older	32	FFR _{ENV_F0_Quiet}	-0.307	[-0.661, 0.048]	-1.765	0.088	1.000	3.116 (0.088)
	Young	18	FFR _{ENV_F0_Quiet}	-0.508	[-1.001, -0.014]	-2.192	0.045*	1.161	3.304 (0.065)
			PTA _{Wide}	0.478	[-0.016, 0.973]	2.062	0.057	1.161	
	Older	14	FFR _{ENV_F0_Quiet}	-0.471	[-1.010, 0.069]	-1.919	0.081	1.068	3.375 (0.072)
			FFR _{TFS_H2_Quiet}	-0.534	[-1.074, 0.005]	-2.179	0.052	1.068	
SRT _{BbN}	Young + older	32	FFR _{ENV_F0_Quiet}	-0.410	[-0.720, -0.099]	-2.698	0.012*	1.012	7.425
			PTA _{Low}	0.460	[0.150, 0.771]	3.030	0.005**	1.012	(0.002**)
	Young	18	PTA _{Wide}	0.323	[-0.181, 0.829]	1.363	0.192	1.000	1.858 (0.192)
	Older	14	FFR _{ENV_F0_Quiet}	-0.455	[-0.857, -0.054]	-2.525	0.030*	1.069	7.622
			FFR _{TFS_H2_Quiet}	-0.551	[-0.969, -0.133]	-2.941	0.015*	1.153	(0.006**)
			PTA _{Wide}	0.441	[0.037, 0.845]	2.434	0.035*	1.081	

(A) Relation between SRT and Logit- θ -PLV obtained in corresponding noise types

(B) Relation between SRT_{BbN} and FFR_{ENV_F0} obtained in Quiet

Figure 7. Scatter plots (young + older) for significant partial correlations after controlling for PTAs that visualize the relation between (A) SRT and Logit- θ -PLV obtained in the corresponding noise types (see statistics in **Table 4**); and (B) SRT_{BbN} and FFR_{ENV_F0} magnitude obtained in Quiet (see statistics in **Table 5**). Red and blue dots represent young and older participants, respectively.

3.3.2 Regression results in the young and the older group separately

Best-Subset Regressions were then conducted separately for the young and older groups. In the young group, SRT in SpN was significantly correlated with FFR_{ENV_F0} magnitude obtained in Quiet after PTA_{Low} was controlled for ($t_{(15)} = -2.195$, $p = 0.045$; greater FFR_{ENV_F0} magnitude correlated with better SPiN; **Table 5**). No significant correlations were found between SRTs and EEG signatures in the corresponding noise types nor between SRT in BbN and in EEG signatures obtained in Quiet (**Tables 4 and 5**). In the older group, when SRTs were predicted by EEG signatures in the respective noise types, SRTs were significantly correlated with FFRs (FFR_{ENV_F0} for SpN: $t_{(10)} = -2.480$, $p = 0.033$; FFR_{TFS_H2} for BbN: $t_{(12)} = -3.368$, $p = 0.006$) and Logit- θ -PLV

(SpN: $t_{(10)} = -2.475$, $p = 0.033$; BbN: $t_{(12)} = -7.103$, $p < 10^{-4}$) (greater FFR magnitudes and Logit- θ -PLV correlated with better SPiN) after PTA_{Wide} was controlled for (**Table 4**). When SRTs were predicted by EEG signatures in Quiet, SRT in BbN was significantly correlated with FFR_{ENV_F0} and FFR_{TFS_H2} (FFR_{ENV_F0} : $t_{(10)} = -2.941$, $p = 0.030$; FFR_{TFS_H2} : $t_{(10)} = -2.525$, $p = 0.015$; greater FFR magnitudes correlated with better SPiN) after PTA_{Wide} was controlled for (**Table 5**).

Taken together, the regression analyses showed that: (1) when combining data from both young and older adults, by controlling for the degree of hearing loss (PTAs), SRTs can be predicted by cortical phase-locked responses (Logit- θ -PLV) to speech obtained in noise (greater Logit- θ -PLV associated with better SPiN) and by subcortical phase-locked responses to speech F_0 obtained in Quiet (greater FFR_{ENV_F0} magnitude associated with better SPiN); (2) SRTs are predicted by subcortical and cortical responses (FFR_{ENV_F0} , FFR_{TFS_H2} and Logit- θ -PLV) obtained in noise in the older group, not in the young group, and by subcortical responses obtained in Quiet ($FFR_{ENV_F0}/FFR_{TFS_H2}$) in the young (SRT_{SpN}) and the older group (SRT_{BbN}).

3.4 Cortical-subcortical/peripheral relationship

Correlations between Logit- θ -PLV and FFR signatures were conducted across the young and older adults to assess the cortical-subcortical/peripheral relationship in different noise types (Quiet, SpN and BbN). This was to see how such relationships are linked to SPiN. It was found that Logit- θ -PLV showed significant positive correlations with FFR_{TFS} magnitudes in SpN ($N = 31$; with FFR_{TFS_H2} , $r = 0.441$; $p = 0.013$; and with FFR_{TFS_F2F3} , $r = 0.560$, $p = 0.001$). As Logit- θ -PLV was significantly associated with SRT in SpN (see **Table 4**), FFR_{TFS} may then indirectly impact SPiN. No significant correlations were found between Logit- θ -PLV and FFR_{TFS} in other noise types (Quiet or BbN), nor between Logit- θ -PLV and $FFR_{ENV_F0}/FFR_{PLV_F0}$ in any noise type, even after age or PTAs were controlled for.

4 Discussion

The relationships between age-related phase-locked responses to speech in the auditory sensory systems and SPiN were investigated. Participant groups were young and older adults. The older adult group had peripheral hearing at high frequencies that ranged from normal to mild/moderate hearing loss. Such demographics are typical in normal aging populations (Popinath et al., 2009;

567 Humes et al., 2010). Other studies have looked specifically at normal-hearing older adults
568 (Anderson et al., 2011, 2012; Presacco et al., 2016a; Schoof and Rosen, 2016).

569 Different patterns of age effects on auditory phase-locked responses were revealed at the subcortical
570 and cortical levels: aging is related to decreases in subcortical responses (FFRs) and increases in
571 cortical responses (θ -PLV). Relationships between behavioral and neural performance showed that
572 subcortical responses obtained in quiet and cortical responses obtained in noise had significant
573 positive associations with SPiN, indicating that effects of aging on subcortical (decrease with aging)
574 and cortical (increase with aging) activities make different impacts on SPiN.

575 4.1 Age effect on subcortical phase-locked activity (FFRs)

576 Speech-evoked FFRs originate primarily from the brainstem (Chandrasekaran and Kraus, 2010;
577 Bidelman, 2018). Previous studies, including ones on older adults who had normal audiometric
578 hearing, reported that FFRs declined with age (Anderson et al., 2012; Presacco et al., 2016a). On
579 the other hand, evidence shows that hearing loss can reduce neural inhibition that allows greater
580 encoding of F_0 -rate envelope modulations in both animals (Kale and Heinz, 2010; Henry et al.,
581 2014; Zhong et al., 2014) and humans (Anderson et al., 2013; Goossens et al., 2019).

582 In the present study, no significant differences were found between young and older adults for
583 magnitudes of FFRs (FFR_{ENV_F0} , FFR_{TFS_H2} , or FFR_{TFS_F2F3}). However, additional ANCOVAs
584 showed that FFR_{ENV_F0} magnitude was significantly smaller in older than young adults after PTA_{Low}
585 (0.25 ~ 1 kHz) was controlled for. Also, PTA_{Low} correlated positively with the FFR_{ENV_F0} magnitude
586 (i.e., FFR_{ENV_F0} magnitude increased with low-frequency hearing loss). Therefore, these results are
587 in line with findings that encoding of envelopes at high-gamma frequencies corresponding to the F_0
588 range declines during aging when peripheral hearing is normal but increases when there is hearing
589 loss (Goossens et al., 2016, 2019), indicating the age-related neural declines in phase-locking and
590 reduced neural inhibition related to hearing loss.

591 However, caution is required when interpreting such results. FFR magnitudes are related to
592 participants' sensitivity to the syllable where greater audibility leads to higher magnitudes
593 (Ananthakrishnan et al., 2016). Therefore, the relation between PTA and FFR magnitudes probably
594 depends on the individual's degree of hearing loss. This could possibly explain why PTA_{Low} , but not
595 PTA_{High} , correlated with FFR magnitude, as the older adults in the present study had a much smaller

range of PTA_{Low} (< 30 dB HL) than PTA_{High} (from normal hearing to mild/moderate hearing loss) (see **Figure 1**) and the wider range of PTA_{High} may lead to a greater mix of reduced audibility and neural inhibition on FFRs. Taken together, effects of aging on subcortical phase-locked processing may be due to a combined effect of age-related hearing loss and the aging process that is independent of peripheral hearing.

4.2 Age effect on cortical phase-locked activity (θ -PLV)

θ -PLV was calculated as low-frequency (theta-band) phase-locked activity time-locked to the syllable onset. It reflects the tracking of speech envelopes (Luo and Poeppel, 2007; Howard and Poeppel, 2010; Peelle et al., 2013) and neural excitability in response to acoustic stimuli in the auditory cortex (Ng et al., 2013). The present study found that θ -PLV increased with age, consistent with previous studies (Tlumak et al., 2015; Goossens et al., 2016). This is also in line with findings that older adults have larger magnitudes of cortical auditory-evoked responses (Alain et al., 2014; Herrmann et al., 2013, 2016) and greater cortical tracking of speech envelopes (Presacco et al., 2016a).

Further ANCOVAs showed that, after PTAs (PTA_{Low} and PTA_{High}) were controlled for, the statistical effect of age was maintained. No statistical correlations occurred between PTAs and θ -PLV in ANCOVAs, indicating that the age-related increase cannot be explained by hearing loss. Consequently, the results may be attributable to hyperexcitability of the central auditory system during the aging process (Casparly et al., 2008). Note that in the present study, participants listened passively to a repeatedly-presented syllable that lacked higher-order semantic information. This indicates that the age effect may occur at an early stage in the auditory cortex.

It is noteworthy that the effects of age and PTAs on FFRs and θ -PLV differed, i.e., FFR_{ENV_F0} declined with age when PTA_{Low} was controlled for, whereas θ -PLV increased with age; FFR_{ENV_F0} increased with PTA_{Low} , whereas no association occurred between θ -PLV and PTAs. Thus these results imply that the mechanisms by which aging and peripheral hearing loss affect the phase-locked neural activities differ at subcortical and cortical levels.

4.3 Contributions of age effects on subcortical and cortical activities to SPiN

The central question for the present study concerned how the effects of age on phase-locked

activities contributed to impaired SPiN. This question can be approached by conducting regression analyses on participants' data across a wide range of ages. Here, linear regressions were conducted to study the neural-behavioral relationship using combined data from young and older adults, as well as young and older adults separately. The present study used an approach that differs from previous attempts (Presacco et al., 2016a; Goossens et al., 2018). Presacco et al. (2016a) studied the relation between speech-evoked phase-locked responses (FFRs and cortical tracking of speech envelopes) and SPiN in both young and older adults, but did not find any significant neural-SPiN correlations. A limitation in this study was that different types of background noise were used in the neural recording (single-talker background) and SPiN tasks (four-talker babble). Hence the neural-SPiN association was not appropriately assessed because acoustic and linguistic properties differed between the two types of background noise (see the discussion in Presacco et al., 2016). Goossens et al., (2018) investigated the relation between subcortical/cortical auditory steady-state responses (ASSRs) and SPiN in both normal-hearing and hearing impaired adults across ages and included age itself as an additional predictor when modeling the neural-behavioral relation. This did not fulfil the aim of testing how age-related neural factors contribute to SPiN. In the present study, regressions were conducted with neural signatures and SPiN under the same types of background noise. That is, SPiN in SpN and 16-talker BbN were predicted by neural data obtained in SpN and 16-talker BbN, respectively; neural data obtained in quiet were additionally used as predictors, as FFRs in quiet could be associated with SPiN (Anderson et al., 2011). Also, age itself was not included as a predictor.

Previous studies have argued that declines in subcortical speech-evoked FFRs are an important determinant of SPiN difficulty in older adults (Anderson et al., 2011, 2012; Presacco et al., 2016a). Consistent with this argument, the present study showed the FFR_{ENV_F0} declines with age and that decreased FFR_{ENV_F0} obtained in quiet was associated with decreased SPiN. At the cortical level, θ -PLV, which was shown to increase with age, was associated with increased SPiN. Thus this argues against the view that increased tracking of speech envelopes or auditory cortical excitability reflects a diminished excitation-inhibition balance in the neural network which results in impaired SPiN (Presacco et al., 2016a). An alternative explanation may be that, cortical hyperexcitability *per se* does not impair SPiN, but influences SPiN in an indirect way: increased cortical excitability is significantly related to decreased attention and inhibitory control in older adults which could, in turn, impair SPiN (Presacco et al., 2016b). The current results thus showed that impacts of age

655 effects on SPiN are different at subcortical (decreased $\text{FFR}_{\text{ENV_F0}}$ with age associated with decreased
656 SPiN) and cortical (increased θ -PLV with age associated with increased SPiN) levels.

657 The results also showed potential different mechanisms of neural-behavioral relationships across
658 ages, in which SPiN is better predicted by cortical and subcortical signatures obtained in noise in
659 the older compared to the young group. This could be because higher-level cognitive functions,
660 such as working memory and attention ability (not tested in the present study) that decline with age
661 (see 4.4.2 for further discussions). These factors could play greater roles in modelling the individual
662 variability of SPiN ability in young adults. The results could also arise because individual
663 variability of SPiN was relatively small in young adults (especially SRT_{BbN} , which had significantly
664 lower variability in the young than the older group according to Levene's test, see 3.1). This would
665 make SPiN more difficult to predict using linear regression.

666 Another interesting finding was that cortical, compared to subcortical, responses obtained in noise
667 is a more reliable predictor of SPiN, which is reflected not only in regression analyses, but also by
668 comparisons across noise types. It was shown that SPiN was significantly better in SpN than in
669 BbN, consistent with previous work (Rosen et al., 2013). Correspondingly, θ -PLV was also
670 significantly higher in SpN than in BbN. The explanation may be that, compared to SpN, BbN is an
671 informational masker that causes phonetic interference (Rosen et al., 2013) leading to worse cortical
672 envelope tracking of target speech, hence worse SPiN in BbN than in SpN. On the other hand, no
673 such effect was found for FFRs when SpN was compared with BbN. It could be that, cortical, but
674 not subcortical, phase-locked responses reflect the mechanisms underlying speech perception as a
675 function of noise types. Alternatively, signal quality of FFRs ($\text{FFR}_{\text{ENV_F0}}$) dropped significantly
676 from quiet to SpN/BbN (see **Figures 5 and 6**), which could hinder its predictability for SPiN in
677 different noise types. This might also explain why $\text{FFR}_{\text{ENV_F0}}$ obtained in quiet, compared to that
678 obtained in noise, showed better predictability (although $\text{FFR}_{\text{ENV_F0_SpN}}$ also significantly predicted
679 SRT_{SpN} in the older group, see **Table 4**).

680 **4.4 Limitations and future work**

681 **4.4.1 Participant samples**

682 The present study recruited older participants who covered a wide range of peripheral hearing from
683 normal to mild/moderate hearing loss. Although the degree of hearing loss was controlled for by

using PTAs as covariates during the analyses, a more direct and better approach would be to compare normal-hearing older adults with those who had hearing loss to clarify the respective effects of aging and hearing loss. However, sample size in the present study (< 17 older participants after data rejection) made it difficult to use this approach. Future research will need to recruit participants with bigger sample sizes with better control in hearing loss for older adults.

4.4.2 Possible role of higher-level functions for SPiN

Phase-locked activities were obtained in the present study with a paradigm where participants listened passively to repeated syllables without high-level linguistic features (semantic or syntactic information), hence reflecting early-stage sensory processing of speech in the auditory brainstem and cortex. Therefore, the present study focused mainly on how effects of age on sensory processing influenced SPiN, while higher-level cognitive functions were not measured.

The present results showed that older adults had significantly worse SPiN than young adults in BbN, but not in SpN, consistent with previous findings (Helfer and Freyman, 2008; Schoof and Rosen, 2014). However, no significant [Noise Type × Age] interactions were found for any of the EEG signatures, hence no sensory neural evidence was provided to explain this behavioral phenomenon. This leaves the alternative explanation through age-related declines in higher-level cognitive functions. Aging and age-related hearing loss are associated with declines in cognitive functions, such as working memory (WM) and attention (Lin et al., 2013). WM capacity is related to SPiN in older adults (Schoof and Rosen, 2014), while attention is critical for suppressing neural processing of background noise in SPiN (Rimmele et al., 2015) and such ability deteriorates during aging (Andres et al., 2006; Presacco et al., 2019). WM and attention may be particularly important for SPiN in BbN as they contribute to resisting the informational masking caused by BbN (Schneider et al., 2007; Shinn-Cunningham and Best, 2008).

As well as WM and attention capacity during SPiN, Mild Cognitive Impairment (MCI) could occur in some participants during normal aging (Petersen et al., 1999) and speech-evoked cortical and subcortical activities have been reported to be related to MCI that may have further affected SPiN (Anderson et al., 2013; Bidelman et al., 2017). For example, Bidelman et al. (2017) found that people with MCI showed hyperexcitability of subcortical and cortical responses to speech compared to controls. It is thus not clear how MCI contributes to the neural-behavioral relation found the in

713 the present study. Future studies should include measurements of cognitive functions and screening
714 for cognitive impairment in the model to better predict SPiN.

715 In addition, previous studies have shown that older adults have diminished sensitivity to temporal
716 fine structure (TFS) that leads to degraded SPiN (Hopkins and Moore, 2011; Fullgrabe et al., 2015).
717 However, although TFS processing at the resolved harmonics region ($\text{FFR}_{\text{TFS_H2}}$) was significantly
718 associated with SPiN in older adults (see **Table 4**, modelling SRT_{BbN}), the present study did not find
719 statistical evidence for age effects on TFS processing ($\text{FFR}_{\text{TFS_H2}}$ or $\text{FFR}_{\text{TFS_F2F3}}$). Diminished
720 sensitivity to TFS in SPiN in older adults may at least in part arise from declines in TFS processing
721 that involves higher-level cognitive function, compared to declines at early-stage sensory
722 processing. Such claims may be further tested in the future.

723 4.4.3 Paradigms used for EEG recording

724 In the present study, EEGs were recorded when participants listened to a repeatedly presented
725 vowel over a loudspeaker. The reason for using a loudspeaker rather than inserted earphones was
726 because the present older adult's data were obtained from our previous study (Mai et al., 2018).
727 This study included hearing-aid users as participants, for whom inserted earphones are not
728 appropriate (their data, however, were not included here, see *Footnote 1* for detailed explanations).
729 Changes in head and body positions were controlled by instructions to avoid movements and
730 monitored by a webcam. However, the possibility of slight movements, such as head jitters, cannot
731 be ruled out. Such possibilities can change the ear position and thus interfere with phase-locked
732 responses to high-frequency TFS properties ($\text{FFR}_{\text{TFS_H2}}$ and $\text{FFR}_{\text{TFS_F2F3}}$) in the free field. Future
733 research using inserted earphones thus is needed to confirm the current results relevant to FFR_{TFS} .

734 Furthermore, although arousal effects were controlled in the present study by restricting the data
735 used in analyses to those when participants were in the high arousal states, amount of attention to
736 the acoustic stimuli was not controlled during the EEG recording. While selective auditory attention
737 can modulate auditory cortical (Choi et al., 2013; Kong et al., 2014) and subcortical (Galbraith et
738 al., 2003; Hairston et al., 2013; Lehmann and Schönwiesner, 2014) electrophysiological responses,
739 it is not totally clear how they are affected by unpredictable changes in attention during passive
740 listening as in the present study. Therefore, additional tasks of active listening to target speech
741 stimuli under the corresponding noise types need to be conducted in the future to investigate

742 whether the current neural-behavioral relationships is replicable.

743 **4.4.4 Cortical-subcortical interaction**

744 Relationships between cortical and subcortical/peripheral responses were also tested. It was found
 745 that Logit- θ -PLV had significant positive correlations with $\text{FFR}_{\text{TFS_H2}}$ and $\text{FFR}_{\text{TFS_F2F3}}$ magnitudes
 746 in SPiN, indicating that FFR_{TFS} might indirectly contribute to SPiN via cortical-subcortical
 747 interactions (see 3.4). However, as they are correlations based on data across different participants,
 748 it is not known whether such correlations reflect cortical-subcortical connectivity within
 749 individuals, whether connectivity is afferent or efferent, and how the connectivity is associated with
 750 age or SPiN. It is necessary in future work to develop effective methods to quantify the strength and
 751 direction of cortical-subcortical interactions (directional phase relations between cortical responses
 752 and FFRs, e.g., [Bidelman et al., 2019](#)).

753 **4.4.5 Other factors that may influence the hearing loss**

754 Degree of hearing loss was tested via air-conduction in the present study. The audiogram for
 755 majority of the participants showed hearing loss at high frequencies (≥ 2 kHz) (**Figure 1**),
 756 consistent with the pattern of presbycusis which is usually sensorineural hearing disorders caused
 757 by age-related functional declines in the inner ear. Despite this, we cannot totally exclude that some
 758 of the participants may also suffer from conductive hearing loss. Future work should further include
 759 bone-conduction to test for conductive hearing loss in order to better clarify the roles of different
 760 types of hearing loss.

761 **4.5 Conclusions**

762 Subcortical (FFRs) and cortical (θ -PLV) phase-locked activities reflect how the brain processes
 763 fine-grained acoustic properties (F_0 , temporal fine structures and acoustic envelopes) that are crucial
 764 for SPiN. The present study hypothesized that effects of age on these activities should be associated
 765 with SPiN. Compared to young adults, it was found that older adults have smaller FFR magnitude
 766 (when low-frequency hearing loss was controlled for) and greater θ -PLV, illustrating distinct
 767 mechanisms of age effects at the subcortical and cortical levels. Smaller FFR magnitude reflects
 768 declines in subcortical phase-locking during aging and was associated with decreased SPiN, whilst
 769 greater θ -PLV reflects the neural hyperexcitability in the auditory cortex during aging and was

770 associated with increased SPiN. The current study thus provided evidence for different mechanisms
771 at the sensory subcortical and cortical levels by which age affects speech-evoked phase-locked
772 activities and SPiN. Future work need to be conducted by combining cognitive assessments to study
773 how higher-level cognitive functions influence such mechanisms and contribute to SPiN together
774 with sensory processing during aging.

775

776 **References:**

- 777 Aiken SJ, Picton TW. Envelope and spectral frequency-following responses to vowel sounds.
778 *Hearing Research*. 2008; 245:35-47.
- 779 Alain C. Effects of age-related hearing loss and background noise on neuromagnetic activity from
780 auditory cortex. *Frontiers in Systems Neuroscience*. 2014; 8:8.
- 781 Ananthakrishnan S, Krishnan A, Bartlett E. Human frequency following response: neural
782 representation of envelope and temporal fine structure in listeners with normal hearing and
783 sensorineural hearing loss. *Ear and Hearing*. 2016; 37:e91.
- 784 Anderson S, Parbery-Clark A, Yi HG, Kraus N. A neural basis of speech-in-noise perception in
785 older adults. *Ear and Hearing*. 2011; 32:750.
- 786 Anderson S, Parbery-Clark A, White-Schwdoch T, Kraus N. Aging affects neural precision of
787 speech encoding. *Journal of Neuroscience*. 2012; 32:14156-64.
- 788 Anderson S, White-Schwoch T, Parbery-Clark A, Kraus N. A dynamic auditory-cognitive system
789 supports speech-in-noise perception in older adults. *Hearing Research*. 2013; 300:18-32.
- 790 Andrés P, Parmentier FB, Escera C. The effect of age on involuntary capture of attention by
791 irrelevant sounds: a test of the frontal hypothesis of aging. *Neuropsychologia*. 2006; 44:2564-8.
- 792 Aviyente S, Bernat EM, Evans WS, Sponheim SR. A phase synchrony measure for quantifying
793 dynamic functional integration in the brain. *Human Brain Mapping*. 2011; 32:80-93.
- 794 Bench J, Kowal Å, Bamford J. The BKB (Bamford-Kowal-Bench) sentence lists for partially-
795 hearing children. *British journal of audiology*. 1979; 13:108-12.
- 796 Bidelman GM. Subcortical sources dominate the neuroelectric auditory frequency-following
797 response to speech. *NeuroImage*. 2018; 175:56-69.
- 798 Bidelman GM, Price CN, Shen D, Arnott SR, Alain C. Afferent-efferent connectivity between
799 auditory brainstem and cortex accounts for poorer speech-in-noise comprehension in older adults.
800 *Hearing research*. 2019; 107795:1-12.
- 801 Bidelman GM, Lowther JE, Tak SH, Alain C. Mild cognitive impairment is characterized by
802 deficient brainstem and cortical representations of speech. *Journal of Neuroscience*. 2017;
803 37(13):3610-20.
- 804 Boersma P, Weenink D. Praat: doing phonetics by computer [Computer program] Version 5.3.51,
805 Retrieved 2 June 2013.
- 806 Burnham KP, Anderson DR. Model selection and multimodel inference: a practical information-

807 theoretic approach. 2003; Springer Science & Business Media; 2003.

808 Carpenter AL, Shahin AJ. Development of the N1–P2 auditory evoked response to amplitude rise
809 time and rate of formant transition of speech sounds. *Neuroscience Letters*. 2013; 544:56-61.

810 Caspary DM, Ling L, Turner JG, Hughes LF. Inhibitory neurotransmission, plasticity and aging in
811 the mammalian central auditory system. *Journal of Experimental Biology*. 2008; 211:1781-91.

812 Chandrasekaran B, Kraus N. The scalp-recorded brainstem response to speech: Neural origins and
813 plasticity. *Psychophysiology*. 2010; 47:236-46.

814 Choi I, Rajaram S, Varghese LA, Shinn-Cunningham BG. Quantifying attentional modulation of
815 auditory-evoked cortical responses from single-trial electroencephalography. *Front Hum*
816 *Neurosci*. 2013; 7:115

817 Czisch M, Wehrle R, Kaufmann C, Wetter TC, Holsboer F, Pollmächer T, Auer DP. Functional MRI
818 during sleep: BOLD signal decreases and their electrophysiological correlates. *European Journal of*
819 *Neuroscience*. 2004; 20:566-74.

820 Dajani HR, Purcell D, Wong W, Kunov H, Picton TW. Recording human evoked potentials that
821 follow the pitch contour of a natural vowel. *IEEE Transactions on Biomedical Engineering*. 2005;
822 52:1614-8.

823 Fujihira H, Shiraishi K. Correlations between word intelligibility under reverberation and speech
824 auditory brainstem responses in elderly listeners. *Clinical Neurophysiology*. 2015; 126:96-102.

825 Füllgrabe C, Moore BC, Stone, MA. Age-group differences in speech identification despite
826 matched audiometrically normal hearing: contributions from auditory temporal processing and
827 cognition. *Frontiers in Aging Neuroscience*. 2010; 6:347.

828 Galbraith GC, Olfman DM, Huffman TM. Selective attention affects human brain stem frequency-
829 following response. *Neuroreport*. 2003;14:735-8.

830 Goossens T, Vercammen C, Wouters J, Wieringen AV. Aging affects neural synchronization to
831 speech-related acoustic modulations. *Frontiers in Aging Neuroscience*. 2016; 8:133.

832 Goossens T, Vercammen C, Wouters J, van Wieringen A. Neural envelope encoding predicts speech
833 perception performance for normal-hearing and hearing-impaired adults. *Hearing Research*. 2018;
834 370:189-200.

835 Goossens T, Vercammen C, Wouters J, van Wieringen A. The association between hearing
836 impairment and neural envelope encoding at different ages. *Neurobiology of Aging*. 2019; 74:202-
837 12.

838 Gopinath B, Rochtchina E, Wang JJ, Schneider J, Leeder SR, Mitchell P. Prevalence of age-related
839 hearing loss in older adults: Blue Mountains Study. *Archives of internal medicine*. 2009; 169:415-8.

840 Hairston WD, Letowski TR, McDowell K. Task-related suppression of the brainstem frequency
841 following response. *PLoS One*. 2013; 8:e55215.

842 Harris KC, Eckert MA, Ahlstrom JB, Dubno JR. Age-related differences in gap detection: Effects of
843 task difficulty and cognitive ability. *Hearing research*. 2010; 264:21-9.

844 Helfer KS, Freyman RL. Aging and speech-on-speech masking. *Ear and Hearing*. 2008; 29:87.

845 Henry KS, Kale S, Heinz MG. Noise-induced hearing loss increases the temporal precision of
846 complex envelope coding by auditory-nerve fibers. *Frontiers in Systems Neuroscience*. 2014; 8:20.

847 Herrmann B, Henry MJ, Scharinger M, Obleser J. Auditory filter width affects response magnitude

848 but not frequency specificity in auditory cortex. *Hearing Research*. 2013; 304:128-36.

849 Herrmann B, Henry MJ, Johnsrude IS, Obleser J. Altered temporal dynamics of neural adaptation in
850 the aging human auditory cortex. *Neurobiology of Aging*. 2016; 45:10-22.

851 Hopkins K, Moore BC. The effects of age and cochlear hearing loss on temporal fine structure
852 sensitivity, frequency selectivity, and speech reception in noise. *The Journal of the Acoustical*
853 *Society of America*. 2011; 130:334-49.

854 Howard MF, Poeppel D. Discrimination of speech stimuli based on neuronal response phase
855 patterns depends on acoustics but not comprehension. *Journal of neurophysiology*. 2010; 104:2500-
856 2511.

857 Humes LE, Busey TA, Craig JC, Kewley-Port D. The effects of age on sensory thresholds and
858 temporal gap detection in hearing, vision, and touch. *Attention, Perception, & Psychophysics*. 2009;
859 71:860-71.

860 Humes LE, Dubno JR. Factors affecting speech understanding in older adults. In: *The Aging*
861 *Auditory System*. Springer, New York, NY. 2010; p. 211-257.

862 Humes LE, Kewley-Port D, Fogerty D, Kinney D. Measures of hearing threshold and temporal
863 processing across the adult lifespan. *Hearing Research*. 2010; 264:30-40.

864 Hunter LL, Sanford CA. Tympanometry and Wideband Acoustic Immitance. In: *Handbook of*
865 *Clinical Audiology*, 7th edition. 2015; p.137-163.

866 Kale S, Heinz MG. Envelope coding in auditory nerve fibers following noise-induced hearing loss.
867 *Journal of the Association for Research in Otolaryngology*. 2010; 11:657-673.

868 Kong YY, Mullangi A, Ding N. Differential modulation of auditory responses to attended and
869 unattended speech in different listening conditions. *Hearing research*. 2014; 316:73-81.

870 Lakatos P, Chen CM, O'Connell MN, Mills A, Schroeder CE. Neuronal oscillations and
871 multisensory interaction in primary auditory cortex. *Neuron*. 2007; 53:279-92.

872 Lehmann A, Schönwiesner M. Selective attention modulates human auditory brainstem responses:
873 relative contributions of frequency and spatial cues. *PLoS One*. 2014; 9:e85442

874 Lin FR, Yaffe K, Xia J, Xue QL, Harris TB, Purchase-Helzner E, Satterfield S, Ayonayon HN,
875 Ferrucci L, Simonsick EM, Health ABC Study Group F. Hearing loss and cognitive decline in older
876 adults. *JAMA Internal Medicine*. 2013; 173:293-299.

877 Luo H, Poeppel D. Phase patterns of neuronal responses reliably discriminate speech in human
878 auditory cortex. *Neuron*. 2007; 54:1001-1010.

879 Mai G, Tuomainen J, Howell P. Relationship between speech-evoked neural responses and
880 perception of speech in noise in older adults. *The Journal of the Acoustical Society of America*.
881 2018; 143:1333-1345.

882 Mai G, Schoof T, Howell P. Modulation of phase-locked neural responses to speech during different
883 arousal states is age-dependent. *NeuroImage*. 2019; 189:734-744.

884 Martin N, Lafortune M, Godbout J, Barakat M, Robillard R, Poirier G, Bastien C, Carrier J.
885 Topography of age-related changes in sleep spindles. *Neurobiology of aging*. 2013; 34:468-476.

886 Morillon B, Liégeois-Chauvel C, Arnal LH, Bénar CG, Giraud AL. Asymmetric function of theta
887 and gamma activity in syllable processing: an intra-cortical study. *Frontiers in Psychology*. 2012;
888 3:248.

889 Noguchi Y, Fujiwara M, Hamano S. Temporal evolution of neural activity underlying auditory
890 discrimination of frequency increase and decrease. *Brain Topography*. 2015; 28:437-444.

891 Ng BS, Logothetis NK, Kayser C. EEG phase patterns reflect the selectivity of neural firing.
892 *Cerebral Cortex*. 2012; 23:389-398.

893 Oya H, Gander PE, Petkov CI, Adolphs R, Nourski KV, Kawasaki H, Howard MA, Griffiths TD.
894 Neural phase locking predicts BOLD response in human auditory cortex. *NeuroImage*. 2018;
895 169:286-301.

896 Peelle JE, Gross J, Davis MH. Phase-locked responses to speech in human auditory cortex are
897 enhanced during comprehension. *Cerebral Cortex*. 2013; 23:1378-1387.

898 Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild cognitive
899 impairment: clinical characterization and outcome. *Archives of Neurology*. 1999; 56(3):303-8.

900 Plomp R, Mimpen AM. Speech-reception threshold for sentences as a function of age and noise
901 level. *The Journal of the Acoustical Society of America*. 1979; 66:1333-1342.

902 Portas CM, Krakow K, Allen P, Josephs O, Armony JL, Frith CD. Auditory processing across the
903 sleep-wake cycle: simultaneous EEG and fMRI monitoring in humans. *Neuron*. 2000; 28:991-999.

904 Presacco A, Simon JZ, Anderson S. Evidence of degraded representation of speech in noise, in the
905 aging midbrain and cortex. *Journal of Neurophysiology*. 2016; 116:2346-2355.

906 Presacco A, Simon JZ, Anderson S. Effect of informational content of noise on speech
907 representation in the aging midbrain and cortex. *Journal of Neurophysiology*. 2016; 116:2356-2367.

908 Presacco A, Simon JZ, Anderson S. Speech-in-noise representation in the aging midbrain and
909 cortex: Effects of hearing loss. *PloS One*. 2019; 14(3):e0213899.

910 Rimmele JM, Golumbic EZ, Schröger E, Poeppel D. The effects of selective attention and speech
911 acoustics on neural speech-tracking in a multi-talker scene. *Cortex*. 2015; 68:144-154.

912 Schneider BA, Li L, Daneman M. How competing speech interferes with speech comprehension in
913 everyday listening situations. *Journal of the American Academy of Audiology*. 2007; 18(7):559-72.

914 Schneider BA, Hamstra SJ. Gap detection thresholds as a function of tonal duration for younger and
915 older listeners. *The Journal of the Acoustical Society of America*. 1999; 106:371-380.

916 Schoof T, Rosen S. The role of auditory and cognitive factors in understanding speech in noise by
917 normal-hearing older listeners. *Frontiers in Aging Neuroscience*. 2014; 6:307.

918 Schoof T, Rosen S. The role of age-related declines in subcortical auditory processing in speech
919 perception in noise. *Journal of the Association for Research in Otolaryngology*. 2016; 17:441-460.

920 Shinn-Cunningham BG, Best V. Selective attention in normal and impaired hearing. *Trends in*
921 *Amplification*. 2008; 12(4):283-99.

922 Skoe E, Kraus N. Auditory brainstem response to complex sounds: a tutorial. *Ear and Hearing*.
923 2010; 31:302.

924 Sommers MS, Gehr SE. Auditory suppression and frequency selectivity in older and younger
925 adults. *The Journal of the Acoustical Society of America*. 1998; 103:1067-74.

926 Stine RA. Graphical interpretation of variance inflation factors. *The American Statistician*. 1995;
927 49:53-56.

928 Thatcher RW. Coherence, phase differences, phase shift, and phase lock in EEG/ERP analyses.

929 Developmental neuropsychology. 2012; 37:476-96.

930 Tlumak AI, Durrant JD, Delgado RE. The effect of advancing age on auditory middle-and long-
931 latency evoked potentials using a steady-state-response approach. American journal of audiology.
932 2015; 24:494-507.

933 Tun PA, O'kane G, Wingfield A. Distraction by competing speech in young and older adult
934 listeners. Psychology and Aging. 2002; 17:453.

935 Tun PA, McCoy S, Wingfield A. Aging, hearing acuity, and the attentional costs of effortful
936 listening. Psychology and Aging. 2009; 24(3):761.

937 Waschke L, Wöstmann M, Obleser J. States and traits of neural irregularity in the age-varying
938 human brain. Scientific reports. 2017; 7:17381.

939 Wilf M, Ramot M, Furman-Haran E, Arzi A, Levkovitz Y, Malach R. Diminished auditory
940 responses during NREM sleep correlate with the hierarchy of language processing. PloS One. 2016;
941 11:e0157143.

942 Wong PC, Skoe E, Russo NM, Dees T, Kraus N. Musical experience shapes human brainstem
943 encoding of linguistic pitch patterns. Nature Neuroscience. 2007; 10:420.

944 Zhong Z, Henry KS, Heinz MG. Sensorineural hearing loss amplifies neural coding of envelope
945 information in the central auditory system of chinchillas. Hearing Research. 2014; 309:55-62.