Fibronectin-dependent tissue mechanics regulate the translation of segmentation clock oscillations into periodic somite formation

Patrícia Gomes de Almeida^{1,2,3}, Pedro Rifes^{1,#}, Ana Patrícia Martins-Jesus^{2,3}, Gonçalo G. Pinheiro^{1,2}, Raquel P. Andrade^{2,3,4,*} and Sólveig Thorsteinsdóttir^{1,*,¶}

¹ Centre for Ecology, Evolution and Environmental Change – cE3c, Departamento de Biologia Animal, Faculdade de Ciências, Universidade de Lisboa, 1740-016 Lisboa, Portugal

² Centre for Biomedical Research - CBMR, Universidade do Algarve, 8005-139 Faro, Portugal

³ Algarve Biomedical Center, Campus Gambelas, Edifício 2. Ala Norte, 8005-139 Faro, Portugal

⁴ Department of Medicine and Biomedical Sciences, Universidade do Algarve, 8005-139 Faro, Portugal

[#]Present address: The Novo Nordisk Foundation Center for Stem Cell Biology (DanStem), University of Copenhagen, 2200 Copenhagen N, Denmark

*These authors contributed equally.

[¶]Correspondence to: Sólveig Thorsteinsdóttir (solveig@fc.ul.pt)

Abstract:

Somitogenesis starts with cyclic waves of expression of segmentation clock genes in the presomitic mesoderm (PSM) and culminates with periodic budding of somites in its anterior-most region. How cyclic clock gene expression is translated into timely morphological somite formation has remained unclear. A posterior to anterior gradient of increasing PSM tissue cohesion correlates with increasing fibronectin matrix complexity around the PSM, suggesting that fibronectin-dependent tissue mechanics may be involved in this transition. Here we address whether the mechanical properties of the PSM tissue play a role in regulating the pathway leading to cleft formation in the anterior PSM. We first interfered with cytoskeletal contractility in the chick PSM by disrupting actomyosin-mediated contractility directly or via Rho-associated protein kinase function. Then we perturbed fibronectin matrix accumulation around the PSM tissue by blocking integrin-fibronectin binding or fibronectin matrix assembly. All four treatments perturbed *hairy1* and *meso1* expression dynamics and resulted in defective somitic clefts. A model is presented where a gradient of fibronectin-dependent tissue mechanics participates in the PSM wavefront of maturation by ensuring the correct spatio-temporal conversion of cyclic segmentation clock gene expression into periodic somite formation.

Keywords:

Fibronectin, Tissue mechanics, Segmentation Clock, Cleft Formation, Somitogenesis

Introduction:

Cells in the developing embryo are constantly receiving and integrating information, including mechanical signals generated by the adhesion to neighbor cells and/or the surrounding extracellular matrix (ECM). Cell-cell adhesion molecules and cell-ECM receptors, such as cadherins and integrins, respectively, are linked to the intracellular actomyosin cytoskeleton via intermediate proteins (Campbell and Humphries, 2011; Charras and Yap, 2018; Takeichi, 2014; Wolfenson et al., 2013). These adhesion complexes, called adhesomes, allow cells to perceive and respond to changes in their physical surroundings (Horton et al., 2016; Zaidel-Bar, 2013). Signaling events in adhesomes can impact the actomyosin cytoskeleton through the phosphorylation of non-muscle myosin II (NM II) which binds to actin and converts ATP into mechanical energy (Zaidel-Bar et al., 2015). The resulting actomyosin contractility leads to changes in cell shape and can transmit signals from integrin adhesomes to cadherin adhesomes and vice versa, as well as from the cell surface to the nucleus (Burute and Thery, 2012; Mui et al., 2016; Wolfenson et al., 2019). In this way, through continuous probing of their mechanical environment, cells adjust their shape, functions and behaviors, such as proliferation, differentiation, cell polarity and migration (Burute and Thery, 2012; Mui et al., 2016; Wolfenson et al., 2019). While morphogens have been extensively studied as major chemical regulators of developmental processes (Marek and Kubíček, 1981; Slack, 1987; Tiedemann, 1976), the importance of mechanical forces in embryo development has, until recently, received less attention (Marek and Kubíček, 1981; Slack, 1987; Tiedemann, 1976). It is, however, becoming increasingly clear that the ability of cells to sense and respond to mechanical signals regulates numerous basic developmental processes (e.g. Barriga et al., 2018; Brunet et al., 2013; Hiramatsu et al., 2013; Smutny et al., 2017).

One of the most conspicuous morphogenetic events during early vertebrate embryogenesis is the formation of somites, which are the source of axial skeleton and skeletal muscle precursor cells (Christ et al., 2007). Somites are spheres of epithelioid cells that are formed periodically from the anterior portion of the mesenchymal presomitic mesoderm (PSM), bilateral

to the axial structures (Bailey and Dale, 2015). Temporal control of somite formation is dependent on cyclic waves of expression of segmentation clock genes, many of which are targets of the Notch signaling pathway (Dequéant et al., 2006; Masamizu et al., 2006; Palmeirim et al., 1997). These waves periodically sweep the PSM in a posterior to anterior direction (Aulehla and Pourquié, 2010; Bailey and Dale, 2015) and, as they reach the anterior PSM, oscillations slow down and then arrest (Morimoto et al., 2005; Shih et al., 2015). The transcription factor *Mesp2/Meso1* is upregulated downstream of the segmentation clock in the anterior PSM, leading to Eph/Ephrin signaling and somitic cleft formation (Barrios et al., 2003; Nakajima et al., 2006; Saga, 2012; Watanabe et al., 2009), followed by progressive cell rearrangements into a somite (Martins et al., 2009; Morimoto et al., 2005; Shih et al., 2015).

Fibronectin is essential for somite formation in all vertebrate models studied to date (George et al., 1993; Georges-Labouesse et al., 1996; Goh et al., 1997; Koshida et al., 2005; Kragtorp and Miller, 2007; Rifes et al., 2007; Sato et al., 2007). Fibronectin matrix assembly is a complex cell-dependent process that requires the engagement and unfolding of globular fibronectin by the major fibronectin matrix assembly receptor, the $\alpha 5 \beta 1$ integrin, followed by fibrillogenesis involving fibronectin-fibronectin binding (Mao and Schwarzbauer, 2005; Singh et al., 2010). In the chick, a fibronectin matrix starts being assembled around the caudal PSM tissue and then gets progressively denser as the tissue matures (Rifes et al., 2007; Rifes and Thorsteinsdóttir, 2012). This results in the formation of a gradient of fibronectin matrix complexity along the PSM (Rifes and Thorsteinsdóttir, 2012), which correlates with a posterior to anterior gradient in cell density (Bénazéraf et al., 2010; Lawton et al., 2013; Mongera et al., 2018). At the rostral end, fibronectin is required for the polarization of N-cadherin and epithelialization of peripheral cells to form a somite (Martins et al., 2009; Rifes et al., 2007). Interestingly, adhesion to a fibronectin substrate was noted as a regulator of the oscillations of the segmentation clock gene *Lfng* in cultured mouse tailbud cells (Hubaud et al., 2017; Lauschke et al., 2013). Cell adhesion to fibronectin was linked to dampening and eventual arrest of *Lfng* oscillations (Hubaud et al.,

2017), reminiscent of what is observed in the anterior PSM prior to somite epithelialization. However, whether the mechanical properties of the PSM tissue play a role in the slowing down of segmentation clock oscillations and their conversion into segments remains unknown.

In this study, we addressed the involvement of PSM tissue mechanics in the regulation of both segmentation clock gene expression dynamics and subsequent somite formation using the chick embryo as a model. First, we experimentally perturbed actomyosin contractility by blocking either NM II ATPase activity or Rho-associated protein kinase (ROCK). We then addressed the role of the fibronectin matrix surrounding the PSM by blocking integrin-fibronectin binding through RGD or by perturbing extracellular fibronectin fibrillogenesis. We found that each one of the four treatments resulted in abnormal segmentation clock oscillations, mis-positioning of *meso1* expression in the rostral PSM and perturbations in somite morphogenesis. These results strongly suggest that fibronectin-dependent PSM tissue mechanics play a role in converting segmentation clock oscillations into periodic somite formation.

Results:

Intracellular actomyosin contractility is required for timely segmentation clock oscillations and *meso1* activation

In the chick embryo, sequential pairs of somites bud off from the anterior PSM every 90 min, which corresponds to the period of segmentation clock oscillations (Figure 1 A). To investigate the involvement of intracellular actomyosin contractility in this process, the expression of the segmentation clock gene *hairy1* (Palmeirim et al., 1997) was analyzed in the PSM of embryo half explants cultured in the presence of either Blebbistatin, which directly inhibits the ATPase activity of NM II and consequently all actomyosin contractility (direct inhibition), or RockOut, a chemical inhibitor of ROCK I and II (ROCK I/II) enzymes involved in activating NM II (indirect inhibition) (Figure 1 B, C; Ringer et al., 2017; Straight et al., 2003; Yarrow et al., 2005). The contralateral control sides were cultured with an equal volume of DMSO.

Explants cultured for 6 hours in each experimental condition presented significantly altered *hairy1* expression. *hairy1* expression was either absent or in a different phase of the cycle relative to the contralateral control in 80% of the Blebbistatin- (n=7/9) and RockOut-treated explants (n= 8/10; Figure 1 D, a-c), suggesting that temporal control of *hairy1* oscillations requires the generation of tensional cues mediated by NM II and ROCK I/II activity.

Segmentation clock oscillations are required for the correct spatial and temporal upregulation of *Mesp2* in the anterior PSM (Niwa et al., 2011; Saga and Takeda, 2001; Sato et al., 2002), which regulates downstream targets needed for the formation of the somitic cleft (Saga, 2012). The expression of the chick *Mesp2* homolog, *meso1*, was altered in Blebbistatin- (n=9/11) and RockOut-treated explants (n=10/10; Figure 1 D, d-f). *meso1* expression was either absent, located more rostrally or presented a different number of bands of expression, clearly indicating that the normal cycles of activation and suppression of *meso1* in the rostral PSM were altered. Importantly, *meso1* expression was also perturbed after 3 hours in culture with Blebbistatin or RockOut (n=8/9 and 5/5, respectively; Supplementary Figure 1 A-B), corresponding to an effect within two segmentation clock cycles. Furthermore, timely downregulation of *dll1* in the anterior-most PSM (Palmeirim et al., 1998), which normally occurs downstream of *Meso1/Mesp2* activity (Takahashi et al., 2000; Takahashi et al., 2003), was not observed in either Blebbistatin- (80%, n=8/10) or RockOut-treated (75%, n=6/8) explants after 10.5h of culture (Figure 1 D, g-i). Together these data indicate that interfering with actomyosin contractility perturbs three sequential events: the spatio-temporal expression dynamics of *hairy1*, timely *meso1* expression and the downregulation of *dll1* expression in the anterior-most PSM.

Alterations in somite formation were observed concomitantly with the perturbations in *hairy1* and *meso1* expression. Control explants formed an average of 3.6 somites after 6 hours, consistent with a 90 min periodicity (Figure 1 E), while contralateral RockOut-treated explants formed 3.1 somites, of which only the first two somites were clearly individualized, while subsequent somite-like condensations were poorly defined (Figure 1 E). After 10.5 hours, control explants had formed an average of 7.5 somites, while RockOut-treated explants formed 7.1 somites of which the first two appeared normal, but the remaining ones were ill-defined (Figure 1 E). Importantly, explants cultured with Blebbistatin were unable to form more than 1-2 somite-like aggregates after 6 hours, or even 10.5 hours, of culture (Figure 1 E), evidencing an absolute requirement for NM II ATPase activity in somite formation. These effects were not due to an increase in apoptosis (Supplementary Figure 2 A-D).

Our data reveal a previously unknown role for NM II- and ROCK I/II-mediated cell contractility in the temporal regulation of the segmentation clock, *meso1* expression, *dll1* downregulation and, consequently, in somite formation.

NM II activity is required for somite cleft formation and cell polarization

Somite formation involves a mesenchymal-to-epithelial transition (MET) of anterior PSM cells (Martins et al., 2009; Saga, 2012). To determine to what extent this process is impaired upon NM II or ROCK I/II inhibition, we performed a detailed analysis of the morphology of S0 to SIII in explants after a 6 hour culture period (regions e-b in Figure 1 A).

In control explants, S0 showed apically enriched N-cadherin (Figure 2 A, D) and some *zonula occludens* protein 1 (ZO-1) accumulation was observed apically (Figure 2 B, D, arrowheads). Peripheral cell alignment occurred (Figure 2 C, D) and fibronectin matrix was detected in the nascent somitic clefts (Figure 2 E, F, arrows). In contrast, in explants cultured with Blebbistatin, N-cadherin was homogeneous (Figure 2 G, J), ZO-1 immunostaining was absent (Figure 2 H, J) and neither peripheral cell alignment (Figure 2 I, J) nor fibronectin matrix accumulation within the

188 tissue was observed (Figure 2 K, L). Furthermore, the continuous and dense fibronectin matrix
189 observed surrounding the rostral PSM in control explants was disrupted in Blebbistatin-treated
190 explants (compare Figure 2 E and K, arrowheads). Moreover, the characteristic nuclear alignment
191 and F-actin apical enrichment observed in control SI (Supplementary Figure 3 A-C) was absent in
192 Blebbistatin-treated explants and no signs of somitic boundaries could be detected
193 (Supplementary Figure 3 D-F). We conclude that exposure of the S-IV and S-III regions of the PSM
194 (regions e and d in Figure 1 A) to Blebbistatin for 6 hours completely blocks their capacity to form
195 somites.

196
197 **Figure 2. NM II inhibition abolishes N-cadherin and ZO-1 polarization and impairs fibronectin**
198 **fibrillogenesis.**
199 **(A-X)** Sagittal optical sections of control explants (A-F, M-R) and their Blebbistatin-treated contralateral
200 halves after 6 hours of culture, immunostained for N-cadherin (Ncad), ZO-1, fibronectin (FN), stained for
201 DNA and imaged at S0 (A-L; region e in Figure 1 A) and SIII (M-X; region b in Figure 1 A) levels. S0 of DMSO-
202 treated explants shows apically enriched N-cadherin (A) and ZO-1 (B, arrowheads) and peripheral nuclei are
203 aligned (C, D), while no signs of polarized cell-cell adhesions (G, H) or nuclear alignment (I, J) are found in
204 the contralateral Blebbistatin-treated explant. Apical polarization of N-cadherin and ZO-1 is maintained in
205 SIII cells of DMSO-treated explants (M, N, arrowheads), but no polarized N-cadherin (S) or ZO-1 (T,
206 arrowheads) are observed in contralateral Blebbistatin-treated halves. Nuclear alignment (U, arrowheads)
207 and fibronectin assembly around somites (K, W, arrowheads) are also deficient in Blebbistatin-treated
208 explants compared to contralateral controls (E, Q, arrowheads). Arrows in E show fibronectin assembly in
209 the nascent somitic cleft. Rostral to the left and dorsal on top. Dashed lines mark borders of S0 (B, H) and

SIII (N, T). FN – fibronectin. Ncad – N-cadherin. ZO-1 – *Zonula occludens* 1. Blebb – Blebbistatin. Scale bars: 50 μ m.

We next turned our attention to SII and SIII somites after 6 hours of culture. These were at stage S-II and S-I in the PSM, respectively (regions c and b in Figure 1 A), when the explants were placed in culture and had thus already upregulated *meso1* (Buchberger et al., 1998). As before, in the presence of Blebbistatin, apical enrichment of N-cadherin failed to occur (Figure 2 S, V), ZO-1 was only detected in a few small foci (Figure 2 T, V) and, although a fibronectin matrix was present, it appeared less dense (Figure 2 W, X). An incipient nuclear alignment was sometimes observed (Supplementary Figure 3 K, arrowheads), but cells did not polarize their F-actin into apically enriched adhesion belts (compare Supplementary Figure 3 G-I with J-L). Epithelial tissues other than somites (e.g. ectoderm and neural tube) did not present significant alterations after incubation with Blebbistatin (Supplementary Figure 4). Altogether, these results point to an indispensable role for NM II activity for the MET underlying somite formation.

RockOut-treated explants also showed perturbations in somite formation, although to a lesser extent (Figure 1 E). When compared to control explants (Figure 3 A-D; Supplementary Figure 5 A-C), RockOut treatment resulted in incomplete somitic clefts, such that S0 shared the somitocoel with SI and sometimes also with SII (Fig 3 E-H, arrows, Supplementary Figure 5 D-F, arrows). In contrast to the accumulation of fibronectin in the nascent clefts in controls (Figure 3 B, arrow), no fibronectin was observed in the incipient somitic clefts of RockOut-treated explants (Figure 3 F, arrows). These results suggest that ROCK I/II activity in the S-IV and S-III regions of the PSM (regions e and d in Figure 1 A) is required for the formation of individualized somites. In contrast, when the rostral-most PSM (stage S-I before culture, region b in Figure 1 A) was exposed to RockOut for 6 hours, it was indistinguishable from control explants, showing apical accumulation of ZO-1 (Figure 3 I, M, arrowhead) and N-cadherin (Supplementary Figure 5, G, J), nuclear alignment (Figure 3, K, O, Supplementary Figure 5, H, K) and a complete, fibronectin

236 matrix-containing cleft (Figure 3 J, N). This indicates that ROCK I/II activity is not required for S-I to
 237 develop into a somite.
 238 Altogether, our data indicate that intracellular actomyosin contractility plays a role in
 239 periodic somite cleft formation, and that ROCK I/II-independent NM II activity is essential for
 240 MET.

241 **Figure 3. ROCK I/II inhibition impairs morphological somite formation, leading to deficient ZO-1**
 242 **polarization and fibronectin assembly.**
 243 (A-P) Sagittal sections of explants cultured in control (DMSO) medium (A-D, I-L) and their contralateral
 244 RockOut-treated halves (E-H, M-P) at S0 (A-H; region e in Figure 1 A) and SIII (I-P; region b in Figure 1 A)
 245 levels, immunostained for ZO-1 (first column), fibronectin (second column), stained for DNA (third column)
 246 and the respective merged image (fourth column). Explants were cultured for 6 hours. S0 in control explants
 247 show normal accumulation of ZO-1 (A, arrowhead), fibronectin assembly in the nascent cleft (B, arrow) and
 248 nuclear alignment (C, arrowhead). In contrast, S0 in contralateral RockOut-treated explants fails to form a
 249 clear cleft (E-H, arrows), although ZO-1 is generally polarized (E, arrowhead) and nuclei are aligned (G,
 250 arrowhead). At SIII level, both explants show normal ZO-1 polarization (I, M, arrowheads), fibronectin
 251 assembly (J, N) and nuclear alignment (K, O). Rostral on the left and dorsal on top. FN – fibronectin. ZO-1 –
 252 *Zonula occludens* protein 1. Scale bar: 50 μ m.
 253

Blocking integrin-fibronectin binding perturbs segmentation clock oscillations and somitic cleft formation

Our next aim was to address the requirement for the fibronectin ECM surrounding the PSM in regulating segmentation clock oscillations and somite formation. PSM cells bind to the RGD motif of fibronectin through the $\alpha 5 \beta 1$ integrin, an interaction that plays a crucial role during somitogenesis (Girós et al., 2011; Yang et al., 1993). αv integrins, which have been described to bind the RGD motif and partially compensate for the absence of the $\alpha 5 \beta 1$ integrin in the mouse (Yang et al., 1999), are not detected in the PSM of the chick embryo (Gomes de Almeida et al., 2016). We cultured embryo half explants in the presence of a linear RGD peptide, which competes with fibronectin for integrin binding (Huveneers et al., 2008; Pierschbacher and Ruoslahti, 1984), and compared them to contralateral control explants (Figure 4 A, B). RGD-treated explants formed ill-defined somite-like condensations, although in approximately the same number as the contralateral control (Figure 4 C). This was not due to cell death (Supplementary Figure 2 E-F). Concomitantly, RGD-treated explants displayed alterations in *hairy1* (Figure 4 D, a-c), *meso1* (Figure 4 D, d-f) and *dll1* expression patterns (Figure 4 D, g-i), evidencing that integrin-fibronectin interactions via RGD are required for proper segmentation clock oscillations, *meso1* positioning and timely downregulation of *dll1* in the anterior PSM.

Figure 4. Integrin-fibronectin binding through RGD is required for timely *hairy1* and *meso1* expression and *dll1* downregulation.

(A) Schematic representation of the explant culture system. Posterior explants of HH11-14 chick embryos were bisected along the midline and cultured for 6 to 12 hours. One side of the explant was cultured with RGD, while the contralateral half was cultured in control medium. **(B)** Schematic representation of the action of RGD. RGD competes with the RGD-binding pockets of integrins, interfering with their binding to the ECM. **(C)** Number of somites (black bars) or somite-like condensations (dotted bars) formed in culture in control

and RGD-treated explants. p values were calculated using a paired Student's t-test. *p<0.05. **(D)** Expression of *hairy1* (a, b), *meso1* (d, e) and *dll1* (g, h) in RGD-treated explants and contralateral controls at representative timepoints of culture. Straightened images of respective explant pairs (right) aligned by SIV. Rostral is on top. Percentage of RGD-treated explants with altered *hairy1*, *meso1* and *dll1* expression compared to the contralateral controls is shown in c, f and i, respectively. Impairing integrin-fibronectin binding with RGD alters *hairy1* and *meso1* expression relative to contralateral controls at 6 hours of culture onwards (a-f, arrowheads). *dll1* expression was altered at 9 hours of culture onwards (g-i, arrowheads). Scale bar: 500 μ m.

Figure 5. Inhibition of integrin-fibronectin binding impairs somitic cleft formation.

(A-X) Longitudinal (left) and sagittal (right) sections of explants cultured in control medium (A-F, M-R) and their contralateral RGD-treated halves (G-L, S-X) at S0 (A-L; region e in Figure 1 A) and SII (M-X; region c in Figure 1 A) levels, immunostained for N-cadherin (first column), fibronectin (fourth column) and stained for DNA (second and fifth columns). Third and sixth columns show the respective merged images. Explants were cultured for 6 hours. At the S0 level, control explants show normal apical accumulation of N-cadherin (A, arrowheads) and nuclear alignment (B, E, arrowheads) in peripheral somitic cells as well as fibronectin assembly in the nascent cleft (D, F, arrowheads). All of these are deficient in their RGD-treated contralateral halves (G-L, arrowheads). At the level of SII, N-cadherin accumulation and nuclear alignment occurs normally in both control and RGD-treated explants (M, N, S, T, arrowheads), but fibronectin assembly between adjacent somites (P, V, arrowheads) and cleft formation (Q, W, arrowheads) are deficient in RGD-treated explants compared to contralateral controls. Ncad – N-cadherin; FN – fibronectin. Scale bars: 50 μ m.

When compared to the contralateral control, the area corresponding to S0 after 6 hours of culture with RGD (region e in Figure 1 A) showed deficient nuclear alignment (Figure 5 B, E, H, K, arrowheads) and N-cadherin polarization (Figure 5 A, G, arrowheads), accompanied by deficient fibronectin assembly in the nascent cleft (Figure 5 D, J arrowheads). At the level of SII (region c in Figure 1 A), complete somite individualization was impaired in RGD-treated explants (Figure 5 N, Q, T, W, arrowheads) and, although N-cadherin polarization appeared normal (Figure 5 M, S, arrowheads), cleft formation (Figure 5 Q, W, arrowheads; R, X) and fibronectin assembly between adjacent somites was deficient (Figure 5 P, V, arrowheads; R, X).

These findings implicate cell-ECM interactions, mediated by integrin-fibronectin binding via the RGD motif, in temporal control of *hairy1* expression, correct positioning of *meso1* expression, downregulation of *dll1* in the anterior PSM and somite morphogenesis.

Impaired fibronectin matrix assembly results in altered segmentation clock dynamics, *meso1* expression and defects in somite morphogenesis

Another way to assess the relevance of the extracellular fibronectin matrix on the events leading up to somite formation is to perturb fibronectin assembly in the PSM and somites. To this end, primitive streak-stage embryos were electroporated with a construct expressing the 70kDa fibronectin fragment, a dominant-negative inhibitor of fibronectin matrix assembly (Figure 6 A, B; McKeown-Longo and Mosher, 1985; Sato et al., 2017). 70kDa-electroporated embryos exhibited a disrupted fibronectin matrix, composed of thinner fibrils when compared to control pCAGGs-electroporated embryos (Supplementary Figure 6 A; also see Figure 7 A, B). These embryos displayed multiple morphological defects, including kinked neural tube and detached tissues as well as perturbations in somite morphogenesis, which are all reminiscent of phenotypes obtained in previous studies interfering with fibronectin matrix deposition and/or with fibronectin-integrin binding (Supplementary Figure 6 B, C; Drake et al., 1992; Drake and Little, 1991; George et al., 1993; Girós et al., 2011; Takahashi et al., 2007). Although the average number of somite-like

structures formed in 70kDa-electroporated embryos was similar to the number of somites in controls (Figure 6 C), the former were ill-defined, often appearing fused or crammed (Supplementary Figure 6 B e-f, C), closely resembling the somite-like condensations formed in RockOut- and RGD-treated explants (Figures 3 and 5).

Next, we sought to evaluate the impact of inhibiting fibronectin matrix assembly on the molecular machinery underlying somite formation. Unilateral electroporation of the 70kDa fragment was performed (Figure 6 A) to allow the direct comparison of gene expression patterns within the same embryo. We observed a significant increase in the frequency of perturbations in the expression of embryonic clock genes *hairy1* ($p < 0.01$) and *hairy2* ($p < 0.05$), as well as in *meso1* ($p < 0.01$) and *dll1* ($p < 0.01$) expression, when compared with embryos electroporated with pCAGGs alone (Figure 6 D).

Figure 6. Impairing fibronectin assembly by electroporation with a 70kDa expressing vector perturbs segmentation clock oscillations, timely *meso1* expression and *dll1* downregulation.

(A) Schematic representation of the electroporation strategy. PSM/ectoderm progenitors of primitive-streak stage embryos were electroporated with either a pCAGGs GFP-expressing vector (pCAGGs) alone, or co-electroporated with a 70kDa-expressing vector (70kDa) and were incubated for about 26 hours. (B) Schematic representation of the action of 70kDa. 70kDa disrupts the assembly of fibronectin matrix by competitively binding to the N-terminal self-assembly domains of the protein, impairing fibronectin fibril formation. (C) Number of somites (black bars) or somite-like condensations (dotted bars) formed in pCAGGs- and 70kDa-electroporated embryos after 26 hours. p values were calculated using a paired Student's t-test. (D) Examples of the expression of *hairy1* (a-b), *hairy2* (d-e), *meso1* (g-h) and *dll1* (j-l) in pCAGGs- (top row) and 70kDa-electroporated embryos (middle row). Electroporated side is on right (a, b, g) or left (d, e, h, j, k). Perturbing the assembly of fibronectin on one side of the PSM leads to an asymmetric pattern of *hairy1* (b, arrowheads), *hairy2* (e, arrowheads), *meso1* (h, arrowheads) and *dll1* expression (k, arrowheads) in a higher percentage of embryos than in controls. Percentage of pCAGGs- and 70kDa-electroporated embryos with asymmetric expression of *hairy1*, *hairy2*, *meso1* and *dll1* between the electroporated PSM and the contralateral non-electroporated control PSM is shown in c, f, i and l, respectively. The number of embryos with an asymmetric pattern was significantly higher in 70kDa-electroporated embryos for all four genes studied. p values were calculated using a Chi-square test. *** p<0.01. Rostral is on top. Scale bar: 200 μ m.

Consistent with these data, 70kDa-electroporated embryos have deficiencies in somite morphogenesis which, in severe cases, leads to incomplete somitic clefts (Figure 7 A, B k, w, arrows). Peripheral cells of nascent somites of 70kDa-electroporated embryos did, however, accumulate ZO-1 apically (Figure 7 B e, h, i, l) which was maintained as the somites matured (Figure 7 B q, u, t, x). Nevertheless, these somites were abnormal in shape and appeared smaller in severely affected embryos (Figure 7 B, i-l, u, x). In fact, the SI of the electroporated sides of 70kDa-treated embryos were significantly smaller in width than those of the contralateral control, while SV was significantly shorter in length (n=151; Supplementary Figure 7). In addition to defects in somite morphology, the ectoderm and endoderm were separated from the paraxial mesoderm in 70kDa-electroporated embryos, indicating that their fibronectin matrix was insufficient to hold these tissues together (brackets in Figure 7 B g, k, s, w).

We conclude that proper fibronectin matrix assembly in the PSM is required for timely clock gene expression dynamics, positioning of *meso1* expression and *dll1* downregulation in the rostral-most PSM, as well as for the complete separation and morphogenesis of somites.

Figure 7. Somite morphology of 70kDa-electroporated embryos is severely compromised.

(A) Sagittal sections of embryos electroporated with pCAGGs and 70kDa, the latter with a severe phenotype. Arrows point to deficient somitic clefts. Rostral to the left and dorsal on top. **(B)** Sagittal views of embryos electroporated with pCAGGs (a-d, m-p) and 70kDa (e-l, q-x) with either mild (e-h, q-t) or severe (i-l, u-x) phenotypes, at S0 (a-l) and SII (m-x) levels, immunostained for ZO-1 (first column) and fibronectin (second column) and stained for DNA (third column). Fourth column shows the merge of the respective channels. S0 of pCAGGs- and 70kDa-electroporated embryos all polarize ZO-1 normally (a, e, i, arrowheads) and this is

maintained at SII levels (m, q, u), but the fibronectin matrix surrounding the somites of 70kDa-treated embryos is disrupted compared to pCAGGs-electroporated embryos (b, f, j, v, arrows). Somites of 70kDa-electroporated embryos are also severely detached from both the ectoderm and endoderm compared to embryos electroporated only with pCAGGS (third column, brackets), and the fibronectin matrix connecting the endoderm to the somites is severely compromised (second column, empty arrowheads). Somites of embryos electroporated with 70kDa with more severe defects also fail to fully detach from adjacent somites (k, w, arrows). Rostral to the left and dorsal to the top. Dashed lines indicate altered somite morphology. Experiments were performed in 6 (pCAGGs) and 7 (70kDa) biological replicates. FN – fibronectin. ZO-1 – *Zonula occludens* protein 1. Scale bars: 50 μ m.

Discussion:

Fibronectin-dependent tissue mechanics coordinate segmentation clock dynamics and cleft formation

We have identified fibronectin-dependent tissue mechanics as a regulator of segmentation clock gene expression and the positioning of the presumptive somitic cleft in the chick embryo. Four independent treatments interfering with the mechanical properties of the PSM (Figure 8 A) consistently lead to asymmetric patterns of *hairy1* expression as well as incorrect positioning of *meso1* expression on the experimental versus the control sides of the same embryo. The similarity of the phenotypes obtained in experiments impairing fibronectin fibrillogenesis, cell-fibronectin interactions via RGD and blocking ROCK (Figure 8 B) suggests that the fibronectin matrix, the RGD-binding $\alpha 5 \beta 1$ integrin, and ROCK-dependent actomyosin contractility are part of the same pathway. The $\alpha 5 \beta 1$ integrin can transduce mechanical signals by ROCK activation (Schiller et al., 2013) and this requires the binding of $\alpha 5 \beta 1$ to two sites of fibronectin, namely the RGD and the synergy site (Friedland et al., 2009). Surprisingly, although the RGD site of fibronectin is crucial for somitogenesis (Girós et al., 2011), removing the synergy site of fibronectin (*Fn1^{syn/syn}*) had no effect on mouse embryonic development (Benito-Jardón et al., 2017). However, cells expressing αv -integrins form strong adhesions on fibronectin lacking the synergy site and can compensate for the inability of $\alpha 5 \beta 1$ to mediate adhesion strengthening on this fibronectin (Benito-Jardón et al., 2017), and αv -integrins can partially compensate for the complete absence

of $\alpha 5 \beta 1$ during mouse somitogenesis (Yang et al., 1999). Alternatively, ROCK can be activated indirectly, for example through cadherin engagement and subsequent adherens junction formation (Burute and Thery, 2012; Schwartz and DeSimone, 2008), which occurs upon fibronectin-induced polarization of peripheral PSM cells (Martins et al., 2009).

It is well established that Notch signaling plays a role in the segmentation clock and is also required for timely *meso1* activation (Saga, 2012). Hence, the mechanical environment may be regulating Notch signaling in the PSM. In agreement with our results, chicken embryos electroporated with RNAi constructs against integrin $\beta 1$ showed alterations in *hairy2*, *lnfg* and *meso1* expression in the PSM (Rallis et al., 2010). Mouse embryos where the fibronectin RGD site has been substituted with an RGE sequence (*Fn1*^{RGE/RGE}) also showed asymmetric and/or dampened expression of *Lnfg* and *Hes7* in the PSM (Girós et al., 2011) and *EphA4*, a direct target of *Mesp2* in the anterior PSM (Nakajima et al., 2006), was diffusely expressed or absent (Girós et al., 2011). Finally, combined roles of integrin $\alpha 5 \beta 1$ and Notch are required for zebrafish somitogenesis (Jülich et al., 2005).

Exactly how tissue mechanics regulate Notch signaling is unknown. It is becoming increasingly clear that mechanics play a crucial role in Notch activation (Gordon et al., 2015; Luca et al., 2017; Meloty-Kapella et al., 2012) and sustained Notch signaling in the *Drosophila* notum requires actomyosin contractility in both signal sending and receiving cells (Hunter et al., 2019). Intriguingly, clock oscillations in mouse tailbud PSM explants cultured on fibronectin are sustained in the presence of a ROCK inhibitor, suggesting that ROCK activity must be low for the maintenance of clock oscillations and that an increase in ROCK activity normally stops segmentation clock oscillations in this system (Hubaud et al., 2017). Moreover, in the same study, cell adhesion to fibronectin was linked to nuclear localization of Yes-associated protein (YAP), an intracellular sensor of cell mechanics, and dampening and eventual arrest of *Lnfg* oscillations was found to be YAP-dependent (Hubaud et al., 2017). ROCK-mediated actomyosin contractility is known to promote the nuclear localization of YAP in several cell types (Piccolo et al., 2014) and YAP-null

mouse mutants (Morin-Kensicki et al., 2006) have a phenotype very similar to that of integrin $\alpha 5$ -null mutants (Yang et al., 1993) and *Fnl1*^{RGE/RGE} embryos (Girós et al., 2011), suggesting that they contribute the same processes during early embryo development. Thus, it is conceivable that increased fibronectin-dependent tissue cohesion may translate into increased ROCK activity and actomyosin contractility, promoting sustained Notch signaling and nuclear localization of YAP, leading to the dampening and eventual arrest of clock oscillations. Further studies are needed to test this hypothesis.

Altogether, our results show that perturbation of the normal PSM tissue mechanics leads to a dysregulation of segmentation clock oscillations and the mispositioning of the segmental border, indicating that the mechanical properties of the PSM modulate Notch signaling and coordinate the translation of clock oscillations into periodic segmental border formation.

Somite cleft formation and cell epithelialization have different mechanical requirements

In the rostral PSM, *Mesp2*/*Meso1* activates the expression of *EphA4*, which interacts with *EphrinB2* in cells rostral to the presumptive cleft, causing cell-cell repulsion and the formation of an incipient cleft (Nakajima et al., 2006; Watanabe et al., 2009). Then, fibronectin matrix assembly within the cleft stabilizes it (Jülich et al., 2015; Rifés and Thorsteinsdóttir, 2012) and promotes the epithelialization of cells rostral to the cleft (Martins et al., 2009). This can be defined as the first step of morphological somite individualization. The second step is defined as the complete epithelialization of the remaining cells of the nascent somite and lasts until SII, when all somitic cells have acquired a spindle-like shape and are organized into a rosette (Martins et al., 2009).

RockOut-, RGD- and 70kDa-treated embryos (Figure 8 A, B) all show perturbations in segmentation clock gene expression, abnormal positioning of *meso1* and defects in the somitic clefts. Although somite morphology is also perturbed, the acquisition of the spindle-shape cell morphology which occurs as S0 develops into SII does not appear to be significantly perturbed. Thus, the first step of morphological somite formation is affected but the second step is not. In

contrast, Blebbistatin-treated explants not only have the defects listed above, but cells that had already upregulated *meso1* before the addition of the drug and formed an incipient cleft during culture, were completely unable to epithelialize. In fact, Blebbistatin-treated explants formed only 1 or 2 somites, and their cells did not acquire the elongated, spindle-shape typical of SI and SII somites. Hence, Blebbistatin affects both steps of morphological somite formation.

RockOut targets NM II activity indirectly by inhibiting ROCK I and II, two of the kinases that activate NM II (Newell-Litwa et al., 2015). In contrast, Blebbistatin directly targets the NM II ATPase. Our results thus raise the possibility that the acquisition of the spindle-shaped morphology of cells may be dependent on another NM II activator. Interestingly, Ca^{++} /calmodulin signaling can activate NM II and inhibiting calmodulin was shown to block the acquisition of this morphology during chick somitogenesis (Chernoff and Hilfer, 1982).

A gradient of fibronectin-dependent tissue mechanics as a player in the PSM wavefront of maturation

While the waves of Notch oscillatory activity travel through the entire length of the PSM, they are only translated into segments in the anterior-most region of the tissue. Opposing gradients of Fgf/Wnt and Retinoic acid (RA) in the PSM are thought to define the wavefront, which marks the region where PSM cells become competent for somite formation (Hubaud and Pourquié, 2014). Rostral to the wavefront, segmentation clock oscillations progressively slow down until they reach a halt and cells become part of a somite (Palmeirim et al., 1997).

Herein, we propose a model (Figure 8 C) where a posterior to anterior gradient of fibronectin-dependent tissue cohesion is interpreted by the PSM cells as an increasing gradient of mechanical tension (Figure 8 C). We propose that the fibronectin-dependent tensional state of the PSM at the level of the wavefront acts as a threshold that activates a mechanotransduction signaling cascade, ensuring the correct spatio-temporal conversion of the cyclic expression of segmentation clock genes into periodic *meso1* expression, which in turn defines the next somitic

487 cleft (Figure 8 C). In support of this hypothesis, a gradient of increasing paraxial mesoderm stiffness
 488 from the tail to rostral somites has been identified in the chick embryo (Bénazéraf et al., 2017;
 489 Marrese et al., 2019). During its maturation, the PSM thus integrates a combination of chemical
 490 and mechanical signals, namely gradients of Fgf/Wnt and RA (Aulehla and Pourquié, 2010) and,
 491 simultaneously, a progressive increase in fibronectin-dependent tissue cohesion (Figure 8 C).
 492 Fibronectin matrix-dependent tissue mechanics would thus be a key contributor to the PSM
 493 wavefront of maturation in that it regulates where and when the next somitic cleft is positioned
 494 in the anterior PSM (Figure 8 C). Hence, we propose that a mechanotransduction pathway
 495 downstream of fibronectin plays a major role in the translation of cyclic waves of expression of
 496 segmentation clock genes into the periodic morphogenesis of somites.

497

Figure 8. Working model illustrating a fibronectin-dependent mechanotransduction pathway in chick somite formation.

(A) Schematic representation of the targets of experimental inhibition used in this study. (B) Effect of the different treatments on somitogenesis. Interfering with NM II (Blebb) or ROCK I/II (RockOut) activity, integrin-fibronectin binding (RGD), or fibronectin matrix assembly (70kDa), leads to altered segmentation clock dynamics, misregulated *meso1* expression, impaired or deficient cleft formation and alterations in somite morphology. Interfering directly with NM II further disrupts full somite epithelialization. (C) Working model proposing that an increase of fibronectin matrix complexity and tissue cohesion in the PSM, from caudal to rostral, is sensed by cells as a tensional gradient, with anterior PSM cells being in the stiffest environment. This increased tensional gradient occurs concomitantly with a decrease in Fgf/Wnt levels and an increase in RA. As cells reach the wavefront and sense a threshold of stiffness (due to fibronectin-dependent tissue cohesion), ROCK activity is increased and so is actomyosin contractility. This in turn slows down clock oscillations and stabilizes Notch signaling, which activates *meso1* expression in the S-II/S-I region, setting the stage for cleft formation and *dll1* downregulation in the rostral-most PSM later on. As the clefts form, somites bud off as SI. Finally, ROCK-independent NM II activity is required for full somite epithelialization into SII.

Materials and Methods:

Embryos

Fertilized chicken (*Gallus gallus*) eggs were obtained from commercial sources (Sociedade Agrícola Quinta da Freiria or Pintobar Exploração Avícola, Lda, Portugal) and incubated at 37.5°C in a humidified chamber until the desired HH stage (HH4 or HH11-14; Hamburger and Hamilton, 1992). Somite nomenclature is according to Pourquié and Tam (2001).

Embryo explant culture and chemical treatments

Explant tissues of HH11-14 embryos were collected and cultured as previously described (Palmeirim et al., 1997; Rifes et al., 2007). Embryos were bisected along the midline and then cut transversally rostral to somites IV and Hensen's node. The two contralateral halves thus retained half of the neural tube and notochord as well as the first four somites and the PSM, with all remaining neighboring tissues intact. Explants were placed on top of a polycarbonate filter floating on M199 medium supplemented with 10% chick serum, 5% fetal calf serum and 100 U/ml of penicillin and streptomycin (Palmeirim et al., 1997). Explants were then cultured at 37°C with 5% CO₂ from 6 to 12 hours.

InSolution™Blebbistatin (Calbiochem) and RockOut (Calbiochem) diluted in DMSO were used at a final concentration of 50 µM in culture medium. Equal volumes of DMSO (Sigma-Aldrich) were used as control for both drugs. The linear RGD peptide (GRGDS - G4391, Sigma) was diluted in culture medium and used at 0.9 mM, while control explants were cultured in medium only. RGD peptide efficiency was confirmed in a cell adhesion assay (Danen et al., 2002; Pierschbacher and Ruoslahti, 1984) before using it on explants.

Embryo electroporation and *ex ovo* culture

HH4-5 embryos were electroporated on one (randomly selected) side of the primitive streak in the presumptive PSM and/or ectoderm and cultured *ex ovo* using the Early Chick culture

method (Chapman et al., 2001). The electroporation mixture contained plasmid DNA at 0.5-1 µg/µl mixed with 0.4% Fast Green for visualization. Embryos were submerged in an electroporation chamber filled with Tyrode's saline and three pulses of 6-9 V, 50 ms each, at 350 ms intervals were applied. Control embryos were electroporated with pCAGGs containing a GFP reporter (pCAGGs-GFP; abbreviated pCAGGs). pCAGGs-70kDa qFN1 was kindly provided by Yuki Sato (Sato et al., 2017) and was co-electroporated with the pCAGGs-GFP plasmid in experimental embryos (treatment abbreviated 70kDa). Electroporated embryos were screened for GFP after fixation to select embryos with an intense signal on only one side to process for whole mount morphological analysis, *in situ* hybridization experiments and transverse sectioning. For morphological analysis in sagittal sections, the embryo side electroporated with pCAGGs was compared to the 70kDa-electroporated side of other same stage embryos.

Cryosectioning and immunohistochemistry

Cryosectioning was performed on embryo explants and whole embryos fixed in 4% paraformaldehyde in 0.12 M phosphate buffer containing 4% sucrose and processed for cryoembedding. Fixed samples were washed in 0.12 M phosphate buffer with 4% and 15% sucrose and then embedded in 7.5% gelatin in 0.12 M phosphate buffer containing 15% sucrose, frozen on dry ice-chilled isopentane and stored at -80°C until sectioning. Cryostat sections (10-30 µm) were processed for immunofluorescence as previously described (Gomes de Almeida et al., 2016). Permeabilization of sections was performed with 0.2% Triton-X100 in phosphate buffered saline (PBS). 5% bovine serum albumen (BSA) or a combination of 1% BSA and 10% Normal Goat Serum (NGS) in PBS were used for blocking depending on the presence or absence of anti-fibronectin antibodies, respectively. Primary and secondary antibodies were diluted in 1% BSA in PBS. Sections were incubated with primary antibodies overnight at 4°C and with secondary antibodies for 1 hour at room temperature. For whole-mount immunodetection, explants were fixed in 4% paraformaldehyde in PBS and processed as previously described (Martins et al., 2009; Rifes and

Thorsteinsdóttir, 2012). 1% Triton-X100 in PBS was used for permeabilization and 1% BSA in PBS was used for blocking and antibody dilution. Antibody incubation was performed overnight at 4°C.

The following primary antibodies were used: anti-ZO-1 (Zymed, #40-2200, 1:100 or Invitrogen, #33-9100, 1:100); anti-N-cadherin (BD Biosciences, #610920, 1:100); anti-fibronectin (Sigma, #F-3648, 1:400), anti-activated caspase3 (Cell Signaling, #9661, 1:1000) and anti-GFP (Invitrogen, #A11122, 1:100). For F-actin staining we used Alexa 488-conjugated phalloidin (Invitrogen, 1:40) and for staining DNA we used ToPro3 (Invitrogen, 1:500) in conjunction with ribonuclease A (Sigma, 10 µg/ml), 4% Methyl Green (Sigma, diluted 1:250; Prieto et al., 2015) or 4',6-diamidino-2-phenylindole (DAPI, 5µg/ml in PBS with 0.1% Triton-X100). For detection of the primary antibodies the adequate secondary goat anti-mouse and anti-rabbit Alexa 488-, Alexa 568- or Alexa 546-conjugated F'ab fragments from Invitrogen were used (#A-11017, #A-21069, #A-11071, #A-11019, #A-11070, 1:1000). Immunohistochemistry was performed on at least 6 different explants/embryos and the respective controls for each treatment (Blebbistatin n=13; RockOut n=15; RGD n=13; 70kDa n=7/pCAGGs n=6).

In situ hybridization

In situ hybridization using DIG-labeled RNA probes was performed as described previously (Henrique et al., 1995) with minor alterations (Gomes de Almeida et al., 2016). RNA probes were synthesized from linearized plasmids: *dll1* (Henrique et al., 1995), *meso1* (Buchberger et al., 1998), *hairy1* (Palmeirim et al., 1997) and *hairy2* (Jouve et al., 2000).

Statistical analysis

Paired Student's t-tests were performed to assess for differences in the number of somites formed in Blebbistatin-, RockOut- and RGD-treated explants relative to the respective controls, and in embryos electroporated with pCAGGs only and pCAGGs + 70kDa. Differences in the frequency of morphological and gene expression phenotypes found in 70kDa-electroporated

embryos compared to pCAGGs-electroporated control embryos was tested through a Chi-square test. Differences in somite size between pCAGGs- or 70kDa-electroporated sides compared to the control (non-electroporated) sides of embryos was tested through a nested ANOVA. The side in which the embryo was electroporated (left or right) was nested in the treatment (non-electroporated, pCAGGs-electroporated or 70kDa-electroporated) to account for a potential variability between the two sides. Statistical significance was set at $p < 0.05$. Statistical analyses were performed in Statistica 10 (<https://statistica.software.informer.com/10.0/>), Graphpad Prism 5 (<https://graphpad-prism.software.informer.com/5.0/>) and RStudio (<https://rstudio.com/>).

Sample preparation and imaging

Whole mount explants were gradually dehydrated in methanol and cleared in methylsalicylate (Sigma-Aldrich) as described previously (Martins et al., 2009; Rifes and Thorsteinsdóttir, 2012), except for phalloidin-labelled embryos and explants, where a shorter series of ethanol dehydration series was used. Cryostat sections were mounted in Vectashield (Vector Laboratories) or in 5mg/ml propyl gallate in glycerol/PBS (9:1) with 0.01% azide. Immunofluorescence images were taken on a confocal Leica SPE microscope, following imaging acquisition steps described previously (Rifes and Thorsteinsdóttir, 2012). Imaging of electroporated embryos and explants processed for *in situ* hybridization was performed using a Zeiss LUMAR V12 Stereoscope coupled to a Zeiss AxioCam 503 color 3MP camera. Image analysis was performed using Fiji v. 1.49 (<https://imagej.net/Fiji>) software. Image histogram corrections and, when appropriate, maximum intensity projections of immunofluorescence confocal stacks were produced in Fiji and exported as TIFF files. When applicable, contiguous images were stitched together into a single image using the pairwise stitching Fiji plugin (Preibisch et al., 2009). For the analysis of *in situ* hybridization patterns along the PSM explants, the Fiji plugin Straighten (Kocsis et al., 1991) was used.

Acknowledgements:

We thank Dr. Yuki Sato for generously sharing the pCAGGs-q70kDa construct and Inês Fragata for help with the statistical analysis. This work was supported by Fundação para a Ciência e a Tecnologia (FCT, Portugal) projects PTDC/SAU-OB/103771/2008, PTDC/BEXBID/5410/2014, UID/BIA/00329/2013, UID/BIM/04773/2019 CBMR, and FCT scholarships SFRH/BD/86980/2012 (PGA) and SFRH/BD/37423/2007 (PR). Imaging and image analysis were done in the Microscopy Facility at the Faculty of Sciences of the University of Lisbon and the Light Microscopy Unit of CBMR-UAlg, nodes of the Portuguese Platform for BioImage (reference PPBI-POCI-01-0145-FEDER-022122). Finally, we are grateful to Isabel Palmeirim and all members of our laboratories for their support and helpful discussions.

Competing interests:

The authors declare no competing or financial interests.

Author contributions:

Conceptualization: P.G.A., P.R., R.P.A., S.T.; Methodology: P.G.A., P.R., R.P.A., S.T.; Validation: P.G.A., P.R., A.P.M.J., G.G.P., R.P.A., S.T.; Formal analysis: P.G.A.; Investigation: P.G.A., P.R., A.P.M.J., G.G.P.; Resources: R.P.A., S.T.; Writing - original draft: P.G.A., S.T.; Writing - review & editing: P.G.A., P.R., A.P.M.J., G.G.P., R.P.A., S.T.; Visualization: P.G.A., P.R., R.P.A., S.T.; Supervision: R.P.A., S.T.; Project administration: P.G.A., R.P.A., S.T.; Funding acquisition: R.P.A., S.T.

References:

- Aulehla A, Pourquié O. 2010. Signaling gradients during paraxial mesoderm development. *Cold Spring Harb Perspect Biol* 2:1–17. doi:10.1101/cshperspect.a000869
- Bailey C, Dale K. 2015. Somatogenesis in Vertebrate Development. ELS. John Wiley & Sons, Ltd: Chichester. Chichester, UK: John Wiley & Sons, Ltd. pp. 1–15.

doi:10.1002/9780470015902.a0003820.pub2

Barriga EH, Franze K, Charras G, Mayor R. 2018. Tissue stiffening coordinates morphogenesis by triggering collective cell migration in vivo. *Nature* **554**:523–527. doi:10.1038/nature25742

Barrios A, Poole RJ, Durbin L, Brennan C, Holder N, Wilson SW. 2003. Eph/Ephrin signaling regulates the mesenchymal-to-epithelial transition of the paraxial mesoderm during somite morphogenesis. *Curr Biol* **13**:1571–82. doi:10.1016/j.cub.2003.08.030

Bénazéraf B, Beaupeux M, Tchernookov M, Wallingford A, Salisbury T, Shultz A, Shultz A, Huss D, Pourquoié O, François P, Lansford R. 2017. Multi-scale quantification of tissue behavior during amniote embryo axis elongation. *Development* **144**:4462–4472. doi:10.1242/dev.150557

Bénazéraf B, Francois P, Baker RE, Denans N, Little CD, Pourquoié O. 2010. A random cell motility gradient downstream of FGF controls elongation of an amniote embryo. *Nature* **466**:248–252. doi:10.1038/nature09151

Benito-Jardón M, Klapproth S, Gimeno-LLuch I, Petzold T, Bharadwaj M, Müller DJ, Zuchtriegel G, Reichel CA, Costell M. 2017. The fibronectin synergy site re-enforces cell adhesion and mediates a crosstalk between integrin classes. *Elife* **6**:1–24. doi:10.7554/elife.22264

Brunet T, Bouclet A, Ahmadi P, Mitrossilis D, Driquez B, Brunet A-C, Henry L, Serman F, Béalé G, Ménager C, Dumas-Bouchiat F, Givord D, Yanicostas C, Le-Roy D, Dempsey NM, Plessis A, Farge E. 2013. Evolutionary conservation of early mesoderm specification by mechanotransduction in Bilateria. *Nat Commun* **4**:2821. doi:10.1038/ncomms3821

Buchberger A, Seidl K, Klein C, Eberhardt H, Arnold HH. 1998. cMeso-1, a novel bHLH transcription factor, is involved in somite formation in chicken embryos. *Dev Biol* **199**:201–15. doi:10.1006/dbio.1998.8919

Burute M, Thery M. 2012. Spatial segregation between cell–cell and cell–matrix adhesions. *Curr Opin Cell Biol* **24**:628–636. doi:10.1016/j.ceb.2012.07.003

Campbell ID, Humphries MJ. 2011. Integrin structure, activation, and interactions. *Cold Spring*

671 *Harb Perspect Biol* **3**:a004994–a004994. doi:10.1101/cshperspect.a004994

672 Chan CJ, Heisenberg C-P, Hiiragi T. 2017. Coordination of morphogenesis and cell-fate
673 specification in development. *Curr Biol* **27**:R1024–R1035. doi:10.1016/j.cub.2017.07.010

674 Chapman SC, Collignon J, Schoenwolf GC, Lumsden A. 2001. Improved method for chick whole-
675 embryo culture using a filter paper carrier. *Dev Dyn* **220**:284–9. doi:10.1002/1097-
676 0177(20010301)220:3<284::AID-DVDY1102>3.0.CO;2-5

677 Charras G, Yap AS. 2018. Tensile Forces and Mechanotransduction at Cell–Cell Junctions. *Curr*
678 *Biol* **28**:R445–R457. doi:10.1016/j.cub.2018.02.003

679 Chernoff EA, Hilfer SR. 1982. Calcium dependence and contraction in somite formation. *Tissue*
680 *Cell* **14**:435–49. doi:10.1016/0040-8166(82)90038-6

681 Christ B, Huang R, Scaal M. 2007. Amniote somite derivatives. *Dev Dyn* **236**:2382–2396.
682 doi:10.1002/dvdy.21189

683 Danen EHJ, Sonneveld P, Brakebusch C, Fässler R, Sonnenberg A. 2002. The fibronectin-binding
684 integrins $\alpha 5 \beta 1$ and $\alpha v \beta 3$ differentially modulate RhoA-GTP loading, organization of cell
685 matrix adhesions, and fibronectin fibrillogenesis. *J Cell Biol* **159**:1071–1086.
686 doi:10.1083/jcb.200205014

687 Dequéant M-L, Glynn E, Gaudenz K, Wahl M, Chen J, Mushegian A, Pourquié O. 2006. A complex
688 oscillating network of signaling genes underlies the mouse segmentation clock. *Science*
689 **314**:1595–8. doi:10.1126/science.1133141

690 Drake CJ, Davis LA, Hungerford JE, Little CD. 1992. Perturbation of beta 1 integrin-mediated
691 adhesions results in altered somite cell shape and behavior. *Dev Biol* **149**:327–38.
692 doi:10.1016/0012-1606(92)90288-R

693 Drake CJ, Little CD. 1991. Integrins play an essential role in somite adhesion to the embryonic
694 axis. *Dev Biol* **143**:418–21. doi:10.1016/0012-1606(91)90092-H

695 Friedland JC, Lee MH, Boettiger D. 2009. Mechanically activated integrin switch controls
696 $\alpha 5 \beta 1$ function. *Science* **323**:642–4. doi:10.1126/science.1168441

George EL, Georges-Labouesse EN, Patel-King RS, Rayburn H, Hynes RO. 1993. Defects in mesoderm, neural tube and vascular development in mouse embryos lacking fibronectin. *Development* **119**:1079–91.

Georges-Labouesse EN, George EL, Rayburn H, Hynes RO. 1996. Mesodermal development in mouse embryos mutant for fibronectin. *Dev Dyn* **207**:145–56. doi:10.1002/(SICI)1097-0177(199610)207:2<145::AID-AJA3>3.0.CO;2-H

Girós A, Grgur K, Gossler A, Costell M. 2011. $\alpha 5 \beta 1$ integrin-mediated adhesion to fibronectin is required for axis elongation and somitogenesis in mice. *PLoS One* **6**:e22002. doi:10.1371/journal.pone.0022002

Goh KL, Yang JT, Hynes RO. 1997. Mesodermal defects and cranial neural crest apoptosis in alpha5 integrin-null embryos. *Development* **124**:4309–19.

Gomes de Almeida P, Pinheiro GG, Nunes AM, Gonçalves AB, Thorsteinsdóttir S. 2016. Fibronectin assembly during early embryo development: A versatile communication system between cells and tissues. *Dev Dyn* **245**:520–35. doi:10.1002/dvdy.24391

Gordon WR, Zimmerman B, He L, Miles LJ, Huang J, Tiyanont K, McArthur DG, Aster JC, Perrimon N, Loparo JJ, Blacklow SC. 2015. Mechanical Allostery: Evidence for a Force Requirement in the Proteolytic Activation of Notch. *Dev Cell* **33**:729–36. doi:10.1016/j.devcel.2015.05.004

Hamburger V, Hamilton HL. 1992. A series of normal stages in the development of the chick embryo. *Dev Dyn* **195**:231–272. doi:10.1002/aja.1001950404

Henrique D, Adam J, Myat A, Chitnis A, Lewis J, Ish-Horowicz D. 1995. Expression of a Delta homologue in prospective neurons in the chick. *Nature* **375**:787–790. doi:10.1038/375787a0

Hiramatsu R, Matsuoka T, Kimura-Yoshida C, Han S-W, Mochida K, Adachi T, Takayama S, Matsuo I. 2013. External Mechanical Cues Trigger the Establishment of the Anterior-Posterior Axis in Early Mouse Embryos. *Dev Cell* **27**:131–144. doi:10.1016/j.devcel.2013.09.026

Horton ER, Humphries JD, James J, Jones MC, Askari JA, Humphries MJ. 2016. The integrin

723 adhesome network at a glance. *J Cell Sci* **129**:4159–4163. doi:10.1242/jcs.192054

724 Hubaud A, Pourquié O. 2014. Signalling dynamics in vertebrate segmentation. *Nat Rev Mol Cell*

725 *Biol* **15**:709–721. doi:10.1038/nrm3891

726 Hubaud A, Regev I, Mahadevan L, Pourquié O. 2017. Excitable dynamics and Yap-dependent

727 mechanical cues drive the segmentation clock. *Cell* **171**:668–682.e11.

728 doi:10.1016/j.cell.2017.08.043

729 Hunter GL, He L, Perrimon N, Charras G, Giniger E, Baum B. 2019. A role for actomyosin

730 contractility in Notch signaling. *BMC Biol* **17**:12. doi:10.1186/s12915-019-0625-9

731 Huveneers S, Truong H, Fässler R, Sonnenberg A, Danen EHJ. 2008. Binding of soluble fibronectin

732 to integrin alpha5 beta1 - link to focal adhesion redistribution and contractile shape. *J Cell*

733 *Sci* **121**:2452–62. doi:10.1242/jcs.033001

734 Jouve C, Palmeirim I, Henrique D, Beckers J, Gossler A, Ish-Horowicz D, Pourquié O. 2000. Notch

735 signalling is required for cyclic expression of the hairy-like gene HES1 in the presomitic

736 mesoderm. *Development* **127**:1421–1429. doi:10.1016/S0092-8674(00)80451-1

737 Jülich D, Cobb G, Melo AM, McMillen P, Lawton AK, Mochrie SGJ, Rhoades E, Holley SA. 2015.

738 Cross-scale integrin regulation organizes ECM and tissue topology. *Dev Cell* **34**:33–44.

739 doi:10.1016/j.devcel.2015.05.005

740 Jülich D, Geisler R, Holley SA, Tübingen 2000 Screen Consortium. 2005. Integrin alpha5 and

741 delta/notch signaling have complementary spatiotemporal requirements during zebrafish

742 somitogenesis. *Dev Cell* **8**:575–86. doi:10.1016/j.devcel.2005.01.016

743 Kocsis E, Trus BL, Steer CJ, Bisher ME, Steven AC. 1991. Image averaging of flexible fibrous

744 macromolecules: the clathrin triskelion has an elastic proximal segment. *J Struct Biol* **107**:6–

745 14. doi:10.1093/bioinformatics/btp184

746 Koshida S, Kishimoto Y, Ustumi H, Shimizu T, Furutani-Seiki M, Kondoh H, Takada S. 2005.

747 Integrin alpha5-dependent fibronectin accumulation for maintenance of somite boundaries

748 in zebrafish embryos. *Dev Cell* **8**:587–98. doi:10.1016/j.devcel.2005.03.006

749 Kragtorp KA, Miller JR. 2007. Integrin alpha5 is required for somite rotation and boundary
750 formation in *Xenopus*. *Dev Dyn* **236**:2713–20. doi:10.1002/dvdy.21280

751 Lauschke VM, Tsiairis CD, François P, Aulehla A. 2013. Scaling of embryonic patterning based on
752 phase-gradient encoding. *Nature* **493**:101–105. doi:10.1038/nature11804

753 Lawton AK, Nandi A, Stulberg MJ, Dray N, Sneddon MW, Pontius W, Emonet T, Holley SA. 2013.
754 Regulated tissue fluidity steers zebrafish body elongation. *Development* **140**:573–582.
755 doi:10.1242/dev.090381

756 Luca VC, Kim BC, Ge C, Kakuda S, Wu D, Roein-Peikar M, Haltiwanger RS, Zhu C, Ha T, Garcia KC.
757 2017. Notch-Jagged complex structure implicates a catch bond in tuning ligand sensitivity.
758 *Science (80-)* **355**:1320–1324. doi:10.1126/science.aaf9739

759 Mao Y, Schwarzbauer JE. 2005. Fibronectin fibrillogenesis, a cell-mediated matrix assembly
760 process. *Matrix Biol* **24**:389–399. doi:10.1016/j.matbio.2005.06.008

761 Marek M, Kubíček M. 1981. Morphogen pattern formation and development in growth. *Bull*
762 *Math Biol* **43**:259–70.

763 Marrese M, Antonovaite N, Nelemans BKA, Smit TH, Iannuzzi D. 2019. Micro-indentation and
764 optical coherence tomography for the mechanical characterization of embryos:
765 Experimental setup and measurements on chicken embryos. *Acta Biomater*.
766 doi:10.1016/j.actbio.2019.07.056

767 Martins GG, Rifes P, Amândio R, Rodrigues G, Palmeirim I, Thorsteinsdóttir S. 2009. Dynamic 3D
768 cell rearrangements guided by a fibronectin matrix underlie somitogenesis. *PLoS One*
769 **4**:e7429. doi:10.1371/journal.pone.0007429

770 Masamizu Y, Ohtsuka T, Takashima Y, Nagahara H, Takenaka Y, Yoshikawa K, Okamura H,
771 Kageyama R. 2006. Real-time imaging of the somite segmentation clock: Revelation of
772 unstable oscillators in the individual presomitic mesoderm cells. *Proc Natl Acad Sci*
773 **103**:1313–1318. doi:10.1073/pnas.0508658103

774 McKeown-Longo PJ, Mosher DF. 1985. Interaction of the 70,000-mol-wt amino-terminal

775 fragment of fibronectin with the matrix-assembly receptor of fibroblasts. *J Cell Biol*
776 **100**:364–74. doi:10.1083/jcb.100.2.364

777 Meloty-Kapella L, Shergill B, Kuon J, Botvinick E, Weinmaster G. 2012. Notch Ligand Endocytosis
778 Generates Mechanical Pulling Force Dependent on Dynamin, Epsins, and Actin. *Dev Cell*
779 **22**:1299–1312. doi:10.1016/j.devcel.2012.04.005

780 Merle T, Farge E. 2018. Trans-scale mechanotransductive cascade of biochemical and
781 biomechanical patterning in embryonic development: the light side of the force. *Curr Opin*
782 *Cell Biol* **55**:111–118. doi:10.1016/j.cecb.2018.07.003

783 Mongera A, Rowghanian P, Gustafson HJ, Shelton E, Kealhofer DA, Carn EK, Serwane F, Lucio AA,
784 Giammona J, Campàs O. 2018. A fluid-to-solid jamming transition underlies vertebrate body
785 axis elongation. *Nature* **561**:401–405. doi:10.1038/s41586-018-0479-2

786 Morimoto M, Takahashi Y, Endo M, Saga Y. 2005. The Mesp2 transcription factor establishes
787 segmental borders by suppressing Notch activity. *Nature* **435**:354–359.
788 doi:10.1038/nature03591

789 Morin-Kensicki EM, Boone BN, Howell M, Stonebraker JR, Teed J, Alb JG, Magnuson TR, O’Neal
790 W, Milgram SL. 2006. Defects in Yolk Sac Vasculogenesis, Chorioallantoic Fusion, and
791 Embryonic Axis Elongation in Mice with Targeted Disruption of Yap65. *Mol Cell Biol* **26**:77–
792 87. doi:10.1128/MCB.26.1.77-87.2006

793 Mui KL, Chen CS, Assoian RK. 2016. The mechanical regulation of integrin–cadherin crosstalk
794 organizes cells, signaling and forces. *J Cell Sci* **129**:1093–1100. doi:10.1242/jcs.183699

795 Nakajima Y, Morimoto M, Takahashi Y, Koseki H, Saga Y. 2006. Identification of EphA4 enhancer
796 required for segmental expression and the regulation by Mesp2. *Development* **133**:2517–
797 25. doi:10.1242/dev.02422

798 Newell-Litwa KA, Horwitz R, Lamers ML. 2015. Non-muscle myosin II in disease: mechanisms and
799 therapeutic opportunities. *Dis Model Mech* **8**:1495–515. doi:10.1242/dmm.022103

800 Niwa Y, Shimojo H, Isomura A, Gonzalez A, Miyachi H, Kageyama R. 2011. Different types of

oscillations in Notch and Fgf signaling regulate the spatiotemporal periodicity of
somitogenesis. *Genes Dev* **25**:1115–1120. doi:10.1101/gad.2035311

Palmeirim I, Dubrulle J, Henrique D, Ish-Horowicz D, Pourquié O. 1998. Uncoupling segmentation
and somitogenesis in the chick presomitic mesoderm. *Dev Genet* **23**:77–85.
doi:10.1002/(SICI)1520-6408(1998)23:1<77::AID-DVG8>3.0.CO;2-3

Palmeirim I, Henrique D, Ish-Horowicz D, Pourquié O. 1997. Avian hairy gene expression
identifies a molecular clock linked to vertebrate segmentation and somitogenesis. *Cell*
91:639–648. doi:10.1016/S0092-8674(00)80451-1

Piccolo S, Dupont S, Cordenonsi M. 2014. The Biology of YAP/TAZ: Hippo Signaling and Beyond.
Physiol Rev **94**:1287–1312. doi:10.1152/physrev.00005.2014

Pierschbacher MD, Ruoslahti E. 1984. Cell attachment activity of fibronectin can be duplicated by
small synthetic fragments of the molecule. *Nature* **309**:30–3.

Pourquié O, Tam PPL. 2001. A nomenclature for prospective somites and phases of cyclic gene
expression in the presomitic mesoderm. *Dev Cell* **1**:619–20. doi:10.1016/S1534-
5807(01)00082-X

Preibisch S, Saalfeld S, Tomancak P. 2009. Globally optimal stitching of tiled 3D microscopic
image acquisitions. *Bioinformatics* **25**:1463–1465. doi:10.1093/bioinformatics/btp184

Prieto D, Aparicio G, Machado M, Zolessi FR. 2015. Application of the DNA-specific stain methyl
green in the fluorescent labeling of embryos. *J Vis Exp* e52769. doi:10.3791/52769

Rallis C, Pinchin SM, Ish-Horowicz D. 2010. Cell-autonomous integrin control of Wnt and Notch
signalling during somitogenesis. *Development* **137**:3591–3601. doi:10.1242/dev.050070

Rifes P, Carvalho L, Lopes C, Andrade RP, Rodrigues G, Palmeirim I, Thorsteinsdóttir S. 2007.
Redefining the role of ectoderm in somitogenesis: a player in the formation of the
fibronectin matrix of presomitic mesoderm. *Development* **134**:3155–3165.
doi:10.1242/dev.003665

Rifes P, Thorsteinsdóttir S. 2012. Extracellular matrix assembly and 3D organization during

827 paraxial mesoderm development in the chick embryo. *Dev Biol* **368**:370–381.

828 doi:10.1016/j.ydbio.2012.06.003

829 Ringer P, Colo G, Fässler R, Grashoff C. 2017. Sensing the mechano-chemical properties of the

830 extracellular matrix. *Matrix Biol* **64**:6–16. doi:10.1016/j.matbio.2017.03.004

831 Saga Y. 2012. The mechanism of somite formation in mice. *Curr Opin Genet Dev* **22**:331–338.

832 doi:10.1016/j.gde.2012.05.004

833 Saga Y, Takeda H. 2001. The making of the somite: molecular events in vertebrate segmentation.

834 *Nat Rev Genet* **2**:835–845. doi:10.1038/35098552

835 Sato Y, Nagatoshi K, Hamano A, Imamura Y, Huss D, Uchida S, Lansford R. 2017. Basal filopodia

836 and vascular mechanical stress organize fibronectin into pillars bridging the mesoderm-

837 endoderm gap. *Development* **144**:281–291. doi:10.1242/dev.141259

838 Sato Y, Sato Y, Kasai T, Kasai T, Nakagawa S, Nakagawa S, Tanabe K, Tanabe K, Watanabe T,

839 Watanabe T, Kawakami K, Kawakami K, Takahashi Y, Takahashi Y. 2007. Stable integration

840 and conditional expression of electroporated transgenes in chicken embryos. *Dev Biol*

841 **305**:616–24. doi:10.1016/j.ydbio.2007.01.043

842 Sato Y, Yasuda K, Takahashi Y. 2002. Morphological boundary forms by a novel inductive event

843 mediated by Lunatic fringe and Notch during somitic segmentation. *Development*

844 **129**:3633–44.

845 Schiller HB, Hermann M-R, Polleux J, Vignaud T, Zanivan S, Friedel CC, Sun Z, Raducanu A,

846 Gottschalk K-E, Théry M, Mann M, Fässler R. 2013. β 1- and α v-class integrins cooperate to

847 regulate myosin II during rigidity sensing of fibronectin-based microenvironments. *Nat Cell*

848 *Biol* **15**:625–636. doi:10.1038/ncb2747

849 Schwartz MA, DeSimone DW. 2008. Cell adhesion receptors in mechanotransduction. *Curr Opin*

850 *Cell Biol* **20**:551–556. doi:10.1016/j.ceb.2008.05.005

851 Shih NP, Francois P, Delaune EA, Amacher SL. 2015. Dynamics of the slowing segmentation clock

852 reveal alternating two-segment periodicity. *Development* **142**:1785–1793.

853 doi:10.1242/dev.119057

854 Singh P, Carraher C, Schwarzbauer JE. 2010. Assembly of Fibronectin Extracellular Matrix. *Annu*

855 *Rev Cell Dev Biol* **26**:397–419. doi:10.1146/annurev-cellbio-100109-104020

856 Slack JMW. 1987. We have a morphogen! *Nature* **327**:553–554. doi:10.1038/327553a0

857 Smutny M, Ákos Z, Grigolon S, Shamipour S, Ruprecht V, Čapek D, Behrndt M, Papusheva E, Tada

858 M, Hof B, Vicsek T, Salbreux G, Heisenberg CP. 2017. Friction forces position the neural

859 anlage. *Nat Cell Biol* **19**:306–317. doi:10.1038/ncb3492

860 Straight AF, Cheung A, Limouze J, Chen I, Westwood NJ, Sellers JR, Mitchison TJ. 2003. Dissecting

861 temporal and spatial control of cytokinesis with a myosin II Inhibitor. *Science* **299**:1743–7.

862 doi:10.1126/science.1081412

863 Takahashi Y, Inoue T, Gossler A, Saga Y. 2003. Feedback loops comprising Dll1, Dll3 and Mesp2,

864 and differential involvement of Psen1 are essential for rostrocaudal patterning of somites.

865 *Development* **130**:4259–68. doi:10.1242/dev.00629

866 Takahashi Y, Koizumi K, Takagi A, Kitajima S, Inoue T, Koseki H, Saga Y. 2000. Mesp2 initiates

867 somite segmentation through the Notch signalling pathway. *Nat Genet* **25**:390–6.

868 doi:10.1038/78062

869 Takeichi M. 2014. Dynamic contacts: rearranging adherens junctions to drive epithelial

870 remodelling. *Nat Rev Mol Cell Biol* **15**:397–410. doi:10.1038/nrm3802

871 Tiedemann H. 1976. Pattern formation in early developmental stages of amphibian embryos. *J*

872 *Embryol Exp Morphol* **35**:437–44.

873 Watanabe T, Sato Y, Saito D, Tadokoro R, Takahashi Y. 2009. EphrinB2 coordinates the formation

874 of a morphological boundary and cell epithelialization during somite segmentation. *Proc*

875 *Natl Acad Sci* **106**:7467–7472. doi:10.1073/pnas.0902859106

876 Wolfenson H, Lavelin I, Geiger B. 2013. Dynamic regulation of the structure and functions of

877 integrin adhesions. *Dev Cell* **24**:447–58. doi:10.1016/j.devcel.2013.02.012

878 Wolfenson H, Yang B, Sheetz MP. 2019. Steps in Mechanotransduction Pathways that Control

879 Cell Morphology. *Annu Rev Physiol* **81**:585–605. doi:10.1146/annurev-physiol-021317-
880 121245

881 Yang JT, Bader BL, Kreidberg J a, Ullman-Culleré M, Trevithick JE, Hynes RO. 1999. Overlapping
882 and independent functions of fibronectin receptor integrins in early mesodermal
883 development. *Dev Biol* **215**:264–277. doi:10.1006/dbio.1999.9451

884 Yang JT, Rayburn H, Hynes RO. 1993. Embryonic mesodermal defects in alpha 5 integrin-deficient
885 mice. *Development* **119**:1093–105.

886 Yarrow JC, Totsukawa G, Charras GT, Mitchison TJ. 2005. Screening for cell migration inhibitors
887 via automated microscopy reveals a Rho-kinase inhibitor. *Chem Biol* **12**:385–95.
888 doi:10.1016/j.chembiol.2005.01.015

889 Zaidel-Bar R. 2013. Cadherin adhesome at a glance. *J Cell Sci* **126**:373–378.
890 doi:10.1242/jcs.111559

891 Zaidel-Bar R, Zhenhuan G, Luxenburg C. 2015. The contractome - a systems view of actomyosin
892 contractility in non-muscle cells. *J Cell Sci* **128**:2209–2217. doi:10.1242/jcs.170068

Supplementary Figures

Supplementary Figure 1. Blocking NM II or ROCK I/II activity for 3 hours leads to altered *meso1* expression. (A, B) *meso1* expression evaluated by *in situ* hybridization in explants treated for 3 hours with Blebbistatin (A) or RockOut (B) and in the respective contralateral controls, shows that *meso1* expression is already altered two clock cycles after adding the drugs. Straightened images of the respective explant pairs (right) were aligned by SIV. Rostral is on top.

904

905

906 **Supplementary Figure 2. Apoptosis is not enhanced in the experimental culture conditions used.**

907 **(A, H)** Immunostaining for activated Caspase3 in transverse sections of control and contralateral explants
908 treated with either Blebbistatin **(A, B)**, RockOut **(C, D)**, or RGD **(E, F)** for 6 hours, and in pCAGGs- and 70kDa-
909 electroporated embryos **(G, H)**. DNA (blue), activated Caspase3 (magenta) and GFP (green). Arrows indicate
910 Caspase3-positive cells. Apoptosis levels are not increased by any of the treatments. Blebb – Blebbistatin;
911 nt – neural tube; s – somite; i – intermediate mesoderm; lp – lateral plate mesoderm. Dorsal is on top. Scale
912 bars: 50 μm.

Supplementary Figure 3. NM II inhibition impairs F-actin apical enrichment and nuclear alignment in somites formed during culture.

(A, L) Sagittal optical sections of DMSO-treated explants (A-C, G-I) and contralateral Blebbistatin-treated explants (D-F, J-L) stained for F-actin and DNA. Rostral PSM epithelialization of DMSO-treated explants occurs normally, with F-actin apical enrichment (A) and nuclear alignment (B) in S0 and SI. At the same axial level in the contralateral Blebbistatin-treated explant (D-F), F-actin staining is dispersed (D) and nuclei do not align (E). SII and SIII of DMSO-treated explants are epithelial (G-I), composed of an outer cell layer with aligned nuclei (H) and elongated cells with apically enriched F-actin (G). At the equivalent axial level in the Blebbistatin-treated explant (J-L), the somitic segments were severely affected. There is only a slight nuclear alignment at the prospective inter-somitic border (K, arrowheads) and F-actin aggregates into dispersed and separate foci (J, L). Rostral to the left and dorsal on top. Blebb – Blebbistatin; Phall – phalloidin F-actin staining. Scale bars: 50 μ m.

Supplementary Figure 4. Epithelial structure of the neural tube and ectoderm are unaffected by NM II inhibition.

(A, N) Control explants (A-D and I-K) showed strong apically located ZO-1 labeling (B, I, arrowheads) both in the neural tube (A-D; transverse optical section, medial on top) and in the overlying surface ectoderm (I-K; dorsal view of ectoderm). In the presence of Blebbistatin (E-H, L-N), ZO-1 labeling remained restricted to the apical end of neural tube cells (F, arrowheads) and of surface ectoderm cells (L, arrowheads). Blebb – Blebbistatin; Ncad – N-cadherin. ZO-1 – *Zonula occludens* protein 1. Arrowheads point to ZO-1 labeling. Scale bars: 20 μ m.

Supplementary Figure 5. ROCK I/II inhibition impairs apical polarization of N-cadherin in nascent somites.

(A, L) Longitudinal sections of explants cultured for 6 hours in control (DMSO) medium (A-C, G-I) and their contralateral RockOut-treated halves (D-F, J-L) at SI (A-F) and SIII (G-L) levels, immunostained for N-cadherin (first column) and stained for DNA (second column). Third column shows the respective merge of all stainings. SI in control explants shows normal accumulation of N-cadherin (A, arrowhead) and nuclear alignment (B, arrowhead). In contrast, SI in contralateral RockOut-treated explants fail to form a clear cleft (D-F, arrows) leading to partially fused somites. N-cadherin is, however, partially polarized (D, arrowhead) and nuclei are aligned (E, arrowhead). At SIII level, both explants show normal N-cadherin polarization (G, J) and nuclear alignment (H, L). Rostral on the left and midline on top. Ncad - N-cadherin. Scale bars: 50 μ m.

Supplementary Figure 6. Embryos electroporated with the 70kDa construct exhibit numerous morphological defects.

(A) Representative images of the morphology of embryos electroporated with either pCAGGs only (a) or 70kDa, the latter being a severe phenotype (e). Severe phenotypes included severely detached tissues (e, arrow) and a truncated A-P axis. Transverse sections of pCAGGs- (b-d) and 70kDa-electroporated embryos (f-h) immunostained for fibronectin (b-c, f-g), GFP (d, h) and stained for DNA (c-d, g-h). Electroporated side is on left (a-d) or right (e-h). Detachment of tissues is clearly visible in 70kDa-electroporated embryos (e), which is accompanied by a severe disruption in the fibronectin matrix (compare b-c with f-g, arrows). **(B)** Close up of embryos electroporated with 70kDa showing kinked neural tube (d, arrowheads), fused somites (e, arrowheads) and fewer somites on the electroporated side (f, arrowheads). Electroporated sides are on left. Ventral view and rostral on top. **(C)** Percentage of pCAGGS- (green bars) and 70kDa-electroporated (pink

957 bars) embryos with morphological defects, including detached (pCAGGs: 39/154, 70kDa: 97/144) and
 958 severely detached tissues (pCAGGs: 6/154, 70kDa: 79/144), kinked neural tube (pCAGGs: 15/154, 70kDa:
 959 34/144), truncated A-P axis (pCAGGs: 13/154, 70kDa: 21/144), abnormal somite morphology (pCAGGs:
 960 18/154, 70kDa: 41/144), and fewer somites on the electroporated side compared to the control non-
 961 electroporated side (pCAGGs: 0/154, 70kDa: 15/144). p values were calculated using a Chi-square test.
 962 ***p<0.001. FN – fibronectin. Scale bars: (A, a, e; C, a-c) 500 μ m, (C, d-f) 200 μ m, (A, b-d, f-h) 50 μ m.

963

Length and width of SI and SV of non-electroporated side vs of pCAGGs/70kDa side

964

965

966

967

968

969

970

971

972

973

Supplementary Figure 7. Quantification of somite length and width in pCAGGs- and 70kDa-electroporated embryos.

(A-D) Length of SI (A) and SV (B) and width of SI (C) and SV (D) from pCAGGS- and 70kDa-electroporated embryos compared to the control non-electroporated side. The measurements were made on images from whole mount embryos. Somites from pCAGGs-electroporated embryos did not show a significant difference in either length or width between electroporated vs non-electroporated sides (n=151), but the width of SI and length of SV of the electroporated side of 70kDa-treated embryos were significantly smaller than that of the contralateral non-electroporated control side (n=143). Bars represent the standard error of the mean. p values were calculated using a nested ANOVA. ns – not significant; *** p<0.001.