HUMAN GENES ARE *IN SILICO* POTENTIAL TARGETS FOR RICE miRNA

Aizhan Rakhmetullina, Anna Pyrkova, Dana Aisina, Anatoliy Ivashchenko

¹ Department of Biotechnology, SRI of Biology and Biotechnology Problems, al-Farabi Kazakh National University, Almaty, al-Farabi 71, Almaty, 050040, Kazakhstan

ORCID IDs: 0000-0002-0117-7110 (A.K.); 0000-0001-8483-451X (A.P.); 0000-0002-7137-3408 (D.A.); 0000-0002-7969-2016 (A.I.)

Address correspondence to a.iavashchenko@gmail.com.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Anatoliy Ivashchenko (a.iavashchenko@gmail.com).

Abstract

Exogenous miRNAs enter the human body through food, and their effects on metabolic processes can be considerable. It is important to determine which miRNAs from plants affect the expression of human genes and the extent of their influence. The binding sites of 738 osa-miRNAs that interact with 17508 mRNAs of human genes were determined using the MirTarget program. The characteristics of the binding of 46 single osa-miRNAs to 86 mRNAs of human genes with a value of free energy (ΔG) interaction equal 94% to 100% from maximum ΔG were established. The findings showed that osa-miR2102-5p, osa-miR5075-3p, osa-miR2097-5p, osa-miR2919 targeted the largest number of genes at 38, 36, 23, 19 sites, respectively. mRNAs of 86 human genes were identified as targets for 93 osa-miRNAs of all family osa-miRNAs with ΔG values equal 94% to 98% from maximum ΔG . Each miRNA of the osa-miR156-5p, osa-miR164-5p, osa-miR168-5p, osa-miR395-3p, osa-miR396-3p, osa-miR396-5p, osa-miR444-3p, osa-miR529-3p, osa-miR1846-3p, osa-miR2907-3p families had binding sites in mRNAs of several human target genes. The binding sites of osa-miRNAs in mRNAs of the target genes for each family of osa-miRNAs were conserved when compared to flanking nucleotide sequences. mRNA human genes of osa-miRNAs are candidate genes of cancer, cardiovascular and neurodegenerative diseases.

Introduction

Plant miRNA (pl-miR) and animal miRNA (an-miR) are exogenous miRNA or xeno miRNA (ex-miR or xe-miR) for humans and do not have distinctive features among themselves. Therefore, in the human body, they will be perceived as the general diversity of the endogenous miRNA (en-miR) of a human, and all human genes can potentially be their targets. The effect of

ex-miR on human target genes and the consequences of this effect will depend on their concentration and the duration of their presence in the cells. pl-miR can enter the human body through the gastrointestinal tract and spread with blood in combination with proteins or as part of exosomes (Buck et al., 2014; Escrevente et al., 2011; Montecalvo et al., 2012; van der Grein & Nolte-'t Hoen, 2014). Many studies have shown that miRNAs that enter the gastrointestinal tract with food (dietary miRNA) are then found in various tissues (Chiang et al., 2015; Jonathan et al., 2013; Liang et al., 2015; Stephen & Snow, 2017; Vaucheret & Chupeau, 2012; Zemleni et al., 2017; Zhang et al., 2019; Zhang et al., 2012; Zhao et al., 2018). For example, after consumption of a fresh maize diet most zma-miRNAs were detected in the heart, brain, mammary gland, lung, liver, kidney and serum exosomes of pig (Luo et al., 2017). The concentration of pl-miR in different human organs varies and can be compared with en-miR. Some studies have found minor concentrations of pl-miR in humans and animals (Zhang et al., 2016). The inclusion of pl-miR into the recipient's organism causes reproducible changes of some properties and physiological processes in it (Cui et al., 2017; Javed et al., 2017; Lang et al., 2019; Vaucheret et al., 2012; Zhang et al., 2012). ex-miR involvement in the regulation of recipient gene expression may affect disease (Chin, Fong & Somlo, 2016; Gopinath, 2019; Hou et al., 2018; Jones et al., 2016). The program for searching binding sites in target genes can reliably identify pl-miR and an-miR binding sites regardless of the origin of miRNAs (Ivashchenko et al., 2016), which makes it possible to predict the interaction of pl-miR with hsa-mRNA. We studied the possible interactions of pl-miRs with hsa-mRNA genes after pl-miRs have entered the human body. It was also assumed that pl-miRs can circulate in the blood throughout the body in the absence of features restricting their entry into any cell (Wagner et al., 2015). The basis for this assumption is the diversity of the nucleotide composition of hsa-mRNAs, some of which overlap with diverse pl-miRs (Liang et al., 2013; Pirro et al., 2016).

Recently, the miRNAs of plants ingested for food ex-miR have been actively studied in the regulation of vital processes in humans and animals (Arroyo et al., 2011; Liang et al., 2012; Luo et al., 2017; Vaucheret & Chupeau, 2012; Zhang et al., 2016). It is presumed that ex-miRs regulate cellular function in healthy cells and act as important mediators in the development of animal diseases (Cong et al., 2018; Hoy & Buck, 2012; Makarova et al., 2016; Rutter & Innes, 2018). Ex-miRs can function as evolutionary linkers between different species and contribute to signal transmission both within and between species (Arteaga-Vazquez et al., 2006; Axtell, Westholm & Lai, 2011; Millar & Waterhouse, 2005; Moran et al., 2017; Zhao, Cong & Lukiw, 2018). Based on the available data, the authors suggest that such xe-miRNAs contribute to the beneficial properties of medicinal plants (Lukasik & Zielenkiewicz, 2016; Xie, Weng & Melzig, 2016), contribute to the negative properties of disease-causing or poisonous plants, and cross-

link species between kingdoms of living organisms by participating in many of the mechanisms associated with the occurrence and pathogenesis of various diseases (Mallocci et al., 2018; Melnik, John & Schmitz, 2014; Perge et al., 2017; Pogue et al., 2014).

To determine which pl-miRs affect the mRNAs of human genes, we chose osa-miRNAs because rice has the most miRNAs in the plants, and rice is the most common source of human nutrition. Most plants contain well-known miRNAs, which serve as typical regulators of plant growth and development (Bari, Orazova & Ivashchenko, 2013; Bari et al., 2014, Nair et al., 2010).

Results

Characteristics of the interaction of single osa-miRNAs with mRNA of human genes

Currently, 738 miRNAs encoded by the rice genome are known. For these osa-miRNAs, target genes from among 17508 human genes were searched. A total of 82 miRNAs with one to four target genes were identified (Table S1). miR11339-3p and miR11339-5p; miR1425-3p and miR1425-5p; miR1432-3p and miR1432-5p; miR1870-3p and miR1870-5p; miR2096-3p and miR2096-5p; miR2867-3p and miR2867-5p; and miR390-3p and miR390-5p, originating from the same pre-miRNA, had binding sites in the mRNAs of different genes. The functions of the 162 identified target genes were diverse.

In the group of 49 miRNAs with five or more target genes, there were several miR-3p/miR-5p pairs that originated from the same pre-miRNA (Table S2). The total number of target genes for miRNAs with five or more genes was 479. The number of target genes for miR408-3p, miR5150-3p, miR528-3p, and miR530-3p was comparable to the number of target genes for miR408-5p, miR5150-5p, miR528-5p, and miR530-5p. For miR1847.1-5p, miR1850.1-5p, miR2094-5p, miR2097-5p, miR2102-5p, and miR3979-5p, the set of target genes was significantly larger than that for each corresponding miRNA-3p. Only miR5144-3p had four-fold more target genes compared to the number attributed to miR5144-5p. The miRNAs with the largest number of target genes were miR2102-5p (38 genes), miR5075-3p (36 genes), miR2097-5p (23 genes), and miR2919 (19 genes). Consequently, at high concentrations, these miRNAs could significantly change the metabolism of recipient human cells.

A total of 641 target genes were identified for 131 single miRNAs, which is approximately 3.7% of the total number of studied human genes.

Table 1 shows the characteristics of the binding of some osa-miRNAs with mRNAs of human genes. Each of the 35 miRNAs could bind to mRNAs of one target gene, six miRNAs had targets with two genes, and four miRNAs had three target genes with a value $\Delta G/\Delta G_m$ equal to 94-98%. miR2102-5p had 11 target genes with a value $\Delta G/\Delta G_m$ of 94-100%, and the free energy of the interaction of the miRNAs with the mRNAs of these genes varied from -115 kJ/mole to -121

kJ/mole. The miR2102-5p binding sites were located mainly in the 5'UTR, which suggests that they have a role in the early inhibition of the translation process. However, this property of miR2102-5p indicates the need to control its plant food-derived concentration in the human body. 19 target genes were associated with miR2919. miR5075-3p could bind to three mRNAs at binding sites located in the coding domain sequence and the 5'-untranslated region. The high-affinity binding sites were located in the 5'UTR and CDS of the mRNAs with the $\Delta G/\Delta G_m$ value was 94-98%. 17 miRNA binding sites were located in 5'UTR, 39 in CDS and 33 in 3'-untranslated region. Therefore, monitoring the concentrations of miR2102-5p, miR2919 and miR5075-3p in human biological fluids is also necessary.

Table 1. Characteristics of Interaction single osa-miRNAs with mRNA of Human Genes

Gene	osa-miRNA	Start of site, nt	Region of miRNA	ΔG , kJ/mole	$\Delta G/\Delta G_m$, %	Length, nt
<i>CPA3</i>	miR1320-3p	671	CDS	-98	96	21
<i>PPAP2B</i>	miR1320-3p	2135	3'UTR	-96	94	21
<i>CNPY1</i>	miR1426-5p	2094	3'UTR	-93	94	21
<i>NSL1</i>	miR1440-5p	4601	3'UTR	-98	94	20
<i>PSEN2</i>	miR1847.1-5p	1785	3'UTR	-108	96	21
<i>MDN1</i>	miR1855-3p	13027	CDS	-106	94	21
<i>KLHDC10</i>	miR1860-3p	3134	3'UTR	-108	96	22
<i>TIE1</i>	miR1860-3p	3139	CDS	-106	94	22
<i>OSTM1</i>	miR2093-3p	848	CDS	-93	96	20
<i>ZNF80</i>	miR2099-3p	2562	3'UTR	-96	94	20
<i>SLC36A3</i>	miR2099-5p	2769	3'UTR	-100	94	22
<i>AFAP1</i>	miR2102-5p	144	5'UTR	-115	95	20
<i>C19orf6</i>	miR2102-5p	193	CDS	-115	95	20
<i>CHSY1</i>	miR2102-5p	348	5'UTR	-117	96	20
<i>DIRC2</i>	miR2102-5p	233	CDS	-117	96	20
<i>KATNAL1</i>	miR2102-5p	80	5'UTR	-117	96	20
<i>NR1D2</i>	miR2102-5p	254	5'UTR	-117	96	20
<i>PDAP1</i>	miR2102-5p	29	5'UTR	-115	95	20
<i>PPP2R5C</i>	miR2102-5p	72	5'UTR	-115	95	20
<i>RHOB2</i>	miR2102-5p	158	5'UTR	-115	95	20
<i>UHRF1BP1</i>	miR2102-5p	112	5'UTR	-115	95	20
<i>WT1</i>	miR2102-5p	450	CDS	-121	100	20
<i>ZNF442</i>	miR2866-5p	1020	CDS	-98	96	20
<i>PCDHB15</i>	miR2866-5p	582	CDS	-96	94	20
<i>GPR20</i>	miR2867-3p	455	CDS	-106	94	20
<i>TMEM38A</i>	miR2867-3p	236	CDS	-106	94	20
<i>ATP13A3</i>	miR2867-5p	3035	CDS	-115	95	22
<i>HK2</i>	miR2868-5p	6645	3'UTR	-93	96	20
<i>ZNF395</i>	miR2870-3p	863	CDS	-100	94	21
<i>IQGAP1</i>	miR2876-5p	1234	CDS	-102	94	21
<i>ADAMTS5</i>	miR2919	471	5'UTR	-104	96	19
<i>BDNF</i>	miR2919	2681	3'UTR	-106	98	19
<i>GPBP1L1</i>	miR2919	1015	5'UTR	-104	96	19
<i>KIAA1161</i>	miR2919	3436	3'UTR	-106	98	19
<i>AKAP11</i>	miR2919	7158	3'UTR	-102	94	19
<i>ATG13</i>	miR2919	259	5'UTR	-102	94	19
<i>C1D</i>	miR2919	543	3'UTR	-102	94	19
<i>CDC25B</i>	miR2919	661	5'UTR	-102	94	19
<i>FAM59B</i>	miR2919	257	5'UTR	-102	94	19
<i>FAM83H</i>	miR2919	4348	3'UTR	-102	94	19
<i>MINK1</i>	miR2919	4936	3'UTR	-102	94	19
<i>NEUROD2</i>	miR2919	145	5'UTR	-102	94	19

<i>OTUD4</i>	miR2919	4235	3'UTR	-102	94	19
<i>PRDM11</i>	miR2919	8528	3'UTR	-102	94	19
<i>PTGFRN</i>	miR2919	5633	3'UTR	-102	94	19
<i>RGS9BP</i>	miR2919	2857	3'UTR	-102	94	19
<i>SPRY4</i>	miR2919	1576	3'UTR	-102	94	19
<i>ZNF304</i>	miR2919	2752	3'UTR	-102	94	19
<i>ZNF385A</i>	miR2919	1388	3'UTR	-102	94	19
<i>KPNA4</i>	miR2923-5p	7223	3'UTR	-93	94	22
<i>SHISA6</i>	miR2925-5p	71	CDS	-106	94	19
<i>SPON1</i>	miR2925-5p	93	5'UTR	-106	94	19
<i>ZNHIT2</i>	miR2925-5p	559	CDS	-106	94	19
<i>UFSP1</i>	miR2931-5p	960	3'UTR	-91	96	20
<i>IGSF3</i>	miR394-5p	2007	CDS	-102	94	20
<i>PGAP1</i>	miR394-5p	9153	3'UTR	-102	94	20
<i>ZNF425</i>	miR3979-5p	512	CDS	-102	94	20
<i>RBMS2</i>	miR408-3p	7343	3'UTR	-113	95	21
<i>PPM1F</i>	miR408-5p	2987	3'UTR	-110	95	21
<i>GPBP1L1</i>	miR413-5p	400	5'UTR	-102	94	21
<i>C14orf142</i>	miR414-5p	281	CDS	-104	94	21
<i>LMNA</i>	miR414-5p	1902	CDS	-104	94	21
<i>TMEM30B</i>	miR414-5p	1700	CDS	-104	94	21
<i>PVR</i>	miR415-5p	5452	3'UTR	-104	94	21
<i>SNAPC1</i>	miR417-3p	1169	CDS	-98	94	21
<i>FAM120A</i>	miR418-3p	1029	CDS	-98	94	21
<i>ZNF256</i>	miR5071-5p	348	CDS	-102	94	21
<i>NR2F2</i>	miR5075-3p	350	5'UTR	-117	95	21
<i>PARP2</i>	miR5075-3p	32	CDS	-117	95	21
<i>RPS6KA5</i>	miR5075-3p	261	CDS	-121	98	21
<i>NANOG</i>	miR5077-5p	773	CDS	-100	94	19
<i>PPARGC1A</i>	miR5144-5p	1450	CDS	-104	94	21
<i>FREM2</i>	miR528-3p	1921	CDS	-106	94	21
<i>FXVD6</i>	miR530-5p	788	CDS	-100	94	20
<i>LAMC3</i>	miR530-5p	898	CDS	-100	94	20
<i>NDST1</i>	miR530-5p	6055	3'UTR	-100	94	20
<i>SLC35D1</i>	miR5339-5p	795	CDS	-102	96	21
<i>EML1</i>	miR535-3p	2092	CDS	-106	94	21
<i>LRP5</i>	miR5488-5p	4185	CDS	-102	94	21
<i>SLC25A47</i>	miR5510-5p	678	CDS	-108	94	21
<i>PM20D2</i>	miR5514-5p	1280	CDS	-113	95	21
<i>PPARA</i>	miR5515-3p	1497	CDS	-106	94	21
<i>ZSCAN22</i>	miR5526-3p	2638	3'UTR	-102	94	21
<i>SH3BP2</i>	miR5532-3p	8598	3'UTR	-104	94	22
<i>NANOS1</i>	miR5534a-5p	1967	3'UTR	-106	96	21
<i>DUT</i>	miR5543-5p	1619	3'UTR	-93	94	21
<i>OTUD4</i>	miR5543-5p	3680	3'UTR	-93	94	21
<i>COX20</i>	miR5833-5p	396	CDS	-117	96	21
<i>AKAP11</i>	miR827-3p	2985	CDS	-100	94	21

119

120

121 Table S1-S3 show information about the genes targeted by plant miRNAs that may be
122 involved in the development of various diseases. Most target genes are involved in the
123 development of cancer of various types: *ENAH*, *MAPT*, *PRKCE*, *PRRT2*, *RBMS2*, *RHOBTB2*,
124 *RPS6KA5* and *ZFHX3* - breast cancer (Aisina et al., 2019); *ADAMTS5*, *PRAPRG1A*, *PVR*,
125 *SPRY4* - colorectal cancer; *CHSY1* - colorectal cancer and hepatocellular carcinoma; *PPM1F*,
126 *FXVD6*, *DUT*, *FAM83H* - hepatocellular carcinoma; *HK2* - gallbladder cancer and leukaemia;
127 *C19orf6* - ovarian carcinoma; *AFAP1* - oesophageal adenocarcinoma; *IOGAP1* - pancreatic

ductal adenocarcinoma; *PPP2R5C* - lung adenocarcinoma; *PDAP1* - leukemia; *AKAP11*,
OSTM1, *LRP5* - osteopetrosis; *WT1* - ovarian cancer and myeloid leukaemia; *NR1D2* - various
cancers, including glioblastoma; *CDC25B*, *DIRC2* - renal carcinoma; and *UHRF1BP1* - cell
carcinoma of the head and neck. Some genes are associated with other diseases: *NR2F2* -
metabolic gene regulation and congenital heart defect; *PRAPA*, *UFSP1* - with increased
cardiovascular disease; and *ATP13A3* - pulmonary tumour arterial hypertension and psychiatric
disorder. *BDNF* has an important role in the neurogenesis and neuroplasticity of the brain;
PSEN2 and *LMNA* are associated with Alzheimer's disease; *ZNF442* has a role in psychiatric
disorders; and *NANOS1* is associated with retinoblastoma tumours. The list of oncological
diseases caused by the target genes of the osa-miRNAs indicates that miRNAs can participate in
the development of cancer not only in the gastrointestinal tract but also in other organs.
Therefore, miRNAs that are ingested with food can be transferred to other tissues and organs.

The interaction of miRNAs and mRNA nucleotides of target genes shows how effectively
these molecules bind. The schemes presented in Fig. 1 show the formation of hydrogen bonds
between all the nucleotides of miR5075-3p, miR2866-5p, and miR2919 and the binding sites in
mRNA. Because the MirTarget program takes into account the interaction of the noncanonical
pairs A-C and G-U, it can be seen that the interaction of miRNAs and mRNAs preserves the
spiral structures of both molecules, and therefore, stacking interactions are found between all the
nucleotides of the miRNA and mRNA, which stabilize the duplex. miRNA binding sites are
located in the 5'UTR, CDS, and 3'UTR.

One form of evidence for the reliability of miRNA interaction with mRNA is the
establishment of a conservative nucleotide sequence of binding sites in mRNA of target genes
(Atambayeva et al., 2017; Bari, Orazova & Ivashchenko, 2013; Bari et al., 2014; Yurikova et al.,
2019). The results of the analysis of the similarity of the nucleotide sequences of the binding
sites in mRNA of the target genes miR2102-5p, miR2919 and miR5075-3p are shown in Fig. 2.
For all miR2102-5p, miR2919, and miR5075-3p binding sites, nucleotide conservation is present
compared to the flanking nucleotides of mRNA target genes. With the complete
complementarity of the nucleotides miRNA and mRNA ($\Delta G/\Delta G_m = 100\%$) of the target genes,
absolute conservatism of the site along the entire binding site should be observed, as previously
shown (Atambayeva et al., 2017; Bari, Orazova & Ivashchenko, 2013; Bari et al., 2014;
Yurikova et al., 2019). When the $\Delta G/\Delta G_m$ value changes from 94% to 100% (Table 1), the basis
for the interaction of the nucleotides miR2102-5p, miR2919 and miR5075-3p and mRNA are G-
C pairs.

Characteristics of the interaction of osa-miRNA families with mRNAs of human genes

The total number of osa-miRNAs of all families is 146, and the number of their target genes is equal to 301, which is 1.7% of 17,508 studied human genes. The characteristics of the interaction of 93 osa-miRNAs of all family osa-miRNAs to 86 mRNAs of human genes with values from 94% to 98% were established (Table 2). miRNA of families such as miR156b-3p, miR159a.1,b,f-3p, miR164a,b,c,d,f-5p, miR166a,e-5p, miR166b,c,d,h-5p, miR167a,b,c,d,e,f,g,h,i,j-5p, miR172a,d-3p miR172c-3p, miR396d, miR396a,b-3p, miR531b-5p, miR815a,b,c-3p, miR1428b,c,d,e-3p and miR1858a,b-5p, has one target gene per family. All members of each miRNA family bind at one site due to the homology of their nucleotide sequences. Therefore, the expression of the target gene for each miRNA family will depend on the total concentration of all miRNA families. The miRNA families miR167e,i-3p, miR168b-5p, miR1846a,b,c-5p and miR2907a,b,c,d-3p had two target genes. miR2907a,b,c,d-3p has target genes *IRAK2* and *SLC25A37* with mRNAs of which these miRNAs interact with a large free energy of -123 kJ/mole and -125 kJ/mole, respectively. miR444b.1,c.1-3p had three target genes (*C19orf57*, *KAZN*, *NRG1*).

Table 2. Characteristics of interactions of osa-miRNA of families in mRNA of human genes

Gene	osa-miRNA	Start of site, nt	Region of miRNA	ΔG , kJ/mole	$\Delta G/\Delta G_m$, %	Length, nt
<i>ANKRD27</i>	miR1428b,c,d,e-3p	244	CDS	-98	96	21
<i>AP2A2</i>	miR156a,b,c,d,e,f,g,h,i,j,k-5p	3797	3'UTR	-104	94	21
<i>OBFC1</i>	miR156a,b,c,d,e,f,g,h,i,j-5p	5427	3'UTR	-100	94	20
<i>ZNF652</i>	miR156a,b,c,d,e,f,g,h,i,j-5p	8125	3'UTR	-100	94	20
<i>MROH2B</i>	miR156b-3p	4615	CDS	-106	94	21
<i>PKHD1</i>	miR159a.1,b,f-3p	9966	CDS	-108	98	21
<i>ASXL1</i>	miR164a,b,c,d,f-5p	1923	CDS	-110	95	21
<i>CHST11</i>	miR166a,e-5p	986	CDS	-104	94	21
<i>CCNY</i>	miR166b,c,d,h-5p	367	CDS	-110	95	21
<i>PRICKLE2</i>	miR167a,b,c,d,e,f,g,h,i,j-5p	1959	CDS	-106	94	21
<i>IL17RB</i>	miR167e,i-3p	137	CDS	-102	94	21
<i>KIAA0528</i>	miR167e,i-3p	523	CDS	-102	94	21
<i>APOBR</i>	miR168b-5p	55	CDS	-110	95	21
<i>KLF14</i>	miR168b-5p	623	CDS	-115	98	21
<i>GPR31</i>	miR172a,d-3p	590	CDS	-100	94	21
<i>ATP12A</i>	miR172c-3p	3557	3'UTR	-102	94	21
<i>EIF3B</i>	miR1846a,b,c-5p	2194	CDS	-119	97	21
<i>FASN</i>	miR1846a,b,c-5p	11	5'UTR	-117	95	21
<i>C20orf27</i>	miR1858a,b-5p	921	3'UTR	-115	95	21
<i>IRAK2</i>	miR2907a,b,d-3p	1488	CDS	-123	94	22
<i>SLC25A37</i>	miR2907a,b,d-3p	884	CDS	-125	95	22
<i>EDN1</i>	miR395a-q,t,y-3p	1136	3'UTR	-108	96	21
<i>CD22</i>	miR395b,d,e,g,h-t,y-3p	2684	3'UTR	-104	94	21
<i>CMKLR1</i>	miR396a,b-3p	3970	3'UTR	-98	96	20
<i>SCAMP5</i>	miR396a,b-5p	1727	3'UTR	-102	94	21
<i>SYVN1</i>	miR396a,b-5p	854	CDS	-102	94	21
<i>AKAP13</i>	miR396d	1736	CDS	-100	94	20
<i>RHBDF1</i>	miR444a-3p.1, d.1-3p	1071	CDS	-108	94	21
<i>RUNDC1</i>	miR444a-3p.1, d.1-3p	1191	CDS	-108	94	21
<i>CEP250</i>	miR444a-3p.2, b.2, c.2, d.2,e-3p	1939	CDS	-102	94	21
<i>C19orf57</i>	miR444b.1, c.1-3p	714	CDS	-106	94	21
<i>KAZN</i>	miR444b.1, c.1-3p	4874	3'UTR	-106	94	21
<i>NRG1</i>	miR444b.1,c.1-3p	2444	3'UTR	-106	94	21

<i>ATP6V0A4</i>	miR529a-3p	241	5'UTR	-100	94	20
<i>CCDC94</i>	miR529a-3p	379	CDS	-100	94	20
<i>CHD2</i>	miR529a-3p	6684	3'UTR	-100	94	20
<i>MAP7</i>	miR529a-3p	2137	CDS	-100	94	20
<i>MLL</i>	miR529a-3p	11418	CDS	-100	94	20
<i>PDE4B</i>	miR529a-3p	2270	CDS	-100	94	20
<i>ENAH</i>	miR531b-5p	106	5'UTR	-115	95	20
<i>TIFA</i>	miR815a,b,c-3p	186	5'UTR	-106	94	21

The miRNAs of the large miR395-3p family had *EDN1* and *CD22* target genes, which are involved in the development of diabetes and in the control of immunity, respectively. If these miRNAs get in food in large quantities, then the probability of their impact on human health is high.

miR396a,b-3p can affect the expression of the *CMKLR1* gene, which is involved in cardiovascular disease, and for miR396a,b-5p, the *SCAMP5* and *SYVN1* genes are targeted, the expression of which changes with autism and colon cancer, respectively. miRNAs of the miR444-3p family have binding sites in the mRNA of six genes (Table S4). Their target genes *RHBDF1*, *RUNDC1*, *CEP250*, *C19orf57*, *KAZN*, and *NRG1* are involved in oncogenesis and other diseases.

Therefore, ingestion of these miRNAs with food in humans can significantly affect metabolic processes. miR529a-3p had binding sites in the mRNA of six genes that are involved in the regulation of several physiological processes (Table S5). If a person in the process of evolution consumed this miRNA as a necessary regulator of the expression of its target genes, then for this miRNA there must be target genes.

Fig. 3 shows the interaction patterns of the nucleotide sequences of some representatives of the miRNA families with the mRNA of their target genes. These data indicate a good predictive power for identifying miRNA associations and target genes. The visibility of the interaction of miRNA and mRNA nucleotides in combination with the quantitative characteristics of binding miRNA and mRNA allows us to consider these associations stable and real.

Note that the nucleotide sequences miR156, miR166, miR395, miR396, and miR444 did not have homologous miRNAs among 2565 human miRNAs from the miRBase base. Therefore, these miRNAs do not directly have common binding sites for human miRNAs and can independently regulate the expression of their target genes.

To confirm the conservatism of the interaction of pl-miR with human target genes, we plotted the web logo for mRNA sections containing pl-miR binding sites (Fig. 2). The graphs show the high conservatism of these binding sites compared to flanking nucleotide sequences. The miR156a-j-5 family, consisting of 10 miRNAs, was associated with the mRNA of seven genes (Table S4). The miR156a-j-5 binding sites in each of the genes differed in the number of hydrogen bonds (Fig. 3) and the value of the free interaction energy. Similar results were

obtained for other miRNA families: miR164e-5p, miR168b-5p, miR396c-3p, miR444a-3p, miR164d-1-3p, miR529a-3p, miR815a,b,c-3p, miR1846a,b,c-5p, miR1858a,b-5p, miR2118l-3p, miR2275d-3p, miR2907a,b,d-3p, and miR395a-y-3p (Fig. 2). The conservatism of such bonds between miRNAs and their target genes was established by us for many associations of miRNAs and their target genes in animals and plants (Atambayeva et al., 2017; Bari, Orazova & Ivashchenko, 2013; Bari et al., 2014; Yurikova et al., 2019). These bonds have persisted over tens of millions of years of evolution and indicate the early emergence of the process of regulation by miRNA molecules of target gene expression in animals and plants.

Discussion

Based on the results obtained in this work, the fact of the interaction of pl-miRs with mRNA of human genes is beyond question. It is necessary to establish the possibilities for these miRNAs to enter the human and animal organisms. Several studies have shown that miRNAs in various parts of plants are present in exosomes 30-400 nm in size and are distributed in the body as part of these nanoparticles (Bang & Thum, 2012; Denzer et al., 2000; Xiao et al., 2018). Such compaction of miRNAs in exosomes contributes to their conservation and facilitates the entry of miRNAs into animals through the digestive tract (Redis et al., 2012; Théry, Zitvogel & Amigorena, 2002; Valadi et al., 2007). Further exosomes together with endogenous exosomes with blood move to many tissues and organs. According to the physicochemical properties, plant miRNAs do not differ from animal miRNAs, which makes them competitive when interacting with mRNA target genes. There are no known limitations for the above-described process of ingestion of plant-miRNAs into humans and animals.

Basically, not all pl-miRs will have human target genes, but the most common and vital pl-miRs present in plants can have target genes in animals and humans for a long time eating them. Such pl-miRs usually participate in maintaining the basic physiological functions of plants (productivity, resistance to biotic and abiotic stresses, growth and development). For example, developed rice lines overexpressing *MIR529a* have been shown to have increased resistance to oxidative stress (Chen & Li, 2018; Cimini et al., 2019). The participation of osa-miR159f, osa-miR1871, osa-miR398b, osa-miR408-3p, osa-miR2878-5p, osa-miR528-5p and osa-miR397a in the regulation of a number of physiological processes of rice has been established (Balyan et al., 2017). The expression of miRNA of *Setaria italica* (sit) changed many times: sit-miR1432-3p, sit-miR156a-5p, sit-miR156b-5p, sit-miR164a-5p, sit-miR167b-5p, sit-miR171c-3p, sit-miR2118-3p, sit-miR390-5p, sit-miR394-5p, sit-miR395-3p, sit-miR408-3p, sit-miR529a-3p, sit-miR529b-3p, and sit-miR827, sit-miR159b-3p, sit-miR319c-5p, sit-miR528-5p and sit-miR535-5p under various stresses (Wang et al., 2016). In a broader evolutionary context, miRNAs of *Morus notabilis* were compared to those of seven other plants, including five dicotyledons,

Arabidopsis thaliana, *Glycine max*, *Malus domestica*, *Populus trichocarpa*, *Ricinus communis*, and two monocotyledons, *Oryza sativa* and *Zea mays*. Of the 31 *Morus notabilis* miRNA families, 24 were conserved in the seven plant species. These miRNAs were classified into well-conserved miRNA families. Prominent among them were mulberry miR160b, miR164a, miR167a, miR169a, miR390, and miR396b, which completely matched their counterparts in the seven other plant species, suggesting that those miRNAs were extremely conserved, and might play critical physiological roles in both dicotyledons and monocotyledons. However, seven miRNA families, miR482, miR529, miR858, miR4376, miR4414, miR4995, and miR5523, were found in only one or two plant species. The present data indicated that the conserved miRNA families (miR156, miR166, miR167, miR168, and miR535) miR159, miR160, miR164, miR169, miR171, miR172, miR390, miR396, miR397, miR529 and miR4376, miR162, miR393, miR395, miR398, miR399, miR408 and miR4414 miR319, miR482, miR827, miR828, miR858, miR2111, miR4995 and miR5523 were expressed across a vast range exceeded in all three tissues (Jia et al., 2014). In African rice *Oryza glaberrima* (ogl), some miRNAs such as ogl-miR156l, ogl-miR166c, ogl-miR166k, ogl-miR168a, ogl-miR167i, ogl-miR171f, ogl-miR1846d of the control library and ogl-miR408, ogl-miR528, ogl-miR156, ogl-miR390, and ogl-miR396c of the treated library had higher reads than their complementary strand. This is because miRNA-3p and miRNA-5p may function simultaneously to regulate gene expression (Mondal et al., 2018). It must be understood that the expression of the target gene under the influence of miRNAs can increase with decreasing miRNA concentration below the average physiological level or decrease with increasing miRNA concentration. The data on changes in miRNA concentration in different plants show that the amount of miRNA consumed with food depends on the stage of plant ontogenesis, growing conditions, plant organs, food processing, etc. (Liu et al., 2017). The concentration of pl-miRs after various processing of raw products decreases, but the remaining miR enter the body (Luo et al., 2017; Zhang et al., 2012; Zhou et al., 2015).

Conclusions

As a result of our study, for the first time, among 17508 human genes, 942 target genes for 277 osa-miRNAs were established. The identified target genes account for 5.4% of the total number of studied human genes. miRNA binding sites were found in the CDS, 5'UTR and 3'UTR. The largest number of genes were targeted by osa-miR2102-5p, osa-miR5075-3p, osa-miR2097-5p, and osa-miR2919, which can bind to the mRNA of 38, 36, 23, and 19 genes, respectively. Since osa-miRNAs ingested through plant food have many target genes, they should be controlled in the human body. Most osa-miRNA target genes are involved in the development of diseases, which makes it easier to clarify the role of miRNAs in these processes. Many osa-miRNA target

genes contribute to the development of breast cancer, and other cancer types. The other target genes are involved in cardiovascular and neurodegenerative diseases. Some osa-miRNAs can be effective regulators of human gene expression. The effect of miRNAs can be both positive, contributing to the cure of diseases, and negative, causing a wide range of diseases.

Materials & Methods

The nucleotide sequences of the mRNAs of 17508 targeted genes were downloaded from NCBI GenBank (<http://www.ncbi.nlm.nih.gov>). The nucleotide sequences of the miRNAs were taken from miRBase v.22 (<http://www.mirbase.org/>). The miRNA binding sites in the mRNAs of several genes were predicted using the MirTarget program (Ivashchenko et al., 2016). This program defines the following features of miRNA binding to mRNA: a) the start of the initiation of the miRNA binding to the mRNAs from the first nucleotide of the mRNA's; b) the localization of the miRNA binding sites in the 5'-untranslated region (5'UTR), coding domain sequence (CDS) and 3'-untranslated region (3'UTR) of the mRNAs; c) the free energy of the interaction between miRNA and the mRNA (ΔG , kJ/mole); and d) the schemes of nucleotide interactions between miRNAs and mRNAs. The ratio $\Delta G/\Delta G_m$ (%) is determined for each site (ΔG_m equals the free energy of the miRNA binding with its fully complementary nucleotide sequence). The MirTarget program finds hydrogen bonds between adenine (A) and uracil (U), guanine (G) and cytosine (C), and G and U, A and C. The distances between the bound A and C (1.04 nm) and G and U (1.02 nm) are similar to those between bound G and C and A and U, which are equal to 1.03 nm (Garg & Heinemann, 2018; Kool, 2001; Leontis, Stombaugh & Westhof, 2002). The numbers of hydrogen bonds in the G-C, A-U, G-U and A-C interactions were 3, 2, 1 and 1, respectively. By comparison, MirTarget differs from other programs in terms of finding the binding sites of miRNA on the mRNAs of plant genes (Dai, Zhuang & Zhao, 2011) in that 1) it takes into account the interaction of the miRNA with mRNA over the entire miRNA sequence; 2) takes into account noncanonical pairs G-U and A-C; and 3) calculates the free energy of the interaction of the miRNA with mRNA, and when two or more miRNAs are bound with one mRNA or, if the binding sites of two different miRNAs coincide in part, the preferred miRNA binding site is considered to be the one for which the free binding energy, ΔG , is greater. The MirTarget program does not work directly with the miRBase and NCBI databases. The search for target genes from 17,508 human genes in a special format from NCBI for the known miRNAs from miRBase will be available on request at mirtarget8@gmail.com.

Supplemental Data

Supplemental table 1. osa-miRNA list of 1-4 human target genes.

Supplemental table 2. osa-miRNA list of 5 or more human target genes.

Supplemental table 3. osa-miRNA of target human genes involved in biological processes.

Supplemental table 4. list of osa-miRNA families of human target genes.

Supplemental table 5. osa-miRNA families of target human genes involved in biological processes.

Funding

The work was carried out with the financial support of the Ministry of Education and Science of the Republic of Kazakhstan within the framework of the grant №AP05132460.

AUTHOR CONTRIBUTIONS

A.I., A.R performed the research and analyzed the data. A.P., D.A contributed analytic tools and methods. A.I., A.R., A.P., D.A wrote the article.

References

- Aisina D, Niyazova R, Atambayeva S, Ivashchenko A. 2019.** Prediction of clusters of miRNA binding sites in mRNA candidate genes of breast cancer subtypes. *PeerJ* **7**: e8049 DOI 10.7717/peerj.8049.
- Arroyo JD, Chevillet JR, Kroh EM, Ruf IK, Pritchard CC, Gibson DF, Mitchell PS, Bennett CF, Pogosova-Agadjanyan EL, Stirewalt DL, Tait JF, Tewari M. 2011.** Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 5003-5008 DOI 10.1073/pnas.1019055108.
- Arteaga-Vazquez M, Caballero-Perez J, Vielle-Calzada JP. 2006.** A family of microRNAs present in plants and animals. *Plant Cell* **18**: 3355-3369 DOI 10.1105/tpc.106.044420.
- Atambayeva S, Niyazova R, Ivashchenko A, Pyrkova A, Pinsky I, Akimniyazova A, Labeit S. 2017.** The Binding Sites of miR-619-5p in the mRNAs of Human and Orthologous Genes. *BMC Genomics* **18**: 428. DOI 10.1186/s12864-017-3811-6.
- Axtell MJ, Westholm JO, Lai EC. 2011.** Vive la difference: biogenesis and evolution of microRNAs in plants and animals. *Genome Biology* **12**: 1-13. DOI 10.1186/gb-2011-12-4-221.
- Balyan S, Kumar M, Mutum R, Raghuvanshi U, Agarwal P, Mathur S, Raghuvanshi S. 2017.** Identification of miRNA-mediated drought responsive multi-tiered regulatory network in drought tolerant rice, Nagina 22. *Scientific Reports* **7**: 15446 DOI 10.1038/s41598-017-15450-1.
- Bang C, Thum T. 2012.** Exosomes: new players in cell–cell communication. *The International Journal of Biochemistry & Cell Biology* **44**:2060–2064 DOI 10.1016/j.biocel.2012.08.007.
- Bari A, Orazova S, Ivashchenko A. 2013.** miR156- and miR171-binding sites in the protein-coding sequences of several plant genes. *BioMed Research International* **2013**: 1-7 DOI 10.1155/2013/307145.

349 **Bari A, Sagaidak I, Pinskii I, Orazova S, Ivashchenko A. 2014.** Binding of miR396 to mRNA
350 of Genes Encoding Growth-Regulating Transcription Factors of Plants. *Russian Journal of*
351 *Plant Physiology* **61**: 807-810 DOI 10.1134/S1021443714050033.

352 **Buck AH, Coakley G, Simbari F, McSorley HJ, Quintana JF, Le Bihan T, Kumar**
353 **S, Abreu-Goodger C, Lear M, Marcus Y, Ceroni A, Babayan SA, Blaxter M, Ivens**
354 **A, Maizels RM. 2014.** Exosomes secreted by nematode parasites transfer small RNAs to
355 mammalian cells and modulate innate immunity. *Nature Communications* **5**: 5488 DOI
356 10.1038/ncomms6488.

357 **Chen J, Li L. 2018.** Multiple Regression Analysis Reveals MicroRNA Regulatory Networks in
358 *Oryza sativa* under Drought Stress. *International Journal of Genomics* **2018**: 1-12
359 DOI 10.1155/2018/9395261.

360 **Chiang K, Shu J, Zemleni J, Cui J. 2015.** Dietary MicroRNA Database (DMD): An Archive
361 Database and Analytic Tool for Food-Borne microRNAs. *PLOS One* **10**: 6 DOI
362 10.1371/journal.pone.0128089.

363 **Chin AR, Fong MY, Somlo G. 2016.** Cross-kingdom inhibition of breast cancer growth by
364 plant miR159. *Cell Research* **26**: 217–228 DOI 10.1038/cr.2016.13.

365 **Cimini S, Gualtieri C, Macovei A, Balestrazzi A, De Gara L, Locato V. 2019.** Redox
366 Balance-DDR-miRNA Triangle: Relevance in Genome Stability and Stress Responses in
367 Plants. *Frontiers in Plant Science* **10**: 989 DOI 10.3389/fpls.2019.00989.

368 **Cong L, Zhao Y, Pogue AI, Lukiw WJ. 2018.** Role of microRNA (miRNA) and Viroids in
369 Lethal Diseases of Plants and Animals. Potential Contribution to Human Neurodegenerative
370 Disorders. *Biochemistry (Moscow)* **83**: 1018-1029 DOI 10.1134/S0006297918090031.

371 **Cui J, Zhou B, Ross SA, Zemleni J. 2017.** Nutrition, microRNAs, and Human Health.
372 *Advances in Nutrition* **8**: 105-112 DOI 10.3945/an.116.013839.

373 **Dai X, Zhuang Z, Zhao P. 2011.** Computational Analysis of miRNA Targets in Plants: Current
374 Status and Challenges. *Briefings in Bioinformatics* **12**: 115-121 DOI 10.1093/bib/bbq065.

375 **Denzer K, Eijk MV, Kleijmeer MJ, Jakobson E, Groot CD, Geuze HJ. 2000.** Follicular
376 dendritic cells carry MHC Class II-expressing microvesicles at their surface. *Journal*
377 *of Immunology* **165**:1259–1265 DOI 10.4049/jimmunol.165.3.1259.

378 **Escrevente C, Keller S, Altevogt P, Costa J. 2011.** Interaction and uptake of exosomes by
379 ovarian cancer cells. *BMC Cancer* **11**: 108 DOI 10.1186/1471-2407-11-108.

380 **Garg A, Heinemann U. 2018.** A novel form of RNA double helix based on G·U and
381 C·A⁺ wobble base pairing. *RNA* **24**: 209-218 DOI 10.1261/rna.064048.117.

382 **Gopinath MS. 2019.** Dietary non-coding RNAs from plants: Fairy tale or treasure? Non-coding
383 *RNA Research* **4**: 63-68 DOI 10.1016/j.ncrna.2019.02.002.

384 **Hou D, He F, Ma L, Cao M, Zhou Z, Wei Z, Xue Y, Sang X, Chong H, Tian C, Zheng S, Li**
385 **J, Zen K, Chen X, Hong Z, Zhang CY, Jiang X. 2018.** The potential atheroprotective role
386 of plant MIR156a as a repressor of monocyte recruitment on inflamed human endothelial
387 cells. *The Journal of Nutritional Biochemistry* **57**: 197-205 DOI
388 10.1016/j.jnutbio.2018.03.026.

389 **Hoy AM, Buck AH. 2012.** Extracellular small RNAs: what, where, why? *Biochemical Society*
390 *Transactions* **40**: 886-890 DOI 10.1042/BST20120019.

391 **Ivashchenko AT, Pyrkova AY, Niyazova RY, Alybayeva A, Baskakov K. 2016.** Prediction of
392 miRNA binding sites in mRNA. *Bioinformatics* **12**: 237-240. DOI
393 10.6026/97320630012237.

394 **Javed M, Solanki M, Sinha A, Shukla LI. 2017.** Position Based Nucleotide Analysis of
395 miR168 Family in Higher Plants and its Targets in Mammalian Transcripts. *Microna* **6**: 136-
396 142 DOI 10.2174/2211536606666170215154151.

397 **Jia L, Zhang D, Qi X, Ma B, Xiang Zh, He N, Zhang J. 2014.** Identification of the Conserved
398 and Novel miRNAs in Mulberry by High-Throughput Sequencing. *PLOS One* **9**: 8 DOI
399 10.1371/journal.pone.0104409.

400 **Jonathan WS, Andrew EH, Stephanie KI, Aaron LB, Stephen YC. 2013.** Ineffective delivery
401 of diet-derived microRNAs to recipient animal organisms. *RNA Biology* **10**: 1107-1116
402 DOI 10.4161/rna.24909.

403 **Jones Buie JN, Goodwin AJ, Cook JA, Halushka PV, Fan H. 2016.** The role of miRNAs in
404 cardiovascular disease risk factors. *Atherosclerosis* **254**: 271-281 DOI
405 10.1016/j.atherosclerosis.2016.09.067.

406 **Kool ET. 2001.** Hydrogen bonding, base stacking, and steric effects in DNA replication. *Annual*
407 *Review of Biophysics and Biomolecular Structure* **30**: 1-22 DOI
408 10.1146/annurev.biophys.30.1.1.

409 **Lang C, Karunairetnam S, Lo KR, Kralicek AV, Crowhurst RN, Gleave AP, MacDiarmid**
410 **RM, Ingram JR. 2019.** Common Variants of the Plant microRNA-168a Exhibit Differing
411 Silencing Efficacy for Human Low-Density Lipoprotein Receptor Adaptor Protein 1
412 (LDLRAP1). *Microna* **8**: 166-170 DOI 10.2174/2211536608666181203103233.

413 **Leontis NB, Stombaugh J, Westhof E. 2002.** The non-Watson-Crick base pairs and their
414 associated isostericity matrices. *Nucleic Acids Research* **30**: 3497-3531 DOI
415 10.1093/nar/gkf481.

416 **Liang H, Huang L, Cao J, Zen K, Chen X, Zhang CY. 2012.** Regulation of mammalian gene
417 expression by exogenous microRNAs. *Wiley Interdisciplinary Reviews-RNA* **3**: 733-742 DOI
418 10.1002/wrna.1127.

- 419 **Liang H, Zen K, Zhang J, Zhang CY, Chen X. 2013.** New roles for microRNAs in cross-
420 species communication. *RNA Biology* **10**: 367-370 DOI 10.4161/rna.23663.
- 421 **Liang H, Zhang S, Fu Z, Wang Y, Wang N, Liu Y, Zhao C, Wu J, Hu Y, Zhang J, Chen X,**
422 **Zen K, Zhang CY. 2015.** Effective detection and quantification of dietetically absorbed plant
423 microRNAs in human plasma. *The Journal of Nutritional Biochemistry* **26**: 505-512 DOI
424 10.1016/j.jnutbio.2014.12.002.
- 425 **Liu W, Meng J, Cui J, Luan Yu. 2017.** Characterization and Function of MicroRNAs in
426 Plants. *Frontiers in Plant Science* **8**. DOI 10.3389/fpls.2017.02200.
- 427 **Lukasik A, Zielenkiewicz P. 2016.** Plant microRNAs - novel players in natural medicine? *The*
428 *International Journal of Molecular Sciences* **18**: 1-16 DOI 10.3390/ijms18010009.
- 429 **Luo Y, Wang P, Wang X, Wang Y, Mu Z, Li Q, Fu Y, Xiao J, Li G, Ma Y, Gu Y, Jin L, Ma**
430 **J, Tang Q, Jiang A, Li X, Li M. 2017.** Detection of dietetically absorbed maize-derived
431 microRNAs in pigs. *Scientific Reports* **7**: 1-10 DOI 10.1038/s41598-017-00488-y.
- 432 **Makarova JA, Shkurnikov MU, Wicklein D, Lange T, Samatov TR, Turchinovich AA,**
433 **Tonevitsky AG. 2016.** Intracellular and extracellular microRNA: an update on localization
434 and biological role. *Progress in Histochemistry and Cytochemistry* **51**: 33-49 DOI
435 10.1016/j.proghi.2016.06.001.
- 436 **Mallocci M, Perdomo L, Veerasamy M, Andriantsitohaina R, Simard G, Martinez MC.**
437 **2018.** Extracellular vesicles: mechanisms in human health and disease. *Antioxidants & Redox*
438 *Signaling* **30**: 813-856 DOI 10.1089/ars.2017.7265.
- 439 **Melnik BC, John SM, Schmitz G. 2014.** Milk: an exosomal microRNA transmitter promoting
440 thymic regulatory T cell maturation preventing the development of atopy? *The Journal of*
441 *Translational Medicine* **12**: 1-11 DOI 10.1186/1479-5876-12-43.
- 442 **Millar AA, Waterhouse PM. 2005.** Plant and animal microRNAs: similarities and differences.
443 *Functional & Integrative Genomics* **5**: 129-135 DOI 10.1007/s10142-005-0145-2.
- 444 **Mondal T, Panda A, Rawal H, Sharma T. 2018.** Discovery of microRNA-target modules of
445 African rice (*Oryza glaberrima*) under salinity stress. *Scientific Reports* **8**: 570
446 DOI 10.1038/s41598-017-18206-z.
- 447 **Montecalvo A, Larregina AT, Shufesky WJ, Stolz DB, Sullivan ML, Karlsson JM, Baty**
448 **CJ, Gibson GA, Erdos G, Wang Z, Milosevic J, Tkacheva OA, Divito SJ, Jordan**
449 **R, Lyons-Weiler J, Watkins SC, Morelli AE. 2012.** Mechanism of transfer of functional
450 microRNAs between mouse dendritic cells via exosomes. *Blood* **119**: 756-766 DOI
451 10.1182/blood-2011-02-338004.
- 452 **Moran Y, Agron M, Praher D, Technau U. 2017.** The evolutionary origin of plant and animal
453 microRNAs. *Nature Ecology & Evolution* **1**: 1-22 DOI 10.1038/s41559-016-0027.

- Nair SK, Wang N, Turuspekov Y, Pourkheirandish M, Sinsuwongwat S, Chen G, Sameri M, Tagiri A, Honda I, Watanabe Y, Kanamori H, Wicker T, Stein N, Nagamura Y, Matsumoto T, Komatsuda T. 2010.** Cleistogamous flowering in barley arises from the suppression of microRNA-guided HvAP2 mRNA cleavage. *Proceedings of the National Academy of Sciences of the United States of America* **107**: 490-495 DOI 10.1073/pnas.0909097107.
- Perge P, Nagy Z, Decmann A, Igaz I, Igaz P. 2017.** Potential relevance of microRNAs in inter-species epigenetic communication, and implications for disease pathogenesis. *RNA Biology*. **14**: 391-401 DOI 10.1080/15476286.2016.1251001.
- Pirro S, Minutolo A, Galgani A, Potesta M, Colizzi V, Montesano C. 2016.** Bioinformatics prediction and experimental validation of microRNAs involved in cross-kingdom interaction. *The Journal of Computational Biology* **23**: 976-989 DOI 10.1089/cmb.2016.0059.
- Pogue AI, Clement C, Hill JM, Lukiw WJ. 2014.** Evolution of microRNA (miRNA) structure and function in plants and animals: relevance to aging and disease. *Journal of Aging Science* **2**: 1-12 DOI 10.4172/2329-8847.1000119.
- Redis RS, Calin S, Yang Y, You MJ, Calin GA. 2012.** Cell-to-cell miRNA transfer: from body homeostasis to therapy. *Pharmacology & Therapeutics* **136**:169–174 DOI 10.1016/j.pharmthera.2012.08.003.
- Rutter BD, Innes RW. 2018.** Extracellular vesicles as key mediators of plant–microbe interactions. *Current Opinion in Plant Biology* **44**: 16-22 DOI 10.1016/j.pbi.2018.01.008.
- Stephen Y, Snow J. 2017.** Formidable challenges to the notion of biologically important roles for dietary small RNAs in ingesting mammals. *Genes and Nutrition* **12**: 13 DOI 10.1186/s12263-017-0561-7.
- Thery C, Zitvogel L, Amigorena S. 2002.** Exosomes: composition, biogenesis and function. *Nature Reviews Immunology* **2**:569–579 DOI 10.1038/nri855.
- Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. 2007.** Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nature Cell Biology* **9**:654–659 DOI 10.1038/ncb1596.
- van der Grein SG, Nolte-'t Hoen EN. 2014.** "Small Talk" in the Innate Immune System via RNA-Containing Extracellular Vesicles. *Frontiers in Immunology* **5**: 542 DOI 10.3389/fimmu.2014.00542.
- Vaucheret H, Chupeau Y. 2012.** Ingested plant miRNAs regulate gene expression in animals. *Cell Research* **1**: 3-5 DOI 10.1038/cr.2011.164.
- Wagner AE, Piegholdt S, Ferraro M, Pallauf K, Rimbach G. 2015.** Food derived microRNAs. *Food & Function* **6**: 714-718 DOI 10.1039/c4fo01119h.

- 489 **Wang Yo, Li L, Tang Sh, Liu J, Zhang H, Zhi H, Jia G, Diao X. 2016.** Combined small RNA
490 and degradome sequencing to identify miRNAs and their targets in response to drought in
491 foxtail millet. *BMC Genomics* **17**: 57 DOI 10.1186/s12863-016-0364-7.
- 492 **Xiao J, Feng S, Wang X, Long K, Luo Y, Wang Y, Ma J, Tang Q, Jin L, Li X, Li M. 2018.**
493 Identification of exosome-like nanoparticle-derived microRNAs from 11 edible fruits and
494 vegetables. *PeerJ* **6**: e5186 DOI 10.7717/peerj.5186.
- 495 **Xie W, Weng A, Melzig MF. 2016.** MicroRNAs as new bioactive components in medicinal
496 plants. *Planta Medica* **82**: 1153-1162. DOI 10.1055/s-0042-108450.
- 497 **Yurikova OY, Aisina DE, Niyazova RE, Atambayeva SA, Labeit S, Ivashchenko AT. 2019.**
498 The Interaction of miRNA-5p and miRNA-3p with the mRNAs of Orthologous Genes. *Mol*
499 *Biol (Mosk)* **53**: 692-704 DOI 10.1134/S0026898419040189.
- 500 **Zempleni J, Aguilar-Lozano A, Sadri M, Sukreet S, Manca S, Wu D, Zhou F, Mutai E.**
501 **2017.** Biological Activities of Extracellular Vesicles and Their Cargos from Bovine and
502 Human Milk in Humans and Implications for Infants. *The Journal of Nutrition* **147**: 3-10 DOI
503 10.3945/jn.116.238949.
- 504 **Zhang L, Chen T, Yin Yo, Zhang Ch, Zhang Yo. 2019.** Dietary microRNA-A Novel
505 Functional Component of Food. *Advances in Nutrition* **10**: 711-721.
506 DOI 10.1093/advances/nmy127.
- 507 **Zhang L, Hou D, Chen X, Li D, Zhu L, Zhang Y, Li J, Bian Z, Liang X, Cai X, Yin Y,**
508 **Wang C, Zhang T, Zhu D, Zhang D, Xu J, Chen Q, Ba Y, Liu J, Wang Q, Chen J, Wang**
509 **J, Wang M, Zhang Q, Zhang J, Zen K, Zhang CY. 2012.** Exogenous plant MIR168a
510 specifically targets mammalian LDLRAP1: evidence of cross-kingdom regulation by
511 microRNA. *Cell Research* **22**: 107-126 DOI 10.1038/cr.2011.158.
- 512 **Zhang H, Li Y, Liu Y, Liu H, Wang H, Jin W, Zhang Y, Zhang C, Xu D. 2016.** Role of plant
513 microRNA in cross-species regulatory networks of humans. *BMC Systems Biology* **10**: 1-10
514 DOI 10.1186/s12918-016-0292-1.
- 515 **Zhao Q, Liu Yu, Zhang N, Hu M, Zhang H, Joshi T, Xu D. 2018.** Evidence for plant-derived
516 xenomiRs based on a large-scale analysis of public small RNA sequencing data from human
517 samples. *PLOS One* **13**: 6 DOI 10.1371/journal.pone.0187519.
- 518 **ZhaoY, Cong L, Lukiw WJ. 2018.** Plant and animal microRNAs (miRNAs) and their potential
519 for interkingdom communication. *Cellular and Molecular Neurobiology* **38**: 133-140 DOI
520 10.1007/s10571-017-0547-4.
- 521 **Zhou Z, Li X, Liu J, Dong L, Chen Q, Liu J, Kong H, Zhang Q, Qi X, Hou D, Zhang**
522 **L, Zhang G, Liu Y, Zhang Y, Li J, Wang J, Chen X, Wang H, Zhang J, Chen H, Zen**

523 **K, Zhang CY. 2015.** Honeysuckle-encoded atypical microRNA2911 directly targets
 524 influenza A viruses. Cell research **1**: 39-49 DOI 10.1038/cr.2014.130.

Gene, miRNA, start of site, region, ΔG , $\Delta G/\Delta G_m$	Gene, miRNA, start of site, region, ΔG , $\Delta G/\Delta G_m$
<p><i>RPS6KA5</i>, miR5075-3p, 261, CDS, -121, 98, 21</p> <p>5' -GCGGACGGCGCGGACGGAGGA-3'</p> <p> </p> <p>3' -CGCCUGCCGCCGUGCCUCUU-5'</p>	<p><i>ZNF442</i>, miR2866-5p, 1020, CDS, -98, 96, 20</p> <p>5' -GAUGCUGGACACAAACCAGA-3'</p> <p> </p> <p>3' -CUACGACUUGUGUUUGAUCU-5'</p>
<p><i>ZNHIT2</i>, miR2925-5p, 559, CDS, -106, 94, 19</p> <p>5' -GCGGAGCCCGCGGCCGCCG-3'</p> <p> </p> <p>3' -UGCUUCGGGCGCCGGCGGU-5'</p>	<p><i>C14orf142</i>, miR414-5p, 281, CDS, -104, 94, 21</p> <p>5' -GGACGGUGAUGAUGAAGAUGA-3'</p> <p> </p> <p>3' -CCUGCUACUACUACUCCUACU-5'</p>
<p><i>EML1</i>, miR535-3p, 2092, CDS, -106, 94, 21</p> <p>5' -AGUGACAACGGGAGGAAGUAC-3'</p> <p> </p> <p>3' -UCACUGUUGCCUCUUUCGUG-5'</p>	<p><i>C1D</i>, miR2919, 543, 3'UTR, -102, 94, 19</p> <p>5' -UCUUCCCCCCCCCCCCC-3'</p> <p> </p> <p>3' -AGAAAGGGGGGGGGGAA-5'</p>
<p>Note: Gene; miRNA; start of binding site (nt); ΔG (kJ/mole); $\Delta G/\Delta G_m$ (%), length of miRNA (nt). The upper and lower nucleotide sequences of mRNA and miRNA, respectively. The bold type indicates the nucleotide of non-canonical pairs U-G, A-C.</p>	

Figure 1. Schemes of the interaction of the nucleotide sequences of osa-miRNA with mRNA human genes.

<p><i>NKRD27</i>, miR1428e-3p, 244, CDS, -98, 96 5' -CAAAUCCAUGGCAUUGUCUUA-3' 3' -GUUUAAGUACCGUAAUAGAAU-5'</p> <p><i>AP2A2</i>, miR156k-5p, 3797, 3'UTR, -104, 94 5' -UGUGCUCGUCUCUUCUGUCA-3' 3' -ACACGAG-AGAGAGAAGACAGU-5'</p> <p><i>PKHD1</i>, miR159f-3p, 9966, CDS, -108, 98 5' -UAGAGCUCCUCCAAUCCAAG-3' 3' -AUCUCGAGGGAAGUUAGGUUC-5'</p> <p><i>CHST11</i>, miR166a-5p, 986, CDS, -104, 94 5' -CCUGAACCAGUACAGCAUCC-3' 3' -GGAACUUGGUC-UGUUGUAAGG-5'</p> <p><i>CCNY</i>, miR166h-5p, 367, CDS, -110, 96 5' -CCUCGGGCCAGCACAAUUAUCC-3' 3' -GGAGCUCGGUCG-GUUGUAAGG-5'</p> <p><i>PRICKLE2</i>, miR167d-5p, 1959, CDS, -106, 94 5' -CAGGACAUGCUGGCAGCUUCA-3' 3' -GUCUAGUACGACCGUCGAAGU-5'</p> <p><i>KLF14</i>, miR168b-5p, 623, CDS, -115, 98 5' -UUCCCGGGCUGCACCAGCCU-3' 3' -AAGGGCUCGACGUGGUU-CGGA-5'</p> <p><i>GPR31</i>, miR172a-3p, 590, CDS, -100, 94 5' -AUGCAGGCAUCAUCAGGGCUCU-3' 3' -UACGUC-GUAGUAGUUAAGA-5'</p>	<p><i>EIF3B</i>, miR1846b-5p, 2194, CDS, -119, 97 5' -GGCGGCCCCGGCCUCCACACU-3' 3' -UCGCCGGGGCCGGAGGA-GUGA-5'</p> <p><i>SLC25A37</i>, miR2907a-3p, 884, CDS, -125, 95 5' -CCGGGGCCUCGCGCGCGCCGCC-3' 3' -GGCUCGGGAGCGA-GCCGACGG-5'</p> <p><i>CD22</i>, miR395t-3p, 2684, 3'UTR, -104, 94 5' -GAGUUUCCCAGACACCGCCAC-3' 3' -CUCAAAGGGGUUGUGA-AGUG-5'</p> <p><i>SYVN1</i>, miR396a-5p, 854, CDS, -102, 94 5' -CAGUUCAGAAAGCUGUGACAG-3' 3' -GUCAAGUUCUUCGACACC-UU-5'</p> <p><i>C19orf57</i>, miR444b.1-3p, 714, CDS, -106, 94 5' -GACAGCAAGCCUGAGACAGACA-3' 3' -CCGUCGUUCGAACUCUGU-UGU-5'</p> <p><i>PDE4B</i>, miR529a-3p, 2270, CDS, -100, 94 5' -GAAGGAGGGAGAGGGACACAG-3' 3' -CUUCUUCUCUCUCCC-AUGUC-5'</p> <p><i>ENAH</i>, miR531b-5p, 106, 5'UTR, -115, 95 5' -CGGGCGCGCGGCCCGGGCGGG-3' 3' -GCC-GUGCGUCGGGGCCGCUC-5'</p> <p><i>TIFA</i>, miR815a-3p, 186, 5'UTR, -106, 94 5' -CCCAGUCUCCUGAUUCCCCUC-3' 3' -GGGUUAGAGGAGUUA-GGGGAA-5'</p>
<p>Note: Gene; miRNA; start of binding site (nt); ΔG (kJ/mole); $\Delta G/\Delta G_m$ (%), length of miRNA (nt). The upper and lower nucleotide sequences of mRNA and miRNA, respectively. The bold type indicates the nucleotide of non-canonical pairs U-G, A-C.</p>	

Figure 3. Schemes of the interaction of nucleotide sequences of osa-miRNA families with mRNA human genes.

Parsed Citations

Aisina D, Niyazova R, Atambayeva S, Ivashchenko A. 2019. Prediction of clusters of miRNA binding sites in mRNA candidate genes of breast cancer subtypes. PeerJ 7: e8049 DOI 10.7717/peerj.8049.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Arroyo JD, Chevillet JR, Kroh EM, Ruf IK, Pritchard CC, Gibson DF, Mitchell PS, Bennett CF, Pogosova-Agadjanyan EL, Stirewalt DL, Tait JF, Tewari M. 2011. Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. Proceedings of the National Academy of Sciences of the United States of America 108: 5003-5008 DOI 10.1073/pnas.1019055108.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Arteaga-Vazquez M, Caballero-Perez J, Vielle-Calzada JP. 2006. A family of microRNAs present in plants and animals. Plant Cell 18: 3355-3369 DOI 10.1105/tpc.106.044420.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Atambayeva S, Niyazova R, Ivashchenko A, Pyrkova A, Pinsky I, Akimniyazova A, Labeit S. 2017. The Binding Sites of miR-619-5p in the mRNAs of Human and Orthologous Genes. BMC Genomics 18: 428. DOI 10.1186/s12864-017-3811-6.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Axtell MJ, Westholm JO, Lai EC. 2011. Vive la difference: biogenesis and evolution of microRNAs in plants and animals. Genome Biology 12: 1-13. DOI 10.1186/gb-2011-12-4-221.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Balyan S, Kumar M, Mutum R, Raghuvanshi U, Agarwal P, Mathur S, Raghuvanshi S. 2017. Identification of miRNA-mediated drought responsive multi-tiered regulatory network in drought tolerant rice, Nagina 22. Scientific Reports 7: 15446 DOI 10.1038/s41598-017-15450-1.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Bang C, Thum T. 2012. Exosomes: new players in cell-cell communication. The International Journal of Biochemistry & Cell Biology 44:2060-2064 DOI 10.1016/j.biocel.2012.08.007.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Bari A, Orazova S, Ivashchenko A. 2013. miR156- and miR171-binding sites in the protein-coding sequences of several plant genes. BioMed Research International 2013: 1-7 DOI 10.1155/2013/307145.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Bari A, Sagaidak I, Pinski I, Orazova S, Ivashchenko A. 2014. Binding of miR396 to mRNA of Genes Encoding Growth-Regulating Transcription Factors of Plants. Russian Journal of Plant Physiology 61: 807-810 DOI 10.1134/S1021443714050033.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Buck AH, Coakley G, Simbari F, McSorley HJ, Quintana JF, Le Bihan T, Kumar S, Abreu-Goodger C, Lear M, Harcus Y, Ceroni A, Babayan SA, Blaxter M, Ivens A, Maizels RM. 2014. Exosomes secreted by nematode parasites transfer small RNAs to mammalian cells and modulate innate immunity. Nature Communications 5: 5488 DOI 10.1038/ncomms6488.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Chen J, Li L. 2018. Multiple Regression Analysis Reveals MicroRNA Regulatory Networks in Oryza sativa under Drought Stress. International Journal of Genomics 2018: 1-12 DOI 10.1155/2018/9395261.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Chiang K, Shu J, Zemleni J, Cui J. 2015. Dietary MicroRNA Database (DMD): An Archive Database and Analytic Tool for Food-Borne microRNAs. PLOS One 10: 6 DOI 10.1371/journal.pone.0128089.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Chin AR, Fong MY, Somlo G. 2016. Cross-kingdom inhibition of breast cancer growth by plant miR159. Cell Research 26: 217-228 DOI 10.1038/cr.2016.13.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Cimini S, Gualtieri C, Macovei A, Balestrazzi A, De Gara L, Locato V. 2019. Redox Balance-DDR-miRNA Triangle: Relevance in Genome Stability and Stress Responses in Plants. Frontiers in Plant Science 10: 989 DOI 10.3389/fpls.2019.00989.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Cong L, Zhao Y, Pogue AI, Lukiw WJ. 2018. Role of microRNA (miRNA) and Viroids in Lethal Diseases of Plants and Animals. Potential Contribution to Human Neurodegenerative Disorders. *Biochemistry (Moscow)* 83: 1018-1029 DOI 10.1134/S0006297918090031.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Cui J, Zhou B, Ross SA, Zemleni J. 2017. Nutrition, microRNAs, and Human Health. *Advances in Nutrition* 8: 105-112 DOI 10.3945/an.116.013839.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Dai X, Zhuang Z, Zhao P. 2011. Computational Analysis of miRNA Targets in Plants: Current Status and Challenges. *Briefings in Bioinformatics* 12: 115-121 DOI 10.1093/bib/bbq065.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Denzer K, Eijk MV, Kleijmeer MJ, Jakobson E, Groot CD, Geuze HJ. 2000. Follicular dendritic cells carry MHC Class II-expressing microvesicles at their surface. *Journal of Immunology* 165:1259–1265 DOI 10.4049/jimmunol.165.3.1259.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Escrevente C, Keller S, Altevogt P, Costa J. 2011. Interaction and uptake of exosomes by ovarian cancer cells. *BMC Cancer* 11: 108 DOI 10.1186/1471-2407-11-108.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Garg A, Heinemann U. 2018. A novel form of RNA double helix based on G·U and C·A⁺ wobble base pairing. *RNA* 24: 209-218 DOI 10.1261/rna.064048.117.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Gopinath MS. 2019. Dietary non-coding RNAs from plants: Fairy tale or treasure? *Non-coding RNA Research* 4: 63-68 DOI 10.1016/j.ncrna.2019.02.002.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Hou D, He F, Ma L, Cao M, Zhou Z, Wei Z, Xue Y, Sang X, Chong H, Tian C, Zheng S, Li J, Zen K, Chen X, Hong Z, Zhang CY, Jiang X. 2018. The potential atheroprotective role of plant MIR156a as a repressor of monocyte recruitment on inflamed human endothelial cells. *The Journal of Nutritional Biochemistry* 57: 197-205 DOI 10.1016/j.jnutbio.2018.03.026.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Hoy AM, Buck AH. 2012. Extracellular small RNAs: what, where, why? *Biochemical Society Transactions* 40: 886-890 DOI 10.1042/BST20120019.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ivashchenko AT, Pyrkova AY, Niyazova RY, Alybayeva A, Baskakov K. 2016. Prediction of miRNA binding sites in mRNA. *Bioinformation* 12: 237-240. DOI 10.6026/97320630012237.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Javed M, Solanki M, Sinha A, Shukla LI. 2017. Position Based Nucleotide Analysis of miR168 Family in Higher Plants and its Targets in Mammalian Transcripts. *Microna* 6: 136-142 DOI 10.2174/2211536606666170215154151.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Jia L, Zhang D, Qi X, Ma B, Xiang Zh, He N, Zhang J. 2014. Identification of the Conserved and Novel miRNAs in Mulberry by High-Throughput Sequencing. *PLOS One* 9: 8 DOI 10.1371/journal.pone.0104409.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Jonathan WS, Andrew EH, Stephanie KI, Aaron LB, Stephen YC. 2013. Ineffective delivery of diet-derived microRNAs to recipient animal organisms. *RNA Biology* 10: 1107-1116 DOI 10.4161/rna.24909.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Jones Buie JN, Goodwin AJ, Cook JA, Halushka PV, Fan H. 2016. The role of miRNAs in cardiovascular disease risk factors. *Atherosclerosis* 254: 271-281 DOI 10.1016/j.atherosclerosis.2016.09.067.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kool ET. 2001. Hydrogen bonding, base stacking, and steric effects in DNA replication. Annual Review of Biophysics and Biomolecular Structure 30: 1-22 DOI 10.1146/annurev.biophys.30.1.1.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lang C, Karunairetnam S, Lo KR, Kralicek AV, Crowhurst RN, Gleave AP, MacDiarmid RM, Ingram JR. 2019. Common Variants of the Plant microRNA-168a Exhibit Differing Silencing Efficacy for Human Low-Density Lipoprotein Receptor Adaptor Protein 1 (LDLRAP1). Microna 8: 166-170 DOI 10.2174/2211536608666181203103233.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Leontis NB, Stombaugh J, Westhof E. 2002. The non-Watson-Crick base pairs and their associated isostericity matrices. Nucleic Acids Research 30: 3497-3531 DOI 10.1093/nar/gkf481.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liang H, Huang L, Cao J, Zen K, Chen X, Zhang CY. 2012. Regulation of mammalian gene expression by exogenous microRNAs. Wiley Interdisciplinary Reviews-RNA 3: 733-742 DOI 10.1002/wrna.1127.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liang H, Zen K, Zhang J, Zhang CY, Chen X. 2013. New roles for microRNAs in cross-species communication. RNA Biology 10: 367-370 DOI 10.4161/rna.23663.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liang H, Zhang S, Fu Z, Wang Y, Wang N, Liu Y, Zhao C, Wu J, Hu Y, Zhang J, Chen X, Zen K, Zhang CY. 2015. Effective detection and quantification of dietetically absorbed plant microRNAs in human plasma. The Journal of Nutritional Biochemistry 26: 505-512 DOI 10.1016/j.jnutbio.2014.12.002.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu W, Meng J, Cui J, Luan Yu. 2017. Characterization and Function of MicroRNAs in Plants. Frontiers in Plant Science 8. DOI 10.3389/fpls.2017.02200.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lukasik A, Zielenkiewicz P. 2016. Plant microRNAs - novel players in natural medicine? The International Journal of Molecular Sciences 18: 1-16 DOI 10.3390/ijms18010009.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Luo Y, Wang P, Wang X, Wang Y, Mu Z, Li Q, Fu Y, Xiao J, Li G, Ma Y, Gu Y, Jin L, Ma J, Tang Q, Jiang A, Li X, Li M. 2017. Detection of dietetically absorbed maize-derived microRNAs in pigs. Scientific Reports 7: 1-10 DOI 10.1038/s41598-017-00488-y.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Makarova JA, Shkurnikov MU, Wicklein D, Lange T, Samatov TR, Turchinovich AA, Tonevitsky AG. 2016. Intracellular and extracellular microRNA: an update on localization and biological role. Progress in Histochemistry and Cytochemistry 51: 33-49 DOI 10.1016/j.proghi.2016.06.001.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mallocci M, Perdomo L, Veerasamy M, Andriantsitohaina R, Simard G, Martinez MC. 2018. Extracellular vesicles: mechanisms in human health and disease. Antioxidants & Redox Signaling 30: 813-856 DOI 10.1089/ars.2017.7265.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Melnik BC, John SM, Schmitz G. 2014. Milk: an exosomal microRNA transmitter promoting thymic regulatory T cell maturation preventing the development of atopy? The Journal of Translational Medicine 12: 1-11 DOI 10.1186/1479-5876-12-43.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Millar AA, Waterhouse PM. 2005. Plant and animal microRNAs: similarities and differences. Functional & Integrative Genomics 5: 129-135 DOI 10.1007/s10142-005-0145-2.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mondal T, Panda A, Rawal H, Sharma T. 2018. Discovery of microRNA-target modules of African rice (Oryza glaberrima) under salinity stress. Scientific Reports 8: 570 DOI 10.1038/s41598-017-18206-z.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Montecalvo A, Larregina AT, Shufesky WJ, Stolz DB, Sullivan ML, Karlsson JM, Baty CJ, Gibson GA, Erdos G, Wang Z, Milosevic J,

Tkacheva OA, Divito SJ, Jordan R, Lyons-Weiler J, Watkins SC, Morelli AE. 2012. Mechanism of transfer of functional microRNAs between mouse dendritic cells via exosomes. Blood 119: 756-766 DOI 10.1182/blood-2011-02-338004.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Moran Y, Agron M, Praher D, Technau U. 2017. The evolutionary origin of plant and animal microRNAs. Nature Ecology & Evolution 1: 1-22 DOI 10.1038/s41559-016-0027.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nair SK, Wang N, Turuspekov Y, Pourkheirandish M, Sinsuwongwat S, Chen G, Sameri M, Tagiri A, Honda I, Watanabe Y, Kanamori H, wicker T, Stein N, Nagamura Y, Matsumoto T, Komatsuda T. 2010. Cleistogamous flowering in barley arises from the suppression of microRNA-guided HvAP2 mRNA cleavage. Proceedings of the National Academy of Sciences of the United States of America 107: 490-495 DOI 10.1073/pnas.0909097107.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Perge P, Nagy Z, Decmann A, Igaz I, Igaz P. 2017. Potential relevance of microRNAs in inter-species epigenetic communication, and implications for disease pathogenesis. RNA Biology. 14: 391-401 DOI 10.1080/15476286.2016.1251001.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pirro S, Minutolo A, Galgani A, Potesta M, Colizzi V, Montesano C. 2016. Bioinformatics prediction and experimental validation of microRNAs involved in cross-kingdom interaction. The Journal of Computational Biology 23: 976-989 DOI 10.1089/cmb.2016.0059.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pogue AI, Clement C, Hill JM, Lukiw WJ. 2014. Evolution of microRNA (miRNA) structure and function in plants and animals: relevance to aging and disease. Journal of Aging Science 2: 1-12 DOI 10.4172/2329-8847.1000119.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Redis RS, Calin S, Yang Y, You MJ, Calin GA. 2012. Cell-to-cell miRNA transfer:

from body homeostasis to therapy. Pharmacology & Therapeutics 136:169–174

DOI 10.1016/j.pharmthera.2012.08.003.

Rutter BD, Innes RW. 2018. Extracellular vesicles as key mediators of plant–microbe interactions. Current Opinion in Plant Biology 44: 16-22 DOI 10.1016/j.pbi.2018.01.008.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stephen Y, Snow J. 2017. Formidable challenges to the notion of biologically important roles for dietary small RNAs in ingesting mammals. Genes and Nutrition 12: 13 DOI 10.1186/s12263-017-0561-7.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Thery C, Zitvogel L, Amigorena S. 2002. Exosomes: composition, biogenesis and

function. Nature Reviews Immunology 2:569–579 DOI 10.1038/nri855.

Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. 2007. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic

exchange between cells. Nature Cell Biology 9:654–659 DOI 10.1038/ncb1596.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

van der Grein SG, Nolte-'t Hoen EN. 2014. "Small Talk" in the Innate Immune System via RNA-Containing Extracellular Vesicles. Frontiers in Immunology 5: 542 DOI 10.3389/fimmu.2014.00542.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vaucheret H, Chupeau Y. 2012. Ingested plant miRNAs regulate gene expression in animals. Cell Research 1: 3-5 DOI 10.1038/cr.2011.164.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wagner AE, Piegholdt S, Ferraro M, Pallauf K, Rimbach G. 2015. Food derived microRNAs. Food & Function 6: 714-718 DOI 10.1039/c4fo01119h.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang Yo, Li L, Tang Sh, Liu J, Zhang H, Zhi H, Jia G, Diao X. 2016. Combined small RNA and degradome sequencing to identify miRNAs

and their targets in response to drought in foxtail millet. *BMC Genomics* 17: 57 DOI 10.1186/s12863-016-0364-7.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xiao J, Feng S, Wang X, Long K, Luo Y, Wang Y, Ma J, Tang Q, Jin L, Li X, Li M. 2018. Identification of exosome-like nanoparticle-derived microRNAs from 11 edible fruits and vegetables. *PeerJ* 6: e5186 DOI 10.7717/peerj.5186.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xie W, Weng A, Melzig MF. 2016. MicroRNAs as new bioactive components in medicinal plants. *Planta Medica* 82: 1153-1162. DOI 10.1055/s-0042-108450.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yurikova OY, Aisina DE, Niyazova RE, Atambayeva SA, Labeit S, Ivashchenko AT. 2019. The Interaction of miRNA-5p and miRNA-3p with the mRNAs of Orthologous Genes. *Mol Biol (Mosk)* 53: 692-704 DOI 10.1134/S0026898419040189.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zempleni J, Aguilar-Lozano A, Sadri M, Sukreet S, Manca S, Wu D, Zhou F, Mutai E. 2017. Biological Activities of Extracellular Vesicles and Their Cargos from Bovine and Human Milk in Humans and Implications for Infants. *The Journal of Nutrition* 147: 3-10 DOI 10.3945/jn.116.238949.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang L, Chen T, Yin Yo, Zhang Ch, Zhang Yo. 2019. Dietary microRNA-A Novel Functional Component of Food. *Advances in Nutrition* 10: 711-721. DOI 10.1093/advances/nmy127.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang L, Hou D, Chen X, Li D, Zhu L, Zhang Y, Li J, Bian Z, Liang X, Cai X, Yin Y, Wang C, Zhang T, Zhu D, Zhang D, Xu J, Chen Q, Ba Y, Liu J, Wang Q, Chen J, Wang J, Wang M, Zhang Q, Zhang J, Zen K, Zhang CY. 2012. Exogenous plant MIR168a specifically targets mammalian LDLRAP1: evidence of cross-kingdom regulation by microRNA. *Cell Research* 22: 107-126 DOI 10.1038/cr.2011.158.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang H, Li Y, Liu Y, Liu H, Wang H, Jin W, Zhang Y, Zhang C, Xu D. 2016. Role of plant microRNA in cross-species regulatory networks of humans. *BMC Systems Biology* 10: 1-10 DOI 10.1186/s12918-016-0292-1.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhao Q, Liu Yu, Zhang N, Hu M, Zhang H, Joshi T, Xu D. 2018. Evidence for plant-derived xenomiRs based on a large-scale analysis of public small RNA sequencing data from human samples. *PLOS One* 13: 6 DOI 10.1371/journal.pone.0187519.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

ZhaoY, Cong L, Lukiw WJ. 2018. Plant and animal microRNAs (miRNAs) and their potential for interkingdom communication. *Cellular and Molecular Neurobiology* 38: 133-140 DOI 10.1007/s10571-017-0547-4.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhou Z, Li X, Liu J, Dong L, Chen Q, Liu J, Kong H, Zhang Q, Qi X, Hou D, Zhang L, Zhang G, Liu Y, Zhang Y, Li J, Wang J, Chen X, Wang H, Zhang J, Chen H, Zen K, Zhang CY. 2015. Honeysuckle-encoded atypical microRNA2911 directly targets influenza A viruses. *Cell research* 1: 39-49 DOI 10.1038/cr.2014.130.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)