Multidrug resistance and high prevalence of class 1 integrons in *Escherichia coli* isolated from waters and vegetables in Nsukka and Enugu, Nigeria.

Chinyere B. Chigor^{1,2¶}, Ini-Abasi I. Ibangha^{1,3&}, Nkechinyere O. Nweze^{1,2¶},
Chizoba A. Ozochi^{1,3¶}, Valentino C. Onuora^{1,3&}, Yinka Titilawo^{4¶}, Tatyana N. Chernikova^{5¶},
Peter N. Golyshin⁵ and Vincent N. Chigor^{1,3*}

¹Water and Public Health Research Group (WPHRG), University of Nigeria, Nsukka, Enugu State, Nigeria.

²Department of Plant Science and Biotechnology, Faculty of Biological Sciences, University of Nigeria, Nsukka, Enugu State, Nigeria.

³Department of Microbiology, Faculty of Biological Sciences, University of Nigeria, Nsukka, Enugu State, Nigeria.

⁴Department of Biology/Microbiology/Biotechnology, Alex Ekwueme Federal University, Ndufu-Alike Ikwo, Ebonyi State, Nigeria.

⁵School of Natural Sciences, Bangor University, Bangor Gwynedd, United Kingdom

*Corresponding author

E-mail: vincent.chigor@unn.edu.ng (VNC)

"VNC and PNG are Joint Senior Authors"

Keywords: multidrug resistance; Integrons; pathogenic *E. coli*; wastewater-irrigated vegetables; public health risks

Running title: Multidrug resistance and class 1 integrons in *Escherichia coli*

Abstract

In spite of treated wastewater presenting itself as an attractive alternative to scarce quality water in the developing countries, the associated contamination of fresh produce by irrigation waters leading to outbreak of foodborne illnesses is on the rise. Horizontal transfer of integrons play important role in the spread and maintenance of antimicrobial resistance among strains of *Escherichia coli*. This study assessed the effluents from the University of Nigeria, Nsukka Wastewater Treatment Plant (UNN-WWTP) as well as vegetables irrigated with the effluent, and vegetables sold in selected markets from Nsukka and Enugu cities for the presence of *E. coli* and determined the prevalence integrons in multidrug-resistant isolates. Isolation of *E. coli* was done using eosin methylene blue agar and isolates subjected to Gram staining for identification of presumptive colonies. Confirmation of *E. coli* was achieved by polymerase chain reaction (PCR) technique, targeting beta-glucuronidase (*uidA*). Resistance to antibiotics was determined using the Bauer-Kirby disk diffusion assay and the Clinical and Laboratory Standard Institute criteria. Integrons were detected by multiplex PCR using primers specific for class 1 and 2 integrons. A total of 178 *E. coli* isolates were obtained from WWTP effluent (41), and vegetables from greenhouse (46), farms (55) and market (36). Multi-drug resistance was detected in all the isolates, ranging from five-drug resistance in a single isolate to 16-drug resistance patterns in two different isolates. Of the total isolates, class 1 integrons were abundantly detected in 175 (98.3%) and class 2 in 5 (2.8%). All the class 2 integrons were found in isolates that were positive for class 1. The high detection of *E. coli* in the studied effluent and vegetables pose potential public health hazards heightened by observed multidrug resistance in all the isolates and the high prevalence of class 1 integron. It is concluded that the vegetable samples are significant reservoirs for potentially pathogenic *E. coli*. Therefore, vegetable irrigation farming with unsafe water should be discontinued, while appropriate improvement strategies to ensure compliance should be facilitated without further delay.

Introduction

Pathogenic *Escherichia coli* causes significant morbidity and mortality worldwide [1-3]. Reported risk factors in the developing countries and sub-Saharan African regions include poor hygiene, unsafe water, improper disposal of waste and faeces, and contaminated food, local beverages and vegetables [2, 4, 5]. Vegetables can become contaminated with pathogenic and commensal

bacteria from animals and humans, during growth, harvesting, distribution, storage and processing [6]. Although the contamination of fresh produce by irrigation waters has led to outbreak of foodborne illnesses, yet treated wastewater presents itself as an attractive alternative to scarce quality water in the developing countries.

E. coli has been reported as an aetiological agent of diarrhoea in both the northern and south-western parts of Nigeria [7-12]. In a study that detected *E. coli* in 119 (44.74%) of 270 diarrhoeal stool samples in Enugu and Onitsha cities, south-eastern Nigeria [13], enterotoxigenic *E. coli* (ETEC) was reported as the second most prevalent pathotype (21.57%) after enteropathogenic *E. coli* (EPEC) (49.02%). Likewise, a study conducted in Nsukka, that involved watery stools, drinking water, and some fruits and vegetables collected during the rainy periods (between April and October) over 3 year sampling regime (1996 to 1998), [4] reported that enteropathogenic *E. coli* (EPEC) was detected in 9 (1.8%) of 500 stool samples, whereas no enteric bacterial pathogen was isolated from the fruits and vegetables. There appears to be no reports on ETEC prevalence in humans and on irrigated vegetables in Nsukka.

Excessive and inappropriate usage of antimicrobials in preventing or treating human and veterinary bacterial infectious diseases has led to increased antimicrobial and multidrug resistance (MDR) and the risk of transmission of antibiotic resistant bacteria (ARB) and antibiotic resistant genes (ARGs) from one country to another is a growing global challenge. [14-16] Attention should be given to how anthropogenic activities might be causing evolution of antibiotic resistance in the environment [16], and studies have shown that waste water treatment plants form a significant reservoir of resistance genes and suggested that waste water disposal increases the reservoir of resistance determinants in the environment either by the addition of resistance genes or input of agents selective for resistant phenotypes [17].

Along with transposons and plasmids, integrons, genetic elements commonly found in bacterial genomes that allow efficient acquisition and expression of exogenous genes, are central in the dissemination of antibiotic resistance among Gram-negative bacteria [18,19]. Horizontal transfer of integrons have been shown to play important role in the spread and maintenance of antimicrobial resistance among strains of *E. coli* and ARB can be transferred across borders by human travelers, animal and insect vectors, agricultural products and surface water [15,20]. Not much is known about the risk factors in spreading across local borders.

It is thought that University towns, characterized by regular and significant demographic changes arising from admissions and vacations, could play major role in dissemination of resistance determinants locally, and even internationally where the institution has a good number of internationals. Nsukka, in Southeast Nigeria, is the location of one of Nigeria's biggest universities and one also in which the town developed around the university. This study assessed the effluent from the University of Nigeria, Nsukka Wastewater Treatment Plant (UNN-WWTP) as well as vegetables irrigated with the effluent and vegetables sold in selected markets for the presence of *E. coli* and determined the prevalence integrons in multidrug-resistant isolates.

Methods

Description of study area

The university town of Nsukka (6.8429° N, 7.3733° E) is in Enugu State, southeast Nigeria, with an area of 1,810 km² and a population of 309,633 (NPC 2006). The sewage treatment facility (WWTP) in Nsukka, consisting of a screen, primary settling (Imhoff) tank, sludge drying beds and two oxidation ponds, is situated at the northwest end of the University of Nigeria, Nsukka. The final effluents have been widely utilized for fresh produce irrigation during dry season.

Cultivation of *Amaranthus* in the greenhouse

The most commonly cultivated vegetables in the study area, during the dry season, include the green leafy vegetable amaranth (*Amaranthus* spp), fluted pumpkin leaves (*Telfaria occidentalis*), scarlet eggplant leaf (*Solanum aethiopicum*) and water leaf (*Talinum fruticosum*). In this study, *Amaranthus* was chosen, being the second most produced and sold leafy vegetable, after *Telfaria* [21], and it equally grows very easily and matures faster. Amaranths were grown for 10 weeks (July 26 to October 03, 2014) in earthen pots at the Soil Science Departmental greenhouse. They were irrigated daily using the sprinkler method. A total of 60 earthen pots were used for the cultivation of vegetables, 48 were irrigated with treated wastewater (final effluent of the University of Nigeria, wastewater treatment plant (WWTP) and 12 with tap water. The pots irrigated with tap water served as the control.

Collection of Samples

Samples collected for this study included treated wastewater and vegetables. Sampling was done according to the standard procedure [22]. Effluents were collected with 10 L plastic cans for irrigation of the green house vegetables. Samples of the WWTP effluent were collected using sterile wide-mouthed, screw-capped 250-ml bottles. Vegetables were obtained from the green house, irrigated gardens and local markets in Nsukka and Enugu metropolis, during December 2014. Samples of the major vegetables cultivated during the dry season include fluted pumpkin leaves (*Telfaria occidentalis*), scarlet eggplant leaf (*Solanum aethiopicum*), water leaf (*Talinum fruticosum*) and the green vegetable (*Amaranthus* Spp) were collected. All samples were transported on ice to the laboratory and analysed within 6 h of collection.

Isolation and identification of presumptive *E. coli*

This was carried out at the Water and Public Health Laboratory, University of Nigeria Nsukka. Exactly 5 g of each vegetable sample was homogenized in a clean porcelain mortar, and 1 g of the homogenate diluted into 9ml normal saline [23, 24]. Serial dilutions (10-fold) were made by pipetting out 1ml stock solution into successive 9ml of sterile normal saline bottles. A 1 ml working sample dilution (10^{-1} and 10^{-2}) was spread-plated onto eosin methylene blue (EMB) agar (Oxoid, UK), incubated at 44 °C for 18-24 h. Raised, entire colonies with dark greenish metallic sheen, typical *E. coli* colonies were subjected to Gram-staining [25] and standard biochemical tests (IMViC). All presumptive *E. coli* isolates were sub-cultured in tryptic soy broth (Oxoid, UK) and then stored at -20 °C for further investigations. All media were prepared following the manufacturers' instructions.

Extraction of genomic DNA

Genomic DNA were extracted from a pure culture of each isolate grown overnight on nutrient agar at 37°C, by the conventional boiling method, as described [28]. Briefly, one loopful of bacterial cells was suspended in 1ml of sterile distilled water. The bacterial suspensions were then heated for 5 min at 100°C, cooled to room temperature and centrifuged at 12,000 xg for 5 min to remove the debris. The supernatant was stored at -20°C and used as the template DNA for PCR analysis.

Detection of beta-glucuronidase (*uidA*) gene for confirmation of *E. coli*

The confirmation of *E. coli* was achieved by polymerase chain reaction (PCR) detection of the target beta-glucuronidase (*uidA*). This was done at the School of Natural Sciences, Bangor University, United Kingdom, following the procedures described [29, 30]. The extracted DNA were cleaned using QIAGEN (QIAEX®II) gel extraction kits and kept at -20°C. PCRs were carried out with BIORAD DNA Engine Tetrad®Peltier Thermal Cycler (BIORAD, USA). The PCR reaction mixtures consisted of 25 µl of PCR Master Mix (Thermo Scientific, (EU) Lithuania), 0.5 µl each of oligonucleotide primers (Eurofins Genomics, Ebersberg Germany), 10 µl of template DNA and 14 µl of nuclease free water to constitute a total reaction volume of 50 µl. The PCR cycling conditions, with some modifications, were in accordance with the protocols prescribed elsewhere [31]. *E. coli* strain (NCTC 13353) and *Enterobacter aerogenes* (NCTC 10006) were used as positive and negative controls respectively for *E. coli* genus identification. The oligonucleotide sequence of primers used, target genes and expected amplification products are given in Table 1. For gel electrophoresis, 3 µl of DNA ladder (1 Kb Plus DNA Ladder; Invitrogen), 6 µl of positive control and 10µl of template DNA were ran on 2.5% (w/v) agarose gels in 1x-TBE buffer (0.09 M Tris-borate and 0.002 M EDTA, pH 8.0) at 100V for 25-30 min. The gels were viewed and photographed with BIORAD Molecular Imager® Gel Doc™ XR Imaging System (BIORAD, USA).

Antibiotic susceptibility testing

Isolates were subjected to antibiotic susceptibility testing using the Kirby-Bauer disc diffusion test [26]. Evaluation of results was based on the standards of the Clinical Laboratory Standards Institute (CLSI) [27]. Briefly, isolates grown on nutrient broth were suspended into sterile normal saline (0.9% (w/v) NaCl) with the aid of a sterile wire loop until the turbidity equivalent of 0.5 McFarland standard was reached. Sterile non-toxic cotton swabs were dipped into the standardized inoculum and used to smear the entire surface of the Muller-Hinton agar (Thermo Fisher Scientific, USA) plates. Antibiotic discs were placed aseptically using sterile forceps. All plates were incubated at 35±2oC for 16 to 18 h. The following antibiotics were employed for the test: Amoxycillin (AMX) 10µg, Ampicillin (AMP) 10µg, Metronidazole (MTZ) 5µg, Rifampicin (RIF) 5µg, Vancomycin (VAN) 30µg, Cloxacillin (COX) 5µg, Penicillin G (PNG) 10iu, Streptomycin (STR) 10µg, Erythromycin (ERT) 15µg, Clarithromycin (CLR) 15µg,

Cefuroxime (CXM) 30µg, Chloramphenicol (CHL) 30µg, Imipenem (IPM) 10µg, Tetracycline (TET) 30µg, Ciprofloxacin (CIP) 5µg, Trimethoprim (TMP) 5µg, Norfloxacin (NOR) 10µg, Sulphamethoxazole (SMZ) 25µg. The *E. coli* ATCC 25922 strains was used as control for antibiotic susceptibility testing. Zones showing complete inhibition around the discs were measured and classified as resistant (R), intermediate (I) and susceptible (S) according to the diameters of the zones recorded to the nearest millimetres.

Detection of integrons

The isolates were screened for class 1, 2 and 3 integrons by a multiplex PCR procedure as described by Machado *et al.* [32] and Karger *et al.* [33]. The PCR reactions (a total volume of 50µl reaction mixture) each consisted of consisting of 10 µl Buffer of 5x MyTaq Reaction Buffer (Bioline, with dye), 0.75µl of each the primers *intI1*, *intI2* and *intI3* (Eurofins Genomics, Ebersberg, Germany.), 27.25µl nuclease free water (Sigma-Aldrich), 0.25µl MyTaq DNA polymerase (Bioline) and 5µl DNA template. For gel electrophoresis, 3µl of DNA ladder (1 Kb Plus DNA Ladder; Invitrogen), 6 µl of positive control and 10 µl of samples were ran on 1.5% (w/v) agarose gels in 1x-TBE buffer (0.09 M Tris-borate and 0.002 M EDTA, pH 8.0) at 100 V for 25 min. The gels were viewed and photographed with BIORAD Molecular Imager® Gel Doc™ XR Imaging System.

209 **Table 1: Primers for the detection of *E. coli* and integrons**

Genetic marker	Primer name and sequence (5' to 3')		Amplicon size (bp)	Thermocycling conditions	Reference
<i>uidA</i>	UAL-754	AAAACGGCAAGAAAAAGCAG	147	Initial activation at 95°C for 3 min, followed by 40 cycles consisting of denaturing at 94°C for 1 min, annealing at 65°C for 1 min, extension at 70°C for 1 min and final elongation at 72°C for 7 min	[29, 30]
(beta-glucuronidase)	UAR-900	ACGCGTGGTTACAGTCTTGCG			
<i>intI1</i> (Class 1 integron)	Int1-F Int1-R	GGTCAAGGATCTGGATTTTCG ACATGCGTGTAATCATCGTC	436	Initial activation step at 94°C for 5 min, followed by 32 cycles consisting of denaturing at 94°C for 1 min, annealing at 60°C for 1 min, extension at 72°C for 2 min and final elongation at 72°C for 10 min	[32, 33]
<i>intI2</i> (Class 2 integron)	Int2-F Int2-R	CACGGATATGCGACAAAAAGGT GTAGCAAACGAGTGACGAAATG	788		
<i>intI3</i> (Class 3 integron)	Int3-F Int3-R	AGTGGGTGGCGAATGAGTG TGTTCTTGTATCGGCAGGTG	600		

Results and Discussion

It is worthy of note the amaranths irrigated with wastewater effluent were of higher yields compared to the controls irrigated with tap water. This is attributable to the fact the effluent is rich in nutrients. It also underscores the continued preference of effluents over the scarce treated water by the vegetable farmers. Despite a perceived understanding farmer have on the use of unsafe WWTP effluent, this knowledge seems not bother them.

In the present study, the total number of samples collected were 288 including WWTP effluents (60), greenhouse (60), farm (84) and market (84) vegetables. A total of 178 *E. coli* isolates were confirmed by PCR amplification of the β -glucuronidase *uidA* gene (Fig 1), with 41 isolates from the WWTP effluents, 46 greenhouse from the effluent irrigated vegetables, 55 from vegetables collected from gardens that produce vegetables sold in local markets and 36 from vegetables bought from selected markets in Nsukka and Enugu (Table 2).

Fig 1. PCR products for *E. coli* confirmation by *uidA* gene amplification.

This is the Fig 1 legend: M: Molecular weight marker (1 KB), W: water, NC: Negative control (*Enterobacter aerogenes*), PC: Positive control (*E. coli*; NCTC 13353), Lanes 1-15: Positive isolates

Table 2: Isolation of *E. coli* in effluent wastewater and vegetable samples.

Sample type	No. of samples	No. of <i>E. coli</i> isolates
WWTP effluent	60	41
Greenhouse vegetables	60	46
Farm vegetables	84	55
Market Vegetables	84	36
Total	288	178

Table 3 shows the antibiotics susceptibility profiles of the 178 *E. coli* isolates tested with 18 different antibiotics. Generally, higher resistance percentages were observed in *E. coli* from market vegetables compared with others. This could be attributable to further contamination of vegetables by clinical *E. coli* strains arising directly from handling by sellers. All the *E. coli* isolates, showed susceptibility to imipenem and only 5.6% (10/178) of all the isolates were resistant to cefuroxime (a cephalosporin). Chloramphenicol, ciprofloxacin and norfloxacin were very effective. The most significant resistance phenotypes were detected among sulphamethoxazole (58.4%), amoxicillin (52.8%), tetracycline (47.2%), trimethoprim (44.9%) and streptomycin (37.1%), as these antibiotics are commonly used in the studied communities (Table 2).

260 **Table 3: Antibiotics susceptibility profile of *E. coli* isolates**

S/N	Antimicrobial Agent (disc concentration)	Class (Subclasses)	Code	Number of isolates resistant (%)				
				Effluent (n = 41)	Greenhouse (n = 46)	Farm (n = 55)	Market (n = 36)	TOTAL (n = 178)
1	Cloxacillin (5µg)	β-Lactam (Penicillins)	COX	41 (100)	46 (100)	55 (100)	36 (100)	178 (100)
2	Metronidazole (50µg)	Nitroimidazoles	MTZ	41 (100)	46 (100)	55 (100)	36 (100)	178 (100)
3	Vancomycin (30µg)	Glycopeptides	VAN	41 (100)	46 (100)	53 (96.4)	36 (100)	176 (98.9)
4	Rifampicin (5µg)	Rifamycins	RIF	41 (100)	45 (97.8)	52 (94.6)	34 (94.4)	172 (96.6)
5	Penicillin (10U)	β-Lactam (Penicillins)	PNG	39 (95.1)	45 (97.8)	54 (98.2)	34 (94.4)	172 (96.6)
6	Erythromycin (15µg)	Macrolides	ERT	40 (97.6)	46 (100)	50 (90.9)	32 (88.9)	168 (94.4)
7	Clarithromycin (15µg)	Macrolides	CLR	40 (97.6)	37 (80.4)	47 (85.5)	36 (100)	160 (89.9)
8	Ampicillin (5µg)	β-Lactam (Cephalosporins)	AMP	34 (82.9)	38 (82.6)	50 (90.9)	36 (100)	158 (88.7)
9	Sulphamethoxazole (25µg)	Sulphonamides	SMZ	23 (56.1)	27 (58.7)	26 (47.3)	28 (77.8)	104 (58.4)
10	Amoxicillin (10µg)	β-Lactam (Penicillins)	AMX	23 (56.1)	24 (52.2)	26 (47.3)	21 (58.3)	94 (52.8)
11	Tetracycline (30µg)	Tetracyclines	TET	25 (61.0)	20 (43.5)	16 (29.1)	23 (63.9)	84 (47.2)
12	Trimethoprim (5µg)	Dihydrofolate Reductase (DHFR) inhibitors	TMP	27 (65.9)	20 (43.5)	12 (21.8)	21 (58.3)	80 (44.9)
13	Streptomycin (10µg)	Aminoglycosides	STR	15 (36.6)	21 (45.7)	16 (29.1)	14 (38.9)	66 (37.1)
14	Chloramphenicol (30µg)	Phenicol	CHL	4 (9.8)	6 (13.0)	6 (10.9)	9 (25.0)	25 (14.0)
15	Ciprofloxacin (5µg)	Fluoroquinolones	CIP	5 (12.2)	7 (15.2)	6 (10.9)	7 (19.4)	25 (14.0)
16	Norfloxacin (10µg)	Fluoroquinolones	NOR	2 (4.9)	7 (15.2)	9 (16.4)	5 (13.9)	23 (12.9)
17	Cefuroxime (30µg)	β-Lactam (Cephalosporins)	CXM	2 (4.9)	2 (4.4)	2 (3.6)	4 (11.1)	10 (5.6)
18	Imipenem (10µg)	β-Lactam (Carbapenems)	IMP	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

261

262

263

Multidrug resistance (MDR) has been frequently reported in Nigeria among *E. coli* isolates obtained from human specimens [12, 37, 42, 43], animal sources [36] and environmental samples [38, 42, 44, 45]. In the present study, all the isolates were multidrug resistant, ranging from 5-drug to 16-drug resistance patterns. Although some studies have reported a high removal efficiency for total ARGs in wastewater [46], our data suggest that sewage treatment process at UNN is not effective in reducing ARGs as all the *E. coli* isolated from the effluent were MDR. The spread of AMR often limits the availability of therapeutic options to only a very few efficacious antibiotics [47]. The last-resort drugs, the carbapenems such as imipenem (used in this study) and meropenem, are themselves not only increasingly challenged by emerging resistance, as evident from the data presented here, but are not affordable in the developing regions.

Multidrug resistance (MDR) was detected in all *E. coli* isolates, and although this study did not determine the full virulence potentials of all the isolates subjected to antimicrobial susceptibility testing (AST), irrigational use of WWTP effluent represents a pathway for human exposure to antibiotic-resistant commensal and pathogenic bacteria. Vegetable farming at the site should therefore be discontinued as it presents significant threat to the health of consumers of such vegetables.

It is known that AMR and MDR in *E. coli* are acquired by the transfer of mobile genetic elements, such as plasmids, transposons and integrons [15, 18, 20]. In the present study, of the 178 *E. coli* isolates, class 1 integrons were detected (Fig 2) in 175 (98.3%), and class 2 in 5 (2.8%). All the class 2 integrons were found in isolates that were positive for class 1. Such co-carriage has been previously published on *E. coli* from meat turkeys in Italy, [34].

Fig 2. Multiplex PCR products for detection of class 1 and 2 integrons

This is the Fig 2 legend: M: Molecular weight marker (1 KB Plus ladder), W: water, Lanes 1-15: *E. coli* isolates

The integron carriage rate for the 137 vegetable isolates was 97.8%, whereas the rate for 41 effluent isolates was 100%. Considering that all the isolates were MDR, the detected high

prevalence of class 1 integron is not surprising and compares with a previous study that reported that MDR phenotypes were observed in 96.8% of the integron-positive isolates [35]. These rates portend serious public health risks as it is known that class 1 integron could carry diverse antibiotic resistance genes (ARGs) and conduct horizontal gene transfer among microorganisms [20].

The data presented here shows that class 2 integrons were less frequently detected 5 (2.8%). Similar data have been published for Enterobacteriaceae in Nigeria [36, 38] and elsewhere [34, 39, 40]. Ramírez et al [39], reported that unlike the widespread distribution of class 1 integron within Gram-negative bacilli, only *Acinetobacter baumannii* and *Enterobacter cloacae* harboured class 2 integrons at a high frequency. However, in an earlier study in China [41], Class 2 integrons were present in 25 (80.6%) of the *Shigella sonnei* isolates and 29 (87.9%) of the *S. flexneri* isolates whereas class 1 integrons were found in only 6 (9.4%) of *Shigella* spp. isolates.

Conclusions

The present study revealed high detection of *E. coli* in the studied effluent and vegetable samples and represent potential public health hazards intensified by observed multidrug resistance in all the isolates and the high occurrence of class 1 integrons. It is concluded that UNN-WWTP is a significant reservoir for diarrheagenic *E. coli*. Vegetable farming at the site should therefore be discouraged as it presents significant threat to the health of consumers of such vegetables.

Acknowledgments

The authors are also grateful to Mr. Uchenna Oyeagu of the Water and Public Health Laboratory, University of Nigeria, Nsukka, Nigeria, for laboratory assistance.

References

1. Stacy-Phipps S, Mecca JJ, Weiss JB (1995) Multiplex PCR Assay and Simple Preparation Method for Stool Specimens Detect Enterotoxigenic *Escherichia coli* DNA during Course of Infection. J Clin Microbiol 33: 1054–1059.
2. Okeke IN (2009) Diarrheagenic *Escherichia coli* in sub-Saharan Africa: status, uncertainties and necessities. J Infect Dev Ctries 3: 817–842.
3. Brown JW, Badahdah A, Iticovici M, Vickers TJ, Alvarado DM, Helmerhorst EJ, Oppenheim FG, Mills JC, Ciorba MA, Fleckenstein JM. Bullitt E (2018) A Role for

- Salivary Peptides in the Innate Defense Against Enterotoxigenic *Escherichia coli* J Infect Dis. 217:1435–1441
4. Nzeako B, Okafor N (2002) Bacterial enteropathogens and factors associated with seasonal episodes of gastroenteritis in Nsukka, Nigeria. Br J Biomed Sci 59: 76–79.
5. WHO/UNICEF [World Health Organization and the United Nations Children’s Fund] (2017). Progress on drinking water, sanitation and hygiene: 2017 update and SDG baselines. Geneva; 2017. Licence: CC BY-NC-SA 3.0 IGO.
6. FAO/WHO [Food and Agriculture Organization of the United Nations and World Health Organization]. (2008). Microbiological hazards in fresh leafy vegetables and herbs: Meeting Report. Microbiological Risk Assessment Series No. 14. Rome. 151pp.
7. Ogunsanya TL, Rotimi VO, Adenuga A (1994) Study of the aetiological agents of childhood diarrhoea in Lagos Nigeria. J Med Microbiol 40: 10–14.
8. Okeke IN, Fayinka ST, Lamikanra A (2000) Antibiotic Resistance in *Escherichia coli* from Nigerian Students, 1986–1998. Emerg Infect Dis 6: 393–396.
9. Olorunshola ID, Smith SI, Coker AO (2000) Prevalence of EHEC O157: H7 in patients with diarrhea in Lagos, Nigeria APMIS 108: 761–763.
10. Okeke IN, Steinruck H, Kanack KJ, Elliott SJ, Sundstrom L, Kaper JB, Lamikanra A (2002) Antibiotic-resistant cell-detaching *Escherichia coli* strains from Nigerian children. J Clin Microbiol 40: 301–305.
11. Chigor VN, Umoh VJ, Smith SI, Igbinsola EO, Okoh AI (2010) Multidrug resistance and plasmid patterns of *Escherichia coli* O157 and other *E. coli* isolated from diarrhoeal stools and surface waters from some selected sources in Zaria, Nigeria. Int J Environ Res Public Health 7: 3831–3841.
12. Onanuga A, Igbeneghu O, Lamikanra A (2014) A study of the prevalence of diarrhoeagenic *Escherichia coli* in children from Gwagwalada, Federal Capital Territory, Nigeria. Pan Afr Med J 17: 146.
13. Nweze EI (2010) Aetiology of Diarrhoea and Virulence Properties of Diarrhoeagenic *Escherichia coli* among Patients and Healthy Subjects in Southeast Nigeria. J Health Popul Nutr 28:245–252.
14. Okeke IN, Edelman R (2001) Dissemination of antibiotic-resistant bacteria across geographic borders. CID 33: 363–369.
15. Gaze WH, Zhang L, Abdouslam NA, Hawkey PM, Calvo-Bado L, Royle J, Brown H, Davis S, Kay P, Boxall ABA, Wellington EMH (2011) Impacts of anthropogenic activity on the ecology of class 1 integrons and integron-associated genes in the environment. The ISME J 5: 1253–1261.
16. Wellington EMH, Boxall ABA, Cross P, Feil EJ, Gaze WH, Hawkey PM, Johnson-Rollings AS, Jones DL, Lee NM, Otten W, Thomas CM, Williams AP (2013) The role of the natural environment in the emergence of antibiotic resistance in Gram-negative bacteria. Lancet Infect Dis 13: 155–165.
17. Amos GCA, Zhang L, Hawkey PM, Gaze WH, Wellington EM (2014) Functional metagenomic analysis reveals rivers are a reservoir for diverse antibiotic resistance genes. Vet Microbiol 171: 441–447.
18. Gillings MR (2014) Integrons: Past, Present, and Future. Microbiol Mol Biol Rev 78: 257–277.

19. Strugeon E, Tilloy V, Ploy MC, Da Re S (2016) The stringent response promotes antibiotic resistance dissemination by regulating integron integrase expression in biofilms. *mBio* 7(4):e00868–16.
20. Ma L, Li AD, Yin XL, Zhang T (2017) The Prevalence of Integrons as the Carrier of Antibiotic Resistance Genes in Natural and Man-Made Environments. *Environ Sci Technol* 51: 5721–5728.
21. Asadu A, Enwelu I, Ifejika P, Igbokwe E (2016) Urban crop production in southeast Nigeria: Potentials and constraints. *Afr J Agric Res* 11: 4646–4653.
22. American Public Health Association (APHA) (2005). Standard Methods for the Examination of Water and Wastewater. 21st Edn. American Public Health Association, American Water Works Association, Water Environment Federation, Washington DC, USA.
23. Falomir MP, Gozalbo D, Rico H (2010) Coliform bacteria in fresh vegetables: from cultivated lands to consumers. In: A. Mandez-Vilas (Ed.). Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology. FORMATEX; Badajoz Spain, pp. 1175–1181.
24. Saeed AY, Mazin H, Saadi A, Hussein SO (2013) Detection of *Escherichia coli* O157 in vegetables. *IOSR J Agric Vet Sci* 6: 16–18.
25. Claus D (1992) A standardized Gram staining procedure. *World J Microbiol Biotechnol* 8:451–452.
26. Bauer A, Kirby W, Sherris J, Turck M (1966) Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 45:493–496.
27. CLSI (2012) Performance Standards for Antimicrobial Disk Susceptibility Tests; Approved Standard—Eleventh Edition. Vol 32, No.1.
28. Vankerckhoven V, Van Autgaerden T, Vael C, Lammens C, Chapelle S, Rossi R, Jabes D, Goossens H (2004) Development of a Multiplex PCR for the Detection of *asa1*, *gelE*, *cylA*, *esp*, and *hyl* Genes in Enterococci and Survey for Virulence Determinants among European Hospital Isolates of *Enterococcus faecium*. *J Clin Microbiol* 42: 4473–4479.
29. Bej AK, Dicesare JL, Haff L, Atlas RM (1991) Detection of *Escherichia coli* and *Shigella* spp. in Water by Using the Polymerase Chain Reaction and Gene Probes for *uid*. *Appl Environ Microbiol* 57: 1013–1017.
30. Horakova K, Mlejnkova H, Mlejnek P (2008) Specific detection of *Escherichia coli* isolated from water samples using polymerase chain reaction targeting four genes: cytochrome bd complex, lactose permease, b-D-glucuronidase, and b-D-galactosidase. *J Appl Microbiol* 105: 970–976.
31. Lopez-Saucedo C, Cerna JF, Villegas-Sepulveda N, Thompson R, Velazquez FR, Torres J, Tarr PI, Estrada-García T (2003) Single multiplex polymerase chain reaction to detect diverse loci associated with diarrhoeagenic *Escherichia coli*. *Emerg Infect Dis* 9: 127–131.
32. Machado E, Canton R, Baquero F, Galan JC, Rollan A, Peixe L, Coque TM (2005) Integron Content of Extended-Spectrum--Lactamase-Producing *Escherichia coli* Strains over 12 Years in a Single Hospital in Madrid, Spain. *Antimicrob Agents Chemother* 49: 1823–1829.
33. Kargar M, Mohammadalipour Z, Doosti A, Lorzadeh S, Japoni-Nejad A (2014) High Prevalence of Class 1 to 3 Integrons Among Multidrug-Resistant Diarrheagenic *Escherichia coli* in Southwest of Iran. *Osong Public Health Res Perspect* 5: 193–198.

34. Piccirillo A, Giovanardi D, Dotto G, Grilli G, Montesissa, C, Boldrin C, Salata C, Giacomelli M (2014) Antimicrobial resistance and class 1 and 2 integrons in *Escherichia coli* from meat turkeys in Northern Italy. *Avian Pathol* 43: 396–405.
35. Vasilakopoulou A, Psychogiou M, Tzouvelekis L, Tassios PT, Kosmidis C, Petrikos G, Roma ES, Charvalos E, Passiotou M, Avlami A, Daikos GL (2009) Prevalence and characterization of class 1 integrons in *Escherichia coli* of poultry and human origin. *Foodborne Pathog Dis* 6: 1211–1218.
36. Chah KF, Agbo IC, Eze DC, Somalo S, Estepa V, Torres C (2010) Antimicrobial resistance, integrons and plasmid replicon typing in multiresistant clinical *Escherichia coli* strains from Enugu State, Nigeria. *J Basic Microbiol* 50: S18–S24.
37. Odetoyin BW, Labar AS, Lamikanra A, Aboderin AO, Okeke IN (2017) Class 1 and 2 integrons in faecal *Escherichia coli* strains isolated from mother-child pairs in Nigeria. *PLoS ONE* 12: e0183383.
38. Adelowo OO, Helbig T, Knecht C, Reincke F, Mäusezahl I, Müller JA (2018) High abundances of class 1 integrase and sulfonamide resistance genes, and characterisation of class 1 integron gene cassettes in four urban wetlands in Nigeria. *PLoS ONE* 13: e0208269.
39. Ramírez MS, Pineiro S, Argentinian Integron Study Group, Centron D (2010) Novel Insights about Class 2 Integrons from Experimental and Genomic Epidemiology. *Antimicrob Agents Chemother* 54: 699–706.
40. Malek MM, Amer FA, Allam AA, El-Sokkary RH, Gheith T, Arafa MA (2015) Occurrence of class I and II integrons in Enterobacteriaceae collected from Zagazig University Hospitals, Egypt. *Front Microbiol* 6: 601.
41. Pan JC, Ye R, Meng DM, Zhang W, Wang HO, Liu KZ (2006) Molecular characteristics of class 1 and class 2 integrons and their relationships to antibiotic resistance in clinical isolates of *Shigella sonnei* and *Shigella flexneri*. *J Antimicrob Chemother* 58: 288–296.
42. Chigor VN, Umoh VJ, Smith SI, Igbinosa EO, Okoh AI (2010) Multidrug resistance and plasmid patterns of *Escherichia coli* O157 and other *E. coli* isolated from diarrhoeal stools and surface waters from some selected sources in Zaria, Nigeria. *Int J Environ Res Public Health* 7: 3831–3841.
43. Okeke IN, Lamikanra A, Czeaczulin J, Dubovsky F, Kaper JB, Nataro JP (2000b) Heterogeneous Virulence of Enterohemorrhagic *Escherichia coli* Strains Isolated from Children in Southwest Nigeria. *J Infect Dis* 181: 252–260.
44. Smith SI, Bello OS, Goodluck HA, Omonigbehin EA, Agbogu VN, Odeigah P (2009) Prevalence of EHEC O157:H7 from human and environmental samples from Lagos and Zaria. *Pakistan J Med Sci* 25: 398–403.
45. Titilawo Y, Sibanda T, Obi L, Okoh A (2015a) Multiple antibiotic resistance indexing of *Escherichia coli* to identify high-risk sources of faecal contamination of water. *Environ Sci Pollut Res* 22: 10969–10989.
46. Yang Y, Li B, Zou S, Fang HHP, Zhang T (2014) Fate of antibiotic resistance genes in sewage treatment plant revealed by metagenomic approach. *Water Res* 62: 97–106.
47. Juhas M, Widlake E, Teo J, Huseby DL, Tyrrell JM, Polikanov YS, Ercan O, Petersson A, Cao S, Aboklaish AF, Rominski A, Crich D, Bottger EC, Walsh TR, Hughes D, Hobbie SN (2019) In vitro activity of apramycin against multidrug-, carbapenem- and aminoglycoside-resistant Enterobacteriaceae and *Acinetobacter baumannii*. *J Antimicrob Chemother* 74: 944–952.

Figure 1

Figure 2