Functional redundancy and crosstalk between flavodiiron proteins and NDH-1 in *Synechocystis* sp. PCC 6803

Short title: Crosstalk between flavodiiron proteins and NDH-1

Lauri Nikkanen¹, Anita Santana Sánchez¹, Maria Ermakova^{1*}, Matthias Rögner², Laurent Cournac³, and Yagut Allahverdiyeva^{1#}

Affiliations:

¹Molecular Plant Biology, Department of Biochemistry, University of Turku, Turku, Finland

²Plant Biochemistry, Faculty of Biology & Biotechnology, Ruhr-University Bochum, Bochum, Germany

³Eco&Sols, University of Montpellier, IRD, CIRAD, INRA, Montpellier Supagro, Montpellier, France

One sentence summary: Flavodiiron proteins and NDH-1 complex ensure survival of cyanobacterial cells by cooperatively safeguarding the photosynthetic apparatus against excessive reduction

Author contributions: Y.A., and L.N. conceptualized the research; L.N., A.S.S., M.E., and Y.A. designed the research; M.E. and A.S.S. conducted growth experiments; L.N. performed MIMS and spectrophotometric experiments; A.S.S. performed immunoblotting; all authors analyzed the data; L.N. and Y.A. wrote the article with the input from A.S.S and M.E.

Funding: This work was supported by the Academy of Finland (project #315119 to Y.A. and the Finnish Center of Excellence, project #307335), the Nordforsk Nordic Center of Excellence ‘NordAqua’ (#82845) and SFB480- Germany

***Current affiliation:** Australian Research Council Centre of Excellence for Translational Photosynthesis, Division of Plant Science, Research School of Biology, The Australian National University, Acton, Australia

#Author for contact: Yagut Allahverdiyeva, allahve@utu.fi

ABSTRACT

In oxygenic photosynthetic organisms excluding angiosperms, flavodiiron proteins (FDPs) catalyze light-dependent reduction of O₂ to H₂O. This alleviates electron pressure on the photosynthetic apparatus and protects it from photodamage. In *Synechocystis* sp. PCC 6803, four FDP isoforms function as hetero-oligomers of Flv1 and Flv3 and/or Flv2 and Flv4. An alternative electron transport pathway mediated by the NAD(P)H dehydrogenase-like complex (NDH-1) also contributes to redox hemostasis and the photoprotection of photosynthesis. Four NDH-1 types have been characterized in cyanobacteria: NDH-1₁ and NDH-1₂, which function in respiration; and NDH-1₃ and NDH-1₄, which function in CO₂ uptake. All four types are involved in cyclic electron transport. Along with single FDP mutants ($\Delta flv1$ and $\Delta flv3$) and the double NDH-1 mutants ($\Delta d1d2$, which is deficient in NDH-1_{1,2} and $\Delta d3d4$, which is deficient in NDH-1_{3,4}), we studied triple mutants lacking either one of Flv1 or Flv3, and NDH-1_{1,2} or NDH-1_{3,4}. We show that the presence of either Flv1/3 or NDH-1_{1,2}, but not NDH-1_{3,4}, is indispensable for survival during changes in growth conditions from high CO₂ /moderate light to low CO₂ / high light. Our results suggest functional redundancy and crosstalk between FDPs and NDH-1_{1,2} under the studied conditions, and demonstrate that the functions of FDPs and NDH-1_{1,2} are dynamically coordinated for the efficient oxidation of PSI and for photoprotection under variable CO₂ and light availability.

KEYWORDS: *Alternative electron transfer, Flavodiiron proteins, Flv, Mehler-like reaction, NDH-1, photosynthesis, photoprotection, Synechocystis.*

INTRODUCTION

Photosynthetic organisms have evolved a variety of different regulatory mechanisms which are important for the protection of the photosynthetic machinery during rapid changes in environmental conditions. Fluctuating light intensities present particular risk to the photosystems due to over-reduction of the photosynthetic electron transport chain (PETC), particularly Photosystem (PS) I (Allahverdiyeva et al., 2015; Tiwari et al., 2016; Shimakawa et al., 2016). To counter this, cyanobacteria, algae, and plants (excluding angiosperms) employ C-type flavodiiron proteins (FDPs) as a strong photoprotective electron sink, directing excess photosynthetic electrons from

46 downstream of PSI to O₂ (Helman et al. 2003, Allahverdiyeva et al., 2013; Gerotto et al., 2016; Ilik
47 et al., 2017; Chaux et al., 2017; Jokel et al., 2018; Alboresi et al., 2019). This process is referred to
48 as the Mehler-like reaction and, in contrast to the Mehler reaction (Mehler 1957), does not produce
49 harmful reactive oxygen species (ROS) (Vicente et al., 2002; Brown et al., 2019).

50 The genome of the β -cyanobacterium *Synechocystis* sp. PCC 6803 (hereafter *Synechocystis*)
51 encodes four isoforms of FDPs, namely Flv1-4, which function in O₂ photoreduction mainly as
52 hetero-oligomers consisting of either Flv1 and Flv3 or Flv2 and Flv4 (Zhang et al., 2004; Mustila et
53 al., 2016; Santana-Sanchez et al., 2019). Recently, we showed that under air level [CO₂], Flv1/3 and
54 Flv2/4 hetero-oligomers function in a coordinated and interdependent manner: Flv1/3 stimulates
55 strong, but transient O₂ photoreduction upon the onset of illumination or during increases in light
56 intensity, while Flv2/4 hetero-oligomers catalyze a slower and limited steady state reduction of O₂,
57 also using electrons downstream of PSI (Santana-Sanchez et al., 2019). In elevated [CO₂] and in air
58 [CO₂] at alkaline pH 9, the *flv4-flv2* operon encoding Flv2, Flv4, and the Sll0218 proteins is
59 downregulated (Zhang et al., 2009; Santana-Sanchez et al., 2019). Nevertheless, a low level of
60 Flv1/3 in elevated [CO₂] stimulates strong steady state O₂ photoreduction, whereas in air [CO₂] at
61 pH 9, Flv1/3 is solely responsible for strong but transient O₂ photoreduction (Santana-Sanchez et
62 al., 2019). Whilst *in vitro* assays showed that homo-oligomers of recombinant Flv1, Flv3 and Flv4
63 can reduce O₂ with NADH and/or NADPH (Vicente et al. 2002, Shimakawa et al. 2015, Brown et
64 al. 2019), FNR and reduced Fd are yet to be considered as possible donors to FDPs. Moreover, the
65 use of *Synechocystis* mutants solely overexpressing Flv1 or Flv3 clearly demonstrated that, in
66 contrast to *in vitro* experiments, homo-oligomers of Flv3 or Flv1 are not involved in O₂
67 photoreduction *in vivo* (Mustila et al. 2016). Thus, the electron donor to FDPs remains to be
68 elucidated *in vivo*.

69 In cyanobacteria, photosynthetic and respiratory electron transfer reactions coexist on thylakoid
70 membranes and are intricately linked, necessitating strict regulation (Mullineaux, 2014). The
71 NAD(P)H dehydrogenase-like complex 1 (NDH-1, photosynthetic complex I) is mainly localized in
72 thylakoid membranes and involved in several bioenergetic reactions including cyclic electron
73 transfer around PSI (CET), respiration and CO₂ acquisition via the carbon concentrating mechanism
74 (CCM) (for reviews see Battchikova et al. 2011; Ma and Ogawa 2015; Burnap et al. 2015; Peltier et
75 al. 2016). The NDH-1 complex is present as several forms with distinct physiological roles: the
76 NDH-1₁ type features the subunits NdhD1 and NdhF1 and functions as Complex I of the respiratory
77 electron transfer chain (Zhang et al., 2004; He et al., 2015; Saura and Kaila, 2019); the NDH-1₂

form features the NdhD2 subunit instead of NdhD1, but it remains unclear whether NDH-1₁ and NDH-1₂ serve different physiological functions (Peltier et al., 2016). The NDH-1₃ and NDH-1₄ forms that comprise low-Ci-inducible, high affinity CO₂ acquisition subunits NdhD3, NdhF3, CupA, and CupS, or constitutive, low-affinity CO₂ acquisition subunits NdhD4, NdhF4, and CupB, respectively, function in conversion of CO₂ to HCO₃⁻ (Ohkawa et al., 2000a,b; Shibata et al., 2001; Zhang et al., 2004; Burnap et al., 2013; Han et al., 2017; Schuller et al., 2019). This process is possibly enabled by alkaline pockets at the thylakoid membrane created by NDH-1 proton pumping (Kaplan and Reinhold, 1999). All four NDH-1 types catalyze CET from ferredoxin (Fd) to plastoquinone (PQ), which is coupled to the pumping of 4H⁺/2e⁻ into the thylakoid lumen (He et al., 2015; Laughlin et al., 2019; Schuller et al., 2019; Saura and Kaila, 2019). However, for NDH-1_{3,4} this is to a lesser extent than for NDH-1_{1,2} (Bernat et al., 2011).

In the current study, we aim to elucidate how the electron transport pathways mediated by FDPs and NDH-1 cooperate to allow the maintenance of redox poise between the PETC, respiration, CCM, and CO₂ fixation in the Calvin-Benson-Bassham cycle (CBB) under variable light and C_i availability. To this end, we employed both biophysical and biochemical methods to characterize various *Synechocystis* mutant strains with combined deficiencies of both FDP and NDH-1 pathways. Our results provide convincing evidence for the presence of either Flv1/3 or NDH-1_{1,2}, but not NDH-1_{3,4}, being indispensable for the survival of *Synechocystis* cells under transitions from high [CO₂] conditions to the combined stress conditions of air [CO₂] and high light. We show that a dynamically coordinated and cooperative function of Flv1/3 and NDH-1_{1,2} is required for the photoprotection of the photosynthetic apparatus of *Synechocystis* cells and discuss the molecular mechanisms involved.

RESULTS

Simultaneous inactivation of Flv1/3 and NDH-1_{1,2} is lethal upon shift from high to air [CO₂] and high light

In order to examine whether the simultaneous inactivation of NDH-1 and FDPs has an adverse effect on cell survival, we monitored the growth of *Synechocystis* wild-type (WT) and various mutant strains lacking (i) either Flv1 or Flv3 ($\Delta flv1$ and $\Delta flv3$, respectively, Helman et al., 2003), (ii) both NdhD1 and NdhD2 ($\Delta ndh12$, deficient in NDH-1₁ and NDH-1₂, Ohkawa et al., 2000), (iii) both NdhD3

and NdhD4 ($\Delta d3d4$, deficient in NDH-1₃ and NDH-1₄, Ohkawa et al., 2000), as well as triple mutants with combined deficiencies of both pathways ($\Delta flv1 d1d2$, $\Delta flv3 d1d2$, $\Delta flv1 d3d4$, $\Delta flv3 d3d4$) in conditions of differing CO₂ availability and light intensity. The triple mutants exhibited similar or slightly slower growth compared to the WT and other mutant strains under high [CO₂] (3% CO₂) and moderate light (ML, intensity of 50 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ PAR) (Fig. 1A). We also detected no substantial difference in growth when the 3% CO₂/ML- pre-grown cells were diluted (OD₇₅₀=0.1) and shifted to air [CO₂] conditions under constant light intensity (Fig. 1B, Fig. S1A, Fig. S1C). In contrast, when cultures pre-grown in 3% [CO₂]/ML were diluted (OD₇₅₀=0.1) and subjected simultaneously to air [CO₂] and high light stress conditions (HL, intensity of 220 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$), the growth of both $\Delta flv3 d1d2$ and $\Delta flv1 d1d2$ strains was completely inhibited (Fig. 1D). Importantly, $\Delta flv1$, $\Delta flv3$ and $\Delta d1d2$ strains demonstrated growth similar to WT cells under air [CO₂] /HL conditions.

We next examined whether pre-adaptation to air [CO₂] would rescue the growth arrest. When cells previously adapted to air [CO₂] /ML were diluted (OD₇₅₀=0.1) and shifted to air [CO₂] /HL, no differences in growth patterns between the strains were detected (Fig. 1C). Importantly, the triple mutants lacking either Flv1 or Flv3 and NDH-1_{3,4} ($\Delta flv1 d3d4$ and $\Delta flv3 d3d4$ strains) did not show a lethal phenotype after a shift from 3% [CO₂]/ML to air [CO₂]/HL, although a somewhat reduced growth was observed compared to the control strains (Fig. S1B, Fig. S1D). This suggests that, under the studied conditions, crosstalk between Flv1/3 and NDH-1_{3,4} complexes is not as crucial as with NDH-1_{1,2}. Our attempts to create M55/ $\Delta flv1$ and M55/ $\Delta flv3$ double mutants lacking either Flv1 or Flv3 and the central membrane component of the NDH-1 complex, NdhB, were both unsuccessful. We were unable to obtain any M55/ $\Delta flv1$ colonies and the M55/ $\Delta flv3$ mutant strain demonstrated poor growth and was not segregated, suggesting that the absence of both Flv1/3 and NDH-1, disrupts essential cell metabolism.

We also studied the growth of the mutant strains on agar plates containing BG-11 (Fig. 1E). For these experiments, 3% CO₂/ML-grown cells were diluted and grown on plates either under air [CO₂] / ML or air [CO₂] / HL conditions for 7 days. The $\Delta flv1$ and $\Delta flv3$ and $\Delta d1d2$ mutants did not exhibit any visible differences in growth capacity under these conditions in comparison to the WT. In contrast, the growth of the $\Delta flv1 d1d2$ and $\Delta flv3 d1d2$ triple mutants was strongly reduced under air [CO₂] / ML, while no growth was detected under air [CO₂] / HL (Fig. 1E). The $\Delta flv1 d3d4$ and $\Delta flv3 d3d4$

138 triple mutants demonstrated a slow growth phenotype, similarly to that previously reported for $\Delta d3d4$
139 on agar plates (Ohkawa et al. 2000b).

140 As the growth phenotypes of $\Delta flv1$ and $\Delta flv3$, $\Delta flv1 d1d2$ and $\Delta flv3 d1d2$ were similar under all
141 conditions, we only included $\Delta flv3$ and $\Delta flv3 d1d2$ in the majority of subsequent experiments.

Figure 1. Growth of WT and mutant strains under different experimental conditions. All experimental cultures were started from pre-cultures adjusted to $OD_{750}=0.1$. **(A)** Cells were pre-cultivated under 3% $[CO_2]$ / ML (50 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$), adjusted to $OD_{750}=0.1$ and grown in the same conditions. **(B)** Cells were pre-cultivated under 3% $[CO_2]$ /ML and shifted to air $[CO_2]$ /ML. **(C)** Cells were pre-cultivated under air $[CO_2]$ /ML and shifted to air $[CO_2]$ /HL 220 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$. **(D)** Cells were pre-cultivated under 3% $[CO_2]$ /ML and shifted to air $[CO_2]$ /HL. The values presented in A–D are means of three independent biological replicates $\pm$ SE. **(E)** Growth of WT and mutant cells on agar plates containing BG-11. Cells grown under 3% $[CO_2]$ / ML were harvested and spotted on BG-11 agar plates in ten-fold dilution series starting at $OD_{750}=1$, incubation was under air $[CO_2]$ /ML and air $[CO_2]$ /HL for 7 days. See also Supplemental Fig. S1.

Either Flv1/3 or NDH-1_{1,2} is required for the oxidation of PSI during sudden increases in light intensity

The lethal phenotype observed upon the shift to air $[CO_2]$ / HL which was caused by the simultaneous impairment of the Mehler-like reaction catalyzed by Flv1/3 (Helman et al. 2003, Allahverdiyeva et al. 2011, 2013) and by NDH-1_{1,2} complex mediated respiration and CET (Ohkawa et al., 2000b, Bernat et al. 2011) suggests a functional redundancy between these electron transport pathways. To further investigate this possibility, we used membrane inlet mass spectrometry (MIMS) to examine the kinetics of O_2 and CO_2 exchange upon HL-illumination of dark-adapted WT and mutant cells grown for four days in air $[CO_2]$ / ML. In order to distinguish O_2 uptake from photosynthetic gross O_2 evolution under illumination, we enriched cell suspensions with $^{18}O_2$ prior to measurements. In dark-adapted WT cells, a transient peak in O_2 uptake occurs during the first minute of illumination (Fig. 2A). We showed recently that this transient peak is attributed to the Mehler-like reaction catalyzed predominantly by Flv1/3 hetero-oligomers, while Flv2/4 hetero-oligomers mainly contribute to steady-state light-induced O_2 reduction in an interdependent manner (Santana-Sanchez et al., 2019). In $\Delta flv3$ cells, light-induced O_2 reduction was almost abolished and only a slight impairment of the rate of photosynthetic gross O_2 evolution was observed in comparison to WT cells (Fig. 2B). This result is in agreement with previous studies (Helman et al., 2003; Allahverdiyeva et al., 2013; Santana-Sanchez et al., 2019).

In contrast to the WT, $\Delta lld2$ cells lacked a fast decay phase of light-induced O_2 uptake during the first minute of illumination, demonstrating sustained O_2 photoreduction at high levels (c.a. 75–100 $\mu\text{mol } O_2 \text{ mg Chl}^{-1} \text{ hr}^{-1}$) throughout the illumination period. Meanwhile, gross O_2 evolution was

174 diminished nearly two-fold compared to the WT (Fig. 2C). Thus, sustained O₂ photoreduction in
 175 $\Delta d1d2$ is not due to increased electron flow from PSII, but most likely due to increased activity of
 176 FDPs. Indeed, $\Delta flv3 d1d2$ mutant cells showed only minor light-induced O₂ uptake transiently
 177 during the first minute of illumination (Fig. 2D). The $\Delta flv3 d1d2$ cells demonstrated slower
 178 induction of photosynthetic O₂ evolution following two-component kinetics, and slightly impaired
 179 steady-state gross O₂ evolution compared to WT (Fig. 2D).

180 The initial peak in CO₂ uptake rate at the onset of illumination likely reflects activation of the CCM
 181 (Liran et al., 2018). Accordingly, this initial peak is absent in the M55 mutant deficient in NDH-1_{3,4}
 182 in addition to NDH-1_{1,2}, as well as in WT cells grown at pH 6 (Fig. S2). No salient impairment of
 183 CCM was observed in any of the studied strains under air [CO₂] / ML, with CO₂ uptake rates
 184 peaking at c.a. 1.5 mM CO₂ mg Chl⁻¹ hr⁻¹. In WT as well as in $\Delta flv3$, CO₂ fixation is then induced
 185 and the respiratory compensation point surpassed after ~1 min of illumination (Fig. 2A-B). In
 186 $\Delta d1d2$, however, induction of CO₂ fixation is severely delayed, and the cells only reached the
 187 compensation point at the end of the 5 min illumination period (Fig. 2C). Intriguingly, although a
 188 slight delay was observed also in $\Delta flv3 d1d2$ induction of CO₂ fixation was largely recovered in the
 189 triple mutant (Fig. 2D).

Figure 2. O₂ and CO₂ exchange rates in WT and mutant strains. Cells were grown in air [CO₂] / ML for 4 days, after which the cells were harvested and Chl *a* concentration adjusted to 10 µg ml⁻¹ with fresh BG-11. Cells were dark-adapted for 15 min, and gas exchange was monitored by membrane inlet mass spectrometry (MIMS) over a 5 min illumination period with 500 µmol photons m⁻²s⁻¹ of white actinic light. Prior to the measurements, samples were supplemented with ¹⁸O₂ at an equivalent concentration to ¹⁶O₂ to distinguish O₂ uptake from O₂ evolution, and with 1.5 mM NaHCO₃. The dashed line in each panel indicates the compensation point of CO₂ fixation (uptake rate equals respiratory rate). The experiment was repeated with three independent biological replicates, of which representative measurements are shown.

These MIMS results indicated that combined deficiency of Flv1/3 and NDH-1_{1,2} under standard growth conditions, air [CO₂] / ML, slightly perturbs photosynthetic electron transfer and the redox poise between the PETC and cytosolic sink reactions. Next, we examined the ability of the mutant strains to adjust their photosynthetic activity to variable light conditions by measuring Chl *a* fluorescence simultaneously with P700 redox changes in conditions where light intensity periodically fluctuated between low (25 µmol photons m⁻²s⁻¹, LL) and high irradiance (530 µmol photons m⁻²s⁻¹, HL). Cells lacking Flv3 suffered from transient PSI acceptor side limitation Y(NA) upon sudden increases in light intensity, resulting in delayed oxidation of PSI (Fig. 3A), which is in line with previous studies (Helman et al. 2003, Allahverdiyeva et al. 2013). During subsequent 1 min cycles of LL/HL, however, the ability of $\Delta flv3$ to oxidize PSI in HL improved, suggesting that cells were able to acclimate to the fluctuating light condition via a compensatory mechanism that is distinct from Flv1/3 hetero-oligomers, such as *e.g.* NDH-1 mediated electron transport.

We detected no acceptor-side limitation during the HL phases in $\Delta d1d2$ (Fig. 3A) as PSI was oxidized similarly to the WT (Fig. 3B). Interestingly, however, $\Delta d1d2$ cells exhibited slightly elevated acceptor-side limitation during the LL phases of the experiment (Fig. 3A), suggesting a role for NDH-1_{1,2} in maintaining photosynthetic redox poise in light-limited conditions. In the triple mutant strain $\Delta flv3 d1d2$, transitions from low to high irradiance as well as from dark to light caused severe limitation on the acceptor side of PSI, resulting in an inability to oxidize PSI during periodic 1 min HL illumination. Unlike the $\Delta flv3$ mutant, $\Delta flv3 d1d2$ did not show improvement in the PSI acceptor side limitation during subsequent cycles. This implies a compensating activity of NDH-1 in $\Delta flv3$ mutant under the studied conditions. Interestingly, diminished donor side limitation in $\Delta flv3 d1d2$ resulted in slightly elevated effective yield of PSI during the HL phases (Fig. 3C). This could indicate the attenuation of pH-dependent limitation of electron transfer at Cyt *b₆f*, due to impairment of proton motive force (*pmf*) generation in CET and the Mehler-like reaction. The

223 effective yield of PSII was slightly decreased in HL in $\Delta flv3$ and $\Delta d1d2$, but not in $\Delta flv3 d1d2$ (Fig.
224 3D).

225

226 **Figure 3. Photosynthetic parameters in WT and mutants strains in fluctuating light intensities. (A)**
227 Acceptor side limitation of PSI, Y(NA), (B) donor side limitation of PSI, Y(ND), as well as (C) effective yield
228 of PSI, Y(I) were calculated from absorbance difference changes at 875-830 nm according to (Klughammer
229 and Schreiber, 2008). Chl *a* fluorescence changes were measured simultaneously with PSI redox changes and
230 used to calculate (D) effective yield of PSII. Cells were grown in air [CO₂] / ML for 4 days, after which the
231 cells were harvested and Chl *a* concentration adjusted to 10 μg ml⁻¹ with fresh BG-11. Cells were dark-adapted
232 for 10 min, and then illuminated with red actinic light intensity alternating between 25 and 530 μmol photons
233 m⁻²s⁻¹ in 1 min periods. Saturating pulses (500 ms of 5000 μmol photons m⁻²s⁻¹) were provided every 15
234 seconds. The values shown are averages from 3–4 independent biological replicates ±SE.

235 The differences in photosynthetic electron transport reported above may also be contingent on altered
236 redox states of the NADP⁺/ NADPH pool. Therefore, in order to investigate the effect of Flv3 and/or
237 NDH-1_{1,2} deficiency on NADP⁺/NADPH redox kinetics, we recorded NADPH fluorescence changes
238 from dark-adapted cells simultaneously with Chl *a* fluorescence in conditions where actinic light

intensity fluctuated between 25 (LL) and 530 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ (HL) similarly to Fig. 3. Upon dark-to-LL transitions, NADPH rapidly accumulated close to a maximal amount in all strains. Re-oxidation of NADPH then occurred in WT and $\Delta flv3 dld2$ cells, while in $\Delta flv3$ a slower oxidation phase preceded a transient re-reduction phase during the first minute of illumination (Fig. 4A-B). Strong reduction of the NADP⁺ pool was also detected in $\Delta flv3$ during the second HL phase of the experiment. In $\Delta dld2$ cells, very little oxidation of the NADPH pool occurred during illumination, even at HL-LL transitions (Fig. 4C). Chl *a* fluorescence also remained at an elevated level (Fig. 4C), suggesting a reduced PQ pool. This was possibly due to delayed activation of CO₂ fixation in the CBB cycle (Fig. 2C), and impaired CET or dark respiration. In contrast, NADPH was strongly oxidized in $\Delta flv3 dld2$ cells at HL-LL transitions (Fig. 4D), but was predominantly reduced in dark-adapted cells (Fig. 4D). Upon cessation of illumination, oxidation of the NADPH pool was followed by transient re-reduction after c.a. 10 s. As observed previously (Schreiber and Klughammer, 2009; Holland et al., 2015), the re-reduction peak coincided with a secondary post-illumination rise in Chl *a* fluorescence (PIFR), while oxidation of NADPH paralleled an initial PIFR. The post-illumination re-reduction of NADP⁺ was diminished and both Chl *a* PIFR peaks were missing in $\Delta dld2$ (Fig. 4C). Therefore, the PIFR peaks likely originate from cyclic and respiratory electron transport via NDH-1_{1,2} (Holland et al., 2015). Both Chl PIFR peaks were however observed in $\Delta flv3 dld2$ cells, indicating involvement of a complementary pathway, possibly via succinate dehydrogenase (Cooley and Vermaas, 2001), or a CET pathway dependent on PGR5 and PGRL1-LIKE (Yeremenko et al., 2005; Dann and Leister, 2019). Finally, we monitored NADPH redox kinetics under conditions mimicking the MIMS experiments, where strong O₂ photoreduction occurs in WT and $\Delta dld2$ during the first minute of dark-to-high light transition (Fig. 2). No substantial differences were observed between the strains, as close to maximal reduction of NADP⁺ was obtained within ~0.25 s (Fig. 4F) and maintained throughout a 40 s illumination period (Fig. 4E). This argues against NADPH being a primary electron donor to either Flv1/3 or NDH-1_{1,2}.

Figure 4. NADPH and Chl *a* fluorescence under fluctuating light in WT and mutant strains. WT (A), $\Delta flv3$ (B), $\Delta d1d2$ (C), and $\Delta flv3 d1d2$ cells (D) were grown in air [CO₂] / ML for 4 days, after which the cells were harvested and Chl *a* concentration adjusted to 5 $\mu\text{g ml}^{-1}$ with fresh BG-11. Cells were dark-adapted for 10 min, and then subjected to illumination with red actinic light alternating between 25 and 530 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ in 1 min periods. NADPH redox changes were monitored with the Dual-PAM 100 spectrophotometer and its 9-AA/NADPH accessory module by measuring fluorescence changes between 420 and 580 nm induced by excitation at 365 nm, as described by (Schreiber and Klughammer, 2009) and

(Kauny and Setif, 2014). Chl *a* fluorescence was recorded simultaneously. (E) NADPH redox changes during illumination of dark-adapted cells with 530 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$. Other experimental details are as in A-D. (F) shows a magnification of the first 2 s of illumination in (E). The values are normalized according to minimum and maximum fluorescence changes at the onset and cessation of illumination, respectively. The experiments were repeated with three independent biological replicates, of which representative measurements are shown. r.u. = relative units.

Redox status of PSI donor and acceptor side electron carriers and buildup of *pmf* during dark-to-light transitions depend on both Flv1/3 and NDH-1_{1,2}

To elucidate the molecular mechanism behind the photosynthetic phenotypes of the studied mutant strains, we next utilized a DUAL-Klas-NIR 100 spectrophotometer to distinguish between redox changes of plastocyanin (PC), P700, and ferredoxin (Fd) (Klughammer and Schreiber, 2016; Schreiber, 2017; Setif et al., 2019) upon exposure of dark-adapted cells to high irradiance (503 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$). Upon illumination of WT cells, rapid oxidation of P700 and PC occurred, followed by transient reduction of PC, P700, as well as Fd after ~ 0.2 s (Fig. 5A). Re-oxidation of all three electron carriers then ensued after ~ 0.5 s. In $\Delta flv3$ cells P700 and PC remained mostly reduced after the initial oxidation transient until ~ 3 s, when another transient and partial re-oxidation peak of P700 and PC occurred at ~ 8 s (Fig. 5B). This was followed by re-reduction of P700, similarly to that observed recently by Bulychev et al. (2018). After 15 s in light, P700 and PC became gradually oxidized. In contrast to WT, Fd remained reduced for several seconds in light, and was only slowly re-oxidized over the 30 s illumination period (Fig. 5B). These observations indicate the importance of Flv1/3 as an electron sink to O₂, accepting electrons presumably from reduced Fd (after ~ 0.5 s in light). In $\Delta d1d2$, redox changes of PC, P700 and Fd were similar to WT, except that the reduction of P700⁺ following initial oxidation as well as reduction of Fd occurred already after ~ 50 ms (Fig. 5C). In $\Delta flv3 d1d2$, re-reduction of P700⁺ and reduction of Fd also occurred already after ~ 50 ms (Fig. 5D), suggesting involvement of NDH-1_{1,2} as an acceptor of electrons from Fd at that stage. Fd was reduced more quickly in $\Delta flv3 d1d2$ than in any of the other strains and was slowly re-oxidized over 30 s (Fig. 5D), likely due to shortage of electron acceptors (Fig. 3A). In summary, these NIR-spectroscopic measurements revealed that Flv1/3 and NDH-1_{1,2} control the redox poise between PSI, PC, and primary electron acceptor Fd during specific time frames at transitions to high irradiance. The results are congruent with Fd functioning as the electron donor to both NDH-1_{1,2} and Flv1/3.

303 To inspect whether the phenotypes observed above depend on differential buildup or regulation of
304 *pmf*, we measured *in vivo* changes in the *pmf* by monitoring the absorbance change difference between
305 500-480 nm, which constitutes the electrochromic shift (ECS) in *Synechocystis* (Viola et al., 2019).
306 *In vivo* measurement of light-induced ECS revealed that after 1 s of illumination of dark-adapted WT
307 cells with 500 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$, high *pmf* level was transiently generated, followed by decline
308 during subsequent seconds (Fig. 5E). After ~ 10 s of illumination *pmf* again increased towards a
309 steadier value (Fig. 5E). Congruently with the strong reduction of P700 and Fd (Fig. 5B), the initial
310 *pmf* peak after the first second of illumination was heavily dependent on the presence of Flv1/3. In
311 both $\Delta flv3$ and $\Delta flv3 dld2$ *pmf* and thylakoid proton flux (vH^+) were lower than in WT after 1 s (Fig.
312 5G). In $\Delta flv3$ *pmf* remained drastically lower than in WT during the first seconds of illumination, but
313 differed only slightly from WT thereafter due to diminished conductivity of the thylakoid membrane
314 (gH^+) (Fig. 5F), which is mainly determined by the activity of the ATP synthase (Cruz et al., 2005;
315 Viola et al., 2019). In $\Delta dld2$ thylakoid proton flux was similar to WT (Fig. 5G), likely due to
316 enhanced FDP activity (Fig. 2C) compensating for impaired CET and respiration. However, *pmf*
317 remained lower than in WT throughout the 1 min experiment (Fig. 5E) because of elevated
318 conductivity (Fig. 5F). In contrast, despite drastically lower proton flux (Fig. 5G), $\Delta flv3 dld2$ cells
319 maintained *pmf* close to WT level after the first seconds in light (Fig. 5E) due to diminished
320 conductivity (Fig. 5F). This indicates that the increased acceptor side limitation of PSI in $\Delta flv3 dld2$
321 in comparison to $\Delta flv3$ (Fig. 3A) is not caused by lack of photosynthetic control as a consequence of
322 lowered *pmf* generation in CET. Based on comparison of ECS kinetics in single, double, and triple
323 mutant cells studied here, we conclude that both Flv1/3 and NDH-1_{1,2} contribute to proton flux during
324 transitions from dark to high irradiance. However, deficiency in NDH-1_{1,2} in $\Delta dld2$ is mostly
325 compensated by elevated FDP-activity in terms of proton flux, while ATP synthase activity is
326 increased, possibly as a response to the delay in activation of carbon fixation (Fig. 2C). In contrast,
327 impaired thylakoid proton flux in $\Delta flv3$ during dark-to-light transitions cannot be compensated by
328 NDH-1. Instead, downregulation of ATP-synthase activity lowers thylakoid conductivity and allows
329 maintenance of *pmf* (apart from the first seconds of illumination) in $\Delta flv3$ and, even more
330 dramatically, in $\Delta flv3 dld2$.

Figure 5. Redox changes of PC, P700, and Fd, and buildup of pmf during dark-to-HL transitions. WT (A), $\Delta flv3$ (B), $\Delta d1d2$ (C), and $\Delta flv3 d1d2$ cells (D) were dark-adapted for 10 min, after which absorbance difference changes at 780–800 nm, 820–870 nm, 840–965 nm and 870–965 nm were measured with the Klas-NIR 100 spectrometer during 30 s illumination at $503 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ of red actinic light. The PC, P700, and Fd signals were deconvoluted from the absorbance difference changes using differential model plots (DMPs) determined for this study (Fig. S3). The upper panels in A–D show magnifications of the first 1.5 seconds of illumination from the lower panels. The experiment was repeated with three independent biological replicates, of which representative traces are shown. See also Supplemental Fig. S3. (E) Buildup

340 of *pmf* during transitions from dark to 500 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ of green actinic light. Absorbance difference
 341 between 500-480 nm (ECS) was measured with a JTS-10 spectrometer, and light-induced *pmf* was
 342 determined from the dark interval relaxation kinetics (DIRK) of the ECS signal. Means $\pm$ SEM from 3–4
 343 independent experiments are shown. The small inset shows *pmf* values after 1s of illumination for a clearer
 344 view. (F) Conductivity of the thylakoid membrane (gH^+) after one and 30 s of illumination as in (E). gH^+
 345 was determined as the inverse of the time constant of first-order post-illumination decay kinetics of the ECS
 346 signal. (G) Thylakoid proton flux (vH^+), calculated as *pmf* $\times$ gH^+ . (H) Representative traces of 500-480 nm
 347 absorbance differences used to determine the values in (E) and post-illumination ECS decay kinetics during a
 348 500 ms dark interval after 30 s in light (lower panels). First-order fits are shown in green. Cells, after growth
 349 in air [CO_2] / 50 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ for 4 days, were harvested and Chl *a* concentration adjusted to 10 μg
 350 ml^{-1} with fresh BG-11 for A–D and to 7.5 $\mu\text{g ml}^{-1}$ for E–J.

351

352 **PSI content is diminished in triple mutants deficient in Flv1/3 and NDH-1_{1,2} shifted to air**

353 **[CO₂] and high irradiance**

354 The results from the growth assays, along with the real-time gas exchange, P700 redox change, and
 355 ECS measurements suggested that in $\Delta\text{Id}2$ cells, an increase in the activity of FDPs compensates
 356 for the absence of a functional NDH-1_{1,2} complex, allowing efficient oxidation of PSI under high
 357 irradiance. Conversely, in Δflv1 and Δflv3 mutants, over-reduction of the electron transport chain
 358 occurs initially upon exposure to high irradiance, but another mechanism(s) eventually allow
 359 acclimation and survival in high [CO_2] and high light (Helman et al., 2003). Under air [CO_2] Flv1/3
 360 hetero-oligomers catalyse transient O₂ photoreduction, which is why no strong growth phenotype is
 361 observed in Δflv1 or Δflv3 under constant high irradiance (Allahverdiyeva et al. 2013, Santana-
 362 Sanchez et al. 2019). However, when both Flv1/3 and NDH-1_{1,2} are absent and the electron sink
 363 capacity of the cytosol is not elevated by high [CO_2], cells are unable to oxidize PSI in high light,
 364 possibly resulting in lethal photodamage to PSI. To assess this hypothesis, we employed
 365 immunoblotting to provide estimates of the protein content of PSII (D1) and PSI (PsaB) subunits,
 366 orange carotenoid protein (OCP), large isoform of the ferredoxin-NADP⁺ oxidoreductase (FNR_L),
 367 FDPs, as well as NdhD3 and the bicarbonate transporter SbtA after 24 hours exposure to different
 368 growth conditions.

369 WT and Δflv1 or Δflv3 cells grown at 3% [CO_2] / ML and shifted to air [CO_2] / ML at OD=0.1,
 370 accumulated similar levels of the PSII reaction center protein D1 and the PSI subunit PsaB.

371 However, we detected a moderate decrease in the amount of both D1 and PsaB in the $\Delta flv1 dld2$
372 and $\Delta flv3 dld2$ triple mutants, while an increased amount of both proteins was detected in $\Delta dld2$
373 (Fig. 6A). After 24 hours exposure to air [CO₂] / HL, PsaB amount had decreased even further in
374 $\Delta flv1 dld2$ and $\Delta flv3 dld2$ (Fig. 6C). Furthermore, 77K fluorescence spectra revealed that while
375 $\Delta flv3 dld2$ and WT cells grown under air [CO₂] / ML had similar PSI/PSII ratios (Fig. 6D), the 24-
376 hour exposure to air [CO₂] / HL caused a dramatic decrease in the relative PSI fluorescence cross
377 section (Fig. 6E). This strongly supports the hypothesis that loss of PSI contributes to the lethality
378 of the shift to air [CO₂] / HL. Interestingly, a substantial decrease in D1 content in air [CO₂] / HL
379 was also observed in the triple mutants (Fig. 6B-C).

380 Importantly, the $\Delta flv1 dld2$ and $\Delta flv3 dld2$ strains were unable to induce substantial accumulation
381 of the proteins encoded by the *flv4-2* operon after a shift from 3% to air [CO₂] (Fig. 6A and Fig.
382 6C). Only when pre-grown for 4 days in air [CO₂] / ML before being shifted to higher irradiance,
383 the expression of the operon was induced (Fig. 6B). Similarly to *flv4-2* operon proteins, both
384 NdhD3 and SbtA failed to accumulate after shifts from 3% to air [CO₂] / HL in the triple mutants,
385 which likely impairs CCM and contributes to the lethal phenotype of the triple mutants in those
386 conditions. Only small amounts of NdhD3 and SbtA were detected also in triple mutant cells shifted
387 from 3% [CO₂] to air [CO₂] / ML (Fig. 6A). Closer to WT amount of NdhD3 but only a small
388 amount of SbtA was detected after 24 hours in air [CO₂] / HL (shifted from air [CO₂] / ML) (Fig.
389 6B). In the $\Delta dld2$ mutant NdhD3 content was similar to WT, but Flv3 as well as the *flv4-2* operon
390 proteins (Flv2, Flv4, and Sll0218) were upregulated in all conditions tested (Fig. 6), which likely
391 contributes to the increased rate of O₂ photoreduction in that strain (Fig. 2C). The amounts of OCP
392 and of FNR_L were unchanged in the mutant strains in all conditions (Fig. 6).

Figure 6. Immunodetection of proteins related to photosynthesis and CCM and 77K fluorescence emission spectra in WT and mutant strains. Pre-cultures were grown for 3 days at 3% $[CO_2]$ / ML (50 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$) and then pelleted, resuspended in fresh BG-11 at OD₇₅₀=0.15 and (A) incubated for 24 hours in air $[CO_2]$ / ML, (B) grown for 4 days in air $[CO_2]$ / ML, harvested, resuspended in BG-11 at OD₇₅₀=0.15 and then incubated for 24 hours in air $[CO_2]$ / HL (220 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$), (C) incubated for 24 hours in air $[CO_2]$ / HL. Total protein extracts were separated by SDS-PAGE and probed with specific antibodies. The immunoblots shown are representative of 2–3 independent biological replicates. (D) 77K fluorescence emission spectra from intact WT and $\Delta flv3 d1d2$ cells grown for 4 days in air $[CO_2]$ / ML. (E) After pre-growth at 3% $[CO_2]$ / ML cells were pelleted, resuspended in fresh BG-11 at OD₇₅₀=0.15 and incubated for 24 hours in air $[CO_2]$ / HL. The cells (D and E) were pelleted, adjusted to $[Chl\ a]$ of 7.5 $\mu\text{g ml}^{-1}$, and excited with 440 nm light at 77K. The spectra were normalized to the PSII fluorescence peak at 685 nm.

DISCUSSION

Flv1/3 hetero-oligomers have been shown to be essential for photoreduction of O₂ as a rapid response to excessive reduction of PSI, and consequently, for cell survival under fluctuating light conditions (Allahverdiyeva et al., 2013; Santana-Sanchez et al., 2019). Moreover, we recently showed that in air [CO₂], Flv2/4 hetero-oligomers also catalyze low level light-induced steady-state reduction of O₂ on the acceptor side of PSI in a coordinated manner with Flv1/3 (Santana-Sanchez et al., 2019).

In turn, another alternative electron transport component, the NDH-1 complex, functions in CET around PSI, using Fd as electron donor (Saura and Kaila, 2019; Schuller et al., 2019). Besides CET, the NDH-1_{1,2} complexes, incorporating the NdhD1 or NdhD2 subunit, are involved in respiration, whereas NDH-1_{3,4} complexes, incorporating the NdhD3 or NdhD4 subunit, function in CO₂ uptake. In the current study, we have used *Synechocystis* mutant strains with different combinations of deficiencies in Flv1/3 and NDH-1 in order to examine the physiological significance of these alternative electron transport pathways and the possibility of functional interdependence between them. Our results provide compelling evidence for the concomitant physiological functions of Flv1/3 and NDH-1_{1,2} and their dynamic coordination for the efficient oxidation of PSI (thus protecting it from photodamage) under variable light conditions, and when low [CO₂] limits the consumption of reductants in the CBB cycle. The two pathways can compensate for each other's absence to some extent, but absence of both the Flv1/3 hetero-oligomer and NDH-1_{1,2} is lethal when cells are transferred from elevated [CO₂] to the combined condition of air level [CO₂] and high irradiance (Fig. 1D-E). In addition to the absence of these two important pathways, lethality is due to an inability to accumulate low C_i-inducible photoprotective CCM proteins (Fig. 6C). Combined deficiency of Flv1/3 and NDH-1_{3,4} did not result in a lethal phenotype upon similar shifts, indicating that functional redundancy exists specifically between the Mehler-like reaction and NDH-1_{1,2}.

Coordinated functions of Flv1/3 and NDH-1_{1,2} protect PSI by maintaining redox poise between the PETC and carbon fixation

Cells lacking functional Flv1/3 hetero-oligomers suffer from a transient over-reduction of PSI during sudden increases in light intensity (Fig. 3 and Fig. 5) due to the impairment of the Mehler-

like reaction (Fig. 2, Allahverdiyeva et al., 2013). Whilst NAD(P)H has been proposed as the electron donor to Flv3 (Vicente et al., 2002; Brown et al., 2019) and Flv1 homo-oligomers (Brown et al., 2019), *in vivo* experiments have been unsupportive (Mustila et al. 2016), with exact electron donor(s) to FDP hetero-oligomers yet to be proven. *In vivo* experiments performed in this study demonstrate that the quick re-oxidation of Fd after 0.5 s (Fig. 5A) in light is absent in the $\Delta flv3$ deletion strains, whereby Fd remains strongly reduced (Fig. 5B). This result, together with a lack of impairment in NADP⁺ reduction and oxidation kinetics of $\Delta flv3$ deletion strains (similar conditions, Fig. 4F) provide strong support for Fd, rather than NAD(P)H, being the primary electron donor to the Flv1/3 hetero-oligomer *in vivo*. Accordingly, Fd has been shown to interact with Flv1 and Flv3 by Fd-chromatography (Hanke et al., 2011) and with Flv3 by a two-hybrid assay (Cassier-Chauvat and Chauvat, 2014).

The deficiency of NDH-1_{1,2}, in turn, reduced the cells' ability to maintain oxidized P700 and Fd under high irradiance soon after (~50–200 ms) a dark-to-light transition (Fig. 5C). This brief timescale may be due to PSI-NDH-1 supercomplexes (Gao et al., 2016) where P700 can be oxidized rapidly upon the onset of illumination. It has been shown that formation of the NDH-1-PSI supercomplex is important to keep PSI functional under various stress conditions (Zhao et al. 2017). NDH-1_{1,2}-deficiency also caused slightly elevated acceptor side limitation of PSI under low irradiance (Fig. 3A), which was likely due to a lack of NADPH oxidation during the low light phases of fluctuating light (Fig. 4C). Thus, NDH-1_{1,2} appears to play an important role under low light, as has been previously suggested for chloroplastic NDH in angiosperms (Yamori et al., 2011; Yamori et al., 2015) and bryophytes (Ueda et al., 2012). The delayed activation of CBB in $\Delta d1d2$ did not result in an inability to oxidize PSI (Fig. 3B and Fig. 5C), likely due to significant enhancement of FDP-mediated O₂ photoreduction (Fig. 2C) providing an enlarged electron sink for the photosynthetic electron transport chain. An opposite order of causation is also possible, whereby the excessive funneling of photosynthetic electrons to O₂ would cause the delay in induction of CO₂ fixation in $\Delta d1d2$. However, the observation that there is no NADPH shortage at the onset of illumination in $\Delta d1d2$, and rather the consumption of NADPH during transitions from HL to LL is impaired (Fig. 4C), suggests that the availability of reductant for the CBB is not the limiting factor. Nor is it likely CO₂, as CCM is functioning as in WT (Fig. 2C), or ATP, as ATP synthase activity was even higher than in WT in $\Delta d1d2$ (Fig. 5F). Nevertheless, simultaneous deficiency of FDPs in addition to NDH-1_{1,2} (in $\Delta flv3 d1d2$ triple mutant), with the effect of diminishing the flow of photosynthetic electrons to O₂ photoreduction, mostly rescued the delay in CO₂ fixation (Fig. 2D)

as well as NADPH consumption (Fig. 4D) seen in $\Delta dld2$. Transient O_2 photoreduction at a low rate was still observed in $\Delta flv3 dld2$ during dark-to-HL transitions (Fig. 2D), possibly mediated by the thylakoid terminal oxidases (Ermakova et al., 2016) or by photorespiration (Allahverdiyeva et al., 2011). However, the triple mutants suffered from more severe inability to oxidize PSI than $\Delta flv3$ during sudden increases in irradiance (Fig. 3). These observations suggest that in addition Flv1/3, also NDH-1_{1,2} has a role in contributing to oxidation of PSI during changes in light conditions or carbon availability. Upon deficiency of NDH-1_{1,2} (in $\Delta dld2$), cells prioritize protection of PSI over efficient CO_2 fixation by upregulating the Mehler-like reaction via an unknown mechanism. The triple mutants cannot do this, leading to timely induction of CO_2 fixation at the high cost of inability to oxidize PSI (Fig. 3A and 3) or Fd (Fig. 5D). This results in loss of PSI (Fig. 6), possibly due to photodamage to its FeS clusters (Tiwari et al., 2016; Shimakawa et al., 2016).

On the mechanism of NDH-1-mediated oxidation of PSI

Contribution of NDH-1 to oxidation of PSI during sudden increases in light intensity is not unprecedented. In angiosperms, where FDPs have been lost during evolution (Ilik et al., 2017), *Arabidopsis* mutants lacking the NDH complex show a delay in oxidation of PSI during increases in light intensity in comparison to *Arabidopsis* WT (Nikkanen et al., 2018; Shimakawa and Miyake, 2018a). By definition, canonical CET cannot directly increase the relative proportion of oxidized P700, as electrons from the acceptor side of PSI are shunted back to the intersystem electron transfer chain. However, NDH-1 may enhance PSI oxidation by at least three alternative, but mutually non-exclusive mechanisms.

- i) NDH-1-mediated CET is coupled to translocation of protons from cytosol to the thylakoid lumen with a $2H^+/e^-$ stoichiometry (Strand et al., 2017; Saura and Kaila, 2019). It will therefore contribute to buildup of ΔpH , which limits electron transfer to PSI by inhibiting PQH₂ oxidation at Cyt *b6f* (Shimakawa and Miyake, 2018b) and drives ATP synthesis to accommodate the needs of the CBB, increasing its electron sink capacity. As the Mehler-like reaction also contributes to buildup of ΔpH by consuming H^+ on the cytosolic side of the thylakoid membrane and by supporting linear electron flow (Fig. 5), (Allahverdiyeva et al., 2013), enhanced FDP activity in $\Delta dld2$ (Fig. 2C) partly compensates for the lack of NDH-1_{1,2} in respect to generation of proton flux (Fig. 5G). However, this fails to explain

the exacerbated impairment of P700 and Fd oxidation in $\Delta flv3 d1d2$ in comparison to $\Delta flv3$ (Fig. 3 and Fig. 5) because in the triple mutant *pmf* was not lower than in $\Delta flv3$ (Fig. 5E). Adequate *pmf* is maintained however at the expense of ATP synthase activity (Fig. 5F), and it is likely that diminished ATP production contributes to the increased acceptor side limitation of PSI in $\Delta flv3 d1d2$ (Fig. 3A). NDH-1_{1,2} could therefore contribute to P700 oxidation by enhancing cytosolic sink capacity by providing a more suitable ATP:NADPH ratio for the CBB. Adjustment of the ATP:NADPH ratio closer to the theoretically optimal 3:2 has long been considered a fundamental reason for the existence of CET (Kramer et al., 2004; Yamori and Shikanai, 2016).

ii) NDH-1-mediated respiratory electron transfer, *i.e.* coupling of PQ reduction by NDH-1 to O₂ reduction by thylakoid terminal oxidases (*i.e.* Cyd and Cox) would also contribute to oxidation of PSI by relieving electron pressure in the intersystem chain during illumination (Ermakova et al., 2016), as well as further contribute to ΔpH by consuming H⁺ on the cytosolic side of thylakoids (Cyd) and pumping protons to the lumen (Cox) (Brändén et al., 2006). Interestingly, Liu and colleagues (2012) have shown that the subcellular localization of NDH-1 complexes is dependent on the redox state of the PQ pool. An oxidized PQ pool causes NDH-1 to accumulate at specific clusters in thylakoid membranes where it would be more likely to transfer electrons (via the PQ pool) to a terminal oxidase. A reduced PQ pool, in turn, results in a more even distribution of NDH-1 within thylakoids (Liu et al., 2012). However, it is important to note that terminal oxidases do not have a high electron sink capacity (Ermakova et al., 2016).

iii) NDH-1 has been predicted, albeit not yet experimentally shown in photosynthetic organisms, to be able to function in reverse: to oxidize PQH₂ driven by concomitant release of protons from the thylakoid lumen (Strand et al., 2017). Such reverse activity would constitute a ‘pseudo-linear’ electron transfer pathway that would bypass PSI and thereby prevent its over-reduction. This could occur in conditions where the PQ pool is reduced, *pmf* is high, and the Fd pool is oxidized (Strand et al., 2017). Such conditions likely exist transiently during dark-to-light and LL-to-HL transitions (Fig. 4 and 5) (Strand et al., 2019). Accordingly, fast re-reduction of P700⁺ already after 50 ms was observed in $\Delta d1d2$ and $\Delta flv3 d1d2$ during dark-to-HL transitions (Fig. 5C). As supported by the impaired ability to oxidize Fd in absence of Flv1/3 (Fig. 5), the presence of Flv1/3-catalyzed Mehler-

like reaction would likely be essential in this model in order to maintain the Fd pool in a sufficiently oxidized state to provide electron acceptors for reverse-functioning NDH-1. It is noteworthy, however, that NDH-1 reverse activity would also have the effect of lowering ΔpH , thereby relieving photosynthetic control at Cyt *b6f*. This could counteract the effect of any reverse NDH-1 activity by increasing electron flow to PSI.

Hypothetical mechanisms for coordination of Flv1/3 and NDH-1_{1,2} activities are shown in Fig. 7.

Inability to induce a strong CCM network contributes to lethality upon shifts to air [CO₂] / high irradiance in cells deficient in both Flv1/3 and NDH-1_{1,2}

Exposure of *Synechocystis* cells to low [CO₂] induces expression of high-affinity CCM-related genes such as SbtA, a HCO₃⁻ / Na⁺ symporter on the plasma membrane, and NdhD3, which is part of the NDH-1₃ complex on thylakoids specializing in C_i uptake (Ohkawa et al., 2000; Shibata et al., 2001, Shibata et al., 2002; Zhang et al., 2004). CCM is energetically expensive, but the large C_i flux involved in the operation of CCM contributes to dissipation of excess light energy under stress conditions (Xu et al., 2008; Burnap et al. 2015). NDH-1_{3,4} in particular has a photoprotective role, using reduced Fd to drive CO₂ conversion to HCO₃⁻ with concomitant translocation of protons into the thylakoid lumen.

Despite the low induction level of the CCM proteins NdhD3 and SbtA, the triple mutants (Fig. 6) survive under standard growth conditions of air [CO₂] / ML (Fig. 1B-C). When a decrease in [CO₂] is coupled with an increase in irradiance, however, a low amount of CCM, reflected by low accumulation level of NdhD3 and SbtA in the $\Delta flv3d1d2$ and $\Delta flv3d1d2$ triple mutants (Fig. 6C) fails to dissipate excess energy, thus resulting in photodamage and lethality of those conditions. The regulatory mechanisms controlling the inability of the triple mutants to accumulate low C_i-inducible proteins is unclear and remains to be elucidated. Accumulation of NADP⁺ and α -ketogutarate inhibits the induction of CCM gene expression via interaction with the transcription factor NdhR (CcmR) (Daley et al., 2012). NdhR also controls expression of the Flv4-2 operon encoding Flv2, Flv4, and Slh0218 (Eisenhut et al., 2012), all of which also showed decreased accumulation levels in the triple mutants (Fig. 6). However, at least in the tested conditions, no substantial increase in the relative amount of NADP⁺ was detected in $\Delta flv3d1d2$ cells (Fig. 4). It is important to note however,

that Flv1/3 and NDH-1_{3,4} cooperation alone is not crucial for cell metabolism, since the $\Delta flv1 d3d4$ and $\Delta flv3 d3d4$ mutant cells do not show clear growth phenotypes (Fig. S1).

CONCLUSIONS

In the current study we have shown that FDPs and NDH-1 function cooperatively to maintain redox balance between the PETC and cytosolic carbon assimilation upon sudden changes in light intensity and/or carbon availability. It is likely that both pathways receive electrons primarily from Fd, which enables coordinated regulation of their activities. Cooperation of FDPs and NDH-1 has an essential photoprotective role during shifts to air [CO₂] /HL, contributing to efficient oxidation of PSI, buildup of *pmf*, and induction of expression of low C_i-specific CCM-related genes. Finally, it is worth noting that simultaneous removal of Flv1/3 and NDH-1_{1,2} as competitive electron sinks downstream of PSI may be a useful biotechnological tool for maximizing direction of photosynthetic electrons to desired pathways under controlled environmental conditions (McCormick et al., 2013; Thiel et al., 2019; Jokel et al., 2019). Furthermore, our findings about functional interplay between FDPs and NDH-1 will be highly relevant in projects aiming to enhance crop productivity via introduction of exogenous FDPs to higher plants (Yamamoto et al., 2016; Gómez et al., 2018).

Figure 7. Schematic models for the coordination of the activities of FDPs and NDH-1 in preventing over-reduction of PSI. (A) In conditions where the Fd pool is highly reduced and the PQ pool is not, NDH-1L transfers electrons from Fd to the PQ pool, coupled to O₂ reduction by thylakoid-localised RTOs. NDH-1L pumps 2H⁺/e⁻ into the lumen while FDPs and RTOs consume H⁺ when reducing O₂ to H₂O on the

cytosolic side of thylakoids. This increases transthylakoid *pmf*, which in turn drives ATP synthesis as well as inhibits PQH₂ oxidation at Cyt *b₆f*, thereby preventing excessive electron supply to PSI. Carbonic anhydrase activity of NDH1-MS converts HCO₃⁻ to CO₂, consuming H⁺ in the cytosol, likely also coupled to translocation of H⁺ to the lumen, further increasing *pmf*. Increased availability of CO₂ and ATP/NADPH ratio enhances the electron sink capacity of the CBB on the acceptor side of the PETC. **(B)** If the PQ pool is highly reduced, Fd pool is oxidized, and *pmf* is high, NDH-1L could hypothetically function in reverse, transferring electrons from PQH₂ to Fd⁺, powered by release of 2H⁺/e⁻ from the lumen. A strong Mehler-like reaction catalyzed by Flv1/3 would be required to maintain a sufficient pool of oxidized Fd, with Flv1/3 likely accepting electron directly from Fd⁻. Most components depicted in (A) are omitted from (B) to enhance clarity, not to suggest their absence. Extent of the colour fill for PQ/PQH₂ and Fd depict level of reduction.

MATERIALS AND METHODS

Strains and culture conditions

In the current study we used the glucose-tolerant WT strain *Synechocystis* sp. PCC 6803 (Williams, 1988), single mutant strains $\Delta flv1$ and $\Delta flv3$ (Helman et al., 2003), the M55 mutant ($\Delta ndhB$) (Ogawa 1991), double mutants $\Delta d1d2$ and $\Delta d3d4$ (Ohkawa et al., 2000a), and triple mutants $\Delta flv1 d1d2$, $\Delta flv3 d1d2$, $\Delta flv1 d3d4$, and $\Delta flv3 d3d4$ obtained from the CyanoMutants collection (Nakamura et al., 1999). The triple mutants were constructed by T. Ogawa in $\Delta d1d2$ and $\Delta d3d4$ backgrounds. All the mutants demonstrated complete segregation. Pre-experimental cultures were always grown in 30 ml batches of BG-11 medium pH 7.5 (Williams, 1988) under 3% [CO₂] at 30°C under continuous white light of 50 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ (ML) with agitation. Mutant pre-cultures were supplemented with the appropriate antibiotics. At the logarithmic growth phase, cells were harvested and re-suspended in fresh BG-11 without antibiotics at OD₇₅₀ of 0.1–0.2, as described in the appropriate figure legends. Cells were then shifted to air [CO₂] / 30°C and illuminated continuously with white light of either 50 or 220 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$, as described in the figure legends.

612 Gas exchange measurements

613 Exchange of $^{16}\text{O}_2$ ($m/z=32$), $^{18}\text{O}_2$ ($m/z=36$), CO_2 ($m/z=44$) was measured *in vivo* with membrane-
 614 inlet mass spectrometry (MIMS) as described in Mustila et al. (2016). Harvested cells were
 615 resuspended in fresh BG-11 pH 7.5 and adjusted to $10\ \mu\text{g}$ Chlorophyll (Chl) $a\ \text{ml}^{-1}$, and kept for
 616 one hour in air $[\text{CO}_2]$ / $50\ \mu\text{mol photons m}^{-2}\text{s}^{-1}$. Prior to the measurements, cells were
 617 supplemented with $^{18}\text{O}_2$ at an equivalent concentration to $^{16}\text{O}_2$ and with $1.5\ \text{mM}$ NaHCO_3 . Cells
 618 were dark-adapted for 15 min, after which gas exchange was monitored over a 5 min illumination
 619 period of $500\ \mu\text{mol photons m}^{-2}\text{s}^{-1}$ of white actinic light (AL). The gas exchange rates were
 620 calculated according to Beckmann et al., (2009).

621

622 Measurement of fluorescence and absorbance changes

623 Experimental cultures for all spectroscopic experiments were grown for four days in air $[\text{CO}_2]$ / 50
 624 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ at 30°C in BG-11 pH 7.5. Chl a fluorescence and P700-oxido-reduction (875-
 625 830 nm absorbance difference) were simultaneously recorded (Fig. 3) with the Dual-PAM 100
 626 spectrophotometer (Walz, Effeltrich, Germany). Harvested cells were resuspended in fresh BG-11
 627 pH 7.5 and adjusted to $10\ \mu\text{g}$ Chl $a\ \text{ml}^{-1}$, and kept for one hour in air $[\text{CO}_2]$ / $50\ \mu\text{mol photons}$
 628 $\text{m}^{-2}\text{s}^{-1}$ and dark-adapted for 10 min before being subjected to fluctuating light regime alternating
 629 between 1 min periods of 25 and $530\ \mu\text{mol photons m}^{-2}\text{s}^{-1}$ of red AL. Saturating pulses (500 ms of
 630 $5000\ \mu\text{mol photons m}^{-2}\text{s}^{-1}$) were administered at 15 s intervals. Photosynthetic parameters were
 631 calculated as follows: $Y(\text{NA})=(P_m-P_m')/P_m$; $Y(\text{ND})=P/P_m$; $Y(\text{I})=1-Y(\text{ND})-Y(\text{NA})$ (Klughammer and
 632 Schreiber, 2008); $Y(\text{II})=(F_m'-F)/F_m'$ (Genty et al., 1989).

633 A Dual-Klas NIR 100 spectrophotometer (Walz) was used to measure absorbance difference
 634 changes at 780–800 nm, 820–870 nm, 840–965 nm and 870–965 nm, from which PC, P700, and Fd
 635 redox changes (Fig. 5) were deconvoluted based on differential model plots (DMPs) for PC, P700,
 636 and Fd (Klughammer and Schreiber, 2016; Schreiber, 2017; Setif et al., 2019). In WT
 637 *Synechocystis*, very fast re-oxidation of Fd upon illumination impedes measurement of model
 638 spectra. To circumvent this issue, we measured the DMPs from $\Delta flv3\ dld2$ cells where P700
 639 oxidation was severely delayed in comparison to WT (Fig. 3B). The DMPs (Fig. S3) were measured
 640 using scripts provided with the instrument software. Harvested cells were adjusted to $10\ \mu\text{g}$ Chl a
 641 ml^{-1} with fresh BG-11 pH 7.5, and kept for one hour in air $[\text{CO}_2]$ / $50\ \mu\text{mol photons m}^{-2}\text{s}^{-1}$ and

dark-adapted for 30 min before illumination for 30 s with 503 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ of red AL. It is noteworthy that cytochrome c_6 and ($F_A F_B$) redox changes likely contribute to the PC and Fd signals, respectively (Setif et al., 2019).

NADPH fluorescence changes between 420 and 580 nm, induced by excitation at 365 nm, were measured simultaneously with Chl a fluorescence with the Dual-PAM 100 and its 9-AA/NADPH accessory module (Walz) (Schreiber and Klughammer, 2009; Kauny and Setif, 2014). Experimental samples were prepared as above and [Chl a] was adjusted to 5 $\mu\text{g ml}^{-1}$. Cells were dark-adapted for 10 min and subjected to the same fluctuating light regime as in Fig. 3, but without saturating pulses.

Electrochromic shift (ECS) was measured as absorbance difference between 500 and 480 nm (Viola et al., 2019). Absorbance changes were measured from dark-adapted cell suspensions with a JTS-10 spectrophotometer (BioLogic, Seyssinet-Pariset, France) using 500 and 480 nm CWL 50mm 10nm FWHM bandpass filters (Edmund Optics, Barrington, NJ, USA) and BG39 filters (Schott, Mainz, Germany) protecting the light detectors from scattering effects. Absorbance changes induced by measuring light only were subtracted from changes under ML + AL. ECS values were normalized to ECS change induced by a single turnover flash provided by a XST-103 xenon lamp (Walz). Experimental samples were prepared as above and [Chl a] was adjusted to 7.5 $\mu\text{g ml}^{-1}$, and illuminated with green AL of 500 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ for 1s or 60 s interspersed with 500 ms dark intervals at 2, 7.5, 18, 30, and 45 s. Pmf was calculated as the extent of ECS decrease at the dark intervals. Thylakoid conductivity (gH^+) was calculated as the inverse of the time constant of a first-order fit to ECS relaxation kinetics during a dark interval, and proton flux (vH^+) as $pmf \times gH^+$ (Cruz. et al., 2005).

77K fluorescence emission spectra were measured with a QE Pro spectrophotometer (Ocean Optics, Dunedin, FL, USA). Cells were harvested and [Chl a] was adjusted to 7.5 $\mu\text{g ml}^{-1}$ with fresh BG-11 pH 7.5. Cells were frozen in liquid N_2 and excited at 440 nm. Raw spectra were normalized to PSII-fluorescence peak at 685 nm.

Protein extraction and immunoblotting

Total protein extracts from cultures were isolated as previously described (Zhang et al., 2009), separated by SDS-PAGE on 12 % gels with 6M urea, and blotted on PVDF membranes.

671 Membranes were probed with antibodies raised against D1, PsaB (Agrisera, Vännäs, Sweden, AS10
672 695), PetH (Kindly shared by H. Matthijs), OCP (kindly shared by D. Kirilovsky), SbtA (kindly
673 shared by T. Ogawa), NdhD3 (Eurogentec, Liège, Belgium), Flv2, Flv4, Flv3 (Antiprot, Puchheim,
674 Germany), sll0218 (MedProbe, Oslo, Norway). Horseradish peroxidase –conjugated secondary
675 antibody (GE Healthcare, Chicago, IL, USA) and Amersham ECL (GE Healthcare) were used for
676 detection.

677

678 **Supplemental material**

679 Supplemental Figure S1. Growth of $\Delta flv1 d3d4$ and $\Delta flv3 d3d4$ mutant strains.

680 Supplemental Figure S2. O₂ and CO₂ fluxes in M55 and in WT at pH 6.

681 Supplemental Figure S3. Near-infrared (NIR) differential model blots (DMPs) for deconvolution of
682 PC, P700, and Fd signals with the Klas-NIR 100 spectrometer.

683

684 **ACKNOWLEDGEMENTS**

685 We thank Prof. Teruo Ogawa for sharing the FDP and NDH-1 mutants and Dr. Henna Mustila for
686 construction of the $\Delta M55 flv3$ strain. Dr. Duncan Fitzpatrick is thanked for expertise and assistance
687 with MIMS. We also thank Dr. Pierre Setif and Dr. Gert Schansker for their expert advice on
688 measuring the DMPs for the Dual-Klas NIR spectrophotometer, and Prof. Jens Appel for critical
689 reading of the manuscript. We declare no conflicts of interest.

690

691 **FIGURE LEGENDS**

692 **Figure 1. Growth of WT and mutant strains under different experimental conditions.** All
693 experimental cultures were started from pre-cultures adjusted to OD₇₅₀=0.1. (A) Cells were pre-
694 cultivated under 3% [CO₂] / ML (50 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$), adjusted to OD₇₅₀=0.1 and grown in the
695 same conditions. (B) Cells were pre-cultivated under 3% [CO₂]/ML and shifted to air [CO₂]/ML.

696 (C) Cells were pre-cultivated under air [CO₂]/ML and shifted to air [CO₂]/HL 220 μmol photons m⁻²s⁻¹. (D) Cells were pre-cultivated under 3% [CO₂]/ML and shifted to air [CO₂]/HL. The values
697 2s⁻¹). (D) Cells were pre-cultivated under 3% [CO₂]/ML and shifted to air [CO₂]/HL. The values
698 presented in A–D are means of three independent biological replicates ±SE. (E) Growth of WT and
699 mutant cells on agar plates containing BG-11. Cells grown under 3% [CO₂] / ML were harvested
700 and spotted on BG-11 agar plates in ten-fold dilution series starting at OD₇₅₀=1, incubation was
701 under air [CO₂] /ML and air [CO₂] /HL for 7 days. See also Supplemental Fig. S1.

702 **Figure 2. O₂ and CO₂ exchange rates in WT and mutant strains.** Cells were grown in air [CO₂] /
703 ML for 4 days, after which the cells were harvested and Chl *a* concentration adjusted to 10 μg ml⁻¹
704 with fresh BG-11. Cells were dark-adapted for 15 min, and gas exchange was monitored by
705 membrane inlet mass spectrometry (MIMS) over a 5 min illumination period with 500 μmol
706 photons m⁻²s⁻¹ of white actinic light. Prior to the measurements, samples were supplemented with
707 ¹⁸O₂ at an equivalent concentration to ¹⁶O₂ to distinguish O₂ uptake from O₂ evolution, and with 1.5
708 mM NaHCO₃. The dashed line in each panel indicates the compensation point of CO₂ fixation
709 (uptake rate equals respiratory rate). The experiment was repeated with three independent biological
710 replicates, of which representative measurements are shown.

711 **Figure 3. Photosynthetic parameters in WT and mutants strains in fluctuating light intensities.**
712 (A) Acceptor side limitation of PSI, Y(NA), (B) donor side limitation of PSI, Y(ND), as well as (C)
713 effective yield of PSI, Y(I) were calculated from absorbance difference changes at 875-830 nm
714 according to (Klughammer and Schreiber, 2008). Chl *a* fluorescence changes were measured
715 simultaneously with PSI redox changes and used to calculate (D) effective yield of PSII. Cells were
716 grown in air [CO₂] / ML for 4 days, after which the cells were harvested and Chl *a* concentration
717 adjusted to 10 μg ml⁻¹ with fresh BG-11. Cells were dark-adapted for 10 min, and then illuminated
718 with red actinic light intensity alternating between 25 and 530 μmol photons m⁻²s⁻¹ in 1 min periods.
719 Saturating pulses (500 ms of 5000 μmol photons m⁻²s⁻¹) were provided every 15 seconds. The values
720 shown are averages from 3–4 independent biological replicates ±SE.

721 **Figure 4. NADPH and Chl *a* fluorescence under fluctuating light in WT and mutant strains.**
722 WT (A), *Δflv3* (B), *Δd1d2* (C), and *Δflv3 d1d2* cells (D) were grown in air [CO₂] / ML for 4 days,
723 after which the cells were harvested and Chl *a* concentration adjusted to 5 μg ml⁻¹ with fresh BG-
724 11. Cells were dark-adapted for 10 min, and then subjected to illumination with red actinic light
725 alternating between 25 and 530 μmol photons m⁻²s⁻¹ in 1 min periods. NADPH redox changes were
726 monitored with the Dual-PAM 100 spectrophotometer and its 9-AA/NADPH accessory module by

727 measuring fluorescence changes between 420 and 580 nm induced by excitation at 365 nm, as
728 described by (Schreiber and Klughammer, 2009) and (Kauny and Setif, 2014). Chl *a* fluorescence
729 was recorded simultaneously. (E) NADPH redox changes during illumination of dark-adapted cells
730 with 530 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$. Other experimental details are as in A-D. (F) shows a magnification
731 of the first 2 s of illumination in (E). The values are normalized according to minimum and
732 maximum fluorescence changes at the onset and cessation of illumination, respectively. The
733 experiments were repeated with three independent biological replicates, of which representative
734 measurements are shown. r.u. = relative units.

735 **Figure 5. Redox changes of PC, P700, and Fd, and buildup of *pmf* during dark-to-HL**
736 **transitions.** WT (A), $\Delta flv3$ (B), $\Delta d1d2$ (C), and $\Delta flv3 d1d2$ cells (D) were dark-adapted for 10 min,
737 after which absorbance difference changes at 780–800 nm, 820–870 nm, 840–965 nm and 870–965
738 nm were measured with the Klas-NIR 100 spectrometer during 30 s illumination at 503 μmol
739 $\text{photons m}^{-2}\text{s}^{-1}$ of red actinic light. The PC, P700, and Fd signals were deconvoluted from the
740 absorbance difference changes using differential model plots (DMPs) determined for this study
741 (Fig. S3). The upper panels in A–D show magnifications of the first 1.5 seconds of illumination
742 from the lower panels. The experiment was repeated with three independent biological replicates, of
743 which representative traces are shown. See also Supplemental Fig. S3. (E) Buildup of *pmf* during
744 transitions from dark to 500 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ of green actinic light. Absorbance difference
745 between 500–480 nm (ECS) was measured with a JTS-10 spectrometer, and light-induced *pmf* was
746 determined from the dark interval relaxation kinetics (DIRK) of the ECS signal. Means $\pm$ SEM from
747 3–4 independent experiments are shown. The small inset shows *pmf* values after 1s of illumination
748 for a clearer view. (F) Conductivity of the thylakoid membrane (gH+) after one and 30 s of
749 illumination as in (E). gH+ was determined as the inverse of the time constant of first-order post-
750 illumination decay kinetics of the ECS signal. (G) Thylakoid proton flux (vH+), calculated as *pmf* x
751 gH+. (H) Representative traces of 500–480 nm absorbance differences used to determine the values
752 in (E) and post-illumination ECS decay kinetics during a 500 ms dark interval after 30 s in light
753 (lower panels). First-order fits are shown in green. Cells, after growth in air [CO₂] / 50 μmol
754 $\text{photons m}^{-2}\text{s}^{-1}$ for 4 days, were harvested and Chl *a* concentration adjusted to 10 $\mu\text{g ml}^{-1}$ with fresh
755 BG-11 for A–D and to 7.5 $\mu\text{g ml}^{-1}$ for E–J.

756 **Figure 6. Immunodetection of proteins related to photosynthesis and CCM and 77K**
757 **fluorescence emission spectra in WT and mutant strains.** Pre-cultures were grown for 3 days at
758 3% [CO₂] / ML (50 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$) and then pelleted, resuspended in fresh BG-11 at

OD₇₅₀=0.15 and (A) incubated for 24 hours in air [CO₂] / ML, (B) grown for 4 days in air [CO₂] / ML, harvested, resuspended in BG-11 at OD₇₅₀=0.15 and then incubated for 24 hours in air [CO₂] / HL (220 μmol photons m⁻²s⁻¹), (C) incubated for 24 hours in air [CO₂] / HL. Total protein extracts were separated by SDS-PAGE and probed with specific antibodies. The immunoblots shown are representative of 2–3 independent biological replicates. (D) 77K fluorescence emission spectra from intact WT and *Δflv3 dld2* cells grown for 4 days in air [CO₂] / ML. (E) After pre-growth at 3% [CO₂] / ML cells were pelleted, resuspended in fresh BG-11 at OD₇₅₀=0.15 and incubated for 24 hours in air [CO₂] / HL. The cells (D and E) were pelleted, adjusted to [Chl *a*] of 7.5 μg ml⁻¹, and excited with 440 nm light at 77K. The spectra were normalized to the PSII fluorescence peak at 685 nm.

Figure 7. Schematic models for the coordination of the activities of FDPs and NDH-1 in preventing over-reduction of PSI. (A) In conditions where the Fd pool is highly reduced and the PQ pool is not, NDH-1L transfers electrons from Fd to the PQ pool, coupled to O₂ reduction by thylakoid-localised RTOs. NDH-1L pumps 2H⁺/e⁻ into the lumen while FDPs and RTOs consume H⁺ when reducing O₂ to H₂O on the cytosolic side of thylakoids. This increases transthylakoid *pmf*, which in turn drives ATP synthesis as well as inhibits PQH₂ oxidation at Cyt *b₆f*, thereby preventing excessive electron supply to PSI. Carbonic anhydrase activity of NDH1-MS converts HCO₃⁻ to CO₂, consuming H⁺ in the cytosol, likely also coupled to translocation of H⁺ to the lumen, further increasing *pmf*. Increased availability of CO₂ and ATP/NADPH ratio enhances the electron sink capacity of the CBB on the acceptor side of the PETC. (B) If the PQ pool is highly reduced, Fd pool is oxidized, and *pmf* is high, NDH-1L could hypothetically function in reverse, transferring electrons from PQH₂ to Fd⁺, powered by release of 2H⁺/e⁻ from the lumen. A strong Mehler-like reaction catalyzed by Flv1/3 would be required to maintain a sufficient pool of oxidized Fd, with Flv1/3 likely accepting electron directly from Fd⁺. Most components depicted in (A) are omitted from (B) to enhance clarity, not to suggest their absence. Extent of the colour fill for PQ/PQH₂ and Fd depict level of reduction.

LITERATURE CITED

Alboresi A, Storti M, Cendron L, Morosinotto T (2019) Role and regulation of class-C flavodiiron proteins in photosynthetic organisms. *Biochem J* 476: 2487-2498, doi/10.1042/BCJ20180648.

789 Allahverdiyeva Y, Ermakova M, Eisenhut M, Zhang P, Richaud P, Hagemann M, Cournac L, Aro
790 EM (2011) Interplay between Flavodiiron Proteins and Photorespiration in *Synechocystis* sp. PCC
791 6803. *J Biol Chem* 286: 24007-24014. doi/10.1074/jbc.M111.223289.

792 Allahverdiyeva Y, Mustila H, Ermakova M, Bersanini L, Richaud P, Ajlani G, Battchikova N,
793 Cournac L, Aro EM (2013) Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and
794 photosynthesis under fluctuating light. *Proc Natl Acad Sci USA* 110: 4111-4116,
795 doi/10.1073/pnas.1221194110.

796 Allahverdiyeva Y, Suorsa M, Tikkanen M, Aro EM (2015) Photoprotection of photosystems in
797 fluctuating light intensities. *J Exp Bot* 66: 2427-2436. doi/10.1093/jxb/eru463.

798 Allakhverdiev SI, Murata N (2004) Environmental stress inhibits the synthesis de novo of proteins
799 involved in the photodamage-repair cycle of Photosystem II in *Synechocystis* sp PCC 6803.
800 *Biochim Biophys Acta Bioenerg* 1657: 23-32. doi/10.1016/j.bbabbio.2004.03.003.

801 Beckmann K, Messinger J, Badger MR, Wydrzynski T, Hillier W (2009) On-line mass
802 spectrometry: membrane inlet sampling. *Photosynth Res* 102: 511-522. doi/10.1007/s11120-009-
803 9474-7.

804 Bernat G, Appel J, Ogawa T, Rögner M (2011) Distinct Roles of Multiple NDH-1 Complexes in the
805 Cyanobacterial Electron Transport Network as Revealed by Kinetic Analysis of P700(+) Reduction
806 in Various *ndh*-Deficient Mutants of *Synechocystis* sp. Strain PCC6803. *J Bacteriol* 193: 292-295,
807 doi/10.1128/JB.00984-10.

808 Brändén G, Gennis RB, Brzezinski P (2006) Transmembrane proton translocation by cytochrome c
809 oxidase. *Biochim Biophys Acta* 1757: 1052-1063, doi/10.1016/j.bbabbio.2006.05.020.

810 Brown KA, Guo Z, Tokmina-Lukaszewska M, Scott LW, Lubner CE, Smolinski S, Mulder DW,
811 Bothner B, King PW (2019) The oxygen reduction reaction catalyzed by *Synechocystis* sp. PCC
812 6803 flavodiiron proteins. *Sustain Energy Fuels* 3: 3191-3200, doi/10.1039/c9se00523d.

813 Bulychev AA, Cherkashin AA, Muronets EM, Elanskaya IV (2018) Photoinduction of electron
814 transport on the acceptor side of PSI in *Synechocystis* PCC 6803 mutant deficient in flavodiiron
815 proteins Flv1 and Flv3. *Biochim Biophys Acta Bioenerg* 1859, 1086-1095,
816 doi/10.1016/j.bbabi.2018.06.012.

817 Burnap RL, Hagemann M, Kaplan A (2015) Regulation of CO₂ concentrating mechanism in
818 cyanobacteria. *Life (Basel)* 5: 348-371.

819 Burnap RL, Nambudiri R, Holland S (2013) Regulation of the carbon-concentrating mechanism in
820 the cyanobacterium *Synechocystis* sp PCC6803 in response to changing light intensity and inorganic
821 carbon availability. *Photosynth Res.* 118, 115-124, doi/10.1007/s11120-013-9912-4.

822 Cassier-Chauvat C, Chauvat F (2014) Function and regulation of ferredoxins in the cyanobacterium,
823 *Synechocystis* PCC6803: recent advances. *Life* 4: 666–680.

824 Chaux F, Burlacot A, Mekhalfi M, Auroy P, Blangy S, Richaud P, Peltier G (2017). Flavodiiron
825 Proteins Promote Fast and Transient O₂ Photoreduction in *Chlamydomonas*. *Plant Physiol* 174:
826 1825-1836. doi/10.1104/pp.17.00421.

827 Cooley JW, Vermaas W (2001) Succinate dehydrogenase and other respiratory pathways in
828 thylakoid membranes of *Synechocystis* sp strain PCC 6803: Capacity comparisons and
829 physiological function. *J Bacteriol* 183: 4251-4258, doi/10.1128/JB.183.14.4251-4258.2001.

830 Cruz JA, Avenson TJ, Kanazawa A, Takizawa K, Edwards GE, Kramer DM (2005) Plasticity in
831 light reactions of photosynthesis for energy production and photoprotection. *J Exp Bot* 56: 395-406.

832 Daley SM, Kappell AD, Carrick MJ, Burnap RL (2012) Regulation of the cyanobacterial CO₂-
833 concentrating mechanism involves internal sensing of NADP⁺ and α -ketoglutarate levels by
834 transcription factor CcmR. *PLoS One* 7: e41286. doi/10.1371/journal.pone.0041286.

835 Dann M, Leister D (2019) Evidence that cyanobacterial Sll1217 functions analogously to PGRL1 in
836 enhancing PGR5-dependent cyclic electron flow. *Nat Commun* 10: 5299. doi/10.1038/s41467-019-
837 13223-0.

838 Eisenhut M, Georg J, Klähn S, Sakurai I, Mustila H, Zhang P, Hess WR, Aro EM (2012) The
839 antisense RNA As1_flv4 in the Cyanobacterium *Synechocystis* sp. PCC 6803 prevents premature
840 expression of the flv4-2 operon upon shift in inorganic carbon supply. *J Biol Chem* 287: 33153-
841 33162.

842 Ermakova M, Huokko T, Richaud P, Bersanini L, Howe CJ, Lea-Smith DJ, Peltier G,
843 Allahverdiyeva Y (2016) Distinguishing the Roles of Thylakoid Respiratory Terminal Oxidases in
844 the Cyanobacterium *Synechocystis* sp PCC 6803. *Plant Physiol* 171: 1307-1319.
845 doi/10.1104/pp.16.00479.

846 Gao F, Zhao J, Chen L, Battchikova N, Ran Z, Aro EM, Ogawa T, Ma W (2016) The NDH-1L-PSI
847 Supercomplex Is Important for Efficient Cyclic Electron Transport in cyanobacteria. *Plant Physiol*
848 172: 1451-1464. doi/10.1104/pp.16.00585.

849 Genty B, Briantais JM, Baker NR (1989) The Relationship between the Quantum Yield of
850 Photosynthetic Electron-Transport and Quenching of Chlorophyll Fluorescence. *Biochim Biophys*
851 *Acta* 990: 87-92. doi/10.1016/S0304-4165(89)80016-9.

852 Gerotto C, Alboresi A, Meneghesso A, Jokel M, Suorsa M, Aro EM, Morosinotto T (2016)

853 Flavodiiron proteins act as safety valve for electrons in *Physcomitrella patens*. *Proc Natl Acad Sci*

854 USA 113: 12322-12327, doi/10.1073/pnas.1606685113.

855 Gómez R, Carrillo N, Morelli MP, Tula S, Shahinnia F, Hajirezaei MR, Lodeyro AF (2018) Faster

856 photosynthetic induction in tobacco by expressing cyanobacterial flavodiiron proteins in

857 chloroplasts. *Photosynth Res* 146: 129-136.

858 Hanke GT, Satomi Y, Shinmura K, Takao T, Hase T (2011) A screen for potential ferredoxin

859 electron transfer partners uncovers new, redox dependent interactions. *Biochim Biophys Acta*

860 Proteins Proteomics 1814: 366-374.

861 Han X, Sun N, Xu M, Mi H (2017) Co-ordination of NDH and Cup proteins in CO₂ uptake in

862 cyanobacterium *Synechocystis* sp PCC 6803. *J Exp Bot* 68: 3869-3877. doi/10.1093/jxb/erx129.

863 He Z, Zheng F, Wu Y, Li Q, Lv J, Fu P, Mi H (2015) NDH-1L interacts with ferredoxin via the

864 subunit NdhS in *Thermosynechococcus elongatus*. *Photosynth Res* 126: 341-349,

865 doi/10.1007/s11120-015-0090-4.

866 Helman Y, Tchernov D, Reinhold L, Shibata M, Ogawa T, Schwarz R, Ohad I, Kaplan A (2003)

867 Genes encoding a-type flavoproteins are essential for photoreduction of O₂ in cyanobacteria. *Curr*

868 *Biol* 13: 230-235, doi/10.1016/S0960-9822(03)00046-0.

869 Holland SC, Kappell AD, and Burnap RL (2015) Redox changes accompanying inorganic carbon

870 limitation in *Synechocystis* sp. PCC 6803. *Biochim Biophys Acta* 1847: 355-363.

871 doi/10.1016/j.bbabi.2014.12.001.

872 Ilik P, Pavlovic A, Kouril R, Alboresi A, Morosinotto T, Allahverdiyeva Y, Aro EM, Yamamoto H,
873 Shikanai T (2017) Alternative electron transport mediated by flavodiiron proteins is operational in
874 organisms from cyanobacteria up to gymnosperms. *New Phytol* 214: 967-972.
875 doi/10.1111/nph.14536.

876 Jokel M, Johnson X, Peltier G, Aro EM, Allahverdiyeva Y (2018). Hunting the main player
877 enabling *Chlamydomonas reinhardtii* growth under fluctuating light. *Plant J* 94: 822-835,
878 doi/10.1111/tpj.13897.

879 Jokel M, Nagy V, Tóth SZ, Kosourov S, Allahverdiyeva Y (2019) Elimination of the flavodiiron
880 electron sink facilitates long-term H₂ photoproduction in green algae. *Biotechnol Biofuels* 12: 280.
881 doi/10.1186/s13068-019-1618-1.

882 Kaplan A, Reinhold L (1999) CO₂ concentrating mechanisms in photosynthetic microorganisms.
883 *Annu Rev Plant Physiol Plant Mol Biol* 50: 539-570. doi/10.1146/annurev.arplant.50.1.539.

884 Kauny J, Setif P (2014) NADPH fluorescence in the cyanobacterium *Synechocystis* sp PCC 6803:
885 A versatile probe for in vivo measurements of rates, yields and pools. *Biochim Biophys Acta*
886 *Bioenerg* 1837: 792-801. doi/10.1016/j.bbabbio.2014.01.009.

887 Klughammer C, Schreiber U (2008) Saturation Pulse method for assessment of energy conversion
888 in PSI. *PAM Application Notes* 1: 11-14.

889 Klughammer C, Schreiber U (2016) Deconvolution of ferredoxin, plastocyanin, and P700
890 transmittance changes in intact leaves with a new type of kinetic LED array spectrophotometer.
891 *Photosynth Res* 128: 195-214.

892 Kramer DM, Avenson TJ, Edwards GE (2004) Dynamic flexibility in the light reactions of
 893 photosynthesis governed by both electron and proton reactions. *Trends Plant Sci* 9: 349-357.

894 Laughlin TG, Bayne AN, Trempe JF, Savage DF, Davies KM (2019) Structure of the complex I-
 895 like molecule NDH of oxygenic photosynthesis. *Nature* 566: 411–414.

896 Liran O, Shemesh E, Tchernov D (2018) Investigation into the CO₂ concentrating step rates within
 897 the carbon concentrating mechanism of *Synechocystis* sp PCC6803 at various pH and light
 898 intensities reveal novel mechanistic properties. *Algal Res* 33: 419-429.
 899 doi/10.1016/j.algal.2018.06.020.

900 Liu L, Bryan SJ, Huang F, Yu J, Nixon PJ, Rich PR, Mullineaux CW (2012) Control of electron
 901 transport routes through redox-regulated redistribution of respiratory complexes. *Proc Natl Acad*
 902 *Sci USA* 109: 11431-11436. doi/10.1073/pnas.1120960109.

903 Mehler AH (1957) Studies on reactions of illuminated chloroplasts. I. Mechanism of the reduction
 904 of oxygen and other Hill reagents. *Arch Biochem Biophys* 33: 65–77.

905 Ma W, Ogawa T (2015) Oxygenic Photosynthesis-specific Subunits of Cyanobacterial NADPH
 906 Dehydrogenases. *IUBMB Life* 67: 3-8. doi/10.1002/iub.1341.

907 McCormick AJ, Bombelli P, Lea-Smith DJ, Bradley RW, Scott AM, Fisher AC, Smith AG, Howe
 908 CJ (2013) Hydrogen production through oxygenic photosynthesis using the cyanobacterium
 909 *Synechocystis* sp PCC 6803 in a bio-photoelectrolysis cell (BPE) system. *Energy Environ Sci* 6:
 910 2682-2690. doi/10.1039/c3ee40491a.

911 Mullineaux CW (2014) Co-existence of photosynthetic and respiratory activities in cyanobacterial
912 thylakoid membranes. *Biochim Biophys Acta Bioenerg* 1837: 503-511.
913 doi/10.1016/j.bbabbio.2013.11.017.

914 Mustila H, Paananen P, Battchikova N, Santana-Sanchez A, Muth-Pawlak D, Hagemann M, Aro
915 EM, Allahverdiyeva Y (2016) The Flavodiiron Protein Flv3 Functions as a Homo-Oligomer During
916 Stress Acclimation and is Distinct from the Flv1/Flv3 Hetero-Oligomer Specific to the O-2
917 Photoreduction Pathway. *Plant Cell Physiol* 57: 1468-1483. doi/10.1093/pcp/pcw047.

918 Nakamura Y, Kaneko T, Miyajima N, Tabata S (1999) Extension of CyanoBase. CyanoMutants:
919 repository of mutant information on *Synechocystis* sp strain PCC6803. *Nucleic Acids Res* 27: 66-
920 68. doi/10.1093/nar/27.1.66.

921 Nikkanen L, Toivola J, Trotta A, Guinea Diaz M, Tikkanen M, Aro EM, Rintamäki E (2018)
922 Regulation of cyclic electron flow by chloroplast NADPH-dependent thioredoxin system. *Plant*
923 *Direct* 2: 1-24.

924 Ogawa T (1991) A gene homologous to the subunit-2 gene of NADH dehydrogenase is essential to
925 inorganic carbon transport of *Synechocystis* PCC6803. *Proc Natl Acad Sci USA* 88: 4275-4279.

926 Ohkawa H, Price D, Badger M, Ogawa T (2000a) Mutation of *ndh* genes leads to inhibition of CO₂
927 uptake rather than HCO₃⁻ uptake in *Synechocystis* sp. strain PCC 6803 *J Bacteriol* 182: 2591-2596.

928 Ohkawa H, Pakrasi HB, Ogawa T (2000b) Two types of functionally distinct NAD(P)H
929 dehydrogenases in *Synechocystis* sp strain PCC6803. *J Biol Chem* 275: 31630-31634.
930 doi/10.1074/jbc.M003706200.

931 Peltier G, Aro EM, Shikanai T (2016) NDH-1 and NDH-2 Plastoquinone Reductases in Oxygenic
932 Photosynthesis. *Annu Rev Plant Biol* 67: 55-80.

933 Santana-Sanchez A, Solymosi D, Mustila H, Bersanini L, Aro EM, Allahverdiyeva Y (2019)
934 Flavodiiron proteins 1-to-4 function in versatile combinations in O₂ photoreduction in
935 cyanobacteria. *eLife* 8: e45766. doi/10.7554/eLife.45766.

936 Saura P, Kaila VRI (2019) Molecular dynamics and structural models of the cyanobacterial NDH-1
937 complex. *Biochim Biophys Acta Bioenerg* 1860: 201-208. doi/10.1016/j.bbabi.2018.11.010.

938 Schreiber U, Klughammer C (2009) New NADPH/9-AA module for the DUAL-PAM-100:
939 Description, operation and examples of application. *PAM Application Notes* 2: 1-13.

940 Schreiber U (2017) Redox changes of ferredoxin, P700, and plastocyanin measured simultaneously
941 in intact leaves. *Photosynth Res* 134: 343-360.

942 Schuller JM, Birrell JA, Tanaka H, Konuma T, Wulflhorst H, Cox N, Schuller SK, Thiemann J,
943 Lubitz W, Setif P, Ikegami T, Engel BD, Kurisu G, Nowaczyk MM (2019) Structural adaptations
944 of photosynthetic complex I enable ferredoxin-dependent electron transfer. *Science* 363: 257-260.
945 doi/10.1126/science.aau3613.

946 Setif P, Boussac A, Krieger-Liszkay A (2019) Near-infrared in vitro measurements of photosystem
947 I cofactors and electron-transfer partners with a recently developed spectrophotometer. *Photosynth*
948 *Res* 142: 307-319. doi/10.1007/s11120-019-00665-2.

949 Shibata M, Ohkawa H, Kaneko T, Fukuzawa H, Tabata S, Kaplan A, Ogawa T (2001) Distinct
950 constitutive and low-CO₂-induced CO₂ uptake systems in cyanobacteria: Genes involved and their

951 phylogenetic relationship with homologous genes in other organisms. Proc Natl Acad Sci USA 98:
952 11789-11794. doi/10.1073/pnas.191258298.

953 Shibata M, Katoh H, Sonoda M, Ohkawa H, Shimoyama M, Fukuzawa H, Kaplan A, Ogawa T
954 (2002) Genes essential to sodium-dependent bicarbonate transport in cyanobacteria - Function and
955 phylogenetic analysis. J Biol Chem 277: 18658-18664, doi/10.1074/jbc.M112468200.

956 Shimakawa G, Shaku K, Miyake C (2016) Oxidation of P700 in Photosystem I Is Essential for the
957 Growth of Cyanobacteria. Plant Physiol 172: 1443-1450. doi/10.1104/pp.16.01227.

958 Shimakawa G, Miyake C (2018a) Changing frequency of fluctuating light reveals the molecular
959 mechanism for P700 oxidation in plant leaves. Plant Direct 2: e00073.

960 Shimakawa G, Miyake C (2018b) Oxidation of P700 Ensures Robust Photosynthesis. Front Plant
961 Sci 9: 1617. doi/10.3389/fpls.2018.01617.

962 Strand DD, D'Andrea L, Bock R (2019) The plastid NAD(P)H dehydrogenase-like complex:
963 structure, function and evolutionary dynamics. Biochem J 476: 2743-2756.
964 doi/10.1042/BCJ20190365.

965 Strand DD, Fisher N, Kramer DM (2017) The higher plant plastid NAD(P)H dehydrogenase-like
966 complex (NDH) is a high efficiency proton pump that increases ATP production by cyclic electron
967 flow. J Biol Chem 292: 11850-11860.

968 Thiel K, Patrikainen P, Nagy C, Fitzpatrick D, Pope N, Aro EM, Kallio P (2019) Redirecting
969 photosynthetic electron flux in the cyanobacterium *Synechocystis* sp. PCC 6803 by the deletion of
970 flavodiiron protein Flv3. Microb Cell Fact 18: 189-2. doi/10.1186/s12934-019-1238-2

971 Tikkanen M, Mekala NR, Aro EM (2014) Photosystem II photoinhibition-repair cycle protects
972 Photosystem I from irreversible damage. *Biochim Biophys Acta* 1837: 210-215.
973 doi/10.1016/j.bbabi.2013.10.001.

974 Tiwari A, Mamedov F, Grieco M, Suorsa M, Jajoo A, Styring S, Tikkanen M, Aro EM (2016)
975 Photodamage of iron-sulphur clusters in photosystem I induces non-photochemical energy
976 dissipation. *Nat Plants* 2: 16035.

977 Ueda M, Kuniyoshi T, Yamamoto H, Sugimoto K, Ishizaki K, Kohchi T, Nishimura Y, Shikanai T
978 (2012) Composition and physiological function of the chloroplast NADH dehydrogenase-like
979 complex in *Marchantia polymorpha*. *Plant J* 72: 683-693, doi/10.1111/j.1365-3113.2012.05115.x.

980 Vicente JB, Gomes CM, Wasserfallen A, Teixeira M (2002) Module fusion in an A-type
981 flavoprotein from the cyanobacterium *Synechocystis* condenses a multiple-component pathway in a
982 single polypeptide chain. *Biochem Biophys Res Commun* 294: 82-87. doi/10.1016/S0006-
983 291X(02)00434-5.

984 Viola S, Bailleul B, Yu J, Nixon P, Selles J, Joliot P, Wollman F (2019) Probing the electric field
985 across thylakoid membranes in cyanobacteria. *Proc Natl Acad Sci USA* 116: 21900-21906.
986 doi/10.1073/pnas.1913099116.

987 Williams JGK (1988) Construction of Specific Mutations in Photosystem-II Photosynthetic
988 Reaction Center by Genetic-Engineering Methods in *Synechocystis*-6803. *Meth Enzymol*: 167.
989 766-778.

990 Xu M, Bernát G, Singh A, Mi H, Rögner M, Pakrasi HB, Ogawa T (2008) Properties of mutants of
991 *Synechocystis* sp. strain 6803 lacking inorganic carbon sequestration systems. *Plant Cell Physiol*
992 49: 1672-1677. doi/10.1093/pcp/pcn139.

993 Yamamoto H, Takahashi S, Badger MR, Shikanai T (2016) Artificial remodelling of alternative
 994 electron flow by flavodiiron proteins in Arabidopsis. Nat Plants 2: 16012. doi/
 995 10.1038/nplants.2016.12.

996 Yamori W, Sakata N, Suzuki Y, Shikanai T, Makino A (2011) Cyclic electron flow around
 997 photosystem I via chloroplast NAD(P)H dehydrogenase (NDH) complex performs a significant
 998 physiological role during photosynthesis and plant growth at low temperature in rice. Plant J 68:
 999 966-976.

1000 Yamori W, Shikanai T, Makino A (2015) Photosystem I cyclic electron flow via chloroplast NADH
 1001 dehydrogenase-like complex performs a physiological role for photosynthesis at low light. Sci Rep
 1002 5: 15593.

1003 Yamori W, Shikanai T (2016) Physiological Functions of Cyclic Electron Transport Around
 1004 Photosystem I in Sustaining Photosynthesis and Plant Growth. Annu Rev Plant Biol 67: 81-106.

1005 Yeremenko N, Jeanjean R, Prommeeenate P, Krasikov V, Nixon PJ, Vermaas W, Havaux M,
 1006 Matthijs H (2005) Open reading frame *ssr2016* is required for antimycin A-sensitive photosystem I-
 1007 driven cyclic electron flow in the cyanobacterium *Synechocystis* sp PCC 6803. Plant Cell Physiol
 1008 46: 1433-1436, doi/10.1093/pcp/pci147.

1009 Zhang P, Battchikova N, Jansen T, Appel J, Ogawa T, Aro EM (2004) Expression and functional
 1010 roles of the two distinct NDH-1 complexes and the carbon acquisition complex
 1011 NdhD3/NdhF3/CupA/Sll1735 in *Synechocystis* sp PCC 6803. Plant Cell 16: 3326-3340.
 1012 doi/10.1105/tpc.104.026526.

1013 Zhang P, Allahverdiyeva Y, Eisenhut M, Aro EM (2009) Flavodiiron Proteins in Oxygenic
 1014 Photosynthetic Organisms: Photoprotection of Photosystem II by Flv2 and Flv4 in *Synechocystis* sp
 1015 PCC 6803. PLoS One 4: e5331. doi/10.1371/journal.pone.0005331.

1016 Zhao J, Gao F, Fan DY, Chow WS, Ma W (2017) NDH-1 Is Important for Photosystem I function
 1017 of *Synechocystis* sp. strain PCC 6803 under environmental stress conditions. Front Plant Sci 8:
 1018 2183. doi/10.3389/fpls.2017.02183.

1019

1020