

Mucoidy, a general mechanism for maintaining lytic phage in populations of bacteria

Waqas Chaudhry¹, Esther Lee¹, Andrew Worthy¹, Zoe Weiss¹, Marcin Grabowicz^{2,3,4}, Nicole Vega^{1*}, Bruce Levin^{1,2**}

¹ Department of Biology, Emory University, Atlanta, GA 30322, USA.

² Emory Antibiotic Resistance Center, Emory University School of Medicine, Atlanta, GA 30322

³ Department of Microbiology & Immunology, Emory University School of Medicine, Atlanta, GA 30322, USA.

⁴ Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, GA 30322, USA.

[*nvega@emory.edu](mailto:nvega@emory.edu)

[**blevin@emory.edu](mailto:blevin@emory.edu)

Abstract

We present evidence that mucoidy provides a new and general answer to the longstanding question of how lytic phage are maintained in bacterial populations. In serial transfer culture, populations of mucoid *E. coli* MG1655 that appear to be broadly resistant to lytic phage with different receptors are capable of maintaining these phage with little effect on the total density of bacteria. Based on the results of our analysis of a mathematical model, we postulate that the maintenance of phage in populations dominated by mucoid cells can be attributed primarily to high rates of transition from the effectively resistant mucoid states to susceptible non-mucoid states. Our test of this “leaky resistance” hypothesis with both population dynamic and single cell experiments are consistent with it. We discuss reasons for the generalized resistance of these mucoid *E. coli*, and the genetic and molecular mechanisms responsible for the high rate of transition from mucoid to sensitive states.

Introduction

Bacteriophages are natural predators (parasitoids) of bacteria and are present in natural microbial communities across environments, from marine and aquatic systems [1] to the intestine of the mammalian host [2]. By mutation, bacteria can become resistant [3] or, in the case of CRISPR-Cas, immune to these viruses [4]. Most commonly these resistant or immune cells ascend to dominate the bacterial population, but the phage continue to be maintained [5-9]. How virulent phage are maintained in populations of bacteria dominated by cells upon which they cannot replicate is a longstanding question that has been addressed theoretically as well experimentally [3, 10-13]. Thanks to concerns about the factors responsible for maintaining and stabilizing the structure of bacterial communities, microbiomes, if you prefer [14, 15] and the resurrection of interest in phage therapy [16-19], this

question - and more broadly, studies of the population and evolutionary dynamics of the interactions between bacteria and phage - have become an increasingly important and relevant avenue of research.

Those who work with *E. coli* and other Gram negative bacteria occasionally see shiny, “snotty” looking “mucoïd” colonies on agar that can be attributed to contamination with bacteriophage. By providing a physical barrier between the cell surface and the would-be infecting phage [20-24], the overproduction of the exopolysaccharide responsible for the mucoïd phenotype confers resistance to the phage. While this resistance may not be as complete as that generated by the modification of the phage receptors [25] and may engender a substantial fitness cost [26], it is sufficient to select for mucoïdy [26-28] in populations of bacteria exposed to lytic phage. Moreover, and most importantly from an ecological perspective, the resistance provided by mucoïdy is general rather than specific, like the resistance generated by modification of the phage receptors; mucoïd bacteria are resistant to multiple lytic phages [26].

These observations raise several questions about the population dynamics and ecology of mucoïd-based resistance. Under what conditions will mucoïdy be selected for over surface (envelope) or other resistance or immune mechanisms [23]? How might the ecology of phage-bacterial system differ when resistance is conferred by mucoïdy rather than resistance? Why and how, in the absence of phage, do mucoïd-colony producing bacteria revert to states that produce non-mucoïd colonies? Most importantly, what is the contribution of mucoïdy to the coexistence of bacteria and lytic phage?

To address and answer these questions, we studied the population dynamics of mucoïd colony producing *E. coli* K12 strain MG1655 following exposure to different virulent (lytic) bacteriophage, T3, T4, T5, T7 and λ^{VIR} . We present evidence that mucoïdy provides a sufficient answer to the question of how lytic phage are maintained following the evolution of resistance and a general mechanism for the stable co-existence of bacteria with lytic phage. Based on the results of our analysis of mathematical/computer simulation models, we postulate that lytic phage can stably persist on populations of mucoïd *E. coli* by replicating on minority populations of sensitive bacteria that are maintained primarily if not exclusively by a high rate of transition from “resistant” mucoïd to sensitive non-mucoïd colony types. Using a combination of computer simulations, batch culture experiments, and single-cell microscopy, we present evidence in support of this “leaky resistance” [9, 13, 29] hypothesis for the maintenance of lytic phage in populations of mucoïd bacteria.

Results

I- Selection for, and population dynamics of, mucoïd *E. coli* in phage-infected cultures

First, we sought to determine whether lytic phages varied in their ability to select for mucoïdy in *E. coli* MG1655, and whether these phage-selected mucoïds differed

from one another in their interactions with phage. For this, wild type cultures of *E. coli* MG1655 were exposed to T3, T4, T5, or T7 phage and serially transferred over ten passages (Figure 1). For cultures initiated with T3 or T4, the phage were maintained over all ten transfers and the dominant population of bacteria isolated at the end of the final transfer were mucoid (Figure 1A and 1B). All replicates of the experiments using T5 resulted in the extinction of the phage following the emergence of envelope resistance (Figure 1C). Through genetic mapping we present evidence that this resistance was attributed to mutations in the *fhuA* locus which encodes the T5 receptor, also see [30]. For T7, the phage were maintained in one of the three populations, where mucoid cells were the dominant population of bacteria. In the remaining two T7 cultures, neither mucoidy nor resistance were generated and the bacteria were rapidly driven to extinction, as were the phage that no longer had hosts (Figure 1D).

Second, we explored the capacity of these phage to be maintained in populations of mucoid bacteria. For the first of these serial transfer experiments, we used the mucoids generated by selection with the phage T7. As seen in Figure 2, all four phage were maintained, and the densities of viable bacteria were high, similar to that of the phage-free controls. Stated another way, resources rather than phage limited the densities of these bacterial populations. Moreover, as can be seen in Figure S1, this result obtains in minimal medium as well as the LB used for the Figure 1 experiment. In the minimal medium experiments, however, mutants with envelope resistance arose in mucoid cultures treated with phage T4 and T5.

These results support the initial assumption that phage-selected mucoidy in *E. coli* provides broad resistance to virulent phage and indicate that co-existence of large populations of virulent phage and mucoid *E. coli* is a general phenomenon that does not depend on the phage initially used for selection. They also suggest that the capacity to select for mucoidy and/or the rate of mutation to mucoidy varied among the phages. These results also suggest that mucoidy can prevail over envelope resistance, but may not do so in all cases, depending on the environment and on the lytic phage used.




Figure 1: The population dynamics of T3 (A), T4 (B), T5 (C) and T7 (D) phage with wild-type *E. coli* MG1655. Cultures were diluted 1:100 into fresh LB medium every 24 hours; three replicates were performed per condition. Solid lines show bacterial densities in each replicate (red/green/blue); dotted lines show the phage densities in the corresponding cultures. Phage-free cultures (black and purple lines) are shown as controls, to indicate resource-limited population size for the bacteria.




Figure 2: The population dynamics of phage T3 (A), T4 (B), T5(C) and T7 (D) phage with T7-selected mucooid *E. coli* MG1655 as the host strain in LB. Three independent cultures were diluted 1:100 into fresh LB media every 24 hours, and CFU/PFU estimates of the densities of bacteria and phage were made by serial dilution plating. Solid lines show bacterial densities in each replicate (red/green/blue); dotted lines show the phage densities in the corresponding cultures. Phage-free cultures (black and purple lines) are shown as controls, to indicate resource-limited population density for the bacteria. Phage-free controls lost the mucooid phenotype entirely by the end of the 4th serial transfer. At the end of the 15th serial transfer, three colonies from each phage-infected culture were streaked at least three times to get rid of phage, yielding non-mucooid revertants that were again sensitive to the relevant phage.

II- Mechanism of phage-selected mucooidy

Next, we investigated the ability of defined mutations in the *E. coli* colanic acid synthesis pathway to recapitulate the observed ecological interactions between bacteria and phage, specifically the high-level maintenance of phage on apparently resistant hosts. The “regulator of capsule synthesis” (Rcs) phosphorelay system regulates colanic acid capsule production in *E. coli* [31]. The Rcs phosphorelay is complex, but at its core: RcsF is a sensory lipoprotein that causes activation of the

RcsC sensor kinase in an environment-dependent manner; RcsC causes phosphorylation of the RcsB transcription factor; phosphor-RcsB is a positive regulator of the colanic acid biosynthetic operon at the *cpsB* promoter. Overproduction of colanic acid via this pathway results in the distinctive mucoid colony phenotype, and mucoid strains featuring mutations in this pathway are selected for by exposure to virulent phage in *E. coli* MG1655 [26].

To understand the dynamics of mucoid selection and the role of mucoidy in phage maintenance, we sought to determine whether EPS-impaired strains, which do not produce mucoid colonies, would be able to maintain phage populations during serial passage. To this end, we tested two gene knockout strains in the Rcs signaling pathway ($\Delta rcsB$ and $\Delta rcsF$) as well as one gene knockout strain in the EPS operon ($\Delta cpsB$), all of which are impaired in capsule synthesis. We also tested a strain (*rscC137 ompC::Tn5*) that has an activating A904V substitution in RcsC and is therefore constitutively mucoid [32], to determine if the simple induction of mucoidy via a known mechanism was sufficient to allow maintenance of phage. We found that the *rscC137* constitutive-mucoid mutant was able to maintain large populations of phage and bacteria for all phage tested, while phage exposure in loss-of-function mutants impaired in mucoidy ($\Delta rcsB$, $\Delta rcsF$, $\Delta cpsB$) resulted in extinction of both phage and bacterial populations when T3 phage were used (Fig. 3). Exposure to T4 or T7 phage resulted in bacterial population collapse in some conditions but not others, largely dependent on whether evolution of envelope resistance occurred; in the $\Delta rcsF$ background, mucoid colonies were observed, presumably downstream of a mutation compensating for the mucoidy defect in the knockout. Notably, the inactivating mutation in *ompC* in the *rscC137* constitutive mutant strain, which can confer envelope resistance to some phage [33], had no noticeable effect in these assays. For reasons that are not yet clear, T5 phage were maintained on all host backgrounds regardless of mucoidy, despite the fact that evolution of envelope resistance generally leads to extinction of this phage (Fig. 1).




Figure 3. Constitutive presence or absence of mucoidy affects the dynamics of *E. coli*-phage interactions. T3, T4, T5, or T7 phage were co-cultured with genetically modified strains of *E. coli* MG1655 with known gain or loss of function in the Rcs pathway. Strains used were: $\Delta rcsF::kan$ (pathway loss-of-function knockout of the outer membrane Rcs input), $\Delta rcsB::kan$ (pathway loss-of-function knockout of the RcsB transcription factor), $\Delta cpsB::kan$ (pathway loss-of-function knockout of biosynthesis of exopolysaccharide colonic acid), and $rscC137 ompC::Tn5$ (gain-of-function point mutant that is a defined activating mutation of mucoidy). Cultures were diluted 1:100 into fresh LB medium every 24 hours; two replicates were performed per condition. Solid red and blue lines show bacterial densities, while dotted lines show the phage densities in the corresponding cultures. Phage-free cultures are shown as controls (black and purple lines), to indicate resource-limited population size for the bacteria.

III- A hypothesis for the maintenance of phage in mucoid populations of bacteria.

We now consider the population dynamics of lytic phage and mucoid hosts to address two questions. How are the phage maintained in mucoid populations of bacteria? Under what conditions will envelope resistant mutant bacteria emerge and replace the mucoid population?

We postulate that the maintenance of high densities of phage in populations of mucoid bacteria can be attributed to a substantial density of sensitive cells generated by a high rate of transition from the mucoid to non-mucoid states, a mechanism that has been called “leaky” resistance [9]. To illustrate how this mechanism can account for the observed maintenance of high densities of phage with little effect on the density of bacteria, and for the common failure to see the ascent of mutants with surface resistance, we use a simple mass-action model of the population dynamics of bacteria and phage (Fig. 5, Equations 1-5). In this model, there are three populations of bacteria – sensitive (S), mucoid (M), and resistant (N_R), and a single population of lytic phage (P), where S, M, N_R , and P are the densities, cells or particles per ml, as well as the designations of these populations.

The bacteria grow at maximum rates v_S , v_M and v_{NR} ($\text{cell}^{-1} \text{hr}^{-1}$) respectively (S2 Fig), with the realized growth rates dependent on the concentration of the shared limiting resource (r , $\mu\text{g}/\text{mL}$) via a hyperbolic function $\psi(y)=r/(r+k)$ [34], where the Monod constant k ($\mu\text{g}/\text{mL}$) is the concentration of limiting resource where the growth rates are half their maximum value. The phage adsorb to the sensitive, S, and mucoid, M, cells respectively with rate constants δ_S and δ_M ($\text{hr}^{-1}\text{mL}^{-1}$) but do not adsorb to the envelope-resistant bacteria N_R . Upon adsorption, both mucoid and sensitive cells produce β phage particles (neglecting the latent period, for simplicity). Transition from sensitive to mucoid states $S \rightarrow M$ occurs at a rate x , and from mucoid to sensitive $M \rightarrow S$ at a rate y , both per cell per hour with all rates proportional to $\psi(r)$. As in [9] we assume that the limiting resource is taken up at a rate proportional to the net growth rate of each bacterial population, modified by the resource conversion parameter e (μg), which is the amount of limiting resource needed to produce a new cell [35]. Serial transfer is simulated by diluting the

population of bacteria and phage by a factor $d = 0.01$ every 24 hours, removing the existing resource, and adding $C \mu\text{g}/\text{ml}$ of the limiting resource.




Figure 4: Diagram of the mass-action model, showing transitions between states. Bacteria may be mucoid (M), sensitive (S) or envelope-resistant (N_R). Mucoid and sensitive cells adsorb phage (P), but resistant cells do not. See the text and methods for full definitions of the parameters, variables, and assumptions behind the construction of this model. The resource, r , is not considered in this diagram, but is included in the numerical solutions to these equations, which simulates a serial transfer mode of population maintenance.

$$\frac{dr}{dt} = \underbrace{-\psi(r) \cdot e \cdot (v_S \cdot S + v_M \cdot M + v_R \cdot N_R)}_{\text{Resource Uptake}} \quad (1)$$

$$\frac{dS}{dt} = \underbrace{v_S \cdot S \cdot \psi(r)}_{\text{Growth of } S} - \underbrace{\delta_S \cdot P \cdot S \cdot \psi(r)}_{\text{Loss of } S \text{ by phage infection}} + \underbrace{\psi(r) \cdot (-y \cdot S + x \cdot M)}_{\text{Transitions of } S \text{ to } M \text{ and from } M \text{ to } S} \quad (2)$$

$$\frac{dM}{dt} = \underbrace{v_M \cdot M \cdot \psi(r)}_{\text{Growth of } M} - \underbrace{\delta_M \cdot M \cdot P \cdot \psi(r)}_{\text{Loss of } M \text{ by phage infection}} + \underbrace{\psi(r) \cdot (y \cdot S - x \cdot M)}_{\text{Transitions of } M \text{ to and from different states}} \quad (3)$$

$$\frac{dN_R}{dx} = \underbrace{v_R \cdot \psi(r) \cdot N_R}_{\text{Growth of } N_R} \quad (4)$$

$$\frac{dP}{dt} = \underbrace{\delta_S \cdot P \cdot S \cdot \beta \cdot \psi(r)}_{\text{Phage replication by infecting } S} + \underbrace{\delta_M \cdot P \cdot M \cdot \beta \cdot \psi(r)}_{\text{Phage replication by infecting } M} \quad (5)$$

where $\psi(r) = \frac{r}{(r+k)}$

These equations were solved numerically and the dynamics were simulated with Berkeley MadonnaTM. The values of the phage infection parameters δ_s and β for the sensitive cells used in these simulations are in the range estimated in [9, 36] and the growth rates of the bacteria in the ranges estimated in this study (Figure S4). For the mucoids, we postulate the adsorption rates of the phage, which is too low to estimate directly because the rate of decline in phage density due to adsorption to mucoids is within the range of phage deactivation due to other processes in that media. We also postulate different transition rates between the states. Save for the phage λ , we are unable to directly estimate the rate of transition from mucoid to sensitive, $M \rightarrow S$, the parameter γ . In our analysis, we explore the sensitivity of the dynamics and predictions of the model to the values of the postulated parameters. Copies of the programs and instructions for their use are available from blevin@emory.edu.

Simulation Results.

In Figures 5A and 5B, we follow the changes in the densities of the bacterial and phage populations in simulated serial transfer populations initiated with phage and sensitive bacteria, the theoretical analogs of the serial transfer presented in in Figures 1 and 3. In Figure 5 C, D, E and F we follow the changes in the density of bacteria and phage in populations initially dominated by mucoid bacteria; the theoretical analog of the experiments presented in Figures 2. In all of these simulations we are assuming that sensitive cells are the most fit, resistant cells have the next highest fitness, and the mucoid cells have the lowest fitness, respectively $v_S=2.1$, $v_{NR}=2.0$, and $v_M=1.8$. This assumption is supported by our estimates of the maximum growth rates of these different cell lines, and consistent with the evidence present in [26] about mucoidy engendering a substantial fitness cost.

At the start of the simulations in Figure 5A, there are 10^6 phage, 10^7 sensitive cells, a single mucoid cell, no resistant mutants, and the resource is at its maximum level of $r=1000$. Within a single transfer the mucoid bacteria ascend to become the dominant population of bacteria and by the third transfer the population of bacteria and phage are at equilibrium. If mucoids are not present and cannot be generated ($\gamma=0$) but resistant mutants are present, they will ascend to become the dominant bacterial strain and the phage will be lost (Figure 5B). Although the dynamics are different than those observed in this simple model, there is qualitative agreement between the theory and experiments. In Figure 1 with T3 and T4, where mucoid bacteria ascended to dominate, the phage were maintained at high densities. In experiments with T5, resistance evolved instead of mucoidy and phage were lost. In one of the three experiments with T7, mucoidy evolved and phage were maintained. In two experiments with T7, neither mucoidy nor resistance evolved; the cells were killed by phage and the phage were washed out in subsequent transfers.

To get a better idea of the conditions required for mucoid bacteria to maintain the phage, and why resistant mutants may not have ascended to dominance in the serial

transfer experiments presented in Figure 2, we simulated situations where phage are introduced into populations of mucoid bacteria. At the start of these simulations there are 10^6 phage, 10^7 mucoid bacteria, single sensitive and resistant cells ($S=1$ and $N_R=1$), and the resource is present at its maximum level $r=1000$. The results of these simulations illustrate the critical role that the rate of transition between mucoids and sensitive bacteria, the parameter γ , plays in these dynamics. When $\gamma=10^{-3}$, within short order the phage established a stable population in a community dominated by mucoid bacteria (Figure 5C). When $\gamma=10^{-4}$, the phage become established, but their densities oscillate before the population of bacteria and phage becomes stable (Figures 5D and S2A). When $\gamma=10^{-5}$, the oscillations in the densities of the phage are greater (Figure 5E) and stability is not achieved (Figure S2C). Because the resistant mutants in these simulations are more fit than the mucoids, their densities increase at each transfer. However, during the course of these simulations, these resistant mutants may not reach levels where they affect the dynamics of the phage and are unlikely to be detected in the experiments. While this is consistent with some of the experiments where phage are introduced into mucoid populations, as noted in the legend to Figure S1 for both T4 and T5, by the end of the experiments, mutants resistant to these phage ascended and mucoidy declined. If the adsorption rate constant δ_M is increased to 10^{-10} , the phage are not maintained and the higher fitness resistant population ascends to dominate the bacterial population (Figure 5F).




Figure 5: The anticipated population dynamics of a lytic phage in serial transfer populations of sensitive, mucoid and resistant bacteria, simulation results. Changes in the densities of mucoid, M (purple), sensitive, S (blue), and envelope-resistant, N_R (green) bacteria and virulent phage, P (red) in populations initially dominated by mucoids. Standard parameters, $C=1000$, $e=5 \times 10^7$, $v_S=2.1$, $v_M=1.8$, and $v_R=2.0$, $k=1$, $\beta=50$, $x=10^{-5}$, $\delta_S=10^{-7}$. (A) Establishment of the mucoids in a population of sensitive bacteria and phage in the absence of resistance, $\gamma=10^{-3}$, $\delta_S=10^{-11}$. (B) Establishment of resistant mutants in a population that is unable to generate the mucoids. (C) Establishment of phage in a population dominated by mucoid cells and single

sensitive and resistant bacteria, $y=10^{-3}$, $\delta_S=10^{-11}$. (D) Establishment of phage in a population dominated by mucoid cells and single sensitive and resistant bacteria, $y=10^{-4}$, $\delta_S=10^{-11}$. (E) Establishment of phage in a population dominated by mucoid cells and single sensitive and resistant bacteria, $y=10^{-5}$, $\delta_S=10^{-11}$. (F) Establishment of phage in a population dominated by mucoid cells and single sensitive and resistant bacteria, $y=10^{-3}$, and $\delta_S=10^{-10}$.

In the simulations presented in Figure 5, two not mutually exclusive mechanisms can be responsible for the maintenance of phage. Both of these mechanisms involve the continuous production of populations of sensitive bacteria of sufficient density to support the replication of the phage population. In the case of resistance, the sensitive population is maintained because they are more fit than the resistant [6, 10, 30]. In the case of mucoidy, the sensitive cells are maintained because of a continuous transition from the mucoid to sensitive states, the “leaky mechanism”. The dynamics anticipated by these mechanisms are, however, different. This is illustrated in Figure S2.

With the standard parameters and initial conditions used in the 10-transfer simulations presented in Figure 5D, in the longer term the initial oscillations observed in the densities of the phage are damped and the phage are maintained by a minority population of sensitive cells sustained by the fitness cost of resistance, $v_R=2.0$ and $v_S=2.1$ (Figure S2A). If, however, the parameters are changed so that the phage cannot adsorb to the mucoid cells, $\delta_M=0$ rather than 10^{-11} , the oscillations in the densities of sensitive cells and phage continue but do not increase in amplitude (Figure S2B). If the rate of transition from sensitive to mucoid is reduced to $y=10^{-5}$, the amplitude of the oscillations in the densities of phage increases to the point where the phage are lost (Figure S2C). This instability of communities where phage are maintained by the fitness cost of resistance can be seen when the initial density of phage is less than that predicted for the equilibrium (Figure S2D). These oscillations of increasing amplitude do not obtain when the phage are maintained by the transition between mucoid and sensitive cells (Figure S2E). Finally and most critically, the leaky resistance mechanism for the maintenance of phage in a mucoid population does not require a fitness cost of the mucoid cells (Figure S2F).

IV. Test of the leaky resistance hypothesis

In accord with the model, for the sustained density of sensitive cells to be sufficiently great to maintain the phage at the high densities observed, the rate of transition or reversion from mucoid to sensitive non-mucoid bacteria has to be high, on the order of 10^{-4} to 10^{-3} per cell. The phage adsorption rate and fitness cost parameters also have to be in the range modeled for resistant cells not to ascend to dominate the culture and lead to the loss of the phage. More information about the range of parameter values needed to account for the population dynamics observed can be seen in Figure S2. What is clear from this sensitivity analysis is that for mucoids to dominate and the phage to be maintained, the rate parameter of adsorption of the phage to mucoids has to be low, $\delta_M < 10^{-11}$. In the following, we present two lines of evidence in support of the hypothesis that there is a high rate of transition from mucoid to sensitive, on the order of 10^{-3} per cell per hour.

(i) Population dynamic evidence for a high rate of reversion from mucoids to non-mucoids: One line of evidence for a high rate of transition from mucoid to sensitive non-mucoid states comes from population dynamics experiments using a method similar to that employed in [9] with genetically marked λ temperate phage, λ^{CAM} . This method is based on the premise that lysogens will only be formed in populations of resistant mucoid bacteria by infecting cells in minority population of susceptible non-mucoid bacteria. We first performed experiments demonstrating that like the T-phages, λ^{VIR} maintains high densities in mucoid populations of different types (Fig. S3); notably, as for T5 phage (Fig. 1), mucoidy does not arise in an initially sensitive wild-type population in this medium. We then introduced λ^{CAM} to cultures of T3- and T7-selected mucoids and the *rcsC137* constitutive-mucoid mutant. We followed the changes in the densities of free phage, the total densities of cells and, by selective plating (see the Materials and Methods), the lysogens and non-mucoid (sensitive) cells generated during the course of the experiment (Figure 6, A, B, and C). While the rates of accumulation of lysogens varied, these data indicate that reversion from mucoid to non-mucoid occurred frequently during growth for all strains.




Figure 6: Short-term population dynamics of lysogen formation. A, B and C: experimental results, changes in the densities of total cells NT, free λ^{CAM} phage P, lysogens L, and sensitive non-lysogens (revertants) S. (A) T3 selected mucoids, (B) T7 selected mucoids, (C) genetically constructed mucoid *E. coli* MG1655 (*rcsC137 ompC::Tn5*). SA, SB, and SC: simulations, changes in population density, standard parameters, $v_S=2.1$, $v_L=2.1$, $\delta = 2E10^{-7}$, $\delta_M = 10^{-11}$, $x=10^{-5}$, $C=1000$, $k=1$, $e=5 \times 10^{-7}$, probability of lysogeny, $\lambda = 0.01$, $i = 10^{-3}$. (SA) Standard parameters with $y=10^{-3}$,

-4

$v_M=1.8$, (SB) standard parameters with $y=10^{-3}$, $v_M=1.8$, (SC) Standard parameters with $y=10^{-3}$ and $v_M=1.5$.

To obtain a rough estimate of the rates of formation of lysogens, we used a simulation similar to that employed in [9] (see the supplemental material for the equations and the description of the model) as seen in Figure 6 SA, SB and SC. It is clear that this simple model is not a strictly accurate analogue of the short-term population dynamics of lysogen formation for several reasons. First, in the simulations, λ^{CAM} phage does not increase in density at a rate even close to that observed, nor do they reach as high a density. Next, save for the *rsc137* MUC strain, the predicted density of revertants (S) in simulation is lower than observed in experiments. Most importantly, the predicted rate of increase in the density of lysogens for the T3 and T7 selected mucoids is, with the M->S transition rate of 10^{-3} per hour, less than that observed experimentally and, for the *rsc137* mucoids, similar to that observed. Nevertheless, these simulations demonstrate that a high rate of transition from mucoidy is necessary to even approximate the observed trends in lysogen formation in an initially mucoid population of cells.

Additional evidence for a high rate of transition from mucoid to non-mucoid and presumably sensitive cells is presented in Figure S4A-D. In this experiment, we followed the numbers of mucoid and non-mucoid colonies in serial transfer culture. It is clear that for the T7 and T4-generated mucoids, there is a high rate of transition to non-mucoid. This is not the case for the T3-generated mucoids or the genetically constructed mucoids (*rscC137*). In Figure S4E we present the estimates of the maximum growth rates of wild-type MG1655, various mucoid lineages, and mutants with envelope resistance against T5 and λ^{VIR} . These data indicate that mucoidy does incur a substantial growth rate cost, as indicated by the model analysis. Furthermore, envelope resistance to T5 and λ^{VIR} incurs no such defect under these culture conditions, and may well be the reason why mucoidy is not observed when *E. coli* K12 is challenged with these phage (Figure 1, Figure S4 E-F).

Notably, we observed that these revertants to a non-mucoid phenotype were in some cases able to return to mucoidy upon re-exposure to phage (Figure S5).

(ii) Single-cell observations of mucoidy, reversion-induced heterogeneity, and phage infection. The second line of evidence for a high rate of transition from mucoid to normal colony sensitive cells comes from single-cell observation of phage infection in mucoid and non-mucoid cultures. Prior work has shown that reversion from phage-induced mucoidy occurs through secondary mutations within the Rcs system rather than through direct reversion of the initial mutation [26], and that these mutations restore expression of Rcs to roughly wild-type levels. If this is the case, it should be possible to identify individual mucoid cells based on expression levels in the Rcs pathway. To this end, we used a plasmid encoding a fluorescent reporter for Rcs expression (pBAD18Cm::P_{rprA}-mCherry) to visualize heterogeneity within initially mucoid populations (Figure 7, Figure S4). Mucoid and non-mucoid cells do not appear morphologically distinct on a single-cell level, and mucoids do not clump or agglomerate in well-mixed liquid culture, which might have protected

these cells against phage [37]. However, fluorescence expression from the Rcs promoter is clearly different between wild type and mucoid lineages.

If reversion to non-mucoid eliminates over-expression of Rcs and consequently increases sorption by virulent phage, this should be visible as differences in single-cell sorption to phage which anti-correlate with expression of the Rcs pathway. Using cells expressing the mCherry Rcs fluorescent reporter as described above, we infected cultures with GFP-fluorescent T7 phage [37]; the T7-derived mucoid lineage was chosen due to the rapid reversion to non-muoidy and high expected levels of heterogeneity in these cultures, and wild-type non-mucoid *E. coli* MG1655 was used for comparison. We observed that, in wild-type cultures of *E. coli* MG1655 where expression of Rcs is uniformly low, most to all individual cells adsorb phage and become infected (Figure 7A,B; Figure S6). By contrast, in T7-induced mucoid *E. coli*, a substantial minority population of low-Rcs cells, apparently corresponding to the revertants expected in this culture, become infected with T7 phage, while cells strongly expressing Rcs do not become infected (Fig. 7 C, D, S6 Fig).




Figure 7: Rcs pathway induction is negatively correlated with T7 phage infection on the single-cell level. In MG1655 wild-type non-mucoid culture (top), expression of Rcs (red) is low, and T7 phage infection (green) is universal. By contrast, in T7-induced mucoid culture (bottom), Rcs expression is high in the bulk of the population and low in a minority of putative revertants, and only the low-Rcs sub-

population is infected with T7 phage. Images represent separate aliquots from 1 mL LB cultures incubated at 37°C, sampled approximately 1 hour (left panels) and 2.5 hours (right panels) after inoculation. Images are an overlay of DIC, TRITC, and GFP channels, acquired using a 40X objective (total magnification 400X). Scale bar is 25 μm .

Discussion

A longstanding, fundamental question about the ecology, population and evolutionary biology of bacteria and phage is how are these viruses maintained in bacterial communities following the ascent and dominance of populations of bacteria upon which they cannot replicate [3, 6, 8, 10-13, 38]? The results of this experimental study with *E. coli* K12 and five of its lytic phage, T3, T4, T5, T7 and λ^{VIR} , provide a new and general answer to this question in the form of a long recognized mechanism of phage resistance, mucoidy. We present evidence that when serial transfer populations of *E. coli* are confronted with some but not all of these lytic phage, mucoid colony producing bacteria ascend and become the dominant population of bacteria. The phage continue to be maintained for extensive periods at high densities with the dominant mucoid population of bacteria at densities not much less than that of phage-free cultures with those bacteria. As demonstrated by Wielgloss and colleagues [26], the resistance provided by mucoidy is general; mucoid colony producing bacteria are resistant to phages with different receptor sites [39]. The results presented here indicate that whether the phage select for the mucoid bacteria they are cultured with or not, or whether these mucoid colony-producing *E. coli* are produced by genetic manipulation, all five lytic phage studied maintain stable populations with these mucoid bacteria.

Based on the results of our analysis of the properties of a mathematical model, we postulate that a sufficient, if not the unique, mechanism by which the phage are maintained in these mucoid populations of *E. coli* is a high rate of transition from the resistant, mucoid state to the susceptible state, “leaky resistance” [9, 13, 29]. We cannot, however, rule out some contribution of the fitness cost of mucoidy contributing to the maintenance of the phage [10, 30]. Central to the leaky resistance hypothesis is that the rate of transition from the mucoid to the susceptible state is high, on the order of 10^{-4} to 10^{-3} per cell per hour.

A priori, a transition rate between phenotypic states of bacteria of this great a magnitude by classical genetic processes is highly unlikely. To be sure, the large size of the Rcs regulon provides an abundance of mutational targets [26]. Nevertheless, *E. coli* is only expected to experience roughly one mutation per thousand cells per generation overall [40], which, no matter how big the target may be, means that a rate of 10^{-4} is too high to explain by base substitution mutation. Be that as it may, taken at large, the results of our experiments provide support for transition rates of this magnitude for at least some mucoid *E. coli*.

Consistent with other studies, we found that some mucoid phage-resistant mutants revert quickly to a non-mucoid, phage-sensitive state when the phage are removed, probably due to additional mutations elsewhere in the Rcs pathway rather than through reversion of the original mucoidy-inducing mutation [26]. Reversion

appeared to occur at different rates in *E. coli* mucoids obtained through exposure to different T phage, such that T3-derived mucoids appeared to be relatively stable and T4- and T7-derived mucoids reverted more rapidly. Observations of the growth rate defect in mucoid cells (Figure S5.) indicated that T3-derived mucoids had a strong growth defect, suggesting that the difference in observed reversion was not due to a lack of selective pressure for improved growth. It may therefore be the case that the specific mutation(s) that produce mucoidy in our T3-derived lineages are more difficult to reverse than the mutations that arise in other phage-selected mucoid lineages, or that direct reversion of the original mutation is required.

Also consistent with the postulated high transition from mucoid to susceptible are the results of the experiments we have done to estimate rate of production of lysogens for the phage λ . When genetically marked temperate λ are mixed with λ resistant mucoids populations generated in different ways, the rate of production of lysogens is consistent with the transition rate from mucoid to susceptible being on the order of 10^{-4} per cell per hour. While not providing estimates of the transition rate, additional support for this rate being high, comes from the experiments we did with single-cell microscopy. We demonstrate that propagation of phage in cultures of mucoid *E. coli* relies on infection and lysis of these revertant cells, which have restored wild-type levels of expression through the Rcs capsule synthesis system.

We observed that, in some cases, susceptible revertants from mucoidy were able to re-gain the mucoid phenotype upon re-exposure to phage. This is in contrast with a previous study which found that mutations leading to reversion of mucoidy prevented a return to the mucoid phenotype [26]. In that study, the tested non-mucoid revertants had acquired large gene deletions spanning the *rcs* operon or an IS insertion in *rscC*. The nature of these mutations likely explains why subsequent return to the mucoid phenotype was not possible: deletions are a loss of genetic information that cannot be reverted, and excision of an IS element occurs infrequently. We therefore suggest that reversion of mucoidy occurs via other means in our study, most likely nucleotide changes that cause gene loss-of-function that can subsequently be reverted to wild type. As mentioned above, given the high rates of observed reversion from mucoidy, it is possible that our mucoid lineages are not (all) genetic mutants; if there is a phenotypic component, both the high rate of reversion and the ready re-emergence of mucoidy are easily accounted for.

While we have focused on the role of resistance generated by mucoidy, the results of our experiments do not rule out a substantial role of surface, envelope, resistance in the dynamics of *E. coli* and its lytic phage. Indeed, if envelope resistant mutants are less costly than mucoid, any envelope resistant mutants that do enter a community of bacteria and phage will increase in frequency. Consistent with this prediction is that in some cases we observed envelope resistance arising within mucoid cultures. Most notably this occurs with T5 phage, where envelope resistance is expected to have minimal or no cost [30]. There is also evidence that mucoidy and other resistance mechanisms can arise together within single cultures [26]. Our results indicate that the evolution and utility of mucoidy, as compared with other phage resistance mechanisms, will depend heavily on the strains involved and the environmental context of the interaction. Under these conditions (laboratory *E. coli*

exposed to single strains of phage in liquid media), exponential growth rate was a reasonably informative measure of fitness, but this is known to be untrue in other scenarios [41], and particularly in more complex environments [8, 42, 43]. Understanding the selective forces that are likely to dominate in different contexts will therefore be relevant for a global understanding of mucoidy as a part of phage-bacteria interactions. Finally, our models and experimental results suggest that if mucoidy evolves first, mutants with more specific resistance mechanisms are, with the exceptions like T5, unlikely to ascend.

Caveats:

At one level, the population dynamic and experimental and modeling results presented here are mechanistic, they explain how mucoid *E. coli* can maintain high-density populations of lytic phage. However, as suggested in the preceding, at the genetic, physiological and molecular levels, these results and explanations raise mechanistic questions, which we allude to, but do not fully answer. Included among those unanswered or insufficiently questions are: (1) How does mucoidy provide the broad “resistance” to phages? (2) Why (really how) is the rate of transition from mucoid to non-mucoid so high? (3) What are the genetic, physiological, and molecular basis of the observed mucoid phenotypes? (4) Why does exposure to some phage, such as T5 in our experiments, not select for mucoidy, but instead selects for envelope resistance? (5) What might the consequences of mucoidy be for microbial populations in natural environments?

These studies were performed in well-mixed liquid media in serial transfer culture. Under these conditions, wall populations are unlikely to provide a refuge for stationary phase cells and allow the maintenance of the phage in this way [44]. Moreover, under these conditions multicell collectives are not anticipated to be formed and were not observed. In stationary cultures where surface adhesion and biofilm formation occur, these collectives could allow exopolysaccharide production to be used as a common good [37, 45]. Bacteria-phage dynamics in surface-adhered conditions might therefore differ from those observed here; for example, creation of surplus extracellular matrix by mucoid bacteria might provide some protection to phage-sensitive bacteria, either by impeding free diffusion of phage or by providing a scaffolding of phage-sorbing cellular material, allowing maintenance of a sensitive sub-population alongside mucoids and phage [37]. As the Rcs pathway in *E. coli* is a known regulator of biofilm formation [31], it is very plausible that mutations in this pathway that generate mucoidy in the interests of phage resistance will also produce changes in collective behavior.

A practical implication

In themselves, the results of these studies with *E. coli* K12 and its phages are of little practical importance. On the other hand, they may well be relevant for understanding the evolution and maintenance of mucoidy in other more medically or agriculturally relevant systems. Mucoidy has been observed in chronic *Pseudomonas aeruginosa* infections of the cystic fibrosis lung and marks progression of the disease into a more debilitating stage [45, 46]. While it is at present not known whether phage-imposed selection plays a role in this clinical phenomenon,

selection for mucoidy has been observed in *Pseudomonas* exposed to lytic phage [28, 47]. As interest in phage-based therapy for cystic fibrosis lung infection grows [18, 19, 48, 49], clarifying the role of phage in disease-associated mucoidy will be critical in developing safe and useful treatments for these intransigent infections.

Acknowledgements

The authors would like to thank Cary Nadell, Jim Bull, and Sylvain Moineau for generously providing us with the phage used in these experiments and Ole Skovgaard for providing the *E. coli* MG1655 employed here. This research was funded by grants from the U.S. National Institutes of General Medical Science GM098175-17 to B.R. L., 1R35GM133509-01 to M.G., and Emory University start-up funds (MG, NV). We are grateful to Ingrid McCall for managing the logistics of this enterprise and for providing editorial and other suggestions, and to Melony Ivey for technical assistance.

Materials and Methods

Bacteria, phage and growth media

Strains used in these experiments are shown in Table 1. Bacterial cultures were grown at 37°C either in M9 media [M9 salts (248510, Difco) supplemented with 0.4% glucose (6363-53-7, Fisher Scientific), 1 mM MgSO₄ (Sigma Aldrich), 0.1 mM CaCl₂ (JT Baker), and 0.2% Thiamine B1 (Sigma Aldrich)] or LB broth [MgSO₄ 2.5g/L, tryptone (Fisher Bioreagent 10g/L, yeast extract (Bacto) 5g/L, sodium chloride (Fisher Chemical) 10g/L]. All *E. coli* strains used in our experiments were derivatives of the parent strain K12 MG1655. Rcs pathway knockout strains were constructed by the method described in [50]. Gain-of-function point mutant for mucoid phenotype strain was constructed by the method described in [51].

Phage lysates were prepared from single plaques at 37°C in LB medium alongside wild-type *E. coli* MG1655. Chloroform was added to the lysates and the lysates were centrifuged to remove any remaining bacterial cells.

The λ^{VIR} strain used in these experiments was obtained from Sylvain Moineau. The construction of temperate phage λ^{KAN} and λ^{CAM} is described in [36].

Table 1. Bacterial strains and phage used in this study.

#	Bacterial Strain	Properties	Reference
1	<i>E.coli</i> K12 (MG1655)	Wild Type	Ole Skovgaard at Roskilde University in Denmark
2	T7 Mucoïd <i>E. coli</i> MG1655	T7 phage selected mucoïd	This study
3	T3 Mucoïd <i>E. coli</i> MG1655	T3 phage selected mucoïd	This study
4	<i>E.coli</i> K12	$\Delta rcsF::kan$	This study
5	<i>E.coli</i> K12	$\Delta rcsB::kan$	This study
6	<i>E.coli</i> K12	$\Delta cpsB::kan$	This study
7	<i>E.coli</i> K12	<i>rscC137 ompC::Tn5</i>	This study
#	Phage	Properties	Reference
1	T3	Wild Type	Jim Bull Lab
2	T4	Wild Type	Jim Bull Lab
3	T5	Wild Type	Jim Bull Lab
4	T7	Wild Type	Jim Bull Lab
5	T7	GFP-labelled T7	[37]
6	λ^{vir}	Virulent mutant of λ	Sylvain Moineau Lab
7	λ^{KAN}	Wild type λ with Kanamycin antibiotic cassette	[36]
8	λ^{CAM}	Wild type λ with chloramphenicol antibiotic cassette	[36]

Sampling bacterial and phage densities

Bacteria and phage densities were estimated by serial dilutions in 0.85% NaCl solution followed by plating. The total density of bacteria was estimated on LB (1.6%) agar plates. To estimate the densities of λ -KAN or λ -CAM lysogens, cultures were plated on LB (1.6%) agar with 25 $\mu\text{g}/\text{mL}$ kanamycin (AppliChem Lot# 1P0000874) or chloramphenicol (Merck Lot# 2430B61), respectively.

To estimate the densities of free phage, chloroform was added to suspensions before serial dilutions. These suspensions were mixed with 0.1 mL of overnight LB grown cultures of wild-type MG1655 (about 5×10^8 cells per mL) in 3 mL of LB soft (0.65%) agar and poured onto semi-hard (1%) LB agar plates.

Serial Transfer

All serial transfer experiments were carried out in 10-mL cultures grown at 37°C with vigorous shaking. The cultures were initiated by 1:100 dilution from 10-mL overnight cultures grown from single colonies. Phage were added to these cultures to reach the initial density of approximately 10^6 PFU/mL. At the end of each transfer, 100 μL of each culture was transferred into flasks with fresh medium (1:100 dilution). Simultaneously, 100 μL samples were taken for estimating the densities of colony forming units (CFU) and plaque forming units (PFU), by serial dilution and plating on solid agar, with selection as needed as described above.

Bacteriophage and bacteria parameter determination

The parameters critical for the interaction of λ phages and *E. coli* MG1655 used in this study were estimated in independent experiments in LB medium. The maximum growth rate of *E. coli* MG1655 was measured by Bioscreen as described in [41]. Phage burst sizes (β) were estimated with one-step growth experiments [52] in a manner similar to [53]. Adsorption of λ to *E. coli* was estimated as described in [52]. The procedure for estimating the probability of lysogeny and the rate of spontaneous lysogen induction are presented in [36].

Short term dynamics experiments

Overnight cultures of *E. coli* MG1655 were grown in LB media. The cultures were diluted in a 1:1000 ratio on 20 mL of medium in 100 mL flasks to which λ_{KAN} or λ_{CAM} phage was added to an initial density of 1×10^7 PFU/mL. Short term dynamics were conducted culturing the mixture at 37 °C in shaking conditions. Total bacteria count was conducted by sampling 100 μL at regular intervals and plating dilutions on LB agar plates. Lysogen counts were obtained by sampling 100 μL and plating dilutions on LB agar with 25 $\mu\text{g}/\text{mL}$ kanamycin or chloramphenicol. Phage counts were obtained by sampling 1 mL at regular intervals, adding chloroform to kill bacteria, and plating for PFU quantification as previously described.

Resistance Testing

Liquid culture: Colonies from the bacteria plates were streaked thrice to ensure removal of phage. Colonies from the streaked plates were used to establish overnight cultures in 2 mL LB broth incubated at 37°C in a shaker. The overnight

cultures were diluted 1/100 and $\sim 1e5$ phage were added to the cultures. After 24 hours of incubation, free phage in these cultures were measured. Those unable to support the phage replication were considered as resistant [7].

Phage agar: Phage plates were made as previously described, using 0.1 mL of overnight LB grown cultures of the isolates being tested. Dense phage stocks (10^8 - 10^9 PFU/mL) were spotted onto the agar surface; susceptibility was scored according to the presence of visible plaques.

Single Cell Microscopy

Overnight cultures of WT and mucoid *E.coli* MG1655 carrying plasmid pBAD18Cm::P_{rprA}-mCherry (reporter for induction of the Rcs pathway [54]) were grown in LB media. These cultures were diluted 1/1000 in fresh LB and incubated for 30 minutes with vigorous shaking (300 RPM) at 37°C. We then added GFP-labelled T7 phage [37] at 0.1 MOI and returned to incubation in a stationary incubator at 37°C. The phage free control and phage infected culture slides were sampled at 15-30 minute intervals and observed on an inverted fluorescent microscope (Leica DMI8 with LasX Premium software). Images are an overlay of DIC, TRITC, and GFP channels, acquired using a 40X objective (total magnification 400X).

Quantification was performed using Image J. Object thresholds were established using DIC images, and particle analysis was performed on all particles exceeding 0.1 μM^2 area. Resulting plots show mean pixel intensity in each fluorescent channel for all identified objects.

References

1. Chibani-Chennoufi, S., et al., *Phage-host interaction: an ecological perspective*. Journal of bacteriology, 2004. **186**(12): p. 3677-3686.
2. Abeles, S.R. and D.T. Pride, *Molecular bases and role of viruses in the human microbiome*. Journal of molecular biology, 2014. **426**(23): p. 3892-3906.
3. Campbell, A., *Conditions for the existence of bacteriophage*. Evolution, 1961. **15**: p. 153-165.
4. Barrangou, R., et al., *CRISPR provides acquired resistance against viruses in prokaryotes*. Science, 2007. **315**(5819): p. 1709-1712.
5. Horne, M.T., *Coevolution of Escherichia coli and bacteriophages in chemostat culture*. Science, 1970. **168**(3934): p. 992-3.
6. Chao, L., B.R. Levin, and F.M. Stewart, *A complex community in a simple habitat: an experimental study with bacteria and phage*. Ecology, 1977. **58**: p. 369-378.
7. Wei, Y., P. Ocampo, and B.R. Levin, *An experimental study of the population and evolutionary dynamics of Vibrio cholerae O1 and the bacteriophage JSF4*. Proceedings of the Royal Society B: Biological Sciences, 2010. **277**(1698): p. 3247-3254.
8. Gomez, P. and A. Buckling, *Bacteria-phage antagonistic coevolution in soil*. Science, 2011. **332**(6025): p. 106-9.
9. Chaudhry, W.N., et al., *Leaky resistance and the conditions for the existence of lytic bacteriophage*. PLoS Biol, 2018. **16**(8): p. e2005971.
10. Levin, B.R., F.M. Stewart, and L. Chao, *Resource-limited growth, competition, and predation: a model and experimental studies with bacteria and bacteriophage*. The American Naturalist, 1977. **111**(977): p. 3-24.
11. Jover, L.F., M.H. Cortez, and J.S. Weitz, *Mechanisms of multi-strain coexistence in host-phage systems with nested infection networks*. J Theor Biol, 2013. **332**: p. 65-77.
12. Jover, L.F., J. Romberg, and J.S. Weitz, *Inferring phage-bacteria infection networks from time-series data*. R Soc Open Sci, 2016. **3**(11): p. 160654.
13. Weissman, J.L., et al., *Immune Loss as a Driver of Coexistence During Host-Phage Coevolution*. bioRxiv, 2017: p. 105908.
14. Reyes, A., et al., *Gnotobiotic mouse model of phage-bacterial host dynamics in the human gut*. Proc Natl Acad Sci U S A, 2013. **110**(50): p. 20236-41.
15. Manrique, P., et al., *Healthy human gut phageome*. Proc Natl Acad Sci U S A, 2016. **113**(37): p. 10400-5.
16. Bull, J.J., B.R. Levin, and I.J. Molineux, *Promises and Pitfalls of In Vivo Evolution to Improve Phage Therapy*. Viruses, 2019. **11**(12).

17. Schmidt, C., *Phage therapy's latest makeover*. Nat Biotechnol, 2019. **37**(6): p. 581-586.
18. Cafora, M., et al., *Phage therapy against Pseudomonas aeruginosa infections in a cystic fibrosis zebrafish model*. Sci Rep, 2019. **9**(1): p. 1527.
19. Forti, F., et al., *Design of a Broad-Range Bacteriophage Cocktail That Reduces Pseudomonas aeruginosa Biofilms and Treats Acute Infections in Two Animal Models*. Antimicrob Agents Chemother, 2018. **62**(6).
20. Scholl, D., S. Adhya, and C. Merrill, *Escherichia coli K1's capsule is a barrier to bacteriophage T7*. Appl Environ Microbiol, 2005. **71**(8): p. 4872-4.
21. Bernheimer, H.P. and J.G. Tiraby, *Inhibition of phage infection by pneumococcus capsule*. Virology, 1976. **73**(1): p. 308-9.
22. van der Ley, P., P. de Graaff, and J. Tommassen, *Shielding of Escherichia coli outer membrane proteins as receptors for bacteriophages and colicins by O-antigenic chains of lipopolysaccharide*. J Bacteriol, 1986. **168**(1): p. 449-51.
23. Labrie, S.J., J.E. Samson, and S. Moineau, *Bacteriophage resistance mechanisms*. Nat Rev Microbiol, 2010. **8**(5): p. 317-27.
24. Wilkinson, B.J. and K.M. Holmes, *Staphylococcus aureus cell surface: capsule as a barrier to bacteriophage adsorption*. Infect Immun, 1979. **23**(2): p. 549-52.
25. Harrison, E., et al., *Plasmid carriage can limit bacteria-phage coevolution*. Biol Lett, 2015. **11**(8).
26. Wielgoss, S., et al., *Adaptation to parasites and costs of parasite resistance in mutator and nonmutator bacteria*. Molecular biology and evolution, 2015. **33**(3): p. 770-782.
27. Mizoguchi, K., et al., *Coevolution of bacteriophage PP01 and Escherichia coli O157: H7 in continuous culture*. Appl. Environ. Microbiol., 2003. **69**(1): p. 170-176.
28. Scanlan, P.D. and A. Buckling, *Co-evolution with lytic phage selects for the mucoid phenotype of Pseudomonas fluorescens SBW25*. ISME J, 2012. **6**(6): p. 1148-58.
29. Delbruck, M., *Bacterial viruses or bacteriophages*. Biol Rev Camb Philos Soc, 1946. **21**: p. 30-40.
30. Lenski, R.E. and B.R. Levin, *Constraints on the coevolution of bacteria and virulent phage : a model, some experiments, and predictions for natural communities*. American Naturalist, 1985. **125**: p. 585-602.
31. Clarke, D.J., *The Rcs phosphorelay: more than just a two-component pathway*. Future microbiology, 2010. **5**(8): p. 1173-1184.
32. Stout, V., *Identification of the promoter region for the colanic acid polysaccharide biosynthetic genes in Escherichia coli K-12*. J Bacteriol, 1996. **178**(14): p. 4273-80.

33. Morona, R., J. Tommassen, and U. Henning, *Demonstration of a bacteriophage receptor site on the Escherichia coli K12 outer-membrane protein OmpC by the use of a protease*. Eur J Biochem, 1985. **150**(1): p. 161-9.
34. Monod, J., *The growth of bacterial cultures*. Annual review of microbiology, 1949. **3**(1): p. 371-394.
35. Stewart, F.M. and B.R. Levin, *Partitioning of resources and the outcome of interspecific competition: a model and some general considerations*. The American Naturalist, 1973. **107**(954): p. 171-198.
36. Pleška, M., et al., *Phage–host population dynamics promotes prophage acquisition in bacteria with innate immunity*. Nature ecology & evolution, 2018. **2**(2): p. 359.
37. Vidakovic, L., et al., *Dynamic biofilm architecture confers individual and collective mechanisms of viral protection*. Nature microbiology, 2018. **3**(1): p. 26.
38. Wei, Y., A. Kirby, and B.R. Levin, *The population and evolutionary dynamics of Vibrio cholerae and its bacteriophage: conditions for maintaining phage-limited communities*. Am Nat, 2011. **178**(6): p. 715-25.
39. Wright, R.C.T., et al., *Cross-resistance is modular in bacteria-phage interactions*. PLoS Biol, 2018. **16**(10): p. e2006057.
40. Lee, H., et al., *Rate and molecular spectrum of spontaneous mutations in the bacterium Escherichia coli as determined by whole-genome sequencing*. Proc Natl Acad Sci U S A, 2012. **109**(41): p. E2774-83.
41. Concepción-Acevedo, J., et al., *Malthusian parameters as estimators of the fitness of microbes: a cautionary tale about the low side of high throughput*. PloS one, 2015. **10**(6): p. e0126915.
42. Bjorkman, J., et al., *Effects of environment on compensatory mutations to ameliorate costs of antibiotic resistance*. Science, 2000. **287**(5457): p. 1479-82.
43. Santos-Lopez, A., et al., *Evolutionary pathways to antibiotic resistance are dependent upon environmental structure and bacterial lifestyle*. Elife, 2019. **8**.
44. Schrag, S. and J.E. Mittler, *Host parasite coexistence: the role of spatial refuges in stabilizing bacteria-phage interactions*. American Naturalist, 1996. **148**: p. 428-377.
45. Malhotra, S., et al., *Mixed Communities of Mucoïd and Nonmucoïd Pseudomonas aeruginosa Exhibit Enhanced Resistance to Host Antimicrobials*. MBio, 2018. **9**(2).
46. Li, Z., et al., *Longitudinal development of mucoïd Pseudomonas aeruginosa infection and lung disease progression in children with cystic fibrosis*. JAMA, 2005. **293**(5): p. 581-8.

47. Martin, D.R., *Mucoid variation in Pseudomonas aeruginosa induced by the action of phage*. J Med Microbiol, 1973. **6**(1): p. 111-8.
48. Alemayehu, D., et al., *Bacteriophages phiMR299-2 and phiNH-4 can eliminate Pseudomonas aeruginosa in the murine lung and on cystic fibrosis lung airway cells*. MBio, 2012. **3**(2): p. e00029-12.
49. Debarbieux, L., et al., *Bacteriophages can treat and prevent Pseudomonas aeruginosa lung infections*. J Infect Dis, 2010. **201**(7): p. 1096-104.
50. Baba, T., et al., *Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection*. Molecular systems biology, 2006. **2**(1).
51. Gottesman, S., P. Trisler, and A. Torres-Cabassa, *Regulation of capsular polysaccharide synthesis in Escherichia coli K-12: characterization of three regulatory genes*. Journal of bacteriology, 1985. **162**(3): p. 1111-1119.
52. Ellis, E.L. and M. Delbrück, *The growth of bacteriophage*. The Journal of general physiology, 1939. **22**(3): p. 365-384.
53. Abedon, S.T., T.D. Herschler, and D. Stopar, *Bacteriophage latent-period evolution as a response to resource availability*. Appl. Environ. Microbiol., 2001. **67**(9): p. 4233-4241.
54. Konovalova, A., et al., *Inhibitor of intramembrane protease RseP blocks the σE response causing lethal accumulation of unfolded outer membrane proteins*. Proceedings of the National Academy of Sciences, 2018. **115**(28): p. E6614-E6621.

SUPPLEMENTAL MATERIAL

Mucoidy, a general mechanism for maintaining lytic phage in populations of bacteria

Waqas Chaudhry¹, Esther Lee¹, Andrew Worthy¹, Zoe Weiss¹, Marcin Grabowicz^{2,3,4}, Nicole Vega^{1*}, Bruce Levin^{1,2**}

¹ Department of Biology, Emory University, Atlanta, GA 30322, USA.

² Emory Antibiotic Resistance Center, Emory University School of Medicine, Atlanta, GA 30322

³ Department of Microbiology & Immunology, Emory University School of Medicine, Atlanta, GA 30322, USA.

⁴ Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, GA 30322, USA.

[*nvega@emory.edu](mailto:nvega@emory.edu)

[**blevin@emory.edu](mailto:blevin@emory.edu)




Figure S1: The population dynamics of T3 (A), T4 (B), T5 (C) and T7 (D) phage with T7 generated mucoid *E. coli* MG1655 as the host strain. Cultures were diluted 1:100 into fresh M9 + 0.4% glucose every 24 hours, and CFU/PFU measurements were taken via dilution plating. Solid lines show bacterial densities in each replicate (red/green/blue); dotted lines show the phage densities in the corresponding cultures. Phage-free cultures (black and purple lines) are shown as controls, to

indicate resource-limited population density for the bacteria. The phage-free controls lost the mucoid phenotype. At the end of the last serial transfer, three colonies from each phage-infected culture were streaked at least three times to get rid of co-evolving phage. These isolated clones were tested against the phage used in the experiment; envelope resistance arose in T4 and T5-infected cultures, accompanied by a loss of mucoidy.




Figure S2: Changes in the densities of bacteria and phage in serial transfer culture. Standard parameters, $C=1000$, $e=5 \times 10^7$, $v_S=2.1$, $v_R=2.0$, $k=1$, $\beta=50$, $x=10^{-5}$, $\delta_S=10^{-7}$. (A) Long term dynamics of a population with mucoid, sensitive, and resistant bacteria $v_M=1.8$, $\delta_M=10^{-11}$, $y=10^{-3}$, continuation of the dynamics in Figure 5B. (B) Long term dynamics of a population with mucoid, sensitive and resistant bacteria with the mucoid bacteria fully resistant to the phage, $\delta_M=0$, $v_M=1.8$, $y=10^{-3}$. (C) Long term dynamics of a population with mucoid, sensitive, and resistant bacteria with a lower rate of transition from S to M, $y=10^{-5}$, $\delta_M=10^{-11}$, $\delta_S=10^{-7}$, a continuation of the dynamics presented in Figure 5C. (D) Long term dynamics of a community where phage are maintained by the fitness cost of resistance and no mucoid bacteria can be generated. Two simulations, one where the initial densities of phage are the 0.01 of the equilibrium, $N_R=2.0 \times 10^7$, $P=1.40 \times 10^5$, $S=1.67 \times 10^4$ (see Figure S2A) and one where the initial density of phage is 1.4×10^3 . (E) Dynamics of a community where the phage are maintained by the transitions of mucoid to sensitive (leaky resistance) $v_M=1.8$, $\delta_M=10^{-11}$, $y=10^{-3}$, no resistant bacteria; effect of the initial densities of mucoid bacteria and phage. (F) Dynamics of a community where the phage are maintained by the transitions from sensitive to mucoid, but there is no fitness cost of mucoidy, $v_M=v_S=2.1$, $\delta_M=10^{-11}$, $y=10^{-3}$, $\delta_S=10^{-7}$, no resistant cells are present or can be generated.




Figure S3: Population dynamics of λ^{VIR} phage with T3-generated mucoid, T7-generated mucoid, *rcsC137 ompC::Tn5* (constitutive mucoid) and WT *E.coli* MG1655. Cultures were inoculated into LB medium with λ^{VIR} phage (1:1) and diluted 1:100 into fresh LB medium every 24 hours; two replicates were performed per condition. The decline of phage in wild-type cultures corresponds to the emergence of envelope resistance against λ^{VIR} . Solid red and blue lines show bacterial densities while dotted line show the phage densities in the corresponding cultures. Phage-free cultures (black lines) are shown as controls, to indicate resource-limited population size for the bacteria.




Figure S4: Reversion from mucoid and the growth rates of the different mucoid cell lines. To estimate the reversion rate of mucoid to non-mucoid phenotype, single mucoid colonies were picked from the 9th serial transfer of (A) T3, (B) T4, (C) T7 phage infected cultures of *E. coli* MG1655 (Figure 1) and (D) genetically constructed mucoid *E. coli* MG1655 (*rcsC137 ompC::Tn5*) and grown in liquid culture in the absence of phage, with dilution 1:100 into fresh LB broth every 24 hours. At the end of every transfer, the culture was serially diluted and plated on LB agar for CFU/ml counts and colony morphology determination. Red, green and blue lanes represent three independent replicates (E) Mucoid (MUC) cultures suffer a substantial growth rate defect as compared with wild-type *E. coli* MG1655, while envelope resistance (R) to T5 and λ phage incurs no growth rate penalty under these conditions. Note that the “T4 Muc” and “T7 Muc” cultures immediately produce a mixed culture of mucoid and revertant colonies, as seen during CFU plating of 24-hour growth curve cultures (F), and the estimated growth rate for the mixed culture is likely to reflect the more rapid growth rate of the substantial revertant fraction. Data were obtained from OD₆₀₀ growth curves in LB at 37°C. Error bars represent mean \pm SD of ten (E) or three (F) replicates.




Figure S5: The proportion of mucoid colonies is shown for populations of the ancestral revertant. Mucoidy was scored after exposure to phages on the phage plate. Three individual revertant non-mucoid colonies were picked from each ancestor mucoid background (bacteria from transfer 8 derived from mucoid ancestors, Figure S4 A, B, C, D). 100 μ l of overnight cultures of these clones was mixed with 100 μ l of a high titer of each respective phage along with 3ml soft agar and poured on the plate; rscC137 revertants were exposed to T7 phage.




Figure S6: Quantification of red (TRITC) and green (GFP) channels for the images shown in Figure 7. Points represent mean pixel intensity for each channel for objects identified as particles in DIC channel. Image quantification was performed in ImageJ.

Model of the dynamics of lysogen formation with leaky resistance

There are three populations of bacteria sensitive non-lysogens, S, resistant (Mucoïd), M non-lysogens, and lysogens, L and one population of phage, P, where S, M, L and P are the densities, cells and particles per ml as well as the designations of these populations. The bacteria grow at maximum rates, v_S , v_M , and v_L with the realized rate of growth proportional to the concentration of the limiting resource,

$$Z_i(r) = v_i \cdot \frac{r}{(r+k)}$$

r $\mu\text{g/ml}$, where i is S, M or L, and k is the concentration of the resource where the realized growth rate is half its maximum value. For the production of a new cell, e μg of the resource has to be consumed.

The phage adsorb to the bacteria at a rate proportional to the product of their densities and a rate constant δ . A fraction, λ ($0 < \lambda < 1$) of the adsorptions of the temperate phage to sensitive cells, P to S, results in the production of lysogens. Phage that adsorb to lysogens are lost. We assume that the mucoïd population is completely resistant to the phage. The $(1-\lambda)$ fraction of infections of sensitive cells by temperate phage result in the production of β , the burst size, of free phage. With a probability i ($0 < i < 1$) per cell per hour lysogens are induced and produce β phage. With a rate y per cell per hour, the mucoïd, M produce sensitive cells, $M \rightarrow S$, and x per cell per hour, the sensitive cells generate mucoïds, $S \rightarrow M$. With these definitions and assumptions, the rate of change in densities of the bacteria and phage and concentration of the limiting resource are given by the following set of coupled differential equations.

$$\frac{dr}{dt} = -\psi(r) \cdot e \cdot (v_S \cdot S + v_M \cdot M + v_L \cdot L)$$

$$\frac{dS}{dt} = v_S \cdot S \cdot \psi(r) - \delta \cdot S \cdot P \cdot \psi(r) + (-x \cdot S + y \cdot M + i \cdot L) \cdot \psi(r)$$

$$\frac{dL}{dt} = v_L \cdot L \cdot \psi(r) + \delta \cdot S \cdot P \cdot (1-\lambda) \cdot \psi(r) - i \cdot L \cdot \psi(r)$$

$$\frac{dM}{dt} = v_M \cdot M \cdot \psi(r) + (x \cdot S - y \cdot M) \cdot \psi(r)$$

$$\frac{dP}{dt} = \delta \cdot S \cdot P \cdot (1-\lambda) \cdot \beta \cdot \psi(r) + i \cdot L \cdot \beta \cdot \psi(r)$$

$$\text{where } \psi(r) = \frac{r}{(r+k)}$$