Rotten-skin disease significantly changed giant spiny frog(*Paa spinosa*) gut microbiota

Tuoyu He¹、 Yun Jiang¹、 Pengpeng Wang^{1,3}、 Jianguo Xiang^{1*}、 Wangcheng Pan²

¹ College of Animal Science and Technology, Hunan Agricultural University, Changsha, China

² Changde Dabeinong Ltd, Changde, China

³ Hunan Fisheries Science Institute, Changsha, China

Running title: Gut microbiota about rotten-skin disease

* Correspondence author: jianguo Xiang, College of Animal Science and Technology, Hunan Agricultural University, Changsha, 410125, China. e-mail: jgxianghau@163.com;

Abstract: The composition and abundance of gut microbiota is essential for host health and immunity. Gut microbiota is symbiotic with the host, so changes in the host diet,

development, and health will lead to changes in the gut microbiota. Conversely, changes in the gut microbiota also affect the host conditions. In this experiment, 16S rRNA high-throughput sequencing was used to compare the gut microbiota composition of 5 healthy *Paa spinosa* and 6 *P. spinosa* with rotten-skin disease. Results: the gut microbiota composition was significant difference between diseased *P. spinosa* and the healthy *P. spinosa*; LEfSe analysis showed that the relative abundance of *Methanocorpusculum*, *Parabacteroides*, *AF12*, *PW3*, *Epulopiscium*, and *Oscillospira* were significantly higher in the diseased *P. spinosa*, while the relative abundance of *Serratia*, *Eubacterium*, *Citrobacter*, and *Morganella* were significantly lower. Conclusion: Rotten-skin disease changed *P. spinosa* gut microbiota significantly; The relative abundance of *Epulopiscium* and *Oscillospira* might be related to the health conditions of the host skin and gallbladder; The relative abundance of *Serratia* and *Eubacterium* might be important for maintaining the gut microbiota ecosystem.

Keywords: microbiota structure; gut microbiota; rotten-skin disease; *Paa spinosa*; high-throughput sequencing

Introduction

The giant spiny frog (*Paa spinosa*) is a large edible frog distributed in the

mountains of China and Vietnam (1). It is favored by people for its great economic and medicinal value (2). In recent years, due to the increase of market demand and the destruction of habitats, the wild *P. spinosa* have been declined sharply. The Chinese Red Animal List has listed *P. spinosa* as a “vulnerable” species(3) (4). The artificial breeding has provided an effective way to meet market demand and protect wild *P. spinosa*. However, frequent disease problems seriously restrict the development of the *P. spinosa* industry (1) (5).

The rotten-skin disease is a common disease of *P. spinosa*, which is characterized by a dull epidermis and white spots appearing at the beginning, and then the epidermis falls off and begins to rot until the bone is exposed(6). Many pathogens cause rotten-skin disease, such as *Proteus mirabilis* and *Yersinia kristensenii*(6). Some diseased frogs will not die immediately, but growth is affected (7), some pathogens not only cause diseased frogs to show signs of rot but also cause a large number of deaths, such as *Chytridiomycosis*. The diversity of pathogens of rotten-skin disease bring difficulties to routine methods of preventing the disease, a new method is needed immediately.

Animal gut microbial communities as host “microbial organs” play an important role in host immunity and health, such as promoting the absorption of nutrients(8) (9) (10), impeding pathogens colonization in the gut (11) and regulating host immunity to maintain host health. Gut microbiota is symbiotic with the host, so changes in the host diet, development, and health can lead to changes in the gut microbiota (12)(13)(14). Conversely, changes in the gut microbiota also affect the host conditions. The normal microbiota composition is the guarantee for maintaining the physiological function of

the host (15). When the physiological function of the host is abnormal (such as the disease involving), the gut microbiota composition will change as well(16). In recent years, a new understanding of aquatic animal diseases has been gained by comparing the gut microbiota of diseased aquatic animals with healthy ones (Table 1). So we hypothesized that gut microbiota changed significantly between the healthy *P. spinosa* and the rotten-skin diseased *P. spinosa*. And comparing the composition of gut microbiota of healthy *P. spinosa* and the rotten-skin diseased *P. spinosa* we will know the microbiota change, which may be vital to discover the antagonistic bacteria of *P. spinosa* rotten-skin disease and explore new methods to prevent and control rotten-skin disease.

In this paper, the 16S rRNA amplicon high-throughput sequencing technology was used to investigate the effects of rotten-skin disease on gut microbiota composition in *P. spinosa*. The potential probiotics and antagonistic bacteria were screened out in the healthy *P. spinosa* through comparing the composition of gut microbiota between healthy and diseased *P. spinosa*, and expected to enrich the theories about regulating gut microbiota structure by using microbial ways and realizing the microecological prevention of rotten-skin disease in *P. spinosa*.

Materials and methods

Sample Collection

All animal experiments were conducted in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institute of

Health(NIH). The experimental animals were approved by the experimental animal ethic committee of Hunan Agriculture University. All samples were collected from Shimen County, Hunan Province Weixin Town *P.spinosa* farm (110°29'-111°33'E, 29°16'-30°08 N). There were 11 *P.spinosa* in the experiment, including 5 Healthy *P.spinosa* (H) and 6 diseased *P.spinosa* (D). The frog was anesthetized and dissected under sterile conditions, collected gut contents, transferred to 2 ml sterile EP tubes and stored at -80 ° C, used for subsequent DNA extraction.

DNA extraction and high-throughput sequencing

The gut microbiota DNA was extracted using a DNeasy PowerSoil Kit (QIAGEN, Germany). The V4-V5 hypervariable region of the prokaryotic 16S rRNA gene was amplified using the universal primer pair 515F and 909R, with a 12-nt sample-specific barcode sequence, including at the 5'-end of the 515F to distinguish samples (17) (18). PCR was performed, and amplicons were sequenced using a MiSeq system at Guangdong Meilikang Bio-Science, Ltd. (China), as described previously (1). The raw sequences were merged using FLASH-1.2.8 software and processed using the QIIME pipeline 1.9.0 as described previously(1)(19). Chimeric sequences were identified and removed using the Uchime algorithm and the no-chimeric sequences were clustered into OTUs at 97% identity using UPARSE software (20) (21). The RDP classifier was used to detect the taxonomic assignments of each OTU (22).

The merged sequences were submitted to the NCBI SRA database (accession number SRR10765290-SRR10765300).

Data Analysis

The results for each parameter were presented as the mean $\pm$ standard error for each group. Principal coordinate analysis (PCoA) based on unweighted Unifrac distance was applied to evaluate the differences in different groups. Principal component analysis (PCA) was performed by R vegan package. Non-parametric ANOVA (PERMANOVA) was performed using the R vegan package(23) to analyze the significance of differences between groups; Welch's T-test(STAMP) was performed to analyze significantly different phylum in gut microbiota between groups; Prism 6 was used for box plot production; T tests was conducted by SPSS16.0 to analyze the significance of differences in diversity indicators; the p-value less than 0.05 were significant difference, p-value less than 0.01 were significant extremely.

Results

The skin of diseased *P. spinosa* had the white spot or large area of decay. After anatomy, there were obvious lesions in the liver and gallbladder. The liver blackened obviously. The gallbladder was enlarged or discolored. The symptoms of the diseased *P. spinosa* are shown in the picture (Fig. 1).

A total of 466,541 high-quality sequences were obtained from the 11 gut microbiota samples. To avoid the influence of sequencing depth, 25,520 sequences were randomly selected from each sample for further analysis, and 5,793 OTUs from 704 genera were identified. The results of gut microbiota diversity indicated that the diversity index, such as Chao1, Observed-species, PD-whole-tree, Shannon index, and Simpson index, in healthy *P. spinosa* were significantly lower than the diseased *P. spinosa*, but the

Good-coverage had no difference (Fig.2). Removing the unclassified sequences (<0.001%), 19 of the 56 phyla dominated the gut microbiota. *Bacteroides* and *Firmicutes* were the dominant microbiota in the gut of all samples (Fig. 3A), which is consistent with a previous study (1). Among them, the average relative abundance of *Bacteroides*, *Firmicutes*, *Proteobacteria*, *Tenericutes*, and *Euryarchaeota* was more than 1% in all samples (Fig. 3A). STAMP based on relative abundance of top 10 phyla in the gut microbiota showed that *Proteobacteria* was significantly higher in the healthy *P. spinosa*, while the relative abundance of *Euryarchaeota* and *Spirochaetes* were significantly lower (Fig. 3B).

PCA based on the relative abundance of all gut microbiota genera and PCoA based on the relative abundance of the all gut microbiota genera showed that there were significant differences in gut microbiota composition between diseased and healthy *P. spinosa* (PERMANOVA, $F = 3.0464$, $p = 0.008$) (Fig. 4A and 4B). Unweighted Pair-Group Method with Arithmetic means UPGMA analysis shown that microbiota composition were familiar between the groups (Fig. 4C).

Lefse analyzed the difference of gut microbiota at genus level showed that relative abundance of *Serratia*, *Eubacterium*, *Citrobacter*, and *Morganella* were significantly higher in healthy *P. spinosa*, while the relative abundance of *Methanocorpusculum*, *Parabacteroides*, *AF12*, *PW3*, *Epulopiscium*, and *Oscillospira* were significantly lower (Fig. 5).

Discussion

Recent researches have shown that gut microbiota has participated in various disease

processes through the gut-brain axis (24) (25), the gut-lung axis (26) (27) the gut-vascular axis(28, 29) , the gut-bone axis (30) (31),the gut-Hepatic axis (32) (33) and other axis (34). The concept of “core microbiota” indicated that the core microbiota in the gut of healthy hosts could maintain the stability of gut microbiota composition and function, and positively regulated the host through these axis to maintain host health (35). The gut microbiota function was destroyed because of the destruction of the core microbiota, and the host might become sick or aggravate the lesion (16). In this study, the gut microbiota of diseased and healthy *P. spinosa* was compared, and the results revealed significant differences in the gut microbiota composition of healthy and diseased *P. spinosa*. The composition of microbiota was destroyed because of pathogen invading.

Current researches on gut microbiota focused on gut microbiota diversity and gut microbiota composition. According to the diversity resistance hypothesis, the more diverse that the microbial community was and the more possible that the host resisted to pathogen invasion (36). Studies in the largemouth bronze gudgeon (*Coreius guichenoti*) (37), crucian Carp (*Carassius auratus*) (14), and ayu (*Plecoglossus altivelis*) (38) showed that the microbiota diversity was significantly higher in healthy samples. However, this study found that gut microbiota diversity was significantly higher in diseased *P. spinosa*. The results were consistent with the results of grass carp (*Ctenopharyngodon idellus*) (39). And found that the amino acid metabolism, carbohydrate metabolism, and immune-related pathway genes of diseased grass carp were more abundant through microbiota gene prediction (39). The increased

microbiota diversity in the gut of the diseased host may because the microbial homeostasis in the gut of the diseased host has not been broken immediately. To maintain the health of the host, the gut microbiota diversity was increased to protect against pathogen invasion. The results of this study and previous studies have shown that the use of gut microbiota diversity to assess host health is limited.

The relative abundance of *Methanocorpusculum*, *Parabacteroides*, *AF12*, *PW3*, *Epulopiscium*, and *Oscillospira* in the gut of rotten-skin *P.spinosa* were significantly higher than healthy *P. spinosa*. Although *Methanocorpusculum* is not a pathogen, it is abundant in diseased samples (40). It can effectively convert heavy metals or metalloids into more toxic derivatives than compounds, which was harmful to host health (41); *Parabacteroides goldsteinii* in *Parabacteroides* can cause bacteraemia ; Previously studied in the gut microbiota of wild and cultured *P. spinosa* found that the cultured *P. spinosa* with more potential pathogens had more *AF12* in the gut (1); Studies on human gallstones indicated that the relative abundance of *Oscillospira* was positive correlation with the gallstones (42, 43); The relative abundance of *Epulopiscium* was significantly increased in the gut of rotten-skin diseased *P. spinosa*, and also significantly increased in the feces of children with eczema(44). In summary, the gut microbiota that was significantly increased in the gut of rotten-skin disease *P. spinosa* was mostly opportunistic pathogen; *Oscillospira* and *Epulopiscium* were significantly increased in the gut of the diseased host when lesions occurred in the skin and gallbladder. It was speculated that these two species may be indicator microbiota in the pathogenesis of skin and gallbladder.

The relative abundance of *Serratia*, *Eubacterium*, *Citrobacter*, and *Morganella* in the gut of healthy *P. spinosa* was significantly higher than diseased *P. spinosa*. Some species in *Serratia* produced Prodigiosin and β -lactam antibiotic carbapenem to inhibit the growth of pathogens in the host, thereby inhibiting the disease (45, 46); *Bifidobacteria* and *Eubacterium hallii* promoted acetate, butyrate, propionate, and formate to form, potentially contributing to gut SCFA formation with potential benefits for the host and for microbiota colonization of the infant gut (47). *E. hallii* was also capable of metabolizing glycerol to 3-hydroxypropionaldehyde with antibacterial properties (48); The relative abundance of *Citrobacter* and *Morganella* in the gut of healthy *P. spinosa* was significantly higher than diseased *P. spinosa*, however *Citrobacter rodentium* and *Morganella morganii* are common opportunistic pathogen (49) (50). In summary, *Serratia* and *Eubacterium* might be the main gut microbiota in the healthy *P. spinosa* that maintained the health of *P. spinosa*; *Citrobacter* and *Morganella* in the gut of healthy *P. spinosa* were significantly increased without causing disease. It might be that non-pathogenic strains of *Citrobacter* and *Morganella* appeared in the gut of healthy frogs, or there were pathogenic strains in these two species, but due to the inhibition of beneficial microbiota in the gut of healthy hosts, the host still maintained a healthy state of gut microbiota homeostasis.

Conclusion

Rotten-skin disease significantly changed *P. spinosa* gut microbiota; the relative abundance of *Methanocorpusculum*, *Parabacteroides*, *AF12*, *PW3*, *Epulopiscium*, and

Oscillospira were significantly higher in the diseased *P. spinosa* , while the relative abundance of *Serratia*, *Eubacterium*, *Citrobacter*, and *Morganella* were significantly lower; The relative abundance of *Epulopiscium* and *Oscillospira* might be related to the healthy condition of the host skin and gallbladder; The relative abundance of *Serratia* and *Eubacterium* might be important for maintaining the gut microbiota ecosystem.

Acknowledgements

Financial support: This work was supported by Sichuan Provincial Department of Education Key Scientific Research Project 18ZA0443 and Xichang University PhD Project 2017BS009.

Conflicts of interest: All authors: No potential conflicts of interest.

References

1. Xiang J, He T, Wang P, Xie M, Xiang J, Ni J. 2018. Opportunistic pathogens are abundant in the gut of cultured giant spiny frog (*Paa spinosa*). *Aquaculture*

- 246 Research 49:2033-2041.
- 247 2. Zhao W, Li C, Zhang D, Wang R, Zheng Y, Zou H, Li W, Wu S, Wang G, Li M.
248 2018. *Balantidium grimi* n. sp. (Ciliophora, Litostomatea), a new species
249 inhabiting the rectum of the frog *Quasipaa spinosa* from Lishui, China. *Parasite*
250 25:29.
- 251 3. Chan H-K, Shoemaker KT, Karraker NE. 2014. Demography of *Quasipaa* frogs
252 in China reveals high vulnerability to widespread harvest pressure. *Biological*
253 *Conservation* 170:3-9.
- 254 4. Dong BJ, Zhan ZG, Zheng RQ, Chen W, Min JJ. 2015. cDNA cloning and
255 functional characterisation of four antimicrobial peptides from *Paa spinosa*. *Z*
256 *Naturforsch C J Biosci* 70:251-6.
- 257 5. Long J, Xiang J, He T, Zhang N, Pan W. 2020. Gut microbiota differences
258 during metamorphosis in sick and healthy giant spiny frogs (*Paa spinosa*)
259 tadpoles. *Lett Appl Microbiol* 70:109-117.
- 260 6. Shu Xinhua Jin Xieli Xiao Keyu Chen Keyi. 1994. W Studies on the Pathogenic
261 Bacteria of the Rotten Skin and Red leg Disease of the Bullfrog I Virulence and
262 Biological Features of *Aeromonas hydrophila*. *Journal of Natural Science of*
263 *Hunan Normal University*1994-5.
- 264 7. WU Ke-bang, DIAO Xiao-ping, HAN Xin-chou, NI Cheng-jie,WU Ying-hua.
265 2003. Study on treating skin disease of Hainan frog with Chinese herbs complex
266 agents. *Journal of Traditional Chinese Veterinary Medicine* 2003-5.
- 267 8. Hao YT, Wu SG, Xiong F, Tran NT, Jakovlic I, Zou H, Li WX, Wang GT. 2017.
268 Succession and Fermentation Products of Grass Carp (*Ctenopharyngodon*
269 *idellus*) Hindgut Microbiota in Response to an Extreme Dietary Shift. *Front*
270 *Microbiol* 8:1585.
- 271 9. Nicholson JK, Holmes E, Wilson ID. 2005. Gut microorganisms, mammalian
272 metabolism and personalized health care. *Nat Rev Microbiol* 3:431-8.
- 273 10. Leadbetter EA, Rifkin IR, Hohlbaum AM, Beaudette BC, Shlomchik MJ,
274 Marshak-Rothstein A.2002. Chromatin-IgG complexes activate B cells by dual
275 engagement of IgM and Toll-like receptors. *Nature* 416(6881):603.

- 276 11. Ringo E, Jutfelt F, Kanapathippillai P, Bakken Y, Sundell K, Glette J, Mayhew
277 TM, Myklebust R, Olsen RE. 2004. Damaging effect of the fish pathogen
278 *Aeromonas salmonicida* ssp. *salmonicida* on intestinal enterocytes of Atlantic
279 salmon (*Salmo salar* L.). *Cell Tissue Res* 318:305-11.
- 280 12. Li J, Ni J, Li J, Wang C, Li X, Wu S, Zhang T, Yu Y, Yan Q. 2014. Comparative
281 study on gastrointestinal microbiota of eight fish species with different feeding
282 habits. *J Appl Microbiol* 117:1750-60.
- 283 13. Ni J, Yan Q, Yu Y, Zhang T. 2014. Factors influencing the grass carp gut
284 microbiome and its effect on metabolism. *FEMS Microbiol Ecol* 87:704-14.
- 285 14. Li X, Zhou L, Yu Y, Ni J, Xu W, Yan Q. 2017. Composition of Gut Microbiota
286 in the Gibel Carp (*Carassius auratus gibelio*) Varies with Host Development.
287 *Microb Ecol* 74:239-249.
- 288 15. Andoh A. 2016. Physiological Role of Gut Microbiota for Maintaining Human
289 Health. *Digestion* 93:176-81.
- 290 16. Li T, Li H, Gatesoupe FJ, She R, Lin Q, Yan X, Li J, Li X. 2017. Bacterial
291 Signatures of "Red-Operculum" Disease in the Gut of Crucian Carp (*Carassius*
292 *auratus*). *Microb Ecol* 74:510-521.
- 293 17. Ni J, Li X, He Z, Xu M. 2017. A novel method to determine the minimum
294 number of sequences required for reliable microbial community analysis. *J*
295 *Microbiol Methods* 139:196-201.
- 296 18. Huang R, Li T, Ni J, Bai X, Gao Y, Li Y, Zhang P, Gong Y. 2018. Different Sex-
297 Based Responses of Gut Microbiota During the Development of Hepatocellular
298 Carcinoma in Liver-Specific *Tsc1*-Knockout Mice. *Front Microbiol* 9:1008.
- 299 19. Ni J, Huang R, Zhou H, Xu X, Li Y, Cao P, Zhong K, Ge M, Chen X, Hou B,
300 Yu M, Peng B, Li Q, Zhang P, Gao Y. 2019. Analysis of the Relationship
301 Between the Degree of Dysbiosis in Gut Microbiota and Prognosis at Different
302 Stages of Primary Hepatocellular Carcinoma. *Front Microbiol* 10:1458.
- 303 20. Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial
304 amplicon reads. *Nat Methods* 10:996-8.
- 305 21. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME

- 306 improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194-
307 200.
- 308 22. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive Bayesian classifier for
309 rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl*
310 *Environ Microbiol* 73:5261-7.
- 311 23. Dixon P.2003.VEGAN, a package of R functions for community ecology.
312 *Journal of Vegetation Science* 14(6):927-930..
- 313 24. Baruch K, Schwartz M. 2016. Circulating Monocytes in between the Gut and
314 the Mind. *Cell Stem Cell* 18:689-91.
- 315 25. Kennedy PJ, Cryan JF, Dinan TG, Clarke G. 2017. Kynurenine pathway
316 metabolism and the microbiota-gut-brain axis. *Neuropharmacology* 112:399-
317 412.
- 318 26. Budden KF, Gellatly SL, Wood DL, Cooper MA, Morrison M, Hugenholtz P,
319 Hansbro PM. 2017. Emerging pathogenic links between microbiota and the gut-
320 lung axis. *Nat Rev Microbiol* 15:55-63.
- 321 27. Shukla SD, Budden KF, Neal R, Hansbro PM. 2017. Microbiome effects on
322 immunity, health and disease in the lung. *Clinical & Translational Immunology*
323 6.
- 324 28. Zhu W, Gregory JC, Org E, Buffa JA, Gupta N, Wang Z, Li L, Fu X, Wu Y,
325 Mehrabian M, Sartor RB, McIntyre TM, Silverstein RL, Tang WHW, DiDonato
326 JA, Brown JM, Lusa AJ, Hazen SL. 2016. Gut Microbial Metabolite TMAO
327 Enhances Platelet Hyperreactivity and Thrombosis Risk. *Cell* 165:111-124.
- 328 29. Li R, Yang J, Saffari A, Jacobs J, Baek KI, Hough G, Larauche MH, Ma J, Jen
329 N, Moussaoui N, Zhou B, Kang H, Reddy S, Henning SM, Campen MJ, Pisegna
330 J, Li Z, Fogelman AM, Sioutas C, Navab M, Hsiai TK. 2017. Ambient Ultrafine
331 Particle Ingestion Alters Gut Microbiota in Association with Increased
332 Atherogenic Lipid Metabolites. *Sci Rep* 7:42906.
- 333 30. Ohlsson C, Sjogren K. 2015. Effects of the gut microbiota on bone mass. *Trends*
334 *Endocrinol Metab* 26:69-74.
- 335 31. Villa CR, Ward WE, Comelli EM. 2017. Gut microbiota-bone axis. *Crit Rev*

- Food Sci Nutr 57:1664-1672.
32. Nicolas S, Blasco-Baque V, Fournel A, Gilleron J, Klopp P, Waget A, Ceppo F, Marlin A, Padmanabhan R, Iacovoni JS, Terce F, Cani PD, Tanti JF, Burcelin R, Knauf C, Cormont M, Serino M. 2017. Transfer of dysbiotic gut microbiota has beneficial effects on host liver metabolism. *Mol Syst Biol* 13:921.
33. Xue L, He J, Gao N, Lu X, Li M, Wu X, Liu Z, Jin Y, Liu J, Xu J, Geng Y. 2017. Probiotics may delay the progression of nonalcoholic fatty liver disease by restoring the gut microbiota structure and improving intestinal endotoxemia. *Sci Rep* 7:45176.
34. Feng Q, Chen WD, Wang YD. 2018. Gut Microbiota: An Integral Moderator in Health and Disease. *Front Microbiol* 9:151.
35. Yao Z, Yang K, Huang L, Huang X, Qiuqian L, Wang K, Zhang D. 2018. Disease outbreak accompanies the dispersive structure of shrimp gut bacterial community with a simple core microbiota. *AMB Express* 8:120.
36. Fargione JE, Tilman D. 2005. Diversity decreases invasion via both sampling and complementarity effects. *Ecology Letters* 8:604-611.
37. Li T, Long M, Ji C, Shen Z, Gatesoupe FJ, Zhang X, Zhang Q, Zhang L, Zhao Y, Liu X, Li A. 2016. Alterations of the gut microbiome of largemouth bronze gudgeon (*Coreius guichenoti*) suffering from furunculosis. *Sci Rep* 6:30606.
38. Nie L, Zhou QJ, Qiao Y, Chen J. 2017. Interplay between the gut microbiota and immune responses of ayu (*Plecoglossus altivelis*) during *Vibrio anguillarum* infection. *Fish Shellfish Immunol* 68:479-487.
39. Tran NT, Zhang J, Xiong F, Wang GT, Li WX, Wu SG. 2018. Altered gut microbiota associated with intestinal disease in grass carp (*Ctenopharyngodon idellus*). *World J Microbiol Biotechnol* 34:71.
40. Horz HP, Conrads G. 2010. The discussion goes on: What is the role of Euryarchaeota in humans? *Archaea* 2010:967271.
41. Gevers D, Kugathasan S, Denson LA, Vazquez-Baeza Y, Van Treuren W, Ren B, Schwager E, Knights D, Song SJ, Yassour M, Morgan XC, Kostic AD, Luo C, Gonzalez A, McDonald D, Haberman Y, Walters T, Baker S, Rosh J, Stephens

- 366 M, Heyman M, Markowitz J, Baldassano R, Griffiths A, Sylvester F, Mack D,
367 Kim S, Crandall W, Hyams J, Huttenhower C, Knight R, Xavier RJ. 2014. The
368 treatment-naïve microbiome in new-onset Crohn's disease. *Cell Host Microbe*
369 15:382-392.
- 370 42. Keren N, Konikoff FM, Paitan Y, Gabay G, Reshef L, Naftali T, Gophna U.
371 2015. Interactions between the intestinal microbiota and bile acids in gallstones
372 patients. *Environ Microbiol Rep* 7:874-80.
- 373 43. Konikoff T, Gophna U. 2016. Oscillospira: a Central, Enigmatic Component of
374 the Human Gut Microbiota. *Trends Microbiol* 24:523-524.
- 375 44. Wang HP,Wang YL,Zheng YJ,LI P,Deng JK.2016. High-throughput sequencing
376 of intestinal flora of infants with eczema. *Chinese Journal of Microecology*
377 28(7):751-5.
- 378 45. Thomson NR, Crow MA, McGowan SJ ,Cox A, Salmond GP.2000.
379 Biosynthesis of carbapenem antibiotic and prodigiosin pigment in *Serratia* is
380 under quorum sensing control. *Molecular microbiology* 36(3):539-556..
- 381 46. Thomson NR, Cox A,Bycroft BW, Stewart GS, Williams P, Salmond GP. 1997.
382 The Rap and Hor proteins of *Erwinia*, *Serratia* and *Yersinia* a novel subgroup in
383 a growing superfamily of proteins regulating diverse physiological processes in
384 bacterial pathogens. *Molecular microbiology* 26(3):531-544..
- 385 47. Bourassa MW, Alim I, Bultman SJ, Ratan RR. 2016. Butyrate, neuroepigenetics
386 and the gut microbiome: Can a high fiber diet improve brain health? *Neurosci*
387 *Lett* 625:56-63.
- 388 48. Engels C, Ruscheweyh HJ, Beerenwinkel N, Lacroix C, Schwab C. 2016. The
389 Common Gut Microbe *Eubacterium hallii* also Contributes to Intestinal
390 Propionate Formation. *Front Microbiol* 7:713.
- 391 49. Nakajima M, Shirokawa M, Miyakuni Y, Nakano T, Goto H. 2017. Giant
392 Iliopsoas Abscess Caused by *Morganella Morganii*. *Am J Case Rep* 18:395-398.
- 393 50. Higgins LM, Frankel G, Douce G, Dougan G, MacDonald TT.1999. *Citrobacter*
394 *rodentium* Infection in Mice Elicits a Mucosal Th1 Cytokine Response and
395 Lesions Similar to Those in Murine Inflammatory Bowel Disease . *Infection and*

immunity 67(6):3031-3039.

51. Yang HT, Zou SS, Zhai LJ, Wang Y, Zhang FM, An LG, Yang GW. 2017. Pathogen invasion changes the intestinal microbiota composition and induces innate immune responses in the zebrafish intestine. *Fish Shellfish Immunol* 71:35-42.
52. She R, Li TT, Luo D, Li JB, Yin LY, Li H, Liu YM, Li XZ, Yan QG. 2017. Changes in the Intestinal Microbiota of Gibel Carp (*Carassius gibelio*) Associated with Cyprinid herpesvirus 2 (CyHV-2) Infection. *Curr Microbiol* 74:1130-1136.
53. Ding ZF, Cao MJ, Zhu XS, Xu GH, Wang RL. 2017. Changes in the gut microbiome of the Chinese mitten crab (*Eriocheir sinensis*) in response to White spot syndrome virus (WSSV) infection. *J Fish Dis* 40:1561-1571.

Figure legends

Fig.1

A and B are the frogs of the early stage of the rotten-skin disease; C and D are the later stages; E and F are the liver and gallbladder of healthy *P.spinosa*; G and H are the lesions in the liver and gallbladder of diseased *P.spinosa*.

Fig.2

Diversity analysis of the gut microbiota *P. spinosa* between healthy and diseased groups: (A) Chao1; (B) Goods-coverage; (C) observed-species; (D) PD-whole-tree; (E) Shannon index; (F) Simpson index. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. Disease, the gut microbiota from six diseased *P. spinosa*. Health, the gut microbiota from five healthy *P. spinosa*. P values show the difference between the groups. $P>0.05$ represents the little difference between groups, $p<0.05$ indicates the significant difference between the groups, $p<0.01$ indicates the extremely different between the groups. Data are the mean $\pm$ SE

Fig.3

The inner circular diagram (A) shows the relative abundance of different phyla in *P.spinosa* gut samples. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. D, the gut microbiota from six diseased *P. spinosa*. H, the gut microbiota from five healthy *P. spinosa*. the significant difference in phylum between health and disease groups (B). The STAMP based on the top 10 phyla of the gut microbiota compositions analyzed the

significantly different($p < 0.05$) phylum between the groups.

Fig.4

PCA profile(A). PCA was conducted based on the all genus microbial communities showing the differentiation of the *P.spinosa* gut microbiota communities between the health and disease group. PCoA profile(B)、 and UPGMA cluster graph (C) based on the unweight unifracs distance showing the differentiation of the *P.spinosa* gut microbiota communities between each individual. The PCoA was conducted based on the all genus microbial communities. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. Disease, the gut microbiota from 6 diseased *P. spinosa*. Health, the gut microbiota from 5 healthy *P. spinosa*.

Fig.5

LEfSe profile showing differences in healthy and diseased *P.spinosa* gut microbial communities. LEfSe analysis was conducted based on the top 40 genus compositions of the *P. spinosa* gut microbiota. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. Disease, the gut microbiota from 6 diseased *P. spinosa*. Health, the gut microbiota from 5 healthy *P. spinosa*.

Table 1. Recent studies about gut microbiota between healthy and diseased samples

Species	Disease	Finding	Authors
Grass carp (<i>Ctenopharyngodon idellus</i>)	Enteritis	The association between changes of the gut microbiota and enteritis in grass carp	(39)
Crucian Carp (<i>Carassius auratus</i>)	Red-Operculum Disease	The surge of some potential pathogens as bacterial signatures that were associated with “red-operculum” disease in crucian carps	(16)
largemouth bronze gudgeon (<i>Coreius guichenoti</i>)	Furunculosis	The presence of healthy carriers of pathogenic <i>Aeromonas salmonicida</i> among the farmed fish, and the gut appeared as a probable infection source for furunculosis in largemouth bronze gudgeon.	(37)
Ayu (<i>Plecoglossus altivelis</i>)	<i>Vibrio anguillarum</i> infection	<i>Vibrio anguillarum</i> infection substantially disrupted the compositions and interspecies interaction of ayu gut bacterial community.	(38)
Zebrafish(<i>Barc hydanio rerio</i> var)	<i>Aeromonas hydrophila</i> infected	the invasion of pathogen could change the gut microbiota composition and induce gut innate immune responses in zebrafish	(51)
Gibel Carp (<i>Carassius gibelio</i>)	<i>Cyprinid herpesvirus 2</i> (CyHV-2) Infection	The composition was dramatically altered following CyHV-2 infection ; <i>Plesiomonas</i> was highly abundant in infected samples, and could be used as a microbial biomarker for CyHV-2 infection	(52)
Chinese mitten crab (<i>Eriocheir sinensis</i>)	White spot syndrome virus (WSSV) infection	Changes in gut microbiome were closely associated with the severity of WSSV infection and that indicator taxa could be used to evaluate the crab health status.	(53)
Shrimp (<i>Litopenaeus vannamei</i>)	Acute hepatopancreatic necrosis disease	Shrimp health is highly relevant to the homeostasis of its gut bacterial community.	(35)

Figures

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Table 1. Recent studies about gut microbiota between healthy and diseased samples

Species	Disease	Finding	Authors
Grass carp (<i>Ctenopharyngodon idellus</i>)	Enteritis	The association between changes of the gut microbiota and enteritis in grass carp	(39)
Crucian Carp (<i>Carassius auratus</i>)	Red-Operculum Disease	The surge of some potential pathogens as bacterial signatures that were associated with “red-operculum” disease in crucian carps	(16)
largemouth bronze gudgeon (<i>Coreius guichenoti</i>)	Furunculosis	The presence of healthy carriers of pathogenic <i>Aeromonas salmonicida</i> among the farmed fish, and the gut appeared as a probable infection source for furunculosis in largemouth bronze gudgeon.	(37)
Ayu (<i>Plecoglossus altivelis</i>)	<i>Vibrio anguillarum</i> infection	<i>Vibrio anguillarum</i> infection substantially disrupted the compositions and interspecies interaction of ayu gut bacterial community.	(38)
Zebrafish(<i>Barc hydanio rerio</i> var)	<i>Aeromonas hydrophila</i> infected	the invasion of pathogen could change the gut microbiota composition and induce gut innate immune responses in zebrafish	(51)
Gibel Carp (<i>Carassius gibelio</i>)	<i>Cyprinid herpesvirus 2</i> (CyHV-2) Infection	The composition was dramatically altered following CyHV-2 infection ; <i>Plesiomonas</i> was highly abundant in infected samples, and could be used as a microbial biomarker for CyHV-2 infection	(52)
Chinese mitten crab (<i>Eriocheir sinensis</i>)	White spot syndrome virus (WSSV) infection	Changes in gut microbiome were closely associated with the severity of WSSV infection and that indicator taxa could be used to evaluate the crab health status.	(53)
Shrimp (<i>Litopenaeus vannamei</i>)	Acute hepatopancreatic necrosis disease	Shrimp health is highly relevant to the homeostasis of its gut bacterial community.	(35)

Figures

Fig.1

A and B are the frogs of the early stage of the rotten-skin disease; C and D are the later stages; E and F are the liver and gallbladder of healthy *P.spinosa*; G and H are the lesions in the liver and gallbladder of diseased *P.spinosa*.

Fig.2

Diversity analysis of the gut microbiota *P. spinosa* between healthy and diseased groups: (A) Chao1; (B) Goods-coverage; (C) observed-species; (D) PD-whole-tree; (E) Shannon index; (F) Simpson index. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. Disease, the gut microbiota from six diseased *P. spinosa*. Health, the gut microbiota from five healthy *P. spinosa*. P values show the difference between the groups. $P > 0.05$ represents the little difference between groups, $p < 0.05$ indicates the significant difference between the groups, $p < 0.01$ indicates the extremely different between the groups. Data are the mean $\pm$ SE

Fig.3

A

B

The inner circular(A) diagram shows the relative abundance of different phyla in *P. spinosa* gut samples. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. D, the gut microbiota from six diseased *P. spinosa*. H, the gut microbiota from five healthy *P. spinosa*. the significant difference in phylum between health and disease groups(B). The STAMP based on the top 10 phyla of the gut microbiota compositions analyzed the significantly different($p < 0.05$) phylum between the groups.

Fig.4

PCA profile(A). PCA was conducted based on the all genus microbial communities showing the differentiation of the *P.spinosa* gut microbiota communities between the health and disease group. PCoA profile(B)and UPGMA cluster graph (C) based on the unweight unifrac distance showing the differentiation of the *P.spinosa* gut microbiota communities between each individual. The PCoA was conducted based on the all genus microbial communities. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. Disease, the gut microbiota from 6 diseased *P. spinosa*. Health, the gut microbiota from 5 healthy *P.*

spinosa.

Fig.5

LefSe profiles showing differences in healthy and diseased *P. spinosa* gut microbial communities. LefSe analysis was conducted based on the top 40 genus compositions of the *P. spinosa* gut microbiota. The gut microbiota of *P. spinosa* was collected from approximately 0.3g samples of the hindgut of each individual. Disease, the gut microbiota from 6 diseased *P. spinosa*. Health, the gut microbiota from 5 healthy *P. spinosa*.