Archives of Virology, Annotated Sequence Record

Isolation and complete genome sequencing of *Pectobacterium* phage Jarilo, representing a novel genus of bacteriophages within the subfamily *Autographivirinae*

Julie Stenberg Pedersen^{1,*}, Alexander Byth Carstens¹, Amaru Miranda Djurhuus¹, Witold Kot¹, Lars Hestbjerg Hansen¹

¹Department of Plant and Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg, Danmark

*Corresponding author:

Julie Stenberg Pedersen
e-mail: hzf242@alumni.ku.dk
Tel.: +4551847142

Key words: *Pectobacterium* phage, *Pectobacterium carotovorum*, biocontrol agents, genome analysis

Word counts: 125 words (Abstract); 2030 words (Main text)

Figures: 2

Supplementary figures: 1

Abstract

Pectobacterium carotovorum is the causative agent of bacterial soft rot on various plant species. The use of phages for plant disease control have gained increased awareness over the past years. We here describe the isolation and characterization of *Pectobacterium* phage Jarilo, representing a novel genus of bacteriophages within the subfamily *Autographivirinae*. Jarilo possesses a double-stranded DNA genome of 40557 bp with a G+C% content of 50.08% and 50 predicted open reading frames (ORFs). Gene synteny and products seem to be somewhat conserved between *Pectobacterium* phage Jarilo and Enterobacteria phage T7, but limited nucleotide similarity is found between Jarilo and other phages within the subfamily *Autographivirinae*. We propose *Pectobacterium* phage Jarilo as the first member of a new genus of bacteriophages within the subfamily *Autographivirinae*.

Introduction

Pectobacterium carotovorum is part of the family *Pectobacteriaceae*, which members include important bacterial plant pathogens involved in a wide range of diseases[1]. *Pectobacteriaceae* are Gram-negative, facultative anaerobes, that are non-spore forming and produce extracellular enzymes, involved in their pathogenicity[2]. *P. carotovorum* (formerly *Erwinia carotovora*) is able to form soft rot disease symptoms in various plant species. In potatoes *P. carotovorum* is the causative agent of black leg and tuber rot during storage, affecting postharvest loss[3,4]. Control mechanism of bacterial soft rot is limited and is primarily based on sanitary conditions and good agricultural practice[5]. Due to limited control mechanism, increased attention have been directed towards biological control agents[6]. Phages have been proposed as promising biocontrol agents towards *Pectobacteriaceae* in several studies[7–12]. In this report, we describe the isolation and characterization of *Pectobacterium* phage Jarilo, representing a new genus of bacteriophages within the subfamily *Autographivirinae*.

Materials and methods

Jarilo was isolated from an organic waste sample as described previously[12], using *Pectobacterium carotovorum* DSM 30170 as host. Prior to isolation virus particles in the organic waste sample was concentrated by polyethylene glycol precipitation, as described elsewhere[13]. Phage DNA was isolated and sequencing libraries prepared using direct plaque sequencing protocol[14], with the modifications described previously[15]. Reads from sequencing were trimmed and *de novo* assembled using CLC

Genomic Workbench (10.1.1). Open reading frame prediction were automatically executed using DNA Master (version 5.0.2), with GeneMark[16] and Glimmer[17] as gene caller, and were corrected manually. Putative gene functions were manually curated using BLASTp[18] and HHpred[19] with the databases Pfam (version 32.0), SCOP70 (version 1.7.5) and pdb70. tRNA sequences were searched for using tRNAscan-SE[20].

Results

Pectobacterium phage Jarilo represents a new genus within the subfamily *Autographivirinae*, with limited DNA sequence similarity (<50%) with other phages within the same subfamily (Fig. 1)[21]. Pectobacterium phage Jarilo has a double-stranded DNA genome of 40557 bp with a G+C% content of 50.08%. In the genome there is 50 predicted open reading frames (ORFs), whereas the majority (30/50) could be assigned with a putative function due to homology with other known phages within the subfamily *Autographivirinae*. No tRNAs were detected.

Based on bioinformatics predictions, ORFs were categorized into five putative gene functions; hypothetical protein (grey), DNA replication and metabolism (red), lysis proteins (yellow), others/unknown (green) and morphogenesis (blue) (Fig. 2). The gene synteny of the genome of Jarilo is conserved among phages within the subfamily *Autographivirinae*, as well as many of the structural genes (Fig. 2). An insertion of a putative endonuclease VII appear to have happened within the putative DNA polymerase, in the genome of Jarilo. This insertion separates the DNA polymerase into two ORFs, both ORFs share high sequence similarity with the DNA polymerase of related phages (Fig. 2). Yersinia phage YpP-G, also part of the *Autographivirinae* subfamily, encodes a endonuclease VII, which gives the highest score against the putative endonuclease VII encoded by Jarilo, using blastp (Suppl. fig. 1)[18,22]. YpP-G display a similar gene arrangement with an endonuclease VII within the gene encoding the DNA polymerase (Suppl. fig. 1). Furthermore is the same phenomenon described in another phage within the subfamily *Autographivirinae*, Pf-WMP3[23]. A endonuclease VII insertion into the DNA polymerase have been proven as part of a self-splicing intron[24], but have not been demonstrated as part of a selfsplicing intron within the *Autographivirinae* subfamily. The gene encoding the major capsid protein as well as the gene following, categorized as a hypothetical gene, in Jarilo, resembles the capsid gene in T7 with some similarity. The capsid gene in T7 has two products, a major (10A) and a minor (10B), produced by frameshifting into the -1 frame near the end of 10A[25]. The gene encoding head to tail

joining protein appear to be divided in two (Fig. 2), which could also be a result of frameshifting, as in the capsid protein of T7, or due to a nonsense mutation. Genes categorized as morphogenesis, in Jarilo, are conserved among phages within the subfamily *Autographivirinae*, however the tail-fiber protein is conserved to a lesser extent, which could represent different receptor preferences (Fig. 2). Interestingly, Jarilo, T7 and AP10 shares a conserved region at the terminal of the genome, not representing an ORF (Fig. 2). Like other phages within the subfamily *Autographivirinae*, Jarilo encodes a RNA polymerase[26] and do not encode any detectable integrase.

Based on the findings we propose Pectobacterium phage Jarilo as the first representation of a new genus of bacteriophages belonging to the subfamily *Autographivirinae*, family *Podoviridae*, order *Caudovirales*.

Nucleotide sequence accession number

The GenBank accession number for Pectobacterium phage Jarilo is MH059637.1.

References

1. Li X, Ma Y, Liang S, Tian Y, Yin S, Xie S, et al. Comparative genomics of 84 *Pectobacterium* genomes reveals the variations related to a pathogenic lifestyle. *BMC Genomics*. 2018;19: 889. doi:10.1186/s12864-018-5269-6
2. Pérombelon MCM. Potato diseases caused by soft rot erwinias: an overview of pathogenesis. *Plant Pathol*. 2002;51: 1–12. doi:10.1046/j.0032-0862.2001.Shorttitle.doc.x
3. de Haan EG, Dekker-Nooren TCEM, van den Bovenkamp GW, Speksnijder AGCL, van der Zouwen PS, van der Wolf JM. *Pectobacterium carotovorum* subsp. *carotovorum* can cause potato blackleg in temperate climates. *Eur J Plant Pathol*. 2008;122: 561. doi:10.1007/s10658-008-9325-y
4. Kang HW, Kwon SW, Go SJ. PCR-based specific and sensitive detection of *Pectobacterium carotovorum* ssp. *carotovorum* by primers generated from a URP-PCR fingerprinting-derived polymorphic band. *Plant Pathol*. 2003;52: 127–133. doi:10.1046/j.1365-3059.2003.00822.x
5. Rahman MM, Ali ME, Khan AA, Akanda AM, Uddin MK, Hashim U, et al. Isolation, characterization, and identification of biological control agent for potato soft rot in Bangladesh. *ScientificWorldJournal*. 2012;2012: 723293. doi:10.1100/2012/723293
6. Byrne JM, Dianese AC, Ji P, Campbell HL, Cuppels DA, Louws FJ, et al. Biological control of bacterial spot of tomato under field conditions at several locations in North America. *Biol Control*. 2005;32: 408–418. doi:https://doi.org/10.1016/j.biocontrol.2004.12.001
7. Lim J-A, Shin H, Lee DH, Han S-W, Lee J-H, Ryu S, et al. Complete genome sequence of the *Pectobacterium carotovorum* subsp. *carotovorum* virulent bacteriophage PM1. *Arch Virol*. 2014;159: 2185–2187. doi:10.1007/s00705-014-2005-7
8. Lee DH, Lee J-H, Shin H, Ji S, Roh E, Jung K, et al. Complete genome sequence of *Pectobacterium carotovorum* subsp. *carotovorum* bacteriophage My1. *J Virol*. 2012;86: 11410–11411. doi:10.1128/JVI.01987-12
9. Lim J-A, Jee S, Lee DH, Roh E, Jung K, Oh C, et al. Biocontrol of *Pectobacterium carotovorum* subsp. *carotovorum* using bacteriophage PP1. *J Microbiol Biotechnol*. 2013;23: 1147–1153.

doi:10.4014/jmb.1304.04001

10. Lim J-A, Lee DH, Heu S. Isolation and Genomic Characterization of the T4-Like Bacteriophage PM2 Infecting *Pectobacterium carotovorum* subsp. *carotovorum*. *plant Pathol J.* 2015;31: 83–89. doi:10.5423/PPJ.NT.09.2014.0099
11. Lim J-A, Heu S, Park J, Roh E. Genomic characterization of bacteriophage vB_PcaP_PP2 infecting *Pectobacterium carotovorum* subsp. *carotovorum*, a new member of a proposed genus in the subfamily Autographivirinae. *Arch Virol.* 2017;162: 2441–2444. doi:10.1007/s00705-017-3349-6
12. Carstens AB, Djurhuus AM, Kot W, Hansen LH. A novel six-phage cocktail reduces *Pectobacterium atrosepticum* soft rot infection in potato tubers under simulated storage conditions. *FEMS Microbiol Lett.* 2019;366. doi:10.1093/femsle/fnz101
13. Sambrook J, Fritsch EF, Maniatis T. *Molecular cloning: A laboratory manual*. 2nd ed. Analytical Biochemistry. Cold Spring Harbor, USA: Cold Spring Harbor Laboratory Press; 1990.
14. Kot W, Vogensen FK, Sørensen SJ, Hansen LH. DPS - A rapid method for genome sequencing of DNA-containing bacteriophages directly from a single plaque. *J Virol Methods.* 2014;196: 152–156. doi:10.1016/j.jviromet.2013.10.040
15. Carstens AB, Kot W, Lametsch R, Neve H, Hansen LH. Characterisation of a novel enterobacteria phage, CAjan, isolated from rat faeces. *Arch Virol.* 2016;161: 2219–2226. doi:10.1007/s00705-016-2901-0
16. Besemer J, Lomsadze A, Borodovsky M. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res.* 2001;29: 2607–2618. doi:10.1093/nar/29.12.2607
17. Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics.* 2007;23: 673–679. doi:10.1093/bioinformatics/btm009
18. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

19. Hildebrand A, Remmert M, Biegert A, Soding J. Fast and accurate automatic structure prediction with HHpred. *Proteins*. 2009;77 Suppl 9: 128–132. doi:10.1002/prot.22499
20. Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res*. 2016;44: W54-7. doi:10.1093/nar/gkw413
21. Adriaenssens E, Brister JR. How to Name and Classify Your Phage: An Informal Guide. *Viruses*. 2017;9. doi:10.3390/v9040070
22. Rashid MH, Revazishvili T, Dean T, Butani A, Verratti K, Bishop-Lilly KA, et al. A *Yersinia pestis*-specific, lytic phage preparation significantly reduces viable *Y. pestis* on various hard surfaces experimentally contaminated with the bacterium. *Bacteriophage*. 2012;2: 168–177. doi:10.4161/bact.22240
23. Liu X, Kong S, Shi M, Fu L, Gao Y, An C. Genomic analysis of freshwater cyanophage Pf-WMP3 Infecting cyanobacterium *Phormidium foveolarum*: the conserved elements for a phage. *Microb Ecol*. 2008;56: 671–680. doi:10.1007/s00248-008-9386-7
24. Ceyssens P-J, Hertveldt K, Ackermann H-W, Noben J-P, Demeke M, Volckaert G, et al. The intron-containing genome of the lytic *Pseudomonas* phage LUZ24 resembles the temperate phage PaP3. *Virology*. 2008;377: 233–238. doi:10.1016/j.virol.2008.04.038
25. Condrón BG, Atkins JF, Gesteland RF. Frameshifting in gene 10 of bacteriophage T7. *J Bacteriol*. 1991;173: 6998–7003. doi:10.1128/jb.173.21.6998-7003.1991
26. Eriksson H, Maciejewska B, Latka A, Majkowska-Skrobek G, Hellstrand M, Melefors O, et al. A suggested new bacteriophage genus, “Kp34likevirus”, within the Autographivirinae subfamily of Podoviridae. *Viruses*. 2015;7: 1804–1822. doi:10.3390/v7041804
27. Agren J, Sundstrom A, Hafstrom T, Segerman B. Gegenees: fragmented alignment of multiple genomes for determining phylogenomic distances and genetic signatures unique for specified target groups. *PLoS One*. 2012;7: e39107. doi:10.1371/journal.pone.0039107
28. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. *Bioinformatics*. 2011;27: 1009–1010. doi:10.1093/bioinformatics/btr039

Figures

Organism	1	2	3	4	5	6
1: Enterobacteria phage T7	100.0	1.69	2.53	2.83	5.24	6.32
2: Pectobacterium phage Jarilo	1.64	100.0	4.05	2.03	2.13	2.4
3: Yersinia phage AP10	2.68	4.29	100.0	8.4	5.13	6.42
4: Klebsiella phage KOX3	2.32	1.47	8.72	100.0	2.7	3.35
5: Enterobacter phage E-3	7.43	2.83	6.04	3.92	100.0	72.43
6: Citrobacter phage SH2	6.01	1.97	6.61	3.17	58.49	100.0

Figure 1. Pectobacterium phage Jarilo possesses limited DNA sequence similarity with other known *Autographivirinae* phages

All vs. all nucleotide comparison between six genomes of bacteriophages within the subfamily *Autographivirinae*, using the Gegenees software (fragment size: 500 bp, step size: 500 bp)[27]. Phages showing highest scores, using blastn[18], against the genome of Jarilo were chosen together with phage T7 as a reference genome. Colors indicate similarity between phages, from similar (green) to less similar (red). Similarity scores are based on blastn. Following phage genomes were used (with genbank accession numbers); Yersinia phage AP10 (KT852574.1), Klebsiella phage KOX3 (MN101216.1), Enterobacter phage E-3 (KP791806.1), Citrobacter phage SH2 (KU687348.1), Enterobacteria phage T7 (V01146.1).

Figure 2. Comparative genomics reveals conserved gene synteny among phages within the subfamily *Autographivirinae*, as well as a high degree of sequence homology among structural proteins

Comparative genomics of six genomes of bacteriophages within the subfamily *Autographivirinae* using blastn in the Easyfig software (version 2.2.3)[28]. Phages showing highest scores, using blastn[18], against the genome of Jarilo were chosen together with phage T7 as a reference genome. Putative gene functions of the genome of Jarilo were divided into five categories; hypothetical proteins (grey), DNA replication and metabolism (red), lysis proteins (yellow), others/unknown (green) and morphogenesis (blue). Following phage genomes were used (with Genbank accession numbers); Yersinia phage AP10

(KT852574.1), Klebsiella phage KOX3 (MN101216.1), Enterobacter phage E-3 (KP791806.1), Citrobacter phage SH2 (KU687348.1), Enterobacteria phage T7 (V01146.1).