Competitive approach of invasive cocklebur (*Xanthium strumarium*) with native weed species diversity in Northeast China

Short Title: Invasive weed competitions with native diversity

Iqbal, M. F.¹, Liu, M. C.¹, Iram, A.², Feng, Y. L.^{1*}

¹Liaoning Key Laboratory for Biological Invasions and Global Changes, College of Bioscience and Biotechnology, Shenyang Agricultural University Shenyang, 110866, Liaoning Province, China

²Government of the Punjab Education Department, Gujranwala 52250, Punjab, Pakistan

*Corresponding author: e-mail: yl_feng@tom.com; fyl@syau.edu.cn;

Phone: +86-24-8848-7163; fax: +86-24-8849-2799

Abstract

Xanthium strumarium is yearly weed local to North America and is presently an obtrusive species. The intrusive weed can contend with local decent variety may turn into a hazardous weed for the agrarian profitability and rangeland biological systems. The present examination researched the challenge of intrusive and local weed populaces inside two straight out variables in matched quadratic rings (treatment/invasive with control/non-invasive). The present investigation was led at four unique areas of northeast China to discover competition of *Xanthium strumarium* on 40 paired matched vegetations in same condition and living space conditions. The number of species (NOS) occurred in quadratic ring; abundance (A); Simpsons Diversity Index (SDI); Margalef's Richness Index (MRI) compared between invasive and non-invasive quadrates by t-test was recorded significant ($P < 0.05$) suggestion of competitions between plant communities. The abundance in communities decreased significantly in invasive compared to non-invasive quadrate gave an indication about low productivity of plant species due to *Xanthium strumarium*. Rarefaction bend with respect to coefficient of determination (R^2) explored in the overviewed network (0.86) proposed that there is a solid positive polynomial connection between various weed families. Greatest difference list (87.06%) recorded in Huailai province followed by Yangyuan (44.43%), Zhangjiakou (40.13%) and at Fushun (29.02%). Significant ($P < 0.05$) maximum global R demonstrated high species decent variety was found in Huailai area (0.943) trailed by Zhangjiakou. Significant ($P < 0.05$) density of native weed was recorded in non-invasive quadrate which was comparable to the invasive quadrate. Finally invasive *Xanthium strumarium* compete with native weeds diversity created significant threat to the natural diversity. Most extreme thickness of weed species gave cautioning that the predominant edaphic and natural states of the uneven regions are profoundly favorable for the dispersion and development of the weed in future.

Keywords: Species density; Ecological indices; Evenness; community ecosystem; Richness index

Introduction

Invasive weeds are presenting critical danger to the biodiversity and biological system working [1-3]. Biological invasions assumed a crucial job in all biological systems [4] however, *X. strumarium* is creating hindrance in the natural vegetations by its competition for food, shelter and light [5]. Biological invasions are considered as succeeding possible hazard to the natural diversity [6]. The invasive weeds reckoned one of the most terrible invaders in the planet characterized to their massive biomass [7,8]. The loss in biodiversity plays a vital role in global climatic change scenario [9] through significant effects on ecosystem [10,11]. These effects adapted drastically by biotic relationship between trophic levels [12,13].

Cocklebur (*Xanthium strumarium* L.) belongs to family Asteraceae are monoecious, annual herbaceous broadleaved, tap rooted, ridged, rough and hairy plant (Venodha, 2016 [14,15] native to North America and Argentina [16-18]. *Xanthium* germinates at broad spectrum temperatures

and can invade great areas of marshland and drought resistant species [19]. This weed compete with native weed populations in China are known invader and high potential for colonizing in new areas [20]. *Xanthium* is a harmful invasive weed introduced in Beijing [21] and spread drastically into six provinces [22,23] at China. These provinces are located on the coastal and border area which is previously described the hypothesis that this invasive weed was transported to China by means of intercontinental traffic [22].

This weed basically present in the northeastern territories (Liaoning, Hebei, Beijing and Shandong) having calm mainland overwhelming precipitation with common climate, the southeastern area (Guangdong and Guangxi) is subtropical and northwestern Territory (Xinjiang) having unmistakable mild semiarid particular climatic conditions. Because of wide range climatic conditions, this weed has the ability to adjust essentially to assorted natural condition. Natural displaying has anticipated that this invasive weed have propensity to spread different zones in China [23].

It is a troublesome and problematic noxious weed in many agricultural crops such as maize; common bean; mustard, teff; bread wheat; sweet potato; peanut; barley; chickpea; lentil; Soybean and cotton in many part of the world [5,24]. However the infestation of *X. strumarium* causing the utmost reduction in yield [25]. *X. strumarium* yield losses 31% upto 39% in groundnut crop having four plants in one quadratic ring however 27% yields losses was recorded in maize density 4.7m⁻² in row planted crop [25]. It invades economic impact on fallow lands resulting in decline the fodder production ultimately affecting in livestock production especially on pig, cattle and sheep [26-28]. The dispersal mechanism of *X. strumarium* was investigated by seed and flood, animals, wind, vehicles and human clothing however its long dormancy period may be helpful for its pressure [5].

Native plants are important in rangeland ecosystems that evolved to grow in local conditions, however then even do not require any specific irrigations and foliar nutrients. The leaves of the native plants have high amounts of inexpensive source of nutrients for human as well as animal consumptions in developing countries [29].

The aim of this study is to identify invasive and native weeds and to predict the appropriate habitation and ecological distribution of *X. strumarium*.

The research highlights and hypothesis are as under:

1. Impact of invasive cocklebur in native diversity of weed in natural ecosystem
2. Maximum weed density gave the indication that prevailing environmental conditions of the vicinity are conducive for the distributions of invasive weeds.
3. The impact of *Xanthium strumarium* invasion was investigated at different locations.
4. Co-efficient of determination (R^2) suggested strong positive relationship between different families in ecosystem.

The field survey conducted to evaluate the impact and competitions of invasive and non-invasive weeds present in selected quadratic rings at different sites in same habitat and environmental cues under Liaoning Key Laboratory for Biological Invasion and Global Changes, College of Bioscience and Biotechnology, Shenyang Agricultural University, Shenyang, Northeast China.

Materials and Methods

The present study was conducted during peak months (August-September) in four different locations of northeast China by line transects ecological method [30] during 2018 to investigate competitions of *Xanthium strumarium* and native weeds present in invasive and non invasive quadrates (Figure 1). These locations (Fushun County, Zhangjiakou, Huailai County, and Yangyuan County), however, Fushun County located in Liaoning Province with comparatively

high precipitation (mm); however, the other three locations situated in Hebei Province located between Beijing and Inner Mangolia having comparatively 50% low precipitation (Figure 2). Due to dynamic climate conditions and combination of hilly areas, these regions are rich biodiversity of native and invasive populations of weeds.

The two factors were studied (Invasive versus nearby site non-invasive) keeping in view identical ecological circumstances having 20m in a radius [31]. On visual observations where the *Xanthium strumarium* showed its dominance considered as treatment (Invasive) and 2nd vegetation plot selected near invasive treatment considered as the non-invasive/control [19]. Ten pairs of plots selected from each site and sum 40-paired plots investigated. In each selected site, ten quadrates each having 100x100cm size selected and all the individual weed populations recorded in both invasive and non-invasive quadrates (Table 1). This procedure was repeated three times in a site to reduce confounding factors. The geographic coordinates of four different paired sites (Figure 1) were recorded by GPS system and ArcGIS version 9.2 software (ESRI) used to map out the distribution of studied sites [32].

Bar graph regarding rarefaction described the number of individual species at species level suggested species richness in the studied sites calculated by results of samplings. After collecting the species population data were focused on multivariate analysis with invasive and non invasive quadrate (used factor) using PRIMER 7 software [33], this data sets was used to ordinate the similarity between these quadrates based on Bray-Curtis dissimilarity [31]. The competition was calculated by similarity analysis and percentage [34,35]. According to these techniques the impact of each weed species contributing to dissimilarity between groups were calculated [36]. Margalef's Richness Index (MRI) was calculated by counting one and sum total of all species in ten quadrates (Sinha, 2017), Simpson Diversity Index (SDI) was also calculated at species level [31,37]. The abundance and density of each weed recorded in studied quadratic rings by dividing it with sum total of quadratic rings studied, however number of species were calculated directly by counting method [31,38-42]. Comparison of number of weed species present in invasive and non invasive quadrate was calculated [43]. The total weed species present in a selected community subjected to ANOVA and the mean difference between Invasive and non-invasive quadrate calculated by t-test at 5% level of significance.

Results

The rarefaction curve indicated that the individuals of Asteraceae family are more diversified showed highly positive polynomial *Coefficient of determination* $R^2 = 0.88$ compared to other species studied (Figure 3). During this survey sum of 15 plant species from 9 genera were found, however 10 species were recorded in non invasive quadrate and 12 from the plots with invasive quadrate with varying number of individuals (Table 1). Significant high ($P<0.05$) number of species investigated at Fushun (Figure 4A); Zhangjiakou (Figure 4C) and Yangyuan (Figure 4D) locations in non invasive quadratic ring compared to invasive. However, significant ($P<0.05$) maximum weed species abundance was recorded at Fushun (4.00); Huailai (9.00) and Yangyuan county (3.00) in non invasive quadratic ring. Other factor regarding Simpson's Diversity Index (SDI); Margalef's Richness Index (MRI) gave the clear-cut indication that *Xanthium strumarium* present in invasive quadrate compete positively. Maximum SDI investigated higher diversity index or probability between invasive and non invasive quadratic rings, however, 0.99 and 0.97 values indicated more diversified species in the study areas. Significant ($P<0.05$) result in weed density was recorded in non-invasive quadratic ring. On the other side invasive quadrate investigated non-significant ($P<0.05$) effect and recorded high competition in density between two groups of weeds at Zhangjiakou (Figure 5A). In Huailai county significant ($P<0.05$) result

was investigated in non-invasive quadratic ring. The density of weed populations in invasive quadratic ring differed non significantly (Figure 5B). In yangyuan, the density of non-invasive quadrate differed significantly ($P<0.05$) with invasive ring (Figure 5C). Same trend but highly significant density was investigated in non-invasive quadratic ring, which was comparable to the invasive quadrate. Maximum dissimilarity index (87.06%) recorded in Huailai county followed by Yangyuan (44.43%), Zhangjiakou (40.13%) and at Fushun (29.02%). Significant ($P<0.05$) maximum global R indicated high species diversity found in Huailai county (0.943) followed by Zhangjiakou. Similarity (%) data analysis described that weeds computed average dissimilarity with invasive and non-invasive quadrates (Table 2). SIMPER analysis pooled data based on average abundance of each weed species illustrated that *Malva verticilata* (10.40) contributed maximum in non-invasive quadrate followed by *Chenopodium album* (9.00); *Avena sativa* (7.10) and *Seteria viridis* (6.80) compared to invasive quadrate at Yangyuan County. However, *Malva verticilata* (23.94%) contributed maximum both quadratic rings compared to *Seteria viridis* (20.80%). *Seteria viridis* (14.00%; 40.82%) contributed maximum at Huailai and Fushun area followed by *Seteria febrei* (24.40%) at Zhangjiakou. A total of 12 weeds species in different families documented from invasive and non invasive forty paired quadratic rings in different studied area. These results clearly indicate that at first level Primer divided the weed populations of the whole study area into one major community at association level and that further divided into two sub communities at sub association level (Table 1). The similarity analysis found significant degree of variation between species compositions in invasive & non invasive quadrates having global statistics R values of 0.943 ($P<0.01$), 0.779 ($P<0.01$), 0.62 ($P<0.01$), 0.553 ($P<0.01$) for Huailai, Zhangjiakou, Yangyuan and Fushun. Similarity (%) data analysis described that weeds contributing average dissimilarity in invasive and non-invasive quadrates (Table 2). A shade plot showing the multivariate analysis and magnitudes of the invasive and non-invasive weeds distributions. However, the each color represents the species present in an ecosystem. Hierarchical data showed gave an indication of total number of species studied in four different dimensional community (Figure 6).

Discussion

The data of study showed that invasive weed (*Xanthium strumarium*) investigated at these locations having maximum competition with native weeds. Significant competition of native weeds recorded in non-invasive quadrate recorded high significant abundance at Huailai County. However, on all other three sites native weeds compete with invasive weed that was comparable to non invasive quadratic ring. Simpson's Diversity Index (SDI); Margalef's Richness Index (MRI) gave the clear-cut indication that *Xanthium strumarium* present in invasive quadrate compete positively within non-invasive weeds. Maximum SDI investigated higher diversity index or probability between invasive and non-invasive quadratic rings. These results were in line with the scientists who reported that *Xanthium strumarium* affected severally to the natural vegetations. Seventy species investigated in the non-invasive plots, however, thirty-one recorded in *Xanthium strumarium* toxic area. About 55.71% species reduced in invasive compared to non-invasive range. SDI, MRI reduced upto 70.33%, and 69.39% in the invasive plot [28,31]. The results of our study was also in line with the researchers who studied that the effect of invasive species on native population was usually stronger than vice versa [44]. Comparatively significant species diversity were present in Invasive quadrate in Huailai (4B) followed by Zhenjiakou (4C); Fushun (4A) and Yangyuan (4D) than non invasive quadrat that prove the hypothesis that Invasive species compete with the native species abundance because they have high phenotypic plasticity and have maximum ability to adopt and enhance vegetative growth in an ecosystem.

These results are in accordance to the researchers who reported that *X. strumarium* is an invasive possessing negative impacts on natural ecosystem [45]. This weeds can alter abundances of native weeds in a terrestrial food web, but the pattern is complex [28,46,47]. *Xanthium strumarium* having a big canopy and tap root system caused considerable impact on native plant communities; however, its population abundance causes serious threat to the native plant communities. Maximum evenness recorded in invasive quadrat investigated good competitor in ecosystem processes. The plant invasions severely influence on the size and composition of native vegetations through interference of biotic interactions in abiotic network [48,49]. *X. strumarium* caused maximum damage to the native weeds might continue in near future [5,50]. Invasive species are characterized in wide-ranging, rapacious, assertively persistent, elastic, grow rapidly, capability to travel in extensive areas and rich reproduction as these flora emerge to have particular qualities that permit them to compete with native plant species [51,52]. The coefficient of determination (R^2) recorded in the surveyed community of different weeds families suggested that there is a strong positive polynomial relationship investigated at different locations. These finding are in accordance to the other studies on invasive weeds of Asteraceae family in which the researcher indicated the strong effect of invaders on the ecosystem properties [31,53]. The researchers deliberated that *Xanthium strumarium* became most brutal Invasive Alien plant dominated on rangelands of Ethiopia [54]. The researchers previously conducted paired wise survey in invasive and non invasive quadrat suggested that invasive plant species compete and affect positively on native plant species [55]. Our hypothesis is true that Invasive weeds have a potential to compete with of non-invasive/native weeds species in a community. These results were in line to the researchers who reported heterogeneous population of weed community in their experiments [38]. Maximum dissimilarity index (87.06%) was recorded in Huailai county followed by Yangyuan (44.43%), Zhangjiakou (40.13%) and at Fushun (29.02%). *Xanthium strumarium* affected on the species density in our experimental study which were in line with the researchers who described the Jaccard's similarity index between invasive and non-invasive ranges in weed communities which gave indication about 38.4% loss of weed species due to *Xanthium strumarium* infested plot resulted 61.6% dissimilarity index [28]. In our experiment significantly ($P < 0.01$) maximum global test for sample statistics R values were recorded in Huailai County (0.943) followed by Zhangjiakou, Yangyuan and Fushun area investigated by analysis of similarities using one way analysis of variance taking 999 number of permutations. The larger value of R indicated that higher effect of species diversity upon studied variables. Huailai County significantly more affected due to *Xanthium strumarium* invasions compared to Zhangjiakou and Yangyuan areas. The lowest invasion was investigated in Fushun because of highest dissimilarity or beta diversity in invasive and non invasive/control plots. The result of our experiment suggested that there is a strong negative relationship ($R^2 = 0.741$, or 74.1%) found between the *Xanthium strumarium* (%) and Species Richness. These results are in line with the scientists who reported that regression equation and Pearson correlation (-0.861) indicated the existence of strong negative linear relationship between *Xanthium strumarium* and Richness however with the increase in *Xanthium* species (%) the richness decreases significantly [28]. In our study, maximum density of natural weed was gave indication that the prevailing edaphic and environmental conditions of the area are highly conducive for the distribution and growth of the weed species. These work is in line with the researchers who reported same results in their experiments [53,56].

Conclusion

It was reasoned that *Xanthium strumarium* rival local weeds species in a network. Noteworthy challenge in thickness recorded in non invasive quadrat that demonstrate the speculation that invasive weed influenced the local weeds decent variety thought about a huge risk to the characteristic biological system. Greatest SDI at Huailai District gave a sign that weed networks were increasingly heterogeneous in non-invasive locales contrasted with invasive ones. The bigger estimation of global R in measurements demonstrated that higher impact of species decent variety upon examined factors. The most influenced site found by *Xanthium strumarium* attack was Huailai Area followed by Zhangjiakou and Yangyuan separately. The most minimal attack impacts were examined in Fushun in view of most elevated difference or beta assorted variety. Most extreme thickness of weed species gave cautioning that the predominant edaphic and ecological states of the territory are exceptionally favorable for the conveyance and development of the weed species. At long last it is an admonition circumstance to the scientists that there is a critical need of appropriate intending to oversee and dispose of the spread of this weed attack. We also encouraged researchers to explore the defensive mechanism, nutrient cycling, role of pseudomonas bacteria, resistance mechanism of invasive and native weeds in future.

Author Contributions: The following statements stated that “conceptualization, Y.L.F. and M.F.I.; methodology, Y.L.F.; software, M.F.I.; validation, Y.L.F., M.F.I.; formal analysis, M.F.I.; investigation, M.F.I., L. M. C.; resources, Y.L.F.; data curation, M.F.I., A. I.; writing—original draft preparation, M.F.I.; writing—review and editing, M.F.I.; Y.L.F.; visualization, M.F.I.; supervision, Y.L.F.; project administration, Y.L.F.; funding acquisition, Y.L.F.”,

Funding: “The survey conducted with the support of National key R&D Program of China (2017YFC1200101), the National Natural Science Foundation of China (31470575, 31670545, and 31971557)”.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Parker IM, Simberloff D, Lonsdale WM, Goodell K, Wonham M, et al. (1999) Impact: toward a framework for understanding the ecological effects of invaders. *Biological invasions* 1: 3-19.
2. Ehrenfeld JG (2010) Ecosystem consequences of biological invasions. *Annual review of ecology, evolution, and systematics* 41: 59-80.
3. Simberloff D (2011) How common are invasion-induced ecosystem impacts? *Biological invasions* 13: 1255-1268.
4. Khan MA, Qureshi RA, Gillani SA, Ghufuran MA, Batool A, et al. (2010) Invasive species of federal capital area Islamabad, Pakistan. *Pak J Bot* 42: 1529-1534.
5. Nigussie ST, Amare SA, Manaye MM, Edget MB, Ashenafi AH, et al. (2017) Invasion and impacts of *Xanthium strumarium* in Borena Zone of Oromia Region, Ethiopia. *Journal of Coastal Life Medicine* 5: 350-355.
6. Qureshi H, Arshad M, Bibi Y (2014) Invasive flora of Pakistan: a critical analysis. *International Journal of Biosciences* 4: 407-424.
7. Holm LG, Plucknett DL, Pancho JV, Herberger JP (1977) *The world's worst weeds. Distribution and biology*: University press of Hawaii.
8. Holm LG (1991) *Avena fatua* L. and other members of the "wild oats" group. *The World's worst weeds*.
9. Hooper DU, Adair EC, Cardinale BJ, Byrnes JEK, Hungate BA, et al. (2012) A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* 486: 105.
10. Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, et al. (2012) Biodiversity loss and its impact on humanity. *Nature* 486: 59.

11. Naeem S, Duffy JE, Zavaleta E (2012) The functions of biological diversity in an age of extinction. *Science* 336: 1401-1406.
12. Lefcheck JS, Byrnes JEK, Isbell F, Gamfeldt L, Griffin JN, et al. (2015) Biodiversity enhances ecosystem multifunctionality across trophic levels and habitats. *Nature communications* 6: 6936.
13. Soliveres S, Van Der Plas F, Manning P, Prati D, Gossner MM, et al. (2016) Biodiversity at multiple trophic levels is needed for ecosystem multifunctionality. *Nature* 536: 456.
14. Dekker J (2018) The EEW 2nd Edition Narrative Book.
15. Dekker J (2011) Evolutionary ecology of weeds. Agronomy Department, Iowa State University, Ames, IA.
16. Stešević D, Petrović D (2010) Preliminary list of plant invaders in Montenegro. *Biologica Nyssana* 1.
17. Nel JL, Richardson DM, Rouget M, Mgidi TN, Mdzeke N, et al. (2004) A proposed classification of invasive alien plant species in South Africa : towards prioritizing species and areas for management action : working for water. *South African Journal of Science* 100: 53-64.
18. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, et al. (2017) Vegan: Community Ecology Package. R package version 2.4–4.
19. Arifa Z, Ahmad SS, Sardar AA, Zaheer-ud-Din K (2014) Evaluation of ecological aspects of natural vegetation of Pakpattan District using multivariate techniques. *Journal of Biodiversity and Environmental Sciences (JBES)* 5: 230-238.
20. Harrison SK, Regnier EE, Schmoll JT (2003) Postdispersal predation of giant ragweed (*Ambrosia trifida*) seed in no-tillage corn. *Weed Science* 51: 955-964.
21. Liu Q, Yu M, Zhou Y (2002) A preliminary study on the invasive plants in Beijing. *Journal of Beijing Normal University (Natural Science)* 38: 399-404.
22. Li N, Zhu LN, Zhai Q, Zhu MW, Chen XH, et al. (2010) A new alien invasive plant—*Xanthium italicum* Moretti in Liaoning Province. *Plant Quar* 24: 49-52.
23. Wang R, Wan F (2010) Prediction of the potential survival area of *Xanthium italicum* in China. *Acta Prataculturae Sinica* 19: 222-230.
24. Amini R, Izadkhah S, Mohammadinasab AD, Raei Y (2014) Common cocklebur (*Xanthium strumarium* L.) seed burial depth affecting corn (*Zea mays* L.) growth parameters. *International Journal of Biosciences* 4: 164-170.
25. Hussain Z, Marwat KB, Cardina J, Khan IA (2014) *Xanthium strumarium* L. impact on corn yield and yield components. *Turkish journal of agriculture and forestry* 38: 39-46.
26. Kamboj A, Saluja AK (2010) Phytopharmacological review of *Xanthium strumarium* L.(Cocklebur). *International Journal of Green Pharmacy (IJGP)* 4.
27. Weaver SE, Lechowicz MJ (1983) The biology of Canadian weeds.: 56. *Xanthium strumarium* L. *Canadian Journal of Plant Science* 63: 211-225.
28. Seifu A, Seboka N, Misganaw M, Bekele T, Merawi E (2017) Impact of Invasive Alien Plant, *Xanthium Strumarium*, On Species Diversity and Composition of Invaded Plant Communities in Borena Zone, Ethiopia. *Biodiversity Int J* 1 (1): 00004. DOI: 10.15406/bij. 2017.01. 00004 *Xanthium strumarium* is one of the invasive alien species those posing negative impacts on country's biodiversity. *Xanthium strumarium*.
29. Bhargava A, Shukla S, Dixit BS, Bannerji R, Ohri D (2006) Variability and genotypex cutting interactions for different nutritional components in *Chenopodium album* L. *Hortic Sci* 33: 29-38.
30. Iqbal MF, Feng YL, Liu MC, Lu XR, Nasir M, et al. (2019) Parasitic activity of powdery mildew (Pathogen strain HMLAC 226) on prostrate knotweed (*Polygonum aviculare* L.) at various locations of Shenyang, Northeast China. *Applied Ecology and Environmental Research* 17: 13383-13394.

31. Qureshi H, Arshad M, Bibi Y, Osunkoya OO, Adkins SW (2018) Multivariate impact analysis of *Parthenium hysterophorus* invasion on above-ground plant diversity of Pothwar region of Pakistan. *Applied Ecology and Environmental Research* 16: 5799-5813.
32. Chikuruwo C, Masocha M, Murwira A, Ndaimani H (2017) Predicting the suitable habitat of the invasive *Xanthium strumarium* in southeastern Zimbabwe. *Applied Ecology and Environmental Research* 15: 17-32.
33. Clarke KR, Gorley RN (2015) Getting started with PRIMER v7. PRIMER-E: Plymouth, Plymouth Marine Laboratory.
34. Clarke KR, Gorley RN, Somerfield PJ, Warwick RM (2014) Change in marine communities: an approach to statistical analysis and interpretation: Primer-E Ltd.
35. Osunkoya OO, Akinsanmi OA, Lim LSA, Perrett C, Callander J, et al. (2017) *Parthenium hysterophorus* L.(Asteraceae) invasion had limited impact on major soil nutrients and enzyme activity: Is the null effect real or reflects data insensitivity? *Plant and soil* 420: 177-194.
36. Clarke KR, Warwick RM (2001) Change in marine communities. An approach to statistical analysis and interpretation.
37. Magurran AE (1988) Why diversity? Ecological diversity and its measurement: Springer. pp. 1-5.
38. Mahajan M, Fatima S (2017) Frequency, abundance, and density of plant species by list count quadrat method. *International Journal of Multidisciplinary Research* 3: 1-8.
39. Chao A, Chazdon RL, Colwell RK, Shen TJ (2005) A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecology letters* 8: 148-159.
40. Letourneau DK, Jedlicka JA, Bothwell SG, Moreno CR (2009) Effects of Natural Enemy Biodiversity on the Suppression of Arthropod Herbivores in Terrestrial Ecosystems. *Annual Review of Ecology, Evolution, and Systematics* 40: 573-592.
41. Kilewa R, Rashid A (2014) Distribution of invasive weed *Parthenium hysterophorus* in natural and agro-ecosystems in Arusha Tanzania. *International Journal of Science and Research* 3: 1-4.
42. El-Azazi E-S, Khalifa EA, Belal MMSAH, Eltanger NA (2013) Ecological studies of some *Acacia* species grown in Egyptian Deserts. *Earth* 7: 3-8542.
43. Seid A, Goftishu M, Degebassa L, Mekete T (2016) Occurrence, distribution, and abundance of plant-parasitic nematodes associated with khat (*Catha edulis* Forsk) in East Hararghe Zone, Ethiopia. *Nematropica* 45: 208-214.
44. Vila M, Weiner J (2004) Are invasive plant species better competitors than native plant species?—evidence from pair-wise experiments. *Oikos* 105: 229-238.
45. Lemma B, Tessema T, Fessehaie R (2015) Distribution, abundance and socio-economic impacts of invasive plant species (IPS) in Borana and Guji Zones of Oromia National Regional State, Ethiopia. *Ethiopia Journal of Agricultural Science and Review* 4: 271-279.
46. McCary MA, Mores R, Farfan MA, Wise DH (2016) Invasive plants have different effects on trophic structure of green and brown food webs in terrestrial ecosystems: a meta-analysis. *Ecology Letters* 19: 328-335.
47. Rejmánek M, Richardson DM, Pyšek P (2005) Plant invasions and invasibility of plant communities. *Vegetation ecology* 332355.
48. Mack RN, Simberloff D, Mark Lonsdale W, Evans H, Clout M, et al. (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological applications* 10: 689-710.
49. Levine JM, Hille Ris Lambers J (2009) The importance of niches for the maintenance of species diversity. *Nature* 461: 254.
50. Moustakas A, Katsanevakis S (2018) Data Mining and Methods for Early Detection, Horizon Scanning, Modelling, and Risk Assessment of Invasive Species. *Frontiers in Applied Mathematics and Statistics* 4: 5.

51. Valéry L, Fritz H, Lefeuvre J-C, Simberloff D (2008) In search of a real definition of the biological invasion phenomenon itself. *Biological invasions* 10: 1345-1351.
52. Sujay YH, Sattagi HN, Patil RK (2010) Invasive alien insects and their impact on agroecosystem. *Karnataka Journal of Agricultural Sciences* 23: 26-34.
53. Riaz T, Javaid A (2011) Prevalence of alien weed *Parthenium hysterophorus* L. in grazing and wastelands of District attock, Pakistan. *Journal of Animal and Plant Science* 21: 542-545.
54. Berhanu L, Taye T, Rezene F (2015) Distribution, abundance and socio-economic impacts of invasive plant species in Borana and Guji Zones of Oromia National Regional State, Ethiopia. *Basic Res J Agric Sci Rev* 4: 271-279.
55. Powell KI, Chase JM, Knight TM (2013) Invasive plants have scale-dependent effects on diversity by altering species-area relationships. *science* 339: 316-318.
56. Mehraj G, Khuroo AA, Qureshi S, Muzafar I, Friedman CR, et al. (2018) Patterns of alien plant diversity in the urban landscapes of global biodiversity hotspots: a case study from the Himalayas. *Biodiversity and conservation* 27: 1055-1072.
57. Kumar S, Graham J, West AM, Evangelista PH (2014) Using district-level occurrences in MaxEnt for predicting the invasion potential of an exotic insect pest in India. *Computers and Electronics in Agriculture* 103: 55-62.

Figure 1: Latitudinal map showing different study sites in Northeast China in 2018

Figure 2. Monthly climatic data of different locations at Liaoning and Hebei Province for the year 2018

Figure 3. Rarefaction curve showing cumulative number species studied at four locations showing highly positive polynomial *Coefficient of determination* (R^2)

Figure 4 showing ecological indices regarding mean value/10m² of Invasive with Non Invasive quadratic rings at different sites where A (Fushun); B (Huailai); C (Zhangjiakou); D (Yungyuan). Whereas (A = Abundance; SDI = Simpsons Diversity Index; MRI = Margalef's Richness Index; NOS = Number of Species).

Figure 5 showing weed density/m² of invasive with non invasive quadrate at different locations (A) Zhangjiakou; (B) Huailai County; (C) Yangyuan county; (D) Fushun

406 **Figure 1:** Latitudinal map showing different study sites in Northeast China in 2018 [32,57]

407
408

Figure 2. Monthly climatic data of different locations at Liaoning and Hebei Province for the year 2018

413

414

415

416

417

Figure 3. Rarefaction curve showing cumulative number species studied at four locations with *Coefficient of determination* (R^2)

Figure 4 showing ecological indices regarding mean value 10 m⁻² of Invasive with Non Invasive quadratic rings at different sites.

Figure 5 showing weed density 10 m⁻² of invasive with non invasive/control quadrate at different locations (A) Zhangjiakou; (B) Huailai County; (C) Yangyuan county; (D) Fushun County

Fig. 6. A shade plot showing the multivariate relative magnitudes of the invasive and non-invasive weeds distributions where the color shading represents the population diversity in a community. Hierarchical data showed that total number of species studied during extensive field survey during 2018

440 **Table 1.** Plant species found during extensive field survey with family, habitat, and life form

Sr. No.	Species	Local Name	Families	Abb.	Habitat Range	Life form
1	<i>Xanthium strumarium</i>	Common cocklebur	Asteraceae	Xs	Invasive	Herb
2	<i>Humulus scandens</i>	Wild hop	Cannabaceae	Hs	Native	Herb
3	<i>Setaria feberi</i>	Chinese foxtail	Poaceae	Sf	Native	Grass
4	<i>Chenopodium album</i>	Bathu	Amaranthaceae	Ca	Native	Herb
5	<i>Artemisia annua</i>	Annual wormwood	Asteraceae	Ara	Native	Herb
6	<i>Dioscorea zingiberensis</i>	Yellow ginger	Dioscoreaceae	Dz	Native	Herb
7	<i>Setaria pumila</i>	Yellow foxtail	Poaceae	Sp	Native	Grass
8	<i>Erodium stephianum</i>	Pin clover	Geraniaceae	Es	Native	Herb
9	<i>Seteria viridis</i>	Green foxtail	Poaceae	Sv	Native	Grass
10	<i>Malva verticillata</i>	Chinese mallow	Malvaceae	Mv	Native	Herb
11	<i>Avena sativa</i>	Common oat	Poaceae	As	Native	Grass
12	<i>Cynanchum chinensis</i>	Dog strangling vine	Apocynaceae	Cc	Native	Herb
13	<i>Artemisia selengensis</i>	Wormwood	Asteraceae	Ars	Native	Herb
14	<i>Oxalis corniculata</i>	creeping woodsorrel	Oxalidaceae	Oc	Native	Shrub
15	<i>Cynanchum wilfordii</i>	Baekshuoh	Apocynaceae	Cw	Native	Herb
16	<i>Acalypha australis</i>	Asian copperleaf	Euphorbiaceae	Aca	Native	Shrub
17	<i>Taraxacum mongolicum</i>	Dandelion	Asteraceae	Tm	Native	Herb
18	<i>Myosoton aquaticum</i>	Giant chickweed	Caryophyllaceae	Ma	Native	Herb
19	<i>Polygonum aviculare</i>	Common knotweed	Polygonaceae	Pa	Native	Herb
20	<i>Ambrosia trifida</i>	Giant ragweed	Asteraceae	Amt	Invasive	Herb
21	<i>Ambrosia artemisifolia</i>	Common ragweed	Asteraceae	Ama	Invasive	Herb
22	<i>Axonopus compressus</i>	Cow Grass	Poaceae	Axc	Native	Grass
23	<i>Medicago lupulina</i>	hop clover	Fabaceae	Mel	Native	Herb
24	<i>Erigeron acris</i>	bitter fleabane	Asteraceae	Ea	Native	Herb
25	<i>Euphorbia maculata</i>	prostrate spurge	Euphorbiaceae	Eumc	Native	Shrub
26	<i>Leptochloa chinensis</i>	Chinese sprangletop	Poaceae	Lc	Native	Grass
27	<i>Plantago depressa</i>	broad-leaved plantain	Plantaginaceae	Pd	Native	Herb
28	<i>Rubia cordifolia</i>	common madder	Rubiaceae	Rc	Native	Herb
29	<i>Cyperus cyperoides</i>	Madhana	Cyperaceae	Cc	Native	Sedge
30	<i>Kochia scoparia</i>	Kochia	Chenopodiaceae	Ks	Native	Herb
31	<i>Chenopodium hybridum</i>	Maple-leaved goosefoot	Amaranthaceae	Ch	Native	Herb
32	<i>Chenopodium ficifolium</i>	Fig-leaved goosefoot	Amaranthaceae	Cf	Native	Herb
33	<i>Eleusine indica</i>	yard-grass	Poaceae	Ei	Native	Grass
34	<i>Polygonum longisetum</i>	knotweed	Polygonaceae	Pl	Native	Herb
35	<i>Echinochola crus-galli</i>	cockspur grass	Poaceae	Ecg	Native	Grass
36	<i>Zizania latifolia</i>	Zizania	Poaceae	Zl	Native	Grass
37	<i>Imperata cylindrica</i>	kunai grass	Poaceae	Ic	Native	Grass
38	<i>Amaranthus retroflexus</i>	rough pigweed	Amaranthaceae	Ar	Native	Herb
39	<i>Amaranthus blitum</i>	Guernsey pigweed	Amaranthaceae	Ab	Native	Herb

441

442

Table 2. Pooled data of SIMPER analysis of *Xanthium strumarium* invasive and non-invasive/control quadrat in four locations at Northeast China

Average dissimilarity = 44.43% based on Average abundance (Yangyuan County)					
Species	Invasive	Non-Invasive	Av. Diss.	Diss./SD	Contribution (%)
<i>Malva verticillata</i>	4.00	10.40	10.64	2.54	23.94
<i>Seteria viridis</i>	1.40	6.80	9.24	1.95	20.80
<i>Chenopodium album</i>	4.10	9.00	8.27	2.27	18.62
<i>Avena sativa</i>	2.90	7.10	7.38	1.48	16.62
Average dissimilarity = 87.06% based on Average abundance (Huailai County)					
<i>Seteria viridis</i>	2.10	7.60	12.19	1.29	14.00
<i>Chenopodium album</i>	0.00	6.10	12.05	2.14	13.84
<i>Artemisia annua</i>	0.50	6.10	11.53	1.69	13.24
<i>Setaria pumila</i>	0.90	6.50	11.46	1.78	13.16
Average dissimilarity = 40.13% based on Average abundance (Zhangjiakou)					
<i>Seteria febrei</i>	10.00	16.80	9.79	1.30	24.40
<i>Discorea zingiberensis</i>	0.80	6.50	7.28	1.44	18.13
<i>Artemisia annua</i>	6.30	12.50	7.27	1.81	18.12
Average dissimilarity = 29.02% based on Average abundance (Fushun County)					
<i>Seteria viridis</i>	7.90	17.50	11.84	2.41	40.82
<i>Seteria febrei</i>	12.30	18.10	7.86	1.47	27.10
<i>Chenopodium album</i>	3.60	5.80	4.31	1.58	14.86