A RAPIDLY EVOLVED SHIFT IN LIFE HISTORY TIMING DURING ECOLOGICAL SPECIATION IS DRIVEN BY THE TRANSITION BETWEEN DEVELOPMENTAL PHASES

THOMAS H. Q. POWELL^{1,2*}, ANDREW NGUYEN^{1,3}, QINWEN XIA,¹ JEFFREY L. FEDER⁴,
GREGORY J. RAGLAND⁵, DANIEL A. HAHN¹

¹Entomology and Nematology Department, Steinmetz Hall, University of Florida, Gainesville, FL 32611

²Present Address: Department of Biological Sciences, Science III, Binghamton University (State University of New York), Binghamton, NY 13902

Present Address: Independent Researcher, Jacksonville, FL

⁴Department of Biological Sciences, 100 Galvin Life Sciences, University of Notre Dame, Notre Dame, IN 45665

⁴Department of Integrative Biology; 1151 Arapahoe, University of Colorado Denver, Denver, CO 80217

*To whom all correspondence should be sent – email: powellt@binghamton.edu

Key words: Allochronic isolation, phenology, respirometry, dormancy, Tephritidae

Data accessibility: All data (in the form of tables of all metabolic rate measurements for all individual flies in the study) will be available on DRYAD when the manuscript is published.

Abstract

For insect species in temperate environments, seasonal timing is often governed by the regulation of diapause, a complex developmental program that allows insects to weather unfavorable conditions and synchronize their lifecycles with available resources. Diapause consists of a series of phases that govern initiation, maintenance, and termination of this developmental pathway. The evolution of insect seasonal timing depends in part on how these phases of diapause development and post-diapause development interact to affect variation in phenology. Here, we dissect the physiological basis of a recently evolved phenological shift in *Rhagoletis pomonella* (Diptera: Tephritidae), a model system for ecological divergence. A recently derived population of *R. pomonella* shifted from specializing on native hawthorn fruit to earlier fruiting introduced apples, resulting in a 3-4 week shift in adult emergence timing. We tracked metabolic rates of individual flies across post-winter development to test which phases of development may act either independently or in combination to contribute to this recently evolved divergence in timing. Apple and hawthorn flies differed in a number of facets of their post-winter developmental trajectories. However, divergent adaptation in adult emergence phenology in these flies was due almost entirely to the end of the pupal diapause maintenance phase, with post-diapause development having a very small effect. The relatively simple underpinnings of variation in adult emergence phenology suggest that further adaptation to seasonal change in these flies for this trait might be largely due to the timing of diapause termination unhindered by strong covariance among different components of post-diapause development.

Data accessibility: All data (in the form of tables of all metabolic rate measurements for all individual flies in the study) will be available on DRYAD when the manuscript is published.

INTRODUCTION

Phenological adaptation can play an important role in both the origin and maintenance of biodiversity in seasonal environments. Adaptation to temporally predictable environmental variation may be a potent driver of reproductive isolation during ecological speciation (Taylor and Friesen 2017). Moreover, altered seasonality is a critical consequence of global climate change for many ecological communities (Miller-Rushing et al. 2010). Thus, understanding how evolution shapes the regulation of seasonal timing has important implications for the tempo of ecological diversification, as well as the potential of biotic communities to persist in the face of global change.

Divergent adaptation to novel habitats or niches can generate reproductive isolation between populations through the process of ecological speciation (Schluter 2001; 2009; Rundle & Nosil 2005). Temporal isolation, or allochrony, may be a common component of ecological barriers to gene flow for two reasons. First, in temperate habitats, temporally distinct resource opportunities may pose distinct trade-offs, favoring phenological specialization. Second, for organisms with seasonal reproductive biology, phenological divergence may serve as a ‘magic trait’ for speciation by pleiotropically coupling ecological divergence to assortative mating (Servedio et al. 2011). Indeed, adaptation to differences in seasonal timing has been shown to contribute to reproductive isolation in a wide variety of organisms including plants (Salvolainen et al. 2006; Lowry et al. 2008), insects (Tauber et al. 1977; Wood & Guttman 1982; Horner et al. 1999; Eubanks et al. 2003; Abbot & Withgott 2004; Ordning et al. 2010; Wadsworth et al. 2013; Hood et al. 2015), and vertebrates (Friesen et al. 2007).

Shifts in phenology have also received increasing attention as anthropogenic change

causes alterations in the phenology of plant and animal populations (Parmesan 2006; Van Dyck et al. 2015; Thackeray et al. 2016). Many recent studies provide evidence for fitness consequences of disrupting seasonal phenologies, from high bird nestling mortality due to mismatch with seasonal food supplies (Sanz et al. 2003) to declines in recruitment in plant populations resulting from lack of synchrony between flowering and the activity of pollinators (Rafferty & Ives 2011; Gilman et al. 2012; Kudo & Ida 2013). Cases of recent shifts in phenology during incipient speciation also provide useful windows into how the regulation of seasonal timing evolves more generally, offering insight into the factors that may promote or constrain evolutionary responses to anthropogenic change (Scriber et al. 2014, Taylor and Friesen 2017).

Phenological adaptation is crucial for species with specialist life history strategies. For example, univoltine phytophagous insects typically have one chance to breed per year, with most of these species specializing on a single host plant (Strong et al. 1984). Therefore, suboptimal timing can have severe fitness costs (Miller-Rushing et al. 2010). Many univoltine insects use seasonal dormancy, or diapause, to synchronize their active times for growth and reproduction with seasonal resources. Thus, the decision of when to enter dormancy relative to the timing of unfavorable seasonal conditions, such as the onset of winter, the dry season, or lack of host-plant availability, as well as when to exit dormancy to synchronize with favorable conditions are critical aspects of life history timing (Tauber and Tauber 1981). Phenological mismatches between insect life histories and resource availability therefore lead to low reproductive success and evolutionary dead ends (Van Dyck et al. 2015).

Insect diapause is not just a simple cessation of development resulting directly from physiologically stressful conditions (e.g., cold-induced quiescence) (Denlinger 2002; Hahn and

Denlinger 2011). Rather, diapause is a dynamic, environmentally sensitive developmental program, consisting of multiple developmental phases: preparation, maintenance, and the end of diapause followed by the initiation of post-diapause development (Kostal et al. 2006). Each of these developmental phases is characterized by distinct molecular (Kostal et al. 2017; Ragland et al. 2017), morphological (Bryon et al. 2017), and physiological (Ragland et al. 2009) hallmarks. Selection to produce alternative or divergent phenological strategies may act on the relative duration of or developmental rates during these phases (Andrewartha 1952, Kostal 2006, Kostal et al. 2017). Whether or not variation in each phase of the diapause and post-diapause program represents an independent functional module underlain by distinct genetic architecture or are highly interrelated and covary has important consequences for the evolvability of seasonal timing (Wagner et al. 2007; Wagner & Zhang 2011). If independent, then it is possible that each module can be differentially shaped by natural selection to varying environmental conditions at specific points during development. However, if the timing of different phases of diapause development are highly interrelated, then it may be that selection acts on the phase having greatest significance for matching an insect's life cycle with resource availability, with possible pleiotropic consequences on other phases. While the concept of distinct phases within the diapause developmental trajectory has long been recognized (Andrewartha 1952; Tauber et al. 1986; Kostal 2006), how different phases of diapause development interact and how selection acts on variation within these phases to affect life history timing is not well understood and has received very little attention despite its potential importance for seasonal adaptation.

To address the question of the flexibility of the diapause program to compartmentally adapt to environmental challenges requires first characterizing the different phases of diapause development in an insect and then determining their interactions (correlational structure) to

assess whether they are assembled into independent modules or not. Under the simplest scenario, seasonal divergence in emergence timing is due principally to the degree of metabolic suppression occurring during the diapause maintenance phase (Wipking et al. 1995; Feder & Filchak 1999), with no subsequent significant independent effect of any other developmental phase. Thus, there will be a general 1:1 correspondence between individuals having greater metabolic suppression and later adult eclosion. A competing hypothesis is that the key factor dictating phenology is when diapause maintenance is terminated, regardless of metabolic depth, (Wadsworth et al. 2013) Alternatively, variation in adult emergence phenology may be driven solely by differences in the rate of post-diapause morphogenesis (Poseldovich et al. 2012; 2014). In each of these models, other components of diapause preparation, maintenance, and post-diapause development would be weakly correlated with eclosion time. Conversely, correlated shifts of different components within and between diapause phases can also lead to divergent phenology, especially if driven by directional selection. For example, earlier adult emergence timing may be driven by combinatorial effects of higher metabolic rate during the diapause maintenance phase, earlier termination of the diapause maintenance phase, and more rapid post-diapause development.

Here, we examine the role that these different components of diapause development play in the case of rapidly evolved phenological divergence in the apple maggot fly, *Rhagoletis pomonella* (Walsh), a text-book system for ecological speciation (Dres & Mallet 2002; Berlocher and Feder 2002). Divergent seasonal adaptation of the flies to synchronize adult emergence with their specific host fruits is well understood (Feder and Filchak 1999), making *R. pomonella* an excellent system for studying how different components of diapause development interact in the regulation of seasonal timing (Feder & Dombraski 2007; Ragland et al. 2017). Partially

reproductively isolated populations that show consistent patterns of host-associated genetic differentiation, hereafter host races, are undergoing divergent selection to synchronize themselves with the fruiting times of their respective host plants. The derived host race shifted from infesting fruits of a native plant, the downy hawthorn (*Crataegus mollis*) to infest domesticated apples (*Malus pumila*) ~160 years ago (Walsh 1861; Bush 1966). The apple varieties favored by *R. pomonella* ripen ~1 month earlier in the season compared to hawthorns in the Midwestern US (Feder et al 1993, 1994). Apple and hawthorn flies spend most of the year in pupal diapause. Thus, the difference in adult and larval phenology between the races is determined by how flies transition from this shared state of pupal diapause during winter to completing adult development prior to emergence at different points the following summer and fall.

We investigate how variation in components of diapause maintenance and post-diapause development are associated with shifts in seasonal life history timing in the derived compared to the ancestral host race of *R. pomonella*. Specifically, we test whether the earlier phenology of the derived apple race is driven by alteration to individual components or multiple components of their diapause developmental trajectories. We use respirometric phenotyping to compare the post-winter metabolic rate trajectories of apple and hawthorn flies from sympatric populations under controlled laboratory common-garden conditions to identify the key phases of diapause development. Low metabolic rate is a reliable marker for the diapause maintenance phase that identifies individuals in diapause; metabolic rate then increases in a stereotypical trajectory during post-diapause development following the end of diapause (Ragland et al. 2009). We follow cohorts of flies from both host races in the laboratory from the cessation of simulated winter through to adult emergence, producing a time-series of metabolic rates for each

individual. We fit these time series to mathematical models with parameters describing different aspects of diapause development corresponding with post-winter metabolic trajectories that include parameters modeling: 1) baseline metabolic rate during diapause maintenance, 2) timing of the transition from maintenance to post-diapause development, 3) metabolic rate at a plateau coinciding with early adult morphogenesis, 4) duration of the plateau, and 5) an exponential increase in metabolic rate corresponding with late adult morphogenesis culminating in adult emergence (Figure 1c). Comparisons across these individual models of metabolic trajectories allow us to test how patterns of post-winter diapause development differ between the two host races, and importantly test which component or combination of components is significantly associated with variation in adult emergence phenology. Thus, our approach provides a test of the hypothesis that variation in the duration of one or multiple components of the diapause-maintenance phase or post-diapause development account for earlier emergence timing in the derived apple host race. We also conduct a second experiment to specifically test whether overwintering and immediate post-winter diapause metabolic rates differ between the host races and how variation in these traits influences adult emergence timing.

Materials and Methods

Insect collection and rearing

In the summer and fall of 2013 and 2014, apple and hawthorn race *R. pomonella* flies were collected from infested fruit from wild populations in Urbana, IL during peak infestation for both host plants. The apple and hawthorn sites are located ~1,300 m apart, well within the flight range of these flies (Roitberg et al. 1984). Infested fruit were collected from the ground and trees and transported to the laboratory where they were kept in an environmentally

controlled room under standard *Rhagoletis* rearing conditions (24°C and 14:10 Light:Dark (L:D): Filchak et al. 2000; Dambroski & Feder 2007). Fruit were placed in wire mesh baskets in plastic trays where wandering third instar larvae emerged naturally from the fruit and dropped into the trays. Trays were monitored daily for larvae, which were placed in petri dishes of moist vermiculite to pupariate. Petri dishes were kept at 24°C and 14:10 L:D for ten days. At the end of the ten days, pupae were transferred to clean petri dishes and placed in chambers with 85% humidity maintained by a saturated KCl solution (Winston and Bates 1960). Humidity chambers were then transferred to simulated winter conditions at $3.5^{\circ} \pm 1.0$ C, where they remained for 24 weeks. After removal from the cold, pupae were kept at 24° C, 14:10 L:D, and 85% humidity for the duration of the experiment.

Measuring post-winter metabolic trajectories

Metabolic rates were measured using stop-flow respirometry (Lighton 2008), closely following the methods of Ragland et al. (2009). Upon removal from the cold, 96 pupae from each host race were weighed and placed into individual 5ml Norm-Ject™ syringes (Air-Tite Products, Virginia Beach, VA, USA) fitted with three-way luer valves (Cole-Parmer, Vernon Hills, IL, USA). *Rhagoletis* pupae are small enough (~5-11 mg) to reside within the top arm of the luer valve so that the syringe piston can be fully depressed. Syringes were flushed with CO₂ free air, using an aquarium pump pushing air through a Dririte™ (W.A. Hammond Co. Xenia, OH, USA) - Ascarite II™ (Thomas Scientific, Philadelphia, PA, USA) - Dririte™ scrubber column followed by an acidified (pH < 4.0) water bubbler to rehumidify the air. Syringes were sealed with 1ml scrubbed air, and the time of the syringe purge was recorded using Expedata logging software (Sable Systems International, Las Vegas, NV, USA). Four empty control syringes were also purged and sealed. Syringes were then returned to 25° C, 14:10 L:D

incubators. After approximately 24 hours (exact time recorded for each individual), the full volume of the 1mL chamber was injected into a flow-through system with a Li-Cor 7000 infrared CO₂ analyzer (Lincoln, NE, USA) interfaced with Expedata. The flow rate of the respirometry system was maintained at 150ml/min using an MFC-2 mass flow control unit (Sable Systems International) with a Sierra Instruments (Monterey, CA, USA) mass-flow valve, and the air in the system was scrubbed of CO₂ and water using a Dririte™ column followed by a Dririte™-Ascarite II™-Dririte™ column. Water vapor from the injected sample was removed by an additional magnesium perchlorate column before entering the sample cell of the gas analyzer. Because both the purge and injection times were recorded in Expedata to the second, the exact time that each pupa spent sealed in the chamber was known and used in the subsequent analysis of respiration rates. Calibration of the gas analyzer was maintained during the experiment using pure nitrogen and a certified mixture of 465.8 ppm CO₂ in nitrogen (Airgas, Radnor, PA, USA).

The respiration rate for each individual at each time point was calculated using the manual bolus integration method of Lighton (2008) with

$$R = \frac{\int_{t_i}^{t_f} CF \cdot dt}{e}$$

where R is the rate of respiration in $\mu\text{L CO}_2/\text{h}$, C is the instantaneous concentration of CO₂, F is the flow rate in l/s , e is the elapsed time interval in seconds between syringe purge and injection, and t_i - t_f is the time interval in seconds over which the injection bolus is detected by the gas analyzer. We conducted subsequent analyses on wet-mass equivalent metabolic rates in $\mu\text{L CO}_2/\text{h/mg}$. Note that the expected variation in developmental state among individuals at a given time point precluded us from establishing a more complex allometric relationship between body mass and metabolic rate in this system.

Analysis of metabolic trajectories

The previous study by Ragland et al. (2009) described a biphasic shape to *R. pomonella* post-winter metabolic trajectories, with a depressed baseline respiration rate during diapause punctuated by a logistic increase in CO₂ production following the end of diapause that stabilizes at a plateau level before entering an exponential increase phase during later pharate adult development (Fig. 1b). Initial inspection of our data revealed that this basic biphasic pattern was present, but the majority of individuals in our study also revealed a characteristic increase in metabolic rate after transfer from cold to warm conditions, followed by an attenuation back down to a stable baseline (Fig. 1c). This initial increase in metabolic rate followed by a return to depression 2-3 days after, indicating pupae were still in the diapause maintenance phase, was also confirmed by an additional follow-up experiment described below. To account for this additional feature, we fit and compared three nested non-linear models to each individual fly's time series of respiration rates: 1) a monophasic model consisting of an exponential increase from a single post-winter baseline metabolic rate (Fig. 1a):

$$R = b + \left(\frac{b}{2} e^{(t-x)c} \right)$$

2) a biphasic model similar to the one described by Ragland et al. (2009), but redesigned so that the parameter values are more directly reflective of landmark time points and metabolic rates (Fig. 1b):

$$R = b + \left(\frac{p - b}{1 + e^{m-t}} \right) + \left(\frac{p}{2} e^{(t-x)c} \right)$$

and 3) a triphasic model that also includes an initial post-winter elevated respiration rate followed by an immediate exponential decrease down to a baseline diapause rate (Fig. 1c):

$$R = (I \cdot e^{-t}) + b + \left(\frac{p - b}{1 + e^{m-t}} \right) + \left(\frac{p}{2} e^{(t-X)c} \right)$$

where R is the mass equivalent respiration rate at time t , b is the baseline post-winter metabolic rate, p is the plateau metabolic rate between diapause termination and pharate adult development, X represents the timing of the exponential increase in respiration during pharate adult development (equal to the time at which the metabolic rate is 2x higher than the previous stable rate b or p), m is the timing of the logistic increase during diapause termination, I is the initial elevated post-winter metabolic rate, and c is a scaling factor for the slope of the exponential phase. Initial AIC analysis during model development did not support the inclusion of additional scaling parameters for the logistic increase or initial decrease phase for any individuals.

All three models were fit for each individual in R 3.0.1 (R Development Core, Vienna, Austria), using the *nls* function. Starting values for each parameter were obtained from visual inspection of the metabolic trajectories; each parameter except for c models either a metabolic rate or time, and thus a position on the y or x axis of the metabolic trajectory plot, respectively (Fig. 1). Parameters were constrained to being biologically reasonable (e.g., all time and metabolic rate parameters > 0). Models were compared using AICc in the R package MuMIN (Barton 2015). Goodness of fit of the models selected by AICc was assessed by examining fitted curves and the distribution of residuals.

Testing whether variation in adult emergence timing is associated with variation in the same developmental trajectory model parameters across both host races

If particular phases of diapause or post-diapause development account for differences in adult emergence phenology, then inter-individual variation in parameters from the above models representing these phases should associate with adult emergence timing 1) within, and 2) between fly host races. Due to strong covariance among diapause trajectory parameters, as well as collinearity between some parameter estimates and their errors (as expected in this case where the degrees of freedom in each trajectory model vary exactly with phenology because flies that take longer to emerge are measured more times), we relied primarily on a partial correlation approach rather than multiple linear regression for statistical testing. We conducted partial correlation analyses assessing the relationships of the parameters b , m , p , $X-m$, c to days to adult emergence using the R package ppcor (Kim 2015). Analysis of variation was conducted both across the host races considering host as a binary variable, and within apple and hawthorn flies considered separately.

Analysis of over-wintering and immediate post-winter metabolic rates

The unexpected presence of elevated metabolic rates in the first few post-winter measurements 2-3 days after removal from cold combined with early diapause termination of many apple flies introduced more variability into the initial post-winter baseline measurements than expected in the original experiment. To better control for the rapid change in metabolic rates during this immediate post-winter phase and to further examine the dynamics of the immediate post-winter metabolic increase and subsequent decline, we performed a second metabolic phenotyping experiment the following year. Here, we measured metabolic rates of 40 apple race and 40 hawthorn race pupae from the same populations at Urbana, Illinois (but collected during the 2014 field season). Larvae and pupae were handled as above except that we used a 28-week simulated winter. At the end of this 28-week cold period, we placed the pupae in syringes,

purged the syringes with CO₂ free air as described above and sealed them under continuous simulated winter conditions (3.5 °C) for 24 hours. After 24 hours, syringes were placed in an insulated cooler with ice packs and brought to our room-temperature respirometry system. Sealed syringes were kept in the cooler until immediately before injection, resulting in a measurement of CO₂ produced by each individual under uninterrupted winter temperatures. The post-winter metabolic rate of the flies was then measured for 12 days at 24°C as described above with the exception that measurements were made every 24 hours. Syringes were re-purged immediately after injection. Metabolic rates were calculated as described above. Pupae were then kept in humidity chambers in 0.2 ml perforated tubes and monitored daily for adult emergence.

Statistical analyses of immediate post-winter metabolic rates.

Our analyses of this second experiment had two distinct goals: 1) to determine whether the two host races differ in their metabolic profiles over this dynamic period of immediate post-winter metabolic change, and 2) to determine the extent to which metabolic rates during the overwintering and immediate post-winter phase may influence the timing of adult emergence. To test whether and at which time points the host races differ in their immediate post-winter diapause metabolism, we conducted a repeated-measures ANOVA of daily metabolic rates for each fly as a function of host race to determine metabolic time points that differed significantly between apple and hawthorn flies. Based on those results, we then asked how variation in two key post-winter time points that differentiated the host races, specifically 1 day and 10 days out of overwintering, and variation in the cold overwintering metabolic rates covaried with each other and with adult emergence phenology using a partial correlation analyses, as above.

Results

Metabolic rate trajectories suggest stereotypical developmental trajectories

Apple host race flies emerged as adults significantly earlier than hawthorn host race flies with mean and median eclosion times for the apple race flies at 51d and 52d (SE = 2.33, IQR = 38.25 – 56.75), respectively compared to 81d and 79d (SE = 2.98, IQR = 73 - 87.5) for the hawthorn race (Fig. 2a, KS test, $D = 0.793$, $P < 0.00001$). The majority of pupae followed either a biphasic or triphasic metabolic rate trajectory model based on AICc model selection (Fig. 2b). However, there was a single fly showing the monophasic trajectory that emerged only 19 days after simulated winter. AICc results for each fly are presented in Table S1, and the best-fit models for each fly are presented in a file in the online supplementary material and Figure S1. Apple flies had more metabolic rate trajectories following the biphasic model fit, whereas, hawthorn flies had a roughly equal ratio of bi- and triphasic model fits (G-test, $G_{(2df, N = 68)} = 16.962$, $p < 0.001$), Table S1). The lower frequency of triphasic trajectories in apple flies was likely driven by their shorter time to adult emergence. With fewer data points during the baseline metabolic rate phase of the trajectories, apple flies tended to have less power to fit the initial decrease curve, with the logistic increase of diapause termination often beginning before the decrease phase stabilized (Fig. S1).

The pattern of initial decrease of metabolic rates from an immediate post-winter peak was present in both hawthorn and apple flies (Fig. S1 and see results of second experiment below). Curves from the fitted models (Fig. 2b) generally fit the shape of the data closely (Fig. S1). The mean residual standard errors S were low for the best-fit models, with a mean of 0.0041 $\mu\text{l CO}_2/\text{h}$ across all flies and a range of 0.0012 – 0.0089 $\mu\text{l CO}_2/\text{h}$. We note that R^2 is generally not an appropriate metric of goodness-of-fit for non-linear regressions (Spiess & Neumayer 2010). To use parameter estimates from the best fit models for each fly in subsequent analyses, we tested

whether the subset of nested shared parameter estimates differed between the hierarchical biphasic and triphasic models fit for the same individuals. These shared parameters: baseline metabolic rate during the diapause maintenance module (b), timing of the end of the diapause maintenance module (m), post-diapause plateau metabolism (p), post-diapause plateau duration (X), and exponential scaling (c) did not differ between the bi- and tri-phasic models in paired t -tests (Table S2).

Apple and hawthorn flies differ in key post-winter metabolic trajectory parameters

While the host races share the same basic metabolic trajectory templates, apple and hawthorn flies differed in parameters related to both phase-specific metabolic rates and, most notably, the timing of the transition out of the diapause maintenance phase. Estimates of three parameters: baseline metabolic rate b , the timing of the transition out of the diapause maintenance phase m , and the post-diapause plateau metabolic rate p , differed between host races (Fig. 3a-c; two-tailed t -tests, $t = 4.0787$, $P = 0.0026$; $t = -8.61$, $P = 3.64 \times 10^{-12}$; $t = 3.8012$, $P = 0.000385$, respectively). Neither the relative timing of exponential metabolic increase during adult development $X-m$ nor the exponential scaling parameter c were significantly different between host races (Fig. 3d-e; t -tests, $t = -0.39081$, $P = 0.6973$; $t = 1.444$, $P = 0.1542$, respectively). Compared to hawthorn flies, apple flies had higher baseline metabolic rates, earlier diapause termination times, and higher post-diapause metabolic plateaus (Fig. 3a-c).

Associations between model parameters and emergence timing within host races

Our correlation analysis revealed a strong relationship across both host races between m , the timing of the end of the diapause module, and adult emergence timing across both host races ($r = 0.994$, $P = 1.07 \times 10^{-57}$, Fig. 3G, Fig. 4). Earlier timing of the end of the diapause maintenance

module was tightly correlated with earlier emergence timing (Fig. 3G). Weaker but significant partial correlations with adult emergence timing also existed between post-diapause plateau metabolism (p), metabolic rate duration of the post-diapause plateau ($X-m$) and the slope of the exponential increase phase c ($r = -0.37$; $P = 0.0013$; $r = 0.772$, $P = 5e^{-13}$, and $r = -0.533$, $P = 0.00015$, respectively; Fig. 3I-J; Fig. 4). The baseline metabolic rate b did not show significant direct correlations with adult emergence time (Fig 3F; Fig 4). These patterns were consistent both within and across the two host races (Fig 3F-J; Fig. 4; Fig S2). The strength of the correlation between m and adult emergence timing was nearly identical when the partial correlation analyses were run separately for each host race (apple $r = 0.98$ and hawthorn $r = 0.99$; Fig S2), reinforcing the importance of the timing of the end of the diapause maintenance module as the primary driver of phenological divergence between the apple and hawthorn host races.

Host races diverge in how they respond to initial favorable conditions

The overall shape of the metabolic trajectories for both host races in the second respirometry experiment was consistent with the dynamics found in our first experiment. The end of simulated winter and transfer to 25 °C resulted in a sharp initial metabolic increase that peaked at 24-72 hours after removal from the cold and subsequently declined to a baseline metabolic rate over the course of 10 days (Fig. 5). Overwintering metabolic rates (measured in the cold) were nearly an order of magnitude lower than post-winter baseline metabolic rates from the experiment above, mean 0.00115 vs. 0.01035 $\mu\text{CO}_2/\text{hr}/\text{mg}$, respectively, and did not differ between the host races (Fig. S3). However, the two host races did differ in their immediate response to warming, with apple flies being more metabolically responsive to warming than hawthorn flies (repeated measures ANOVA: time x host, $F_{12,552} = 3.54$ $p < 0.001$; Table S3; Fig.

5). Thus, apple flies were more metabolically responsive to the shift from cool simulated overwintering conditions to warm post-winter conditions, rising to higher metabolic rates than hawthorn flies after simulated overwintering, particularly on days 1 & 2 post-winter, with some individuals appearing to terminate diapause before day10 (Fig. 5).

Partial correlation analysis of overwinter, day 1, and day 10 metabolic rates with adult emergence time in the second experiment showed significant but weaker overall covariance among variables than was found in the first experiment (Fig 6). Overwintering metabolic rate was weakly but significantly positively correlated with metabolic rates at the two other time points day 1 and day 10 after removal from cold, as well as host ($r = 0.37$, $P = 0.008$, $r = 0.346$, $P = 0.013$; $r = 0.315$, $P = 0.026$, respectively, Figs. 6, S4). The only variable with a direct correlation to adult emergence timing was an inverse relationship with day 1 metabolic rate ($r = -0.318$, $P = 0.025$, Figs 6, S4). Pupae of both host races that were more metabolically responsive during the rapid rise in metabolism after shifting from cool overwintering conditions to warm post-winter conditions were more likely to emerge earlier as adults. This correlation was much weaker than that observed between adult emergence timing and the timing of the end of the diapause maintenance module ($r = -0.318$ vs $r = 0.994$), indicating that if day 1 metabolic rate influences adult emergence phenology, it likely does so through its effect on the timing of the end of diapause maintenance.

DISCUSSION

Diapause is an important component of adaptation to temperate environments for most insect species, and it may often serve as a critical step governing the seasonal phenological timing of insect life histories (Tauber & Tauber 1981). As a multi-faceted, alternative

developmental pathway, diapause is not just a static resting state. Diapause is a complex trait, possessing a number of physiologically distinct functional components that are expressed along the course of diapause development (Kostal 2006). Thus, the ability for insect populations to rapidly evolve new phenologies is likely dependent on the complexity with which these different components of diapause development interact to affect overall seasonal timing. Here, we show that although apple and hawthorn flies differ in a number of timing and metabolic rate parameters of post-winter development (Figs 3A-E), variation in adult emergence, both between and within the host races, is primarily due to differences in the timing of the transition between diapause maintenance and post-diapause development (parameter m ; Fig 3F-J; Fig 4). In both host races, adult flies emerged on average 31.6 days after diapause termination, with a standard deviation of 2.4 days. Wadsworth et al. (2013) reported similar results for a primary effect of diapause termination on adult emergence time in the univoltine-Z and bivoltine-E strains of the European corn borer, another system where divergence in seasonal phenology drives allochronic isolation (Dopman et al. 2010), perhaps highlighting the general importance of diapause termination timing in the evolution of insect phenology.

In addition to diapause termination, adult emergence timing in the host races is related, although to a much lesser degree, to two other post-winter development parameters: 1) the duration of the post-diapause plateau ($X-m$) when early pharate adult morphogenesis is occurring (Ragland et al. 2009); and 2) the slope of the exponential increase phase of pharate adult development (c) late in the process of adult morphogenesis (Figs 3H-J). In our second experiment, a weak but significant inverse correlation was also observed between adult emergence and the initial day 1 post-winter metabolic rate, such that pupae that more strongly increase metabolic rates after the transition from cool over-wintering temperatures to warm post-

winter temperatures are more likely to emerge as adults earlier (Fig 6). However, the timing of the end of the diapause maintenance phase appears to be the major factor driving variation in overall adult emergence phenology, with the later-acting post-diapause development parameters having much subtler effects on any residuals variation in emergence timing (Figs 3H-J, Fig. 4).

The overall covariance structure presented in the partial correlation network (Fig. 4) indicates that the major effect of the timing of the transition between the diapause maintenance and post-diapause development phases on phenology is not entirely independent of the other metabolic trajectory parameters. Although the timing of transition between phases appears to be the primary driver of phenology, this variation may not act in a singularly modular fashion, as we would infer if the parameters correlated with adult emergence timing did not otherwise overlap in the partial correlation network. However, most of the correlations among metabolic trajectory parameters were relatively modest (only two were stronger than $r = 0.5$, with the strongest being between m and $X-m$ at $r = -0.76$), implying that their respective responses to selection may not be highly constrained. Thus, the covariance between diapause termination and post-diapause development is not absolute, both the diapause maintenance phase and the post-diapause development phase may be capable of separate responses to selection on adult life history timing, a result that has broader implications for phenological adaptation in many contexts from seasonal synchronization with novel hosts to rapid adaptation to new climatic regimes during introductions or range expansion to shifting seasonality as a result of climate change (e.g. Bradshaw & Holzapfel 2001; Ording & Scriber 2005; Gomi et al. 2007; Wadsworth et al. 2013).

In certain insects, variation in diapause phenotype may have relatively simple genetic architecture. Diapause-mediated allochronic isolation in the European corn borer is under the control of a segregating locus of major effect on the Z-chromosome (Glover et al. 1992; Dopman

et al. 2005; Wadsworth et al. 2015). Similarly, variation in diapause induction appears to be driven largely by a large-effect Z-linked locus in a butterfly species complex comprised of *Papilio glaucus*, *P. canadensis*, and *P. appalacheinsis* (Hagen & Scriber 1989; Kunte et al. 2011). But latitudinal variation in life history timing in *Papilio* butterflies may also be associated with autosomal regions as well (Ryan et al. 2017). In contrast, geographic variation in photoperiodic sensitivity for diapause induction in the mosquito *Wyeomia smithii* appears to be under more complex genetic control, involving multiple loci of moderate effect with strong epistasis (Lair et al. 1997; Mathias et al. 2007). Genetic variation in life history timing in *R. pomonella* appears to be highly quantitative, showing widespread statistical associations with markers across the genome (Filchak et al. 2000; Michel et al. 2010; Egan et al. 2015; Ragland et al. in 2017). Our results suggest that the many genes implied to affect adult emergence phenology primarily do so by their effect on the timing of termination of the diapause maintenance module. Thus, timing of diapause termination is a developmental hub through which quantitative allelic variation flows to modify adult emergence phenology.

The genetic complexity of life history timing in *R. pomonella* might seem at odds with the relative simplicity of diapause termination primarily dictating adult emergence timing. However, there is good reason to suspect that many genes or gene networks may play a role in driving variation in the duration of the diapause maintenance module. The pre-winter initiation of diapause is not simply a switch ceasing development, and the termination of diapause is not simply a switch resuming development. Rather the phases are each characterized by a progression of physiological changes that lead up to each transition along the developmental pathway (Denlinger 2002; Kostal 2006; Kostal et al. 2017). The dynamic nature of this developmental trajectory allows for many different transitions at which allelic variation might

influence diapause timing, and many intermediary developmental avenues exist along which highly polygenic variation could act additively to affect eclosion timing. Precisely how each of these processes and events relate to the detectable metabolic increase during diapause termination is not currently known. Further characterizing variation in these intermediate mechanistic steps to diapause termination will help parse the complex pattern of genomic variation associated with life history timing in these flies.

The earlier diapause termination of apple than hawthorn flies may partially be due to differences in their developmental progression during the overwintering portion of the diapause maintenance phase. Apple pupae may become more primed or potentiated to respond to environmental cues sooner after winter than hawthorn pupae. For example, apple flies may be prone to enter or progress further into the stage of diapause development permissive for termination of the maintenance phase earlier in the winter compared to hawthorn flies. Alternatively, the host races could emerge from winter in the same state of diapause development, with the difference in adult emergence time resulting from variation in their respective responses to subsequent environmental cues. Although the two host races did not show a detectable difference in overwintering metabolic rate, the fact that apple flies were more metabolically responsive to the shift from cool overwintering temperatures to warm post-winter temperatures, and the very rapid nature of diapause termination in many of the apple flies, argues that apple fly pupae are potentiated to respond to diapause-termination cues earlier than hawthorn fly pupae. The finding of higher initial metabolic response to warming in apple flies combined with the association between earlier adult emergence and higher initial metabolic rates just after warming in both host races suggests a modification to the simple hypothesis that basal metabolic rates during diapause themselves are a primary driver of diapause duration (Wipking

et al. 1995; Feder & Filchak 1999). Instead, perhaps the dynamic responsiveness of metabolism to thermal shifts indicates greater potentiation to respond to diapause termination cues both within and between the host races and acts upstream of the timing of the end of diapause maintenance.

A recent RNAseq study comparing the transcriptomes of the host races during simulated overwintering and soon after removal from the cold is consistent with the hypothesis that apple flies are more potentiated to terminate diapause after winter than hawthorn flies (Meyers et al. 2016). The head transcriptomes (including neuro-endocrine tissues) of apple and hawthorn flies were found to already be strongly differentiated before pupae were removed from the cold, with biases towards up-regulation of genes involved in cell proliferation and development in the apple flies (Meyers et al. 2016). Our finding of greater responsiveness of metabolic rate in apple than hawthorn race pupae as they shift from cold, developmentally suppressive temperatures to warm, developmentally permissive temperatures also suggests apple race pupae become more potentiated for diapause termination by the end of simulated winter treatments compared to hawthorn race pupae.

The immediate post-winter metabolic increase and subsequent decline observed in our second experiment (Fig. 5) is a novel finding with implications for how dormancy is maintained in the face of abiotic conditions that are otherwise permissive for non-diapause development. Diapause has been most extensively studied in insects with multivoltine life cycles (e.g. Schmidt et al. 2008; Ragland et al 2010; Emerson et al. 2010; Bao et al. 2011; Lehmann et al. 2014). For multivoltine organisms, trophic resources tend to have broad temporal distributions, and selection favors fast resumption of non-diapause development as soon as abiotic conditions are physiologically permissive. Thus, it is not surprising that for multivoltine species, the diapause

maintenance phase typically terminates during winter, with insects remaining in post-diapause quiescence (dormant due to physiologically restrictive abiotic conditions rather than programmed metabolic and developmental suppression) until physiological temperature thresholds are surpassed, after which time they initiate post-diapause development (Hayward et al. 2005). In contrast, univoltine insects with narrow resource windows like *R. pomonella* derive no benefit from terminating diapause maintenance early during the winter because they must remain dormant even after conditions become permissive the following spring until it is time to initiate the post-diapause developmental phase to synchronize themselves with host fruit phenology.

The response of *R. pomonella* to the end of simulated winter was therefore more metabolically dynamic than we expected (Fig. 5). The shift from overwinter respiration to post-winter diapause respiration was neither a gradual linear increase nor a discrete stair-step up to the post-winter level. Rather, pupae showed a rapid but transient metabolic increase immediately after removal from the cold, peaking around 48 hours at a level 4 to 5 fold higher than the eventual post-winter baseline metabolic rates (Fig. 5). During normal development, metabolic rates of ectotherms increase with temperature (Angiletta 2009). Thus, thermally sensitive metabolic rates can lead to increased energy consumption during warmer conditions, which may prove costly to univoltine insects that need to conserve energy stores for prolonged dormancy (Irwin & Lee 2000; Mercader & Scriber 2004; Kostal et al. 2011; Vrba et al. 2014). However, diapausing insects may be able to mitigate these energetic costs by reducing the thermal sensitivity of their metabolic rates (Williams et al. 2012; 2015), as we saw in our trials when initially high metabolic rates were reduced within days of transfer to warm, permissive thermal conditions. Active suppression of metabolism in the face of physiologically permissive temperatures may play an important role in maintaining optimal overwintering energetics in the

face of intermittent periods of warmth both in the fall and spring (Hahn & Denlinger 2011; Sinclair 2015). Differences in the immediate post-winter metabolic dynamics of apple and hawthorn flies might therefore represent an additional axis of divergent seasonal adaptation in this classic speciation system that may also be important to disentangle in other systems with rapid adaptation to new phenological regimes.

CONCLUSIONS

Specialist insects make up a large proportion of metazoan biodiversity (Jaenike 1990; Bush & Butlin 2004). How insect life history timing evolves has important implications for both the origin and maintenance of a large swathe of biodiversity (Bush 1993). The tempo of seasonal adaptation for many species likely depends on the covariance structure of different components of dormancy regulation and how these components act to influence phenology. Are different facets of diapause development free to evolve independently, or are shifts in life history timing constrained by a need for simultaneous evolution across several phases of diapause development? Here, we found that termination of the diapause maintenance phase plays a particularly important (but not sole) role in synchronizing insect phenology with that of ephemeral resources (Figs 3,4). In the case of *R. pomonella*, complex patterns of quantitative allelic variation appear to act primarily through the hub of diapause termination to affect seasonal timing, with additional input via post-diapause development. The dual roles of these two developmental phases may be a coarse and fine adjustment of phenology, respectively. It is important to note that the magnitude of effects of diapause termination and post-diapause development reported here were measured under a simplified laboratory temperature regime: a constant post-winter temperature of 25 °C. It remains to be determined whether the effects of either of these phases on eclosion phenology are themselves function-valued traits driven by

temperature. One possibility is that the duration of the diapause maintenance phase sets life history timing broadly across years and that the post-diapause development module may allow fine-tuning of adult emergence timing to within year variation in host-plant phenology. For example, apples always fruit earlier than hawthorns in our sympatric field sites. However, in some warm years, apples and hawthorns both fruit earlier and in cooler years both fruit later. Metabolic rates during the post-diapause developmental phase are substantially higher than during the diapause maintenance phase and we speculate that post-diapause development may be more responsive to temperature acting to fine tune synchrony with host plant availability and local abiotic conditions from year to year at any particular location and across a latitudinal gradient of seasonal conditions throughout this fly's range. Future work on the thermal sensitivity of specific modular components across these two modules of the diapause developmental trajectory is needed to test this hypothesis for synchronizing adult emergence timing with weather-related local variation in host fruit phenology.

In *Rhagoletis* flies, variation in seasonal adult emergence timing is associated with on-going divergent adaptation to the novel apple host plant and reproductive isolation, but this shift to earlier seasonal timing is analogous to expectations for phenological shifts under climate change in coming decades. A primary motivation of this research was to determine the extent to which the inherent complexity of diapause development is likely to aid or constrain the rapid evolution of seasonality in univoltine insects. In the case of these two host races of *R. pomonella*, the partially independent major and minor effects related to the timing of the transition out of diapause maintenance and components of post-diapause development bodes well for the continued evolvability of adult eclosion time in this system. If phenological adaptation in these flies is not primarily constrained by strong antagonistic pleiotropy among different facets of

diapause development, it may be limited more by available genetic variation affecting each intermediate phenotype. However, the apple and hawthorn fly speciation story is superimposed on strong local phenological adaptation of these flies along latitudinal clines as well (Dambroski & Feder 2007; Michel et al. 2010; Doellman et al. 2018a, b), which may act as a vast reservoir for standing genetic variation in seasonality traits across the continent (Powell et al. 2013). The combination of relatively simple, major effect basis for variation in adult emergence phenology coupled with abundant genetic variation may mean that natural selection has a fairly free hand with these insects to keep pace with changing seasonal conditions or colonize novel niches. However, the existence of moderate antagonistic covariance among traits with links to phenology may potentially constrain the evolution of novel phenotypes outside the current trait distribution. Furthermore, elucidating the roles of diapause maintenance and post-diapause development as important intermediate phenotypes driving differences in phenology sets the stage for future studies to begin drawing stronger connections between allelic variation across the genome and the complex ecological traits of life history timing, local adaptation, and ecological reproductive isolation. Given that genetic variation in the diapause maintenance phase (Wadsworth et al. 2013) or in post-diapause development (Poseldovich et al. 2012; 2014; Stalhansske et al. 2014) are thought to be independent drivers of adaptive shifts in phenology in other insect systems, we expect that the dual role for diapause and post-diapause development in regulating life history timing may be partially modular across a wide range of insect species. Thus, the idea that phenologies may be tuned to local seasonality by a combination of genetic and environmental effects (GxE) is broadly applicable across diverse groups of insects and germane to understanding persistence versus declines of insect populations in the future as seasonality is further altered by anthropogenic change.

Acknowledgements

We would like to thank Meredith Doellman, Glen Hood, and Pete Meyers for help with field collections. Caroline Williams provided invaluable advice on our respirometry configuration. Help with phenotyping was provided by Jennifer Serviss, Caelen Schreiber, Andre Szejner, Chao Chen, Genevieve Comeau, and Xiaoping Wang. This research was supported by NSF IOS 170773 to GJR and JLF, NSF DEB 1638997 to GJR, NSF DEB 1638951 to JLF, and NSF DEB 1639005 to THQP and DAH, as well as NSF IOS 1257298, the Florida Agricultural Experiment Station, and the joint FAO/IAEA CRP Dormancy Management to Enable Mass-rearing to DAH.

REFERENCES CITED

- Abbot, P., and J. H. Withgott. 2004. Phylogenetic and Molecular Evidence for Allochronic Speciation in Gall-Forming Aphids (pemphigus). *Evolution* 58:539–553.
- Andrewartha, H. G. 1952. Diapause in Relation to the Ecology of Insects. *Biological Reviews* 27:50–107.
- Angilletta, M. J. 2009. *Thermal Adaptation: A Theoretical and Empirical Synthesis*. OUP Oxford.
- Bao, B., and W.-H. Xu. 2011. Identification of gene expression changes associated with the initiation of diapause in the brain of the cotton bollworm, *Helicoverpa armigera*. *BMC Genomics* 12:224.
- Barton, K. 2015. Multi-model inference. R package version 1.13. 4.

628 Berlocher, S.H. and J.L. Feder. 2002. Sympatric speciation in phytophagous insects: moving
629 beyond controversy? *Annual Review of Entomology* 47:773-815.

630 Bradshaw, W. E., and C. M. Holzapfel. 2001. Genetic shift in photoperiodic response correlated
631 with global warming. *Proceedings of the National Academy of Sciences* 98:14509–14511.

632 Bradshaw, W. E., P. A. Zani, C. M. Holzapfel, and J. Fry. 2004. Adaptation to temperate
633 climates. *Evolution* 58:1748–1762.

634 Bryon, A., A. H. Kurlovs, W. Dermauw, R. Greenhalgh, M. Riga, M. Grbić, L. Tirry, et al. 2017.
635 Disruption of a horizontally transferred phytoene desaturase abolishes carotenoid accumulation
636 and diapause in *Tetranychus urticae*. *Proceedings of the National Academy of Sciences*
637 114:E5871–E5880.

638 Bush, G. 1969. Sympatric Host Race Formation and Speciation in Frugivorous Flies of Genus
639 *Rhagoletis* (diptera, Tephritidae). *Evolution* 23:237-.

640 Bush, G. L., and R. K. Butlin. 2004. Sympatric Speciation in Insects. *Adaptive Speciation*, eds.
641 Dieckmann U, Doebeli M, Metz JAJ, & Tuatz D. Cambridge University Press.

642 Comparative transcriptomics support evolutionary convergence of diapause responses across
643 Insecta - Ragland - 2017 - *Physiological Entomology* - Wiley Online Library. n.d.

644 Crozier, A. J. G. 1979. Supradian and infradian cycles in oxygen uptake of diapausing pupae of
645 *Pieris brassicae*. *Journal of Insect Physiology* 25:575–582.

646 Dambroski, H. R., and J. L. Feder. 2007. Host plant and latitude-related diapause variation in
647 *Rhagoletis pomonella*: a test for multifaceted life history adaptation on different stages of
648 diapause development. *Journal of Evolutionary Biology* 20:2101–2112.

649 Denlinger, D. L. 2002. Regulation of Diapause. *Annual Review of Entomology* 47:93–122.

650 Doellman, M.M., S.P. Egan, G.J. Ragland, P.J. Meyers, G.R. Hood, T.H.Q. Powell, P. Lazorchak, D.A.
651 Hahn, S.H. Berlocher, P. Nosil, and J.L. Feder. 2018a. Standing genetic variation in eclosion time and
652 genome wide patterns of geographic and host-related differentiation in *Rhagoletis pomonella* Ecology
653 and Evolution 9. doi: 10.1002/ece3.4758.

654 Doellman, M.M., G.J. Ragland, G.R. Hood, P.J. Meyers, S.P. Egan, T.H.Q. Powell, P. Lazorchak, M.M.
655 Glover, C. Tait, H. Schuler, D.A. Hahn, S.H. Berlocher, J.J. Smith, P. Nosil, and J.L. Feder. 2018b.
656 Genomic differentiation during speciation-with-gene-flow: Comparing geographic and host-related
657 variation in divergent life history adaptation in *Rhagoletis pomonella*. Genes May 18;9(5). pii: E262. doi:
658 10.3390/genes9050262.

659 Dopman, E. B., L. Pérez, S. M. Bogdanowicz, and R. G. Harrison. 2005. Consequences of
660 reproductive barriers for genealogical discordance in the European corn borer. Proceedings of
661 the National Academy of Sciences of the United States of America 102:14706–14711.

662 Dopman, E. B., P. S. Robbins, and A. Seaman. 2010. Components of Reproductive Isolation
663 Between North American Pheromone Strains of the European Corn Borer. Evolution 64:881–
664 902.

665 Dres, M., and J. Mallet. 2002. Host races in plant-feeding insects and their importance in
666 sympatric speciation RID B-5114-2008. Philosophical Transactions of the Royal Society B-
667 Biological Sciences 357:471–492.

668 Dyck, H. V., D. Bonte, R. Puls, K. Gotthard, and D. Maes. 2015. The lost generation hypothesis:
669 could climate change drive ectotherms into a developmental trap? Oikos 124:54–61.

670 Egan, S. P., G. J. Ragland, L. Assour, T. H. Q. Powell, G. R. Hood, S. Emrich, P. Nosil, et al.
671 2015. Experimental evidence of genome-wide impact of ecological selection during early stages
672 of speciation-with-gene-flow. Ecology Letters 18:817–825.

Emerson, K. J., W. E. Bradshaw, and C. M. Holzapfel. 2010. Microarrays Reveal Early
Transcriptional Events during the Termination of Larval Diapause in Natural Populations of the
Mosquito, *Wyeomyia smithii*. PLOS ONE 5:e9574.

Eubanks, M., C. Blair, and W. Abrahamson. 2003. One host shift leads to another? Evidence of
host-race formation in a predaceous gall-boring beetle. *Evolution* 57:168–172.

Feder, J. L., and K. E. Filchak. 1999. It's about time: the evidence for host plant-mediated
selection in the apple maggot fly, *Rhagoletis pomonella*, and its implications for fitness trade-offs in phytophagous insects. Pages 211–225 in *Proceedings
of the 10th International Symposium on Insect-Plant Relationships, Series Entomologica*.
Springer, Dordrecht.

Feder, J.L., T.A. Hunt. and G.L. Bush. 1993. The Effects of Climate, Host Plant Phenology and
Host Fidelity on the Genetics of Apple and Hawthorn Infesting Races of *Rhagoletis pomonella*.
Entomologia Experimentalis et Applicata 69:117-135.

Feder, J. L., T. H. Q. Powell, K. Filchak, and B. Leung. 2010. The diapause response of
Rhagoletis pomonella to varying environmental conditions and its significance for geographic
and host plant-related adaptation. *Entomologia Experimentalis Et Applicata* 136:31–44.

Feder, J., S. Opp, B. Wlazlo, K. Reynolds, W. Go, and S. Spisak. 1994. Host fidelity is an
effective premating barrier between sympatric races. *Proceedings of the National Academy of
Sciences of the United States of America* 91:7990–7994.

Feder, J., J. Roethele, B. Wlazlo, and S. Berlocher. 1997. Selective maintenance of allozyme
differences among sympatric host races of the apple maggot fly. *Proceedings of the National
Academy of Sciences of the United States of America* 94:11417–11421.

695 Filchak, K. E., J. L. Feder, J. B. Roethele, U. Stolz, and J. Mallet. 1999. A field test for host-
696 plant dependent selection on larvae of the apple maggot fly, *Rhagoletis Pomonella*. *Evolution*
697 53:187–200.

698 Filchak, K. E., J. B. Roethele, and J. L. Feder. 2000. Natural selection and sympatric divergence
699 in the apple maggot *Rhagoletis pomonella*. *Nature* 407:739–742.

700 Forbes, A. A., T. H. Q. Powell, L. L. Stelinski, J. J. Smith, and J. L. Feder. 2009. Sequential
701 sympatric speciation across trophic levels. *Science* 323:776–779.

702 Friesen, V. L., A. L. Smith, E. Gómez-Díaz, M. Bolton, R. W. Furness, J. González-Solís, and L.
703 R. Monteiro. 2007. Sympatric speciation by allochrony in a seabird. *Proceedings of the National*
704 *Academy of Sciences of the United States of America* 104:18589–18594.

705 Funk, D., K. Filchak, and J. Feder. 2002. Herbivorous insects: model systems for the
706 comparative study of speciation ecology. *Genetica* 116:251–267.

707 Gilman, R. T., N. S. Fabina, K. C. Abbott, and N. E. Rafferty. 2011. Evolution of plant–
708 pollinator mutualisms in response to climate change. *Evolutionary Applications* 5:2–16.

709 Glover, T. J., P. S. Robbins, C. J. Eckenrode, and W. L. Roelofs. 1992. Genetic control of
710 voltinism characteristics in European corn borer races assessed with a marker gene. *Archives of*
711 *Insect Biochemistry and Physiology* 20:107–117.

712 Gomi, T., M. Nagasaka, T. Fukuda, and H. Hagihara. 2007. Shifting of the life cycle and life-
713 history traits of the fall webworm in relation to climate change. *Entomologia Experimentalis et*
714 *Applicata* 125:179–184.

715 Hagen, R. H., and J. M. Scriber. 1989. Sex-linked diapause, color, and allozyme loci in *Papilio*
716 *glaucus*: linkage analysis and significance in a hybrid zone. *Journal of Heredity* 80:179–185.

717 Hahn, D. A., and D. L. Denlinger. 2007. Meeting the energetic demands of insect diapause:
718 Nutrient storage and utilization. *Journal of Insect Physiology*, VIII European Congress of
719 Entomology - Physiology and Endocrinology 53:760–773.

720 ———. 2011. Energetics of insect diapause. *Annual Review of Entomology* 56:103–121.

721 Hayward, S. A. L., S. C. Pavlides, S. P. Tammariello, J. P. Rinehart, and D. L. Denlinger. 2005.
722 Temporal expression patterns of diapause-associated genes in flesh fly pupae from the onset of
723 diapause through post-diapause quiescence. *Journal of Insect Physiology*, Diapause and
724 Biological Clocks 51:631–640.

725 Hood, G. R., A. A. Forbes, T. H. Q. Powell, S. P. Egan, G. Hamerlinck, J. J. Smith, and J. L.
726 Feder. 2015. Sequential divergence and the multiplicative origin of community diversity.
727 *Proceedings of the National Academy of Sciences* 112:E5980–E5989.

728 Horner, J. D., T. P. Craig, and J. K. Itami. 1999. The influence of oviposition phenology on
729 survival in host races of *Eurosta solidaginis*. *Entomologia Experimentalis et Applicata* 93:121–
730 129.

731 Irwin, J. T., and R. E. Lee. 2000. Mild winter temperatures reduce survival and potential
732 fecundity of the goldenrod gall fly, *Eurosta solidaginis* (Diptera: Tephritidae). *Journal of Insect*
733 *Physiology* 46:655–661.

734 Jaenike, J. 1990. Host specialization in phytophagous insects. *Annual Review of Ecology and*
735 *Systematics* 21:243–273.

736 Karpovich, S. A., Ø. Tøien, C. L. Buck, and B. M. Barnes. 2009. Energetics of arousal episodes
737 in hibernating arctic ground squirrels. *Journal of Comparative Physiology B* 179:691–700.

738 Kim, S. 2015. ppcor: An R Package for a Fast Calculation to Semi-partial Correlation
739 Coefficients. *Communications for statistical applications and methods* 22:665–674.

740 Košťál, V. 2006. Eco-physiological phases of insect diapause. *Journal of Insect Physiology*
741 52:113–127.

742 Košťál, V., J. Korbelová, J. Rozsypal, H. Zahradníčková, J. Cimlová, A. Tomčala, and P. Šimek.
743 2011. Long-term cold acclimation extends survival time at 0°C and modifies the metabolomic
744 profiles of the larvae of the fruit fly *Drosophila melanogaster*. *PLOS ONE* 6:e25025.

745 Košťál, V., T. Štětina, R. Poupardin, J. Korbelová, and A. W. Bruce. 2017. Conceptual
746 framework of the eco-physiological phases of insect diapause development justified by
747 transcriptomic profiling. *Proceedings of the National Academy of Sciences* 201707281.

748 Kudo, G., and T. Y. Ida. 2013. Early onset of spring increases the phenological mismatch
749 between plants and pollinators. *Ecology* 94:2311–2320.

750 Kunte, K., C. Shea, M. L. Aardema, J. M. Scriber, T. E. Juenger, L. E. Gilbert, and M. R.
751 Kronforst. 2011. Sex chromosome mosaicism and hybrid speciation among tiger swallowtail
752 butterflies. *PLOS Genetics* 7:e1002274.

753 Lair, K. P., W. E. Bradshaw, and C. M. Holzapfel. 1997. Evolutionary divergence of the genetic
754 architecture underlying photoperiodism in the pitcher-plant mosquito, *Wyeomyia smithii*.
755 *Genetics* 147:1873–1883.

756 Lehmann, P., S. Piironen, M. Kankare, A. Lyytinen, M. Paljakka, and L. Lindström. 2014.
757 Photoperiodic effects on diapause-associated gene expression trajectories in European
758 *Leptinotarsa decemlineata* populations. *Insect Molecular Biology* 23:566–578.

759 Lighton, J. R. B. 2008. Measuring Metabolic Rates: A Manual for Scientists. Oxford University
760 Press, USA.

761 Lowry, D. B., R. C. Rockwood, and J. H. Willis. 2008. Ecological reproductive isolation of coast
762 and inland races of *Mimulus guttatus*. *Evolution* 62:2196–2214.

763 M J Tauber, and and C. A. Tauber. 1976. Insect seasonality: Diapause maintenance, termination,
764 and postdiapause development. *Annual Review of Entomology* 21:81–107.

765 Mathias, D., L. Jacky, W. E. Bradshaw, and C. M. Holzapfel. 2007. Quantitative trait loci
766 associated with photoperiodic response and stage of diapause in the pitcher-plant mosquito,
767 *Wyeomyia smithii*. *Genetics* 176:391–402.

768 Melo, D., and G. Marroig. 2015. Directional selection can drive the evolution of modularity in
769 complex traits. *Proceedings of the National Academy of Sciences* 112:470–475.

770 Mercader, R. J., and J. M. Scriber. 2008. Asymmetrical thermal constraints on the parapatric
771 species boundaries of two widespread generalist butterflies. *Ecological Entomology* 33:537–545.

772 Meyers, P. J., T. H. Q. Powell, K. K. O. Walden, A. J. Schieferecke, J. L. Feder, D. A. Hahn, H.
773 M. Robertson, et al. 2016. Divergence of the diapause transcriptome in apple maggot flies:
774 winter regulation and post-winter transcriptional repression. *Journal of Experimental Biology*
775 219:2613–2622.

776 Michel, A. P., S. Sim, T. H. Q. Powell, M. S. Taylor, P. Nosil, and J. L. Feder. 2010. Widespread
777 genomic divergence during sympatric speciation. *Proceedings of the National Academy of*
778 *Sciences of the United States of America* 107:9724–9729.

779 Miller-Rushing, A. J., T. T. Høye, D. W. Inouye, and E. Post. 2010. The effects of phenological
780 mismatches on demography. *Philosophical Transactions of the Royal Society of London B:*
781 *Biological Sciences* 365:3177–3186.

782 Ordning, G. J., R. J. Mercader, M. L. Aardema, and J. M. Scriber. 2009. Allochronic isolation and
783 incipient hybrid speciation in tiger swallowtail butterflies. *Oecologia* 162:523–531.

784 Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. *Annual*
785 *Review of Ecology, Evolution, and Systematics* 37:637–669.

786 Posledovich, D., T. Toftegaard, J. A. Navarro-Cano, C. Wiklund, J. Ehrlén, and K. Gotthard.
787 2014. Latitudinal variation in thermal reaction norms of post-winter pupal development in two
788 butterflies differing in phenological specialization. *Biological Journal of the Linnean Society*
789 113:981–991.

790 Posledovich, D., T. Toftegaard, C. Wiklund, J. Ehrlén, and K. Gotthard. 2015. Latitudinal
791 variation in diapause duration and post-winter development in two pierid butterflies in relation to
792 phenological specialization. *Oecologia* 177:181–190.

793 Powell, T.H.Q., G.R. Hood, M.O. Murpy, J. Heilveil, S.H. Berlocher, P. Nosil, and J.L. Feder.
794 2013. Genetic divergence along the speciation continuum: the transition from host race to species
795 in *Rhagoletis* (Diptera: Tephritidae). *Evolution* 67: 2561-2576.

796 Ragland, G. J., D. L. Denlinger, and D. A. Hahn. 2010. Mechanisms of suspended animation are
797 revealed by transcript profiling of diapause in the flesh fly. *Proceedings of the National*
798 *Academy of Sciences* 107:14909–14914.

799 Ragland, G. J., M. M. Doellman, P. J. Meyers, G. R. Hood, S. P. Egan, T. H. Q. Powell, D. A.
800 Hahn, et al. 2017. A test of genomic modularity among life-history adaptations promoting
801 speciation with gene flow. *Molecular Ecology* 26:3926–3942.

802 Ragland, G. J., S. P. Egan, J. L. Feder, S. H. Berlocher, and D. A. Hahn. 2011. Developmental
803 trajectories of gene expression reveal candidates for diapause termination: a key life-history
804 transition in the apple maggot fly *Rhagoletis pomonella*. *Journal of Experimental Biology*
805 214:3948–3959.

806 Ragland, G. J., J. Fuller, J. L. Feder, and D. A. Hahn. 2009. Biphasic metabolic rate trajectory of
807 pupal diapause termination and post-diapause development in a tephritid fly. *Journal of Insect*
808 *Physiology* 55:344–350.

809 Roitberg, B. D., R. S. Cairl, and R. J. Prokopy. 1984. Oviposition deterring pheromone
810 influences dispersal distance in tephritid fruit flies. *Entomologia Experimentalis et Applicata*
811 35:217–220.

812 Rundle, H., and P. Nosil. 2005. Ecological speciation. *Ecology Letters* 8:336–352.

813 Ryan, S. F., M. C. Fontaine, J. M. Scriber, M. E. Pfrender, S. T. O’Neil, and J. J. Hellmann.
814 2017. Patterns of divergence across the geographic and genomic landscape of a butterfly hybrid
815 zone associated with a climatic gradient. *Molecular Ecology* 26:4725–4742.

816 Sanz, J. J., J. Potti, J. Moreno, S. Merino, and O. Frias. 2003. Climate change and fitness
817 components of a migratory bird breeding in the Mediterranean region. *Global Change Biology*
818 9:461–472.

819 Savolainen, V., M. Anstett, C. Lexer, I. Hutton, J. Clarkson, M. Norup, M. Powell, et al. 2006.
820 Sympatric speciation in palms on an oceanic island. *Nature* 441:210–213.

821 Schluter, D. 2001. Ecology and the origin of species. *Trends in Ecology & Evolution* 16:372–
822 380.

823 Schluter, D. 2009. Evidence for ecological speciation and its alternative. *Science* 323:737–741.

824 Schmidt, P. S., C.-T. Zhu, J. Das, M. Batavia, L. Yang, and W. F. Eanes. 2008. An amino acid
825 polymorphism in the couch potato gene forms the basis for climatic adaptation in *Drosophila*
826 *melanogaster*. *Proceedings of the National Academy of Sciences* 105:16207–16211.

827 Scriber, J. M., B. Elliot, E. Maher, M. McGuire, and M. Niblack. 2014. Adaptations to “Thermal
828 Time” Constraints in *Papilio*: Latitudinal and local size clines differ in response to regional
829 climate change. *Insects* 5:199–226.

830 Scriber, J. M. 2002. Latitudinal and local geographic mosaics in host plant preferences as shaped
831 by thermal units and voltinism in *Papilio* spp. (Lepidoptera). *European Journal of Entomology*
832 99: 225-239.

833 Scriber, J. M., and G. J. Ordling. 2005. Ecological speciation without host plant specialization;
834 possible origins of a recently described cryptic *Papilio* species. *Entomologia Experimentalis et*
835 *Applicata* 115:247–263.

836 Servedio, M. R., G. S. V. Doorn, M. Kopp, A. M. Frame, and P. Nosil. 2011. Magic traits in
837 speciation: ‘magic’ but not rare? *Trends in Ecology & Evolution* 26:389–397.

838 Sinclair, B. J. 2015. Linking energetics and overwintering in temperate insects. *Journal of*
839 *Thermal Biology*, What sets the limit? How thermal limits, performance and preference in
840 ectotherms are influenced by water or energy balance 54:5–11.

841 Sláma, K., and D. L. Denlinger. 1992. Infradian cycles of oxygen consumption in diapausing
842 pupae of the flesh fly, *Sarcophaga crassipalpis*, monitored by a scanning microrespirographic
843 method. Archives of Insect Biochemistry and Physiology 20:135–143.

844 Spiess, A-N. and N. Neumeyer. 2010. An evaluation of R^2 as an inadequate measure for
845 nonlinear models in pharmacological and biochemical research: a Monte Carlo approach. BMC
846 Pharmacology. 10:6.

847 Stalhansske, S., K. Gotthard, D. Posledovich, and O. Leimar. 2014. Variation in two phases of
848 post-winter development of a butterfly. Journal of Evolutionary Biology. 27:2644-2653.

849 Strong, D. R., J. H. Lawton, and S. R. Southwood. 1984. Insects on plants. Community patterns
850 and mechanisms. Blackwell Scientific Publishing.

851 Tan, Q-Q, Liu, W., Zhu, F., Lei, C-L., Hahn, D.A., and Wang, X-P. 2017. Describing the
852 diapause-preparatory proteome of the beetle *Colaphellus bowringi* and identifying candidates
853 affecting lipid accumulation using isobaric tags for mass spectrometry-based proteome
854 quantification (iTRAQ). Frontiers in Physiology. 8:251

855 Tauber, C. A., M. J. Tauber, and J. R. Nechols. 1977. Two genes control seasonal isolation in
856 sibling species. Science 197:592–593.

857 Tauber, C. A., M. J. Tauber 1981. Insect seasonal cycles: genetics and evolution. Annual Review
858 of Ecology and Systematics 12: 281-308.

859 Tauber, M. J., C. A. Tauber, and S. Masaki. 1986. Seasonal Adaptations of Insects. Oxford
860 University Press.

861 Taylor, R. S., and V. L. Friesen. 2017. The role of allochrony in speciation. Molecular Ecology
862 26:3330–3342.

863 Thackeray, S. J., P. A. Henrys, D. Hemming, J. R. Bell, M. S. Botham, S. Burthe, P. Helaouet, et
864 al. 2016. Phenological sensitivity to climate across taxa and trophic levels. *Nature* 535:241–245.

865 Van Dyck, H., D. Bonte, R. Puls, K. Gotthard, and D. Maes. 2015. The lost generation
866 hypothesis: could climate change drive ectotherms into a developmental trap? *Oikos* 124:54–61.

867 Visser, M. E., and C. Both. 2005. Shifts in phenology due to global climate change: the need for
868 a yardstick. *Proceedings of the Royal Society of London B: Biological Sciences* 272:2561–2569.

869 Vrba, P., M. Dolek, O. Nedvěd, H. Zahradníčková, C. Cerrato, and M. Konvička. 2014.
870 Overwintering of the boreal butterfly *Colias palaeno* in central europe. *Cryoletters* 35:247–254.

871 Wadsworth, C. B., X. Li, and E. B. Dopman. 2015. A recombination suppressor contributes to
872 ecological speciation in *Ostrinia* moths. *Heredity* 114:593–600.

873 Wadsworth, C. B., W. A. Woods, D. A. Hahn, and E. B. Dopman. 2013. One phase of the
874 dormancy developmental pathway is critical for the evolution of insect seasonality. *Journal of*
875 *Evolutionary Biology* 26:2359–2368.

876 Wagner, G. P., M. Pavlicev, and J. M. Cheverud. 2007. The road to modularity. *Nature Reviews*
877 *Genetics* 8:921–931.

878 Wagner, G. P., and J. Zhang. 2011. The pleiotropic structure of the genotype–phenotype map:
879 the evolvability of complex organisms. *Nature Reviews Genetics* 12:204.

880 Walsh, B. D. 1861. On Phytophagic Varieties and Phytophagic Species.

881 Walther, G.-R., E. Post, P. Convey, A. Menzel, C. Parmesan, T. J. C. Beebee, J.-M. Fromentin,
882 et al. 2002. Ecological responses to recent climate change. *Nature* 416:389.

883 Williams, C. M., W. D. Chick, and B. J. Sinclair. 2015. A cross-seasonal perspective on local
884 adaptation: metabolic plasticity mediates responses to winter in a thermal-generalist moth.
885 *Functional Ecology* 29:549–561.

886 Williams, C. M., K. E. Marshall, H. A. MacMillan, J. D. K. Dzurisin, J. J. Hellmann, and B. J.
887 Sinclair. 2012. Thermal variability increases the impact of autumnal warming and drives
888 metabolic depression in an overwintering butterfly. *PLOS ONE* 7:e34470.

889 Winston, P. W., and D. H. Bates. 1960. Saturated solutions for the control of humidity in
890 biological research. *Ecology* 41:232–237.

891 Wipking, W., M. Viebahn, and D. Neumann. 1995a. Oxygen consumption, water, lipid and
892 glycogen content of early and late diapause and non-diapause larvae of the burnet moth *Zygaena*
893 *trifolii*. *Journal of Insect Physiology* 41:47–56.

894 Wolf, J. B., D. Pomp, E. J. Eisen, J. M. Cheverud, and L. J. Leamy. 2006. The contribution of
895 epistatic pleiotropy to the genetic architecture of covariation among polygenic traits in mice.
896 *Evolution & Development* 8:468–476.

897 Wood, T., and S. Guttman. 1982. Ecological and behavioral basis for reproductive isolation in
898 the sympatric *Enchenopa-binotata* complex (*Homoptera: Membracidae*). *Evolution* 36:233–242.

899 Yang, L. H., and V. H. W. Rudolf. 2010. Phenology, ontogeny and the effects of climate change
900 on the timing of species interactions. *Ecology Letters* 13:1–10.

901

Figure Legends

Figure 1. Examples of plotted metabolic trajectories following the A) monophasic, B) biphasic, and C) triphasic models described in the text. To demonstrate how different parameters affect aspects and the shape of metabolic trajectories and reflect different modules of diapause development, the values of the fitted parameters in each model are indicated in the plots with arrows pointing to their relevant position.

Figure 2. Plots showing A) that apple flies (yellow) emerge as adults earlier than hawthorn flies (red), and B) the best-fit non-linear models of the metabolic trajectories for each apple fly (yellow) and hawthorn fly (red) that survived to adult emergence in experiment 1.

Figure 3. Box plots comparing fitted parameter values that show: A) apple flies have a higher baseline metabolic rate (b) than hawthorn flies, B) apple flies have earlier timing of diapause termination (m) than hawthorn flies, C) apple flies have higher post-diapause plateau metabolic rates (p) than hawthorn flies, D) apple and hawthorn flies do not differ in post-diapause plateau duration ($X-m$) nor E) exponential scaling (c). The box plots are followed by scatter plots of the relationships between adult emergence time on the y-axis and F) baseline metabolic rate (b), G) timing of diapause termination (m), H) post-diapause plateau metabolism (p), I) post-diapause plateau duration ($X-m$), and J) exponential scaling (c) with individual apple flies indicated in yellow and hawthorn flies in red, showing that adult emergence timing is strongly correlated with the timing of diapause termination ($r = 0.994$) with much weaker relationships to the other fitted metabolic trajectory parameters.

Figure 4. Partial correlation network showing the major contribution of the timing of diapause termination (m) ($r = 0.994$) along with the minor contribution of post-diapause parameters

describing the plateau metabolic rate (p), the duration of the post-diapause plateau ($X-m$), and scaling of the exponential phase (c) ($r = -0.37$, $r = 0.772$, and $r = 0.533$, respectively) with adult emergence timing as well as additions significant partial correlations among parameters. Edges depict all significant ($P < 0.05$) partial correlations. Edge weights are proportional to the magnitude of Pearson's r , for both positive correlations (solid lines) and negative correlations (dashed lines).

Figure 5. Plots of metabolic rate every 24 hours over 12 days for surviving apple flies (yellow) and hawthorn flies (red) from experiment 2 showing the immediate post-winter metabolic increase over days 1 and 2 exhibited in both host races, with apple fly pupae increasing their metabolic rates substantially higher than hawthorn fly pupae after removal from the cold. After this initial metabolic increase, all hawthorn pupae and most apple pupae enter back into metabolic suppression at baseline post-winter diapause maintenance levels. Time zero on the x-axis represents the overwintering metabolic rate measurement over which flies were purged and measured under continuous simulated winter conditions. See Fig. S4 for a box plot depicting variation in overwintering metabolic rates in more detail.

Figure 6. Partial correlation network showing the relatively weak negative correlation between day 1 metabolic rate and adult emergence timing ($r = -0.318$), as well as additional significant partial correlations among overwintering metabolic rate and post-winter metabolic variables in experiment 2. The three metabolic rate variables over this period are modestly positively correlated, and pupae with particularly strong metabolic increases at day 1 tended to emerge earlier as adults. Edges depict all significant ($P < 0.05$) partial correlations among variables. Line weights are proportional to the magnitude of Pearson's r , for both positive correlations (solid lines) and negative correlations (dashed lines).

Figure S1. Plots of mass adjusted metabolic rates at each time point along with curves fitted for the best-fit non-linear metabolic trajectories for each individual apple fly (orange points) and hawthorn fly (blue points) in experiment 1.

Figure S2. Network of significant ($P < 0.05$) partial correlations between each fitted parameter, representing a different component of diapause development, from the best-fit non-linear models of metabolic trajectories and adult eclosion time for A) all flies, B) hawthorn flies, and C) apple flies from experiment 1, showing that partial correlation relationships among parameters and emergence timing remain largely the same between both host races. Line weights are proportional to the magnitude of Pearson's r , for both positive correlations (solid lines) and negative correlations (dashed lines).

Figure S3. Box plot showing that mean overwintering metabolic rates did not differ between apple flies (yellow) and hawthorn flies (red) from experiment 2. Values come from the measurement of flies over a 24-hour interval during continuous simulating overwintering conditions prior to their removal from the cold. These data are represented as time point zero in Figure 5.

Figure S4. Scatter plots showing A) the significant inverse relationship between ($r = -0.318$) between day 1 metabolic rate and adult emergence time, with higher initial metabolic rate peaks leading to earlier emergence times and B) the non-significant relationship between day 10 metabolic rate and adult emergence timing, with three outlier individuals that have likely exited the diapause maintenance phase by day 10 for each apple fly (yellow) and hawthorn fly (red) in experiment 2.

Figure S5. Networks of significant ($P < 0.05$) partial correlations between overwintering metabolic rate, day 1 metabolic rate (reflective of the initial post-winter metabolic increase), and day 10 metabolic rates (reflective of post-winter diapause baseline metabolic rates) and adult eclosion for A) all flies, B) hawthorn flies, and C) apple flies from experiment 2. Line weights are proportional to the magnitude of Pearson's r , for both positive correlations (solid lines) and negative correlations (dashed lines).

Table S1. AICc values for each of the three hierarchical non-linear models of metabolic trajectories for each fly in experiment 1.

Fly	Host Race	AICc values		
		monophasic	biphasic	triphasic
A1	Apple	-85.42488	-101.16601	-95.87575
A2	Apple	-159.3352	-195.2199	-192.0269
A3	Apple	-90.23488	-93.30185	-86.33917
A4	Apple	-137.7368	-183.5841	-182.2060
A5	Apple	-87.73685	-98.39063	-91.77090
A6	Apple	-100.4272	-133.4496	-128.6462
A7	Apple	-116.0277	-154.3354	-151.7218
A8	Apple	-125.1148	-149.7043	-145.7560
A9	Apple	-96.47984	-126.50280	-120.66406
A10	Apple	-68.01215	-54.19323	-13.90317
A11	Apple	-96.65276	-99.46503	-92.12180
A12	Apple	-120.4935	-134.1151	-129.5280
A13	Apple	-185.0641	-222.9163	-221.6227
A14	Apple	-132.8936	-164.5961	-161.3035
A15	Apple	-117.2497	-146.7242	-142.3276
A16	Apple	-188.1717	-246.0007	-247.0824
A17	Apple	-96.76512	-101.30425	-93.07224
A18	Apple	-175.6545	-221.7109	-218.5474
A19	Apple	-99.23066	-110.27054	-103.61684
A20	Apple	-107.6863	-118.5794	-113.3541

A21	Apple	-126.1644	-145.1380	-141.5845
-----	-------	-----------	-----------	-----------

979 Table S1 continued

Fly	Host Race	AICc values		
		monophasic	biphasic	triphasic
A22	Apple	-217.5604	-291.0552	-288.7704
A23	Apple	-167.1777	-240.5444	-240.7968
A24	Apple	-157.1383	-201.4665	-199.8782
A25	Apple	-104.9526	-136.0942	-132.2242
A26	Apple	-140.7678	-207.0548	-203.6812
A27	Apple	-198.7765	-258.1858	-256.7039
A28	Apple	-156.5090	-213.0256	-210.2791
A29	Apple	-92.64481	-127.87155	-122.72742
A30	Apple	-117.6916	-144.5089	-140.4335
A31	Apple	-214.7300	-275.9143	-275.9728
H1	Hawthorn	-214.3493	-285.7186	-288.1658
H2	Hawthorn	-238.7159	-282.3931	-281.0323
H3	Hawthorn	-180.8670	-233.9812	-231.9984
H4	Hawthorn	-259.3621	-351.8553	-354.2683
H5	Hawthorn	-292.7671	-386.2696	-386.8440
H6	Hawthorn	-395.1055	-476.3699	-499.9463
H7	Hawthorn	-192.0993	-250.6646	-248.1271
H8	Hawthorn	-265.4327	-326.4221	-327.5829
H9	Hawthorn	-240.5597	-293.8475	-297.6247

H10 Hawthorn -326.6583 -408.7803 -410.9211

980

981 Table S1 continued

Fly	Host Race	AICc values		
		monophasic	biphasic	triphasic
H11	Hawthorn	-265.2983	-353.9769	-358.0998
H12	Hawthorn	-244.5326	-308.4879	-310.6853
H13	Hawthorn	-161.4259	-195.7099	-194.0092
H14	Hawthorn	-133.3469	-155.4459	-153.0507
H15	Hawthorn	-254.4221	-343.1827	-354.8459
H16	Hawthorn	-318.0212	-382.7139	-381.1767
H17	Hawthorn	-177.9566	-225.7985	-222.3016
H18	Hawthorn	-376.5118	-504.9609	-513.9705
H19	Hawthorn	-261.0556	-337.7621	-338.1983
H20	Hawthorn	-203.6956	-269.0703	-265.8444
H21	Hawthorn	-315.5219	-386.2596	-387.6650
H22	Hawthorn	-246.0133	-303.8118	-302.6023
H23	Hawthorn	-304.7557	-359.4706	-357.3332
H24	Hawthorn	-287.6338	-397.2828	-399.9917
H25	Hawthorn	-376.0554	-474.7933	-472.3875
H26	Hawthorn	-171.2786	-204.8060	-201.5091
H27	Hawthorn	-301.8461	-350.0046	-350.3595
H28	Hawthorn	-327.8362	-396.3005	-399.3649
H29	Hawthorn	-383.5556	-482.3031	-480.4219

H30	Hawthorn	-259.9859	-315.4500	-314.0465
H31	Hawthorn	-395.3495	-493.7767	-494.3287

982

983 Table S1 continued

		AICc values		
Fly	Host Race	monophasic	biphasic	triphasic
H32	Hawthorn	-233.9627	-311.3159	-315.6969
H33	Hawthorn	-223.9982	-308.7444	-311.5882
H34	Hawthorn	-292.5576	-372.8700	-370.3306
H35	Hawthorn	-217.4432	-267.6473	-265.6989

984

985

Table S2. Results of paired t-tests between parameter coefficients from biphasic and triphasic trajectory models from the same fly.

Parameter	Description	<i>t</i>	p-value	mean of differences
<i>b</i>	Baseline metabolic rate	1.760	0.083	0.00104
<i>p</i>	Plateau metabolic rate	-0.827	0.411	-0.00021
<i>m</i>	Timing of diapause exit	1.152	0.254	0.11743
<i>X</i>	Timing of exponential increase	-0.756	0.453	-0.07246
<i>c</i>	Exponential scaling	-0.639	0.525	-0.00088

Table S3. Summary of repeated-measured ANOVA analysis of metabolic rate over time as a function of host race.

	Df	SS	MS	F-value	p-value
<i>Host</i>	1	1985	1985	21.762	<0.001
<i>Day 1</i>	1	2260	2260.1	24.779	<0.001
<i>Day 9</i>	1	53	52.8	0.578	0.4515
<i>Day 10</i>	1	120	120.2	1.318	0.2579
<i>Day 11</i>	1	17	17.1	0.188	0.667
<i>Host x day 1</i>	1	429	428.9	4.702	<0.05
<i>Host x day 9</i>	1	216	216	2.368	0.1319
<i>Host x day 10</i>	1	564	563.9	2.182	<0.05
<i>Residuals</i>	39	3557	91.2		

A

bioRxiv preprint doi: <https://doi.org/10.1101/2020.01.29.925057>; this version posted January 30, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

B

C

Edge width corresponds to correlation strength:

= $r = 1$

= $r = 0.75$

= $r = 0.5$

Dashed edges represent negative correlations.

Edge width corresponds to correlation strength:

$= r = 1$

$= r = 0.75$

$= r = 0.5$

Dashed edges represent negative correlations

Metabolic Rate ($\mu\text{L CO}_2/\text{hr}/\text{mg}$)

Time (days)

A

B

C

———— = $r = 1$ — — — — = $r = -1$
——— = $r = 0.5$ - - - - = $r = -0.5$

A **B**