1 Detection of hypermucoviscous *Klebsiella pneumoniae* sequence type 86 capsular type
2 K2 in South America as an unexpected cause of a fatal outbreak in captivity marmosets

3

4 Juliana M. Guerra¹, Natália. C.C. de A. Fernandes^{1,2}, Alessandra Loureiro Morales dos
 5 Santos^{1,2}, Joana de Souza Pereira Barrel¹, Bruno Simões Sergio Petri³, Liliane Milanelo³,
 6 Monique Ribeiro Tiba-Casas¹, Alcina Maria Liserre¹, Cláudia Regina Gonçalves¹, Cláudio
 7 Tavares Sacchi¹, José Luiz Catão-Dias², Carlos Henrique Camargo¹.

8

- 9 1. Instituto Adolfo Lutz, São Paulo, Brazil
 10 2. Universidade de São Paulo, São Paulo, Brazil
 11 3. Parque Ecológico do Tietê, São Paulo, Brazil

12

13 Correspondence

14 Carlos Henrique Camargo

15 Instituto Adolfo Lutz - Centro de Bacteriologia - Núcleo de Doenças Entéricas e Infecções
 16 por Patógenos Especiais

17 Avenida Dr. Arnaldo, 351 - 9º Andar

18 Pacaembú - São Paulo, SP – Brasil

19 Phone: +55-11-3068-2896 Fax: +55-11-3068-2898

20 **Email:** carlos.camargo@ial.sp.gov.br

21

22 **First author biographical sketch**

23

24 Dr. Guerra is a veterinary pathologist and scientific researcher in the Center of Pathology of
25 Adolfo Lutz Institute. Her research focuses on the comparative pathology of emerging and
26 reemerging infectious diseases in the context of an integrated One Health approach.

27

28

29 **Running title**

30

31 hvKp ST86 K2 outbreak in marmosets in Brazil

32

33 Abstract

34 After the sudden death of eleven captives marmosets in a rehabilitation center of
 35 wildlife in São Paulo, Brazil, histological and microbiological study was conducted. Liver,
 36 spleen, intestine, central nervous system, lung, thymus, stomach, testicle tissues were
 37 analyzed by light microscopy and microbial cultures were conducted. Environmental cultures
 38 were also performed. Prophylactic antimicrobial therapy, restricted access to marmosets'
 39 cages with dedicated staff, and additional sanitization of animals' fruits were implemented.
 40 Histological findings were compatible with hyperacute septicemia, and microbiological
 41 cultures and molecular tests identified the etiologic agent as hypermucoviscous sequence type
 42 86 capsular type K2 *K. pneumoniae* for the first time in South America. Implementation of
 43 prompt containment measures led to successful control of this outbreak. Detection of a
 44 hypervirulent and zoonotic pathogen, such as hypermucoviscous *K. pneumoniae* ST86 K2, in
 45 an unexpected and human interface reservoir underscores its potential threat in public health
 46 settings.

47

48 **Key-words:** Disease Outbreaks, Primate Diseases, One Health, *Klebsiella pneumoniae*

49

50

1. INTRODUCTION

52

53 *Klebsiella pneumoniae* is a Gram-negative, encapsulated, non-motile, rod-shaped and
 54 opportunistic bacteria present as a normal part of the nasopharynx and gastrointestinal tract
 55 microbiome of humans and animals (1–3). Since the mid-1980s and 1990s, a new
 56 hypervirulent (hypermucoviscous) variant of *K. pneumoniae* (hvKp), initially described in
 57 Southeastern Asia (4–6), has arisen as an important emerging pathogen, affecting young and
 58 healthy individuals from Europe, North, Central and South America, Australia, Africa and the
 59 Middle East (7–13).

60 This variant presents clinical signs and bacterial genotypic and phenotypic features
 61 that allow the differentiation from opportunistic *K. pneumoniae*. hvKp has been frequently
 62 associated with serious clinical complications, including multisystemic pyogenic abscess in
 63 liver, lungs and brain, meningitis, endophthalmitis, osteomyelitis, septicemia and pulmonary
 64 embolism (12–15). Also, hvKp strains usually have a distinguish hypermucoviscosity
 65 phenotype when grown on agar plates, based on a positive string test. The development of
 66 prominent polysaccharide capsules, associated with capsular serotypes K1 or K2, have been
 67 reported as the major virulence determinants for human hvKp in liver abscesses (16–19),
 68 since its seems to protect the bacteria from phagocytosis and preventing destruction by
 69 bactericidal serum factors. In addition, a number of putative virulence genes, such as the
 70 gene *rmpA*, a regulator of the mucoid phenotype gene, located on a plasmid, mucoviscosity-
 71 associated gene *magA*, which encodes a structural outer membrane protein of the K1
 72 serotype, and aerobactin genes have been described (17,20,21).

73 *K. pneumoniae* strains have also been associated with a variety of diseases in animals,
 74 especially in New and Old World primates (22–24). Sudden death or variable clinical signs,
 75 including anorexia, prostration, fever, cough, dyspnea, mucopurulent discharge, meningitis,

76 pneumonia, peritonitis, and sepsis are strongly associated with sporadic infections of
 77 *Klebsiella pneumoniae* in experimental colonies of common marmosets (24,25). A mixed
 78 infection of *Klebsiella* species and *Bordetella bronchiseptica* have also been reported to
 79 cause purulent otitis media in this same species, with frequent extension to the brain, leading
 80 to abscessing encephalitis and meningitis (26). Multiple antibiotic-resistant *Klebsiella* strains
 81 have been isolated from meningitis in *Macaca mulatta* (rhesus macaques) (22). Richard
 82 (1989) described a subcutaneous abscess due to *K. pneumoniae* K5 in a breed of squirrel
 83 monkey (*Saimiri* spp.) at the Pasteur Institute of Cayenne, French Guyana, and a fatal
 84 infection due to *K. pneumoniae* K2 in a colony of lemurs, at Mulhouse zoological garden
 85 (East of France) (27). Positive *K. pneumoniae* bacterial culture related to necrotic and
 86 hemorrhagic enteritis and typhlocolitis were observed in an experimental colony of owl
 87 monkeys (*Aotus nancymai*) (2). More recently, hvKp has been identified in *Cercopithecus*
 88 *aethiops* (Africa green monkeys) causing liver abscesses with bacteremia, meningitis,
 89 endophthalmitis (28–30) and suppurative in a captive gold-handed tamarin (*Saguinus midas*
 90 *midas*) (31). Bilateral mucopurulent nasal discharge, fever, anemia, and swelling of the
 91 inguinal lymph nodes was described in a *Allouatta Clamitans* due to hvKp ST23 serotype K1
 92 (32). However, some species, as cynomolgus and rhesus macaques can act as carriers and
 93 maintain subclinical infection with *K. pneumoniae* (30).

94 Despite the well-recognized zoonotic importance of hvKp and the public health risk
 95 of emerging multidrug-resistant strains (33–35), there is a lack of information regarding the
 96 complete genotyping and phenotyping characterization of the etiologic agent, which is
 97 essential to establish adequate diagnosis and treatment of this pathogen in captive and wild
 98 non-human primates. Thus, the aim of this study was to report an epizootic of common
 99 marmosets in a rehabilitation center of wildlife in Brazil and characterize the serotype,
 100 sequence typing, virulence properties and resistance profile of *K. pneumoniae* strains.

2. MATERIALS AND METHODS

2.1. Study population

All the animals described in this study were maintained in Parque Ecológico do Tietê (23°29'46"S, 46°31'10"O), a center for reception, rehabilitation and referring of wildlife animals, located at São Paulo municipality, São Paulo, Brazil. All callitrichids (*Callithrix jacchus*, *C. penicillata* and hybrids) were housed in a shed of five stainless steel 2 m³ cages and 15 stainless steel 0.4 m³ cages, dedicated only to this species, with a maximum number of eight animals per cage with anti-mosquito screen, feeding platforms, logs and ropes. This center comprises an average daily census of approximately 70 wild-caught callitrichids, that were brought to the facility due to hunting or traffic. These animals have been maintained in captivity for an average period of 140 days for rehabilitation. The cages were cleaned daily by an exclusive employee, except on weekends, when the same worker also cleans the reptilian cages. Standard husbandry procedures included two-times daily feeding program with a commercial food (Megazoo, Minas Gerais, Brazil), diet supplements consisting of a variety of vegetables, fruits, sporadically, *Tenebrio* spp. and crickets, and water *ad libitum*. Routine quarantine procedures consist of physical examination and vermifugation. The animals were also periodically monitored by a veterinarian.

2.2. Animals and sampling

On 10th and 11 February 2019, eleven captive marmosets (eight *C. penicillata*, two *C. jacchus* and one hybrid), without previous clinical signs, suddenly died. They lived in the

same shed and into five different cages with others 72 animals and none of them presented any symptoms. All animals were in captivity from 123 to 399 days. No new animals had been introduced into the cages in the last 25 days.

The necropsy was performed on the same day and macroscopic examination showed no major findings. Tissues samples were preserved in phosphate-buffer formalin 10% and in refrigeration for 12 hours. Since 1999, according to the Non-Human Primates Epizootic Events Surveillance System of Brazilian Ministry of Health (36), which was incorporated into the Yellow Fever Surveillance System. All epizootic event in this population should be investigated in reference laboratories. As so, these samples were sent to Adolfo Lutz Institute for analysis. This study was approved by the Ethics Committee for the Use of Animals (CEUA) of Adolfo Lutz Institute, Brazil (protocol n°11/2016), SISBIO registration no. 50551 for the manipulation of wildlife material and SISGEN registration no. A1A2A72 and A7EB4B6.

2.3 Histological and histochemical examination

For light microscopy small pieces of liver, spleen, intestine, central nervous system, lung, thymus, stomach, and testicle tissue were fixed in neutral 10% phosphate-buffered saline formaldehyde. Fixed tissue specimens were dehydrated by graded ethanol treatment and were routinely embedded in paraffin wax for light microscopy. Embedded sections (3 µm) were stained with haematoxylin and eosin (H&E). Also, additional slides were submitted to histological gram staining. The presence of lesions, intensity, distribution and presence of inflammatory infiltrate were analyzed.

2.4 Isolation and characterization of *Klebsiella* spp.

Liver, brain and pooled tissues were aseptically cultured on sheep blood agar (Oxoid, Basingstoke, UK) and MacConkey agar (Oxoid, Basingstoke, UK) and incubated overnight at 37°C. The plates were examined the day after inoculation for the presence of any growth. Pink-yellow, mucoid, lactose-positive colonies were cultured, and identified as *Klebsiella pneumoniae* Complex based on motility, lysine decarboxylation, citrate utilization and indole non-production. The string test was used to check the hypermucoviscosity of *K. pneumoniae* colonies. The result was defined as positive when an inoculation loop was able to generate a viscous string of 5 mm in length from a colony on a blood agar plate after overnight incubation (8) while a negative sting test was assigned when the string was less than 5 mm in length or no string was present.

2.5 Environmental testing

In order to identify the source of *K. pneumoniae* strains, environmental samples of water from the lake near to the primate cages and drag swabs from their cages were screened for the presence of *K. pneumoniae*. Water samples (100mL) were filtered with 0.22 µm sterile membranes (Jet Biofil, China) and placed onto the surface of a MacConkey Agar plate. Typical colonies (mucoid, lactose-fermenting) were identified by biochemical tests (see above). Drag swabs were streaked onto MacConkey Agar plates and typical *K. pneumoniae* colonies were subjected to the same tests as employed for the water isolates.

2.6 Antimicrobial susceptibility testing

A representative isolate (P04) was subjected to antimicrobial susceptibility testing by using the broth microdilution methodology with Sensititre (Thermo, Waltham, USA) plate,

according to the manufacturer's instructions. Minimal inhibitory concentration values were categorized as susceptible, intermediate or resistance following Clinical and Laboratory Standards Institute (CLSI, Wayne, USA) M100-S29 breakpoints (available at <http://em100.edaptivedocs.net/dashboard.aspx>).

2.7 Pulsed-field gel electrophoresis (PFGE)

All isolates identified as *K. pneumoniae* were submitted to DNA macro-restriction by using 30U of XbaI enzyme followed by Pulsed-field gel electrophoresis in a CHEF-III apparatus, according to the PulseNet International guidelines (<https://www.cdc.gov/pulsenet/pathogens/pfge.html>). The Universal Size Standard Strain H9812 (*Salmonella* Braenderup) (37) was used as reference in all gels. Gel images were analyzed in the BioNumerics v.7.6.2 (Applied Maths, Sint-Martens-Latem, Belgium) with the generation of dendrogram based on unweighted pair group method with arithmetic mean (UPGMA) methods and similarities calculated by Dice coefficient with tolerance and optimization both set at 1.5%.

2.8 DNA extraction and whole genome sequencing

The representative strain P04 was selected based on the PFGE results, and subjected to whole genome sequencing. High quality whole genome was extracted from a pure culture of 16h growth in Luria-Bertani broth (Oxoid, Basingstoke, UK) by using the Wizard Genomic DNA Purification Kit (Promega, Madison, USA) according to the instructions of the manufacturer. DNA was tagged by using the Ion Xpress™ Plus Fragment Library Kit and the libraries constructed were sequenced in an Ion Torrent S5 platform. Reads were de

novo assembled with the Spades algorithm (v.5.12.0.0) and submitted to online platforms. Definitive species identification was carried out by Ribosomal Multilocus Sequence Typing (rMLST) (<https://pubmlst.org/rmlst/>). Antimicrobial resistance genes were sought at the Resfinder tool of the Center for Genomic Epidemiology website (<http://www.genomicepidemiology.org/>). Sequence Type, virulence genes and capsular type data were extracted from the Institute Pasteur MLST and whole genome MLST databases (<http://bigsdbs.pasteur.fr/>). Based on the reference genome CG43 (isolate 1530 on Institute Pasteur database available at https://bigsdbs.pasteur.fr/cgi-bin/bigsdbs/bigsdbs.pl?db=pubmlst_klebsiella_isolates&page=plugin&name=Contigs&format=text&isolate_id=1530&match=1&pc_untagged=0&min_length=&header=1), high quality single-nucleotide polymorphisms (SNPs) were extracted from the P04 genome and from the available ST86 *K. pneumoniae* genomes from the Institute Pasteur MLST and whole genome MLST databases. A phylogenetic tree based on the concatenated alignments was built on the CSI Phylogeny 1.4 (<https://cge.cbs.dtu.dk/services/CSIPhylogeny/>). The genome of the hv ST374 *K. pneumoniae* was included as outgroup. The Newick generated file was uploaded on the Microreact platform along with metadata of such isolates (project V8uV0aPEU) for the generation of a tree with the SNPs and metadata. The whole-genome shotgun project reported here has been deposited at DDBJ/EMBL/GenBank under the accession number SPSP000000000.

RESULTS

2.1 Histological and histochemical examination

Multiple sections of different tissues were microscopically examined and similarly affected. The major microscopic finding consistent to sepsis it was intravascular presence of bacilli in liver (10/10), cerebrum (8/10), lung (3/10), heart (1/7), intestine (1/7), thymus (1/2), and skeletal muscle (1/1). Others relevant microscopic findings of different tissues are summarized on the Table 1. Furthermore, intestine it was the unique organ that it was observed bacteria into the lumen of lymphatic vessels (1/7). Other nonspecific findings consists to tubular proteinosis (4/8) and tubular acute necrosis (1/8) in kidney, hepatic steatosis (1/10), lymph histiocytic admixed or not with neutrophils enteritis (2/7), Peyer plaques hyperplasia (1/7), cardiac congestion (1/7), focus of edema of the cardiomyocytes and necrotizing and lymphohistiocytic myocarditis (1/7). Stomach, tongue, testis, thymus, skin, uterus, lymph-node didn't show microscopic alterations on all the fragments analyzed. Some tissues couldn't be analyzed due to the several autolysis.

2.2 Isolation and characterization of *Klebsiella* spp.

Microbial cultures performed from brain and liver from eight different animals recovered *K. pneumoniae* in all of them, as pure growth, and with hypermucoviscous phenotype, i.e., positive string test. The isolate P04 presented susceptibility to all the antimicrobial agents evaluated (Table 2).

2.3 Environmental testing

K. pneumoniae isolates were recovered from both water and drag swab samples.

2.4 Pulsed-field gel electrophoresis (PFGE)

Molecular typing of *K. pneumoniae* revealed the same XbaI-PFGE profile among the isolates recovered from the dead animals (Figure 2, cluster A, pulsotypes A1 and A2), however, environmental isolates clustered apart from the invasive isolates (Figure 2, clusters B and C).

2.5 Whole genome sequencing

Whole genome sequencing resulted in 1,049,163 reads and the *de novo* assembled genome comprised 5,358,608 bps grouped in 62 contigs, with an average coverage depth of 53. rMLST and kmer tools confirmed the presence of *K. pneumoniae* as the species of isolate P04. Acquired antimicrobial resistant genes were not detected, but the constitutive *bla*SHV-1, *oqxAB* and *fosA* were. *In silico* multilocus sequence typing analysis identified the isolate as the sequence type ST86, with the following combination alleles: *gapA*, 9; *infB*, 4; *mdh*, 2; *pgi*, 1; *phoE*, 1; *rpoB*, 1; *tonB*, 27. The capsular type K2 was identified. According to the virulence database of BIGSdb *K. pneumoniae*, P04 presented the *mrkABCDHFIIJ* cluster (mannose-resistant *Klebsiella*-like type III fimbriae cluster, associated with adhesiveness and fimbrial filament formation to adherence to eukaryotic cells) (38) in the same contig of the *kvgAS* genes. The *iroBCD* genes (salmochelin) were also detected, in the same contig that the *rmpA* gene (regulator of mucoid phenotype A). The *rmpA2* gene was detected within the contig along the *iucABD* with the *iutA* genes (aerobactin). Phylogenetic analysis of high quality SNPs showed that ST86 isolates were closely related, and the P04 strain clustered close to IPEUC340, an isolate recovered in 1975 in France (Figure 3.A). The isolate RJF293, ST374, clustered apart from the ST86 isolates (Figure 3.B).

DISCUSSION

In this study, eleven marmosets suddenly died after habitual management of the captives in one of the largest centers for reception, rehabilitation and referring of wildlife animals in the municipality of São Paulo, the largest city in South America. Histological findings were compatible with a hyperacute septicemia, microbiological cultures and molecular tests identified the etiologic agent as hypermucoviscous, ST86, K2 *K. pneumoniae*. This bacteria represents one of the priority organisms to be controlled according to the World Health Organization, particularly the carbapenem- and polymyxin-resistant strains (39). The emergence of these strains is a concern in human and veterinary medicine due to the potential of these strains to acquire multidrug resistance genes, their capacity to persist in the environment and to infect a wide-range of hosts and the unavailability of vaccines against these strains for humans and animals (40).

The role of *K. pneumoniae* in wildlife has been revisited, especially in callitrichidae species, which are highly susceptible to fatal infections caused by this bacteria in captivity (24,40). The bacteria likely spreads by close contact and aerosol between animals and through fomites between cages (41). Frequently described in association with immunosuppressive conditions in New world monkeys, *Klebsiella* is considered an opportunist agent, leading to pneumonia, peritonitis, cystitis, meningitis, and septicemia (2). The propensity of common marmosets to bacterial infection could be related to the low variability in the MHC class II loci that may compromise anti-bacterial immune response and also explain the high frequency of septicemia and endotoxemia after deep wounds and fighters in this genus (41,42).

We identified that the strains recovered from marmosets that suddenly died were clonally related by PFGE analysis. Since PFGE is known by its high discriminatory power,

we can suppose that a common source was present. In order to investigate the source of such strain, environmental samples (water and drag swabs) were cultured, and despite the growth of *K. pneumoniae*, molecular typing revealed that environmental strains were not related to the clinical isolates. Despite our efforts, we were not able to detect the source of the hypermucoviscous *K. pneumoniae* in this outbreak during our investigation. Other studies also faced this same limitation (2) and suggested that dietary supplements could be the most likely source responsible for the introduction of *K. pneumoniae* in the colonies. Vegetables and fruits, especially bananas and their peels, seems to carry a widely diverse Enterobacteriales microbial community (23). Even we have not found the source of the virulent *K. pneumoniae* isolate causing this outbreak, drag swabs of the living animals were negative for such strain, indicating that the remaining population of marmosets was not carriers of this hvKp isolate. In fact, no other animals died after this event. We can admit that the end of this outbreak can be attributed to implementation of effective measures to contain the dissemination of the virulent *K. pneumoniae* strain. The use of Bactrim as prophylaxis during the first 5 days after the identification of deaths may be one of the successful measures implemented, since the isolate was susceptible to this drug. In fact, the isolate did not present any resistance marker to the evaluated antimicrobial agents (Table 2). In addition, accession to the cages was restricted to a dedicated employee, who wore dedicated clothes during the permanence in the marmosets' shed. Additional sanitization step of fruits with 2% sodium hypochlorite for 15 minutes and dedicated space and staff for preparation of marmosets' meals were also implemented after the epizootic event.

Whole genome sequencing of one representative isolate identified several virulence genes that can be associated with its hypermucoviscous phenotype. Among the several virulence genes detected, the ones associated with invasiveness and with capsule production could contribute to the evasiveness of immune system resulting in the severe and fulminant

clinical conditions observed. Indeed, four out of the five suggested virulence genes recognized to be associated with the hv phenotype were detected in the P04 genome, namely *rmpA*, *rmpA2* (regulators of mucoid phenotype A), *iuc* (aerobactin), and *iroB* (salmochelin) (43).

The hypermucoviscous *K. pneumoniae* P04 strain was identified as sequence type ST86 and capsular typing K2. Previous reports also identified this hypervirulent clone with a wild-type susceptibility profile causing severe infections in both hospital (44) and community-acquired infections in humans (45), highlighting the role of hypermucoviscous *K. pneumoniae* as etiological of human infections. Intriguingly, infections due hvKp ST86 K2 are apparently more severe than the infections due to other hvKp clones (46). As of October 2019, 26 isolates belonging to ST86 were available in the PasteurMLST database (<https://bigsd.b.pasteur.fr/>). All of them isolated from humans, and predominantly from Europe (12/26) and Asia (10/26) (10,47–49). Only two isolates were previously identified in America, one recovered from the blood of a patient with liver abscess in Buffalo, USA, in 2013 (50) and another one recovered from the cerebrospinal fluid of a patient with meningitis in Pointe-a-Pitre, Guadeloupe, in 2014 (51). Therefore, to the best of our knowledge, this is the first description of hvKp ST86 K2 *K. pneumoniae* from South America.

Our phylogenetic analysis demonstrated that ST86 genomes are very closely related, but the origin of the P04 isolate could not be certainly inferred, since it clustered with the RFJ293 isolate, recovered in 1975 from France. The expansion of the emerging pathogen hypermucoviscous *K. pneumoniae* ST86 K2 among different reservoirs should be carefully surveilled, since relates of hypervirulent and multi-drug resistant strains are raising (52).

In summary, we report the detection of hypermucoviscous *K. pneumoniae* ST86 K2 for the first time in South America, responsible for a sudden and fatal outbreak in captive marmosets. Implementation of prompt containment measures led to successful control of this

outbreak. The burden of hypermucoviscous *K. pneumoniae* ST86 K2 in unexpected
reservoirs, including the ones in human interface, deserves further investigation.

ACKNOWLEDGEMENTS

We thank the team of curators of the Institut Pasteur MLST and whole genome MLST
databases for curating the data and making them publicly available at <http://bigsd.b.pasteur.fr/>
and all contributors of Center of Pathology for routine sample processing.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

Tables headings

Table 1. Histologic findings described about the fragments submitted for microscopic evaluation.

Table 2. Antimicrobial susceptibility profile determined by Sensititre (Thermo Scientific) of the epizootic strain hypermucoviscous *K. pneumoniae* ST86 strain P04.

Figure subtitles

Figure 1. Microscopic findings of histological and histochemical examination prepared for light microscopy. A – Spleen. Necro hemorrhagic splenitis in multiple germ centers (H&E 4x). B – Brain (Meningis). Bacterial rods inside vascular lumen (arrows) (H&E 40x). C – Lung. Several interstitial pneumonia (H&E 4x) and alveolar hemorrhage (inset – H&E10x). D – Intestine. Lymphoplasmacytic enteritis (arrowheads) and intravascular bacterias (arrow) (H&E 20x and 40x). E – Liver. Necrosis (arrowheads) associated with the presence of numerous bacterial rods (arrow). Hepatocytes also show steatosis and the sinusoids are filled by neutrophils (H&E 40x). The inset image shows discrete mononuclear portal hepatitis and presence of numerous bacterial rods (H&E 20x). F – Liver. The sinusoids are filled by gram-negative bacterial structures (heads) and neutrophils (Gram 40x).

Figure 2. Dendrogram and PFGE typing of XbaI restricted *K. pneumoniae* strains isolated from infection (blue, cluster A), drag swab (green, cluster B) and water (red, cluster C). PFGE profiles (represented by capital letters) were defined based on 100% Dice similarity cutoff

value of the UPGMA clustering method (1.5% optimization; 1.5% tolerance). NA: not applicable.

Figure 3. Schematic representations of high quality SNPs trees built from publicly available ST86 *K. pneumoniae* genomes. In (A) it is possible to note that the P04 isolate described in this study clustered close to the IPEUC-340 isolate, recovered in 1975 in France. In (B) we observe the high similarity of ST86 *K. pneumoniae* genomes in comparison with the outgroup strain RJF293, an ST374 hypermucoviscous *K. pneumoniae* recovered from invasive infection. Green circles represent the ST86 isolates, and the red one represents the ST374.

REFERENCES

1. Davis GS, Price LB. Recent Research Examining Links Among *Klebsiella pneumoniae* from Food, Food Animals, and Human Extraintestinal Infections. *Curr Environ Heal Reports*. 2016 Jun;3(2):128–35.
2. Gozalo AS, Elkins WR, Lambert LE, Stock F, Thomas ML, Woodward RA. Genetic diversity of *Klebsiella pneumoniae* isolates during an outbreak in a non-human primate research colony. *J Med Primatol*. 2016 Dec;45(6):312–7.
3. Simmons J GS. Bacterial and mycotic diseases of nonhuman primates. In: Bennett BT, Abee CR HR, editor. *Nonhuman Primates in Biomedical Research*. San Diego: Academic Press; 2012. p. 105–72.
4. Liu YC, Cheng DL, Lin CL. *Klebsiella pneumoniae* liver abscess associated with septic endophthalmitis. *Arch Intern Med*. 1986 Oct;146(10):1913–6.
5. cheng D-L. Septic Metastatic Lesions of Pyogenic Liver Abscess. *Arch Intern Med*. 1991 Aug;151(8):1557.
6. Wang J, Liu Y, Lee SS, Yen M, Wang, Yao-Shen Chen J, Wann S, et al. Primary Liver Abscess Due to *Klebsiella pneumoniae* in Taiwan. *Clin Infect Dis*. 1998 Jun;26(6):1434–8.
7. Fang FC, Sandler N, Libby SJ. Liver Abscess Caused by magA+ *Klebsiella pneumoniae* in North America. *J Clin Microbiol*. 2005 Feb;43(2):991–2.
8. Siu LK, Yeh K-M, Lin J-C, Fung C-P, Chang F-Y. *Klebsiella pneumoniae* liver abscess: a new invasive syndrome. *Lancet Infect Dis*. 2012 Nov;12(11):881–7.
9. Shon AS, Bajwa RPS, Russo TA. Hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*. *Virulence*. 2013 Feb;4(2):107–18.
10. Bialek-Davenet S, Criscuolo A, Ailloud F, Passet V, Jones L, Delannoy-Vieillard A-S,

- et al. Genomic Definition of Hypervirulent and Multidrug-Resistant *Klebsiella pneumoniae* Clonal Groups. *Emerg Infect Dis*. 2014 Nov;20(11):1812–20.
11. Yang Z, Liu W, Cui Q, Niu W, Li H, Zhao X, et al. Prevalence and detection of *Stenotrophomonas maltophilia* carrying metallo- β -lactamase blaL1 in Beijing, China. *Front Microbiol*. 2014 Dec;5.
12. Struve C, Roe CC, Stegger M, Stahlhut SG, Hansen DS, Engelthaler DM, et al. Mapping the Evolution of Hypervirulent *Klebsiella pneumoniae*. Segre J, Relman DA, editors. *MBio*. 2015 Jul;6(4).
13. Prokesh BC, TeKippe M, Kim J, Raj P, TeKippe EM, Greenberg DE. Primary osteomyelitis caused by hypervirulent *Klebsiella pneumoniae*. *Lancet Infect Dis*. 2016 Sep;16(9):e190–5.
14. Patel PK, Russo TA, Karchmer AW. Hypervirulent *Klebsiella pneumoniae*. *Open Forum Infect Dis*. 2014 Jun;1(1):ofu028–ofu028.
15. Wang Q, Li B, Tsang AKL, Yi Y, Woo PCY, Liu CH. Genotypic Analysis of *Klebsiella pneumoniae* Isolates in a Beijing Hospital Reveals High Genetic Diversity and Clonal Population Structure of Drug-Resistant Isolates. Manganelli R, editor. *PLoS One*. 2013 Feb;8(2):e57091.
16. Chuang Y, Fang C, Lai S, Chang S, Wang J. Genetic Determinants of Capsular Serotype K1 of *Klebsiella pneumoniae* Causing Primary Pyogenic Liver Abscess. *J Infect Dis*. 2006 Mar;193(5):645–54.
17. Fang C-T, Chuang Y-P, Shun C-T, Chang S-C, Wang J-T. A Novel Virulence Gene in *Klebsiella pneumoniae* Strains Causing Primary Liver Abscess and Septic Metastatic Complications. *J Exp Med*. 2004 Mar;199(5):697–705.
18. Yeh K, Chang F, Fung C, Lin J, Siu LK. Serotype K1 Capsule, Rather than magA Per Se, Is Really the Virulence Factor in *Klebsiella pneumoniae* Strains That Cause

- 445 Primary Pyogenic Liver Abscess. J Infect Dis. 2006 Aug;194(3):403–4.
- 446 19. Yu VL, Hansen DS, Ko WC, Sagnimeni A, Klugman KP, von Gottberg A, et al.
- 447 Virulence Characteristics of Klebsiella and Clinical Manifestations of K. pneumoniae
- 448 Bloodstream Infections. Emerg Infect Dis. 2007 Jul;13(7):986–93.
- 449 20. Yu W-L, Ko W-C, Cheng K-C, Lee H-C, Ke D-S, Lee C-C, et al. Association between
- 450 rmpA and magA Genes and Clinical Syndromes Caused by Klebsiella pneumoniae in
- 451 Taiwan. Clin Infect Dis. 2006 May;42(10):1351–8.
- 452 21. Russo TA, Olson R, MacDonald U, Metzger D, Maltese LM, Drake EJ, et al.
- 453 Aerobactin Mediates Virulence and Accounts for Increased Siderophore Production
- 454 under Iron-Limiting Conditions by Hypervirulent (Hypermucoviscous) Klebsiella
- 455 pneumoniae. Infect Immun. 2014 Jun;82(6):2356–67.
- 456 22. Fox JG, Rohovsky MW. Meningitis caused by Klebsiella spp in two rhesus monkeys. J
- 457 Am Vet Med Assoc. 1975 Oct;167(7):634–6.
- 458 23. Gonzalo A ME. Klebsiella pneumoniae infection in a New World nonhuman primate
- 459 center. Lab Primate Newsl. 1991;30:13–20.
- 460 24. Pisharath HR, Cooper TK, Brice AK, Cianciolo RE, Pistorio AL, Wachtman LM, et al.
- 461 Septicemia and peritonitis in a colony of common marmosets (Callithrix jacchus)
- 462 secondary to Klebsiella pneumoniae infection. Contemp Top Lab Anim Sci. 2005
- 463 Jan;44(1):35–7.
- 464 25. Kindlovits A, Kindlovits L. Clínica e Terapêutica em Primatas Neotropicais. 2nd ed.
- 465 Rio de Janeiro: L.F. Livros; 2009. 535 p.
- 466 26. Chalmers DT, Murgatroyd LB, Wadsworth PF. A survey of the pathology of
- 467 marmosets (Callithrix jacchus) derived from a marmoset breeding unit. Lab Anim.
- 468 1983 Oct;17(4):270–9.
- 469 27. Richard C. [Epidemiology of Klebsiella pneumoniae infections in 2 colonies of

- 470 squirrel monkeys and lemurs]. Bull Soc Pathol Exot Filiales. 1989;82(4):458–64.
- 471 28. Soto E, LaMon V, Griffin M, Keirstead N, Beierschmitt A, Palmour R. Phenotypic and
472 genotypic characterization of *Klebsiella pneumoniae* isolates recovered from
473 nonhuman primates. J Wildl Dis. 2012 Jul;48(3):603–11.
- 474 29. Twenhafel NA, Whitehouse CA, Stevens EL, Hottel HE, Foster CD, Gamble S, et al.
475 Multisystemic Abscesses in African Green Monkeys (*Chlorocebus aethiops*) with
476 Invasive *Klebsiella pneumoniae* —Identification of the Hypermucoviscosity
477 Phenotype. Vet Pathol. 2008 Mar;45(2):226–31.
- 478 30. Burke RL, Whitehouse CA, Taylor JK, Selby EB. Epidemiology of invasive *Klebsiella*
479 *pneumoniae* with hypermucoviscosity phenotype in a research colony of nonhuman
480 primates. Comp Med. 2009 Dec;59(6):589–97.
- 481 31. Guerra MFL, Teixeira RHF, Ribeiro VL, Cunha MPV, Oliveira MGX, Davies YM, et
482 al. Suppurative peritonitis by *Klebsiella pneumoniae* in captive gold-handed tamarin (*Saguinus midas midas*). J Med Primatol. 2016 Feb;45(1):42–6.
- 483
484 32. Anzai EK, de Souza Júnior JC, Peruchi AR, Fonseca JM, Gumpl EK, Pignatari ACC,
485 et al. First case report of non-human primates (*Alouatta clamitans*) with the
486 hypervirulent *Klebsiella pneumoniae* serotype K1 strain ST 23: A possible emerging
487 wildlife pathogen. J Med Primatol. 2017 Dec;46(6):337–42.
- 488 33. Wu F, Ying Y, Yin M, Jiang Y, Wu C, Qian C, et al. Molecular Characterization of a
489 Multidrug-Resistant *Klebsiella pneumoniae* Strain R46 Isolated from a Rabbit. Int J
490 Genomics. 2019 Aug;2019:1–12.
- 491 34. Osman KM, Hassan HM, Orabi A, Abdelhafez AST. Phenotypic, antimicrobial
492 susceptibility profile and virulence factors of *Klebsiella pneumoniae* isolated from
493 buffalo and cow mastitic milk. Pathog Glob Health. 2014 Jun;108(4):191–9.
- 494 35. Boszczowski I, Salomão MC, Moura ML, Freire MP, Guimarães T, Cury AP, et al.

- 495 Multidrug-resistant *Klebsiella pneumoniae*: genetic diversity, mechanisms of
496 resistance to polymyxins and clinical outcomes in a tertiary teaching hospital in Brazil.
497 Rev Inst Med Trop Sao Paulo. 2019;61.
- 498 36. Brasil Ministério da Saúde, Secretaria de Vigilância em Saúde, Doenças D de V das,
499 Transmissíveis. Guia de vigilância de epizootias em primatas não humanos e
500 entomologia aplicada à vigilância da febre amarela / Ministério da Saúde, Secretaria de
501 Vigilância em Saúde, Departamento de Vigilância das Doenças Transmissíveis. 2nd
502 ed. Ministério da Saúde, editor. Brasília; 2014. 100 p.
- 503 37. Hunter SB, Vauterin P, Lambert-Fair MA, Van Duyne MS, Kubota K, Graves L, et al.
504 Establishment of a universal size standard strain for use with the PulseNet standardized
505 pulsed-field gel electrophoresis protocols: converting the national databases to the new
506 size standard. J Clin Microbiol [Internet]. 2005;43(3):1045–50. Available from:
507 <http://www.ncbi.nlm.nih.gov/pubmed/15750058>
- 508 38. Gerlach GF, Clegg S, Allen BL. Identification and characterization of the genes
509 encoding the type 3 and type 1 fimbrial adhesins of *Klebsiella pneumoniae*. J
510 Bacteriol. 1989;171(3):1262–70.
- 511 39. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al.
512 Discovery, research, and development of new antibiotics: the WHO priority list of
513 antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis. 2018 Mar;18(3):318–
514 27.
- 515 40. Cox BL, Schiffer H, Dagget G, Beierschmitt A, Sithole F, Lee E, et al. Resistance of
516 *Klebsiella pneumoniae* to the innate immune system of African green monkeys. Vet
517 Microbiol. 2015 Mar;176(1–2):134–42.
- 518 41. Ludlage E, Mansfield K. Clinical care and diseases of the common marmoset
519 (*Callithrix jacchus*). Comp Med. 2003 Aug;53(4):369–82.

- 520 42. Antunes SG, de Groot NG, Brok H, Doxiadis G, Menezes AAL, Otting N, et al. The
521 common marmoset: A new world primate species with limited Mhc class II variability.
522 Proc Natl Acad Sci. 1998 Sep;95(20):11745–50.
- 523 43. Russo TA, Olson R, Fang C-T, Stoesser N, Miller M, MacDonald U, et al.
524 Identification of Biomarkers for Differentiation of Hypervirulent Klebsiella
525 pneumoniae from Classical K. pneumoniae. Diekema DJ, editor. J Clin Microbiol.
526 2018 Jun;56(9).
- 527 44. Cubero M, Grau I, Tubau F, Pallarés R, Dominguez MA, Liñares J, et al.
528 Hypervirulent Klebsiella pneumoniae clones causing bacteraemia in adults in a
529 teaching hospital in Barcelona, Spain (2007–2013). Clin Microbiol Infect. 2016
530 Feb;22(2):154–60.
- 531 45. Liu Y, Long D, Xiang T-X, Du F-L, Wei DD, Wan L-G, et al. Whole genome
532 assembly and functional portrait of hypervirulent extensively drug-resistant NDM-1
533 and KPC-2 co-producing Klebsiella pneumoniae of capsular serotype K2 and ST86. J
534 Antimicrob Chemother. 2019 May;74(5):1233–40.
- 535 46. Decre D, Verdet C, Emirian A, Le Gourrierec T, Petit J-C, Offenstadt G, et al.
536 Emerging Severe and Fatal Infections Due to Klebsiella pneumoniae in Two
537 University Hospitals in France. J Clin Microbiol. 2011 Aug;49(8):3012–4.
- 538 47. BIALEK-DAVENET S, NICOLAS-CHANOINE M-H, DECRÉ D, BRISSE S.
539 Microbiological and clinical characteristics of bacteraemia caused by the
540 hypermucoviscosity phenotype of Klebsiella pneumoniae in Korea. Epidemiol Infect.
541 2013 Jan;141(1):188–188.
- 542 48. Bialek-Davenet S, Criscuolo A, Ailloud F, Passet V, Nicolas-Chanoine M-H, Decre D,
543 et al. Development of a multiplex PCR assay for identification of Klebsiella
544 pneumoniae hypervirulent clones of capsular serotype K2. J Med Microbiol. 2014

545 Dec;63(Pt_12):1608–14.

546 49. Lin J-C, Koh T, Lee N, Fung C-P, Chang F-Y, Tsai Y-K, et al. Genotypes and
547 virulence in serotype K2 *Klebsiella pneumoniae* from liver abscess and non-infectious
548 carriers in Hong Kong, Singapore and Taiwan. *Gut Pathog.* 2014;6(1):21.

549 50. Russo TA, Gill SR. Draft Genome Sequence of the Hypervirulent *Klebsiella*
550 *pneumoniae* Strain hvKP1, Isolated in Buffalo, New York. *Genome Announc.* 2013
551 Mar;1(2):e0006513.

552 51. Melot B, Brisse S, Breurec S, Passet V, Malpote E, Lamaury I, et al. Community-
553 acquired meningitis caused by a CG86 hypervirulent *Klebsiella pneumoniae* strain:
554 first case report in the Caribbean. *BMC Infect Dis.* 2016;16(1):736.

555 52. Harada S, Doi Y. Hypervirulent *Klebsiella pneumoniae*□: a Call for Consensus
556 Definition and International Collaboration. Diekema DJ, editor. *J Clin Microbiol.* 2018
557 Jun;56(9).

PFGE_Xbal

PFGE_Xbal

kb

60 70 80 90 100

2000 1000 600.00 500.00 400.00 250.00 200.00 150.00 100.00 40.00 20.00

Key

Animal Number

Source

Source_

Outcome

P04

1

Infection

Liver

Death

P08

3

Infection

Liver

Death

P09

4

Infection

Brain

Death

P10

4

Infection

Liver

Death

P11

5

Infection

Brain

Death

P12

5

Infection

Liver

Death

P13

6

Infection

Brain

Death

P14

6

Infection

Liver

Death

P15

8

Infection

Blood

Death

P16

8

Infection

Liver

Death

P17

8

Infection

Intestine

Death

P19

8

Infection

Brain

Death

P06

2

Infection

Liver

Death

B1

NA

Environment

Drag swab

NA

C1

NA

Environment

Drag swab

NA

RC5A1

NA

Environment

Water

NA

RC5B2

NA

Environment

Water

NA

RC5B1

NA

Environment

Water

NA

55.4

74.3

97.3

90.9