MiRNAs profiling and degradome sequencing between the CMS-line N816S and its maintainer line Ning5m during anther development in pepper (*Capsicum annuum* L.)

Hongyuan Zhang, Shuping Huang, Jie Tan, Xia Chen, Min Zhang*

Wuhan Academy of Agricultural Sciences, Wuhan, 430070, Hubei province, China.

*Corresponding author: Min Zhang, email address: zhangmin818@aliyun.com

Abstract

Utilization of cytoplasmic male sterility (CMS) is significant for agriculture. MiRNAs are a class of endogenously non-coding small RNAs (21-24 nt) that play key roles in the regulation of various growth and developmental processes in plants. The knowledge miRNA-guided CMS regulation is rather limited in pepper. To better understand the miRNAs involvement and regulatory mechanism of CMS, miRNA libraries from anther of CMS-line N816S and its maintainer line Ning5m were generated by miRNAome sequencing in pepper. A total of 76 differentially expressed miRNAs were detected, of which 18 miRNAs were further confirmed by quantitative real-time PCR (qRT-PCR). In addition, miRNA targets were identified by degradome sequencing. The result showed that 1292 targets that were potentially cleaved by 321 miRNAs (250 conserved miRNAs and 71 novel miRNAs). Gene Ontology (GO) and KEGG pathway analysis indicated that 35 differentially expressed miRNAs might play roles in the regulation of CMS sterility, by cleaving 77 target transcripts, such as *MYBs*, *SPLs*, and *AFRs*, of which targeted by miR156, miR167, miRNA858 family. Nineteen miRNA-cleaved targets were selectively examined by qRT-PCR, and the results showed that there were mostly negative correlations between miRNAs and their targets on the expression level. These findings provide a valuable information to understand miRNAs mechanism during anther development and CMS occurrence in pepper.

Keywords: CMS; miRNAome; degradome sequencing; miRNA regulation; Pepper

Introduction

MicroRNAs (miRNAs) are a class of endogenous non-coding small RNAs (sRNAs) with 21-24 nucleotides (nt) in animals and plants(Zhou et al., 2020). In plants, mature miRNA sequences were generated from primary miRNA transcripts (pri-miRNA) by Dicer-like (DCL1)(Zhang et al., 2015). The mature miRNAs are

incorporated into the RNA-induced silencing complex (RISC) to guiding the cleavage of specific complementary mRNAs(Schuck et al., 2013). MiRNAs are widely considered as negative-regulated gene expression at the post-transcription or translation process via degrading target mRNAs or repressing mRNA translation(Stepien et al., 2017). Previous studies have manifested that miRNAs involved in various plant growth and developmental processes, such as hormone homeostasis, flower development, embryogenesis and stress responses(Jha and Shankar, 2011;Pei et al., 2013).

Recently, a large number of studies have revealed miRNAs as important regulators of gene expressions to be involved in the developmental transition from vegetative growth to reproductive growth(Lin et al., 2013). MiR156 and miR172, two prime participators in flowering regulation, have been shown to regulate the floral transition in plant(Zhu and Helliwell, 2011;Yu et al., 2012;Diaz-Manzano et al., 2018). MiR156 could target SQUAMOSA Promoter Binding Protein-like (SPL) Transcription Factors (TF) and decreases with increasing life spans of the plant(Zheng et al., 2019). Overexpression of miR156 could delay flowering and prolong vegetative stage in many species, including Arabidopsis, rice and tomato(Zheng et al., 2019). The miR172, which targets APETALA2-like (AP2) transcription factors, has the opposite effect to the miR156 on the regulation of flowering time and increases with phase development and induces floral organogenesis and promote flowering(Tripathi et al., 2018). In addition, miR397 has been reported promoting panicle branching, increasing grain size, and resulting in improving yield in rice(Zhang et al., 2013). These studies showed that miRNAs play important roles in a number of developmental processes and pathways which regulate flower development related process.

Pepper (*Capsicum annuum* L.) is one of the most economically important worldwide vegetable crops(Barrajon-Catalan et al., 2020). Hybrid breeding has made a tremendous contribution to pepper yield, increasing seed production efficiency and protection of the varieties patent(Jifon et al., 2019). The utilization of male sterility in hybrid pepper is mainly based on three-line systems, which including cytoplasmic male sterile line (CMS), a maintainer line and restorer line(Bohra et al., 2016). In the CMS pepper, an ORF named *orf456*, was found at the 3'-end of the *coxII* gene, which is concluded that the *orf456* may represent a candidate gene, from mitochondrial genes, for determining the male-sterile phenotype of CMS(Kim et al., 2007). Furthermore, serval CMS-related sterile genes and fertility restorer genes also have been cloned from various plants(Bohra et al., 2016). In Arabidopsis, miR167 overexpression has been reported leading to male fertility defects(Ru et al., 2006), whereas miR159a overexpression results in decreased expression of *MYB33* and *MYB65*, leading to male sterility and delays flowering time(Anthony, 2005). With the development of miRNAome sequencing technology, differential expression patterns of miRNAs between the cytoplasmic male sterility (CMS) line and its maintainer line have been reported in many vegetable crops, such as Brassica juncea(Yang et al., 2013), Chinese cabbage(Wei et al., 2015) and Radish(Zhang et al., 2016b). A large number of miRNAs related to flowering and flower development have been identified

and characterized in above species. Nevertheless, there are no reports on systematic identification and characterization of CMS-related miRNAs in pepper.

To explore the roles of miRNA in CMS, we identified the miRNAs via a high-throughput sequencing approach from pepper anthers at the early uninucleate stage of the sterile line N816S and its maintainer line Ning5m. Differential expression patterns of miRNAs were analyzed and compared between N816S and Ning5m. Targets were predicted by degradome sequencing. These results may provide insights into clarification of the molecular mechanisms underlying the regulation of miRNAs during pollen development.

Materials and Methods

Pepper materials

The CMS sterile line N816S and its maintainer line Ning5m were used in this study. The plants were grown in greenhouse of the vegetable institute of Wuhan academy of agricultural sciences under normal conditions. In general, anthers at the uni-nucleate stage were manually collected. Anthers were harvested from three individual plants of each cultivar, immediately frozen in liquid nitrogen, stored at -80°C, and then used for RNA isolation. The microspore development was judged by both the floret length as described by Parra- Vega et al (Parra-Vega et al., 2013).

Small RNA library Construction, sequencing, and miRNA analysis

Total RNA was extracted using Trizol reagent (Invitrogen, CA, USA). RNA quantity was detected by Qubit Fluorometer (Invitrogen, CA, USA), and RNA purity was assayed by NanoDrop spectrophotometer (BioRad, PA, USA) and Agilent2100 bioanalyzer (Agilent Technologies, CA, USA). According to the manufacturer's instructions, sRNA libraries were constructed using the Small RNA Sample Prep Kit (Illumina, CA, US) and meanwhile the sRNA passed quality test(Yeri et al., 2018). Briefly, the small RNA (18-30 nt) were ligated to a 3' adaptor and a 5' adaptor sequentially and then converted to cDNA by RT-PCR. The purified cDNAs after reverse transcription reaction were sequenced by the Illumina Hiseq2000 (Illumina, CA, USA). After the Illumina sequencing, the raw sequences were obtained through a quality control process to generate high quality reads, and clean reads were directly used for further bioinformatics analysis with ACGT101-v4.2-miR (LC Sciences, TX, USA) to remove adapter sequences, junk reads, short reads, common RNA families (rRNA, tRNA, snRNA and snoRNA), repeats(Jeyaraj et al., 2019). Because of lacking known miRNA records of pepper in miRNA database (miRBase 21 released), unique sRNA sequences (18-25 nt) were mapped to specific species precursors in the miRBase 21 and the pepper reference sequence to identify conserved miRNAs by BLAST search. New miRNAs were identified by extracting flanking genome sequence of unique sRNAs using MIREAP (<http://sourceforge.net/projects/mireap/>)

(Huang et al., 2010), followed by the prediction of secondary structures by Mfold program (<http://unafold.rna.albany.edu/?q=mfold>) (Reuter and Mathews, 2009).

Degradome Library Construction, Data analysis and Target identification

The degradome library, a mixed samples' library, was constructed according to the method described previously using sliced ends of polyadenylated transcripts (German et al., 2009). In brief, poly-A-containing mRNA was purified from total RNA mixture of N816S and Ning5m, then ligated to 5' RNA adaptor containing a *MmeI* recognition site. Subsequently, RT-PCR was performed to first-strand cDNA, followed by digestion with *MmeI*, and then ligated to 3' adaptor. Finally, ligation product was amplified, purified and subjected to Illumina sequencing.

Raw reads were performed to remove adaptor sequence and low-quality reads resulting in clean reads. The high-quality specific sequences of were collected for subsequent degradome tags analysis. To identified potentially sliced targets of miRNAs, degradome sequence analysis were processed using the Cleveland 3.0 software package and the ACGT301-DGE program (LC Sciences, TX, USA) (Gong et al., 2015). The tags, which mapped to sense cDNA, were used to predict cleavage sites. Height of the degradome peak at each occupied transcript position was placed into five possible categories.

Quantitative real-time PCR (qRT-PCR) validation

Total RNA was extracted from pepper anthers, and RNA-free DNase I (Fermentas, USA) was used to remove DNA contamination for 15 min at 37°C. Stem-loop qRT-PCR were carried out to validate differential expressional levels of miRNAs. The mRNA template for the miRNA target was reverse transcribed using the OligodT₂₀ primer for qRT-PCR. All miRNA detection primers were designed and synthesized based on the mature miRNA sequences. For each miRNA, approximately 1 µg of total RNA was reverse-transcribed by reverse transcriptase using miRNA-specific stem-loop primers and a Fermentas Revert Aid First Strand cDNA Synthesis Kit (Fermentas, USA). Relative expression analysis of the miRNA and its target were performed using the ABI Step One Plus™ Real Time PCR System (Applied Biosystems, USA) and SYBR Green Master Mix (Roche, Germany). All reactions were run with three individual biological replicates, and *18S rRNA* was used as the internal control gene refer to Hwang et al (Hwang et al., 2013). The relative expression of miRNA and mRNA were used quantified the $2^{-\Delta\Delta C_t}$ method to calculate the fold change between N816S and Ning5m (Asha et al., 2016). The primers used are listed in Table S1.

157 Results

158 Overview of Small RNAs Libraries Sequencing Date

159 To determine the involvement regulatory roles of miRNAs in the fertility of sterile
160 and maintainer lines during anther development in pepper. Six small RNA libraries,
161 including three biological replicates from N816S (N816S_1, N816S_2, N816S_3) and
162 Ning5m (Ning5m_1, Ning5m_2, Ning5m_3), were constructed for deep sequencing.
163 An average of 14,567,767 and 14,274,100 raw reads were obtained from N816S and
164 Ning5m anthers, respectively, which after length, Junk reads, Rfam, Repeat, mRNA,
165 rRNA, tRNA, snoRNA and snRNA reads filtering, an average of 10,780,898 (72.21%)
166 valid reads representing 6,882,786 (84.68%) unique sequences and 10,442,445
167 (74.25%) valid reads representing 6,666,094 (86.4%) unique reads, respectively
168 (Table 1). The proportion of the valid reads in the corresponding raw reads was more
169 than 70%, which suggested that the quality of the sequencing data was high (Table1).
170 The length of total sRNAs and ranged from 18 to 25 nt, and in both the N816S and
171 Ning5m libraries, the 24 nt category was most abundant (average of 55.83% and
172 59.92% in N816S and Ning5m libraries, respectively) (Figure 1A-B). The length of
173 unique sRNAs, in both the N816S and Ning5m libraries, the 24 nt category was most
174 abundant (average of 59.15% and 62.90% in N816S and Ning5m libraries,
175 respectively) (Figure 2C-D). This is consistent with the typical lengths of plant
176 sRNAs reported in other studies(Asha et al., 2016;Hu et al., 2016).

177 Conserved miRNAs and novel miRNAs identified in Pepper

178 These unique sequences were subsequently used to identify conserved and novel
179 miRNAs by alignment against miRBase (Version 22), the pepper genome and
180 expressed sequence tag (EST) or pre-miRNA sequences. Unique miRNA transcripts
181 identified from the mappable sequences are divided into five group types (Table
182 S2)(Gong et al., 2015): (1) gp1a type: Unique reads map to specific
183 miRNAs/pre-miRNAs in miRbase and the pre-miRNAs further map to the genome
184 and EST. Total 77 miRNAs from the six samples corresponding to 50 known pepper
185 pre-miRNAs (Table S2-1). (2) gp1a type: Reads map to selected (except for specific)
186 miRNAs/pre-miRNAs in miRbase and the pre-miRNAs further map to the genome
187 and EST. 122 miRNAs, which correspond to 94 known pepper pre-miRNAs, which
188 cannot be mapped to genome or EST (Table S2-2). (3) gp2b type: Unique reads map
189 to selected miRNAs/pre-miRNAs in miRbase. The mapped pre-miRNAs do not map
190 to the genome, but the reads (and of course the miRNAs of the pre-miRNAs) map to
191 genome. The extended genome sequences from the genome loci may form hairpins.
192 261 miRNAs corresponding to 314 others known miRbase plant pre-miRNAs, which
193 are mapped to genome or EST (Table S2-3). (4) gp3a type: Unique reads map to
194 selected miRNAs/pre-miRNAs in miRbase. The mapped pre-miRNAs do not map to
195 the genome, and the reads do not map to the genome; Forty miRNAs corresponding to
196 35 others known miRbase plant pre-miRNAs, which cannot be mapped to genome or
197 EST (Table S2-4). (5) gp4a: Unique reads do not map to selected pre-miRNAs in

miRBase, but the reads map to genome and the extended genome sequences from genome may form hairpins; 411 miRNAs corresponding to 421 candidate pre-miRNAs, which are predict RNA hairpins derived from genome or EST, and these miRNAs are novel miRNA, which are labeled PC (pepper candidate) (Table S2-5).

In the present study, identified mature miRNAs are divided into two type, including conserved miRNA and novel miRNA (Table 1), referring to the research of Maize(Li et al., 2017). The length of the mature miRNAs ranged from 18 to 25 nt. The 24 nt and 21 nt miRNA category was most abundant, 41.85% and 27.41%, respectively (Figure 2A). This is consistent with the typical lengths of plant sRNAs reported in other studies(Gao et al., 2016;Zhang et al., 2016a). Of the conserved miRNA 21-nt miRNAs were most abundant (37.32%) (Figure 2B), representing the dominant size of mature miRNAs in plants. The 5' terminal nucleotides of sRNA sequences influence classification of their AGO complexes and is an important feature affecting function(Schuck et al., 2013). Most miRNAs are incorporated into the AGO1 effector complex, resulting in sequence specificity that either cleaves or translationally represses their targets(Dalmadi et al., 2019). Therefore, we examined the 5' nucleotide distribution of conserved miRNAs, and 42.49% started with uridine at 5' -end, and 26.17% started with adenine (Figure 2C).

Differentially expressed miRNAs analysis between pepper sterile line N816S and its maintainer line Ning5m

MiRNAs are play an important role in plant development and apoptosis(Jovanovic and Hengartner, 2006;Wang et al., 2007). To determine differential expression between N816S and Ning5m anthers. MiRNA expression was normalized to transcripts per million and simplified as normalized expression (Norm)(Gong et al., 2015). A miRNA was considered if the Norm value was greater than one in the all given replicated samples. Based on this criterion, 525 miRNAs were detected (Figure 3A), including a total of 350 conserved miRNAs (38.42% of the total miRNAs) belonging to 55 families were observed in at least one of the group samples (Table S3). In correlation analysis, Norm values of the N816S and Ning5m anthers were found to be highly correlated between repeats ($r>0.99$), indicating good reproducibility of the miRNAome results (Figure S1).

A one-tailed t-test was used to identify differentially expressed miRNAs with p value < 0.05 (Xing et al., 2012). The hierarchical clustering of the differential expression mRNAs was made and showed different expression patterns between the N816S and Ning5m (Figure 5A). As a result, a total of 76 miRNAs (59 conserved miRNAs and 17 novel miRNAs) were found to be differentially expressed between the two phenotypes (Table 3). Compared with the Ning5m, 44 miRNAs (34 conserved miRNAs and 10 novel miRNAs) were found to be down-regulated and 32 miRNAs (25 conserved miRNAs and 7 novel miRNAs) up-regulated in the N816S (Figure 3B). Nta-miR156g_L+1 and stu-miR156a-5p had high expression abundance in MiRNA156 family, were down-regulated in sterile line N816S. However, miRNA390 family, including hex-MIR390a-p5_1ss21GT, hex-MIR390b-p5_2ss10TC21GA and

hex-MIR390b-p5_1ss21GT, were up-regulated in sterile line N816S. Five miRNAs (stu-miR393-3p, stu-miR399j-3p_1ss21GA, nta-miR6149a_L+1R-1_1ss21GC and stu-miR398b-3p ath-miR8175_L+4) were specifically expressed in Ning5m. Five miRNA(NTA-MIR172e-p3_2ss15CG19TA, ppe-MIR399a-p3_2ss5AT18TC, ppe-MIR399a-p5_2ss5AT18TC, stu-miR3627-5p_R-1 and sly-MIR10528-p3_2ss9GA19TC) were specifically expressed in N816S. To validate conserved miRNAs identified and novel miRNAs predicted, we selected 18 differential expressed miRNAs for stem-loop qRT-PCR. The expression trends of these miRNAs were consistent with the high-throughput sequencing results. As showed in Figure 3C, expression of these miRNAs from qRT-PCR displayed a similar tendency with those from small RNA sequencing.

253 **Targets analysis of miRNAs in anthers of Pepper.**

254 To understand the function of the mature miRNAs from anthers of Pepper, 255 targets of miRNAs were detected using degradome sequencing technology. A total of 256 34233673 (99.24%) mappable reads from raw reads were obtained, including 257 12320360(99.19%) unique mappable reads, while 20851831 (60.45%) transcript 258 mapped reads, including 6635426(53.42%) unique transcript mappable reads, were 259 obtained (Table S4). The target transcripts were sorted into five categories according 260 to the relative abundance of the tags at the target mRNA sites(Gong et al., 2015): 261 category “0” is defined as >1 raw read at the position, with abundance at a position 262 equal to the maximum on the transcript and with only one maximum on the transcript; 263 category “1”, the expected cleavage signature was equal to the maximum on the 264 transcript and more than one maximum position on the transcript; category “2”, 265 abundance at the position was less than the maximum but higher than the median for 266 the transcript; category “3”, the abundance at the position equal to or less than the 267 median for the transcript; category “4”, abundance at the position was only one raw 268 read. Figure 6 showed the typical five categories of the target transcripts.

269 A total of 1292 reliable targets ($p_value < 1$) that were potentially cleaved by 270 321 miRNAs (250 conserved miRNAs and 71 novel miRNAs) were identified (Table 271 S5). A total of 77 target genes were identified to be targeted by 35 differentially 272 expressed miRNAs (27 conserved miRNAs and 8 novel miRNAs) (Table S6). Many 273 targets of miR156, miR167, miRNA858 families were identified as transcription 274 factors, such as SPL, ARF, and MYB, respectively, which have been experimentally 275 validated by the previous studies.

276 **Gene Ontology (GO) and KEGG Pathway Analysis of Targets.**

277 The targets were annotated using the GO annotations analysis, which is 278 commonly used to describe the function of genes and gene products, meanwhile the 279 KEGG analysis is used to provide the pathway of annotated targets. Figure 5B

showed the GO functional classification of miRNA targets in the pepper anthers. Targets covered a broad range of functional categories. However, targets of DNA binding, regulation of transcription-, and DNA-templated genes, transcription factor activity-, and sequence-specific DNA binding, transcription-, and DNA-templated genes were mostly enriched. This suggested that those target genes may play an important role in pepper anthers. The pathways of all the miRNAs and targets involved were listed (Table S7). There were 107 pathways that targets involved, while carbohydrate metabolism, translation, folding sorting and degradation pathways owned most targets (Figure 5C). Furthermore, the 77 miRNA targets from 35 differentially expressed miRNAs were found to be involved in multiple pathways, including plant hormone signal transduction, purine metabolism, starch and sucrose metabolism, oxidative phosphorylation, and others (Table S8).

Expression profiles of miRNA targets examined by qRT-PCR

To examine the correlation between the targets and the corresponding miRNAs, the expression levels of 19 selected targets were examined by qRT-PCR analysis. MiRNAome sequencing result showed that miRNA156a/g (stu-miR156a and nta-miR156g_L+1), miRNA167a/h (stu-miR167a-5p, stu-miR167a-5p_1ss21AT and mdm-miR167h_1ss22AT), stu-miR393-5p and nta-miR6149a_L+1R-1_1ss21GC were down-regulated in N816S compare to Ning5m (Table 3). The degradome sequencing results indicate that miRNA156a/g target SPL family genes (SPL3, SPL6 and SPL9), miRNA167a/h (aly-miR167a-5p_R+1_1ss21AT, gma-miR167a_1ss21AT, mdm-miR167h_1ss22AT, stu-miR167a-5p_1ss21AT) targets an AUXIN RESPONSE FACTOR gene (ARF8), MiRNA393 targets two AUXIN SIGNALING F-box genes (TIR1 and AFB2) (Table S8). QRT-PCR analysis showed that the expression profiles of the above miRNA targets were up-regulated in N816S (Figure 6A).

In contrast, miRNA159_R3, nta-MIR169j-p3_2ss8GA19CT, stu-miR319b_L+1R-1, hex-MIR390a-p5_1ss21GT, ppe-MIR399a-p3_2ss5AT18TC, ppe-MIR858-p5_1ss4GA, cst-MIR11334-p5_2ss9TG18TC and PC-5p-154497_29 were up-regulated in N816S (Table 3). MiRNA159_R3 targets a plasma membrane H⁺-ATPase gene (*ACA13*), a γ -TuC Protein 3 (GCP3)-Interacting Protein gene (*GIP1*), MiRNA169j-p3 targets a heat shock protein gene (HSP70), MiRNA319b targets four TEOSINTE BRANCHED/CYCLOIDEA/PCF transcription factor genes (*TCP2*, *TCP4*, *TCP24*) and a Aldehyde dehydrogenase gene (*ALDH6B2*), MiRNA390a-p5 targets a GDP dissociation inhibitor gene (*GDI1*), miRNA399a-p3 targets a ceramide-1-phosphate transfer protein gene (*ACD11*), miRNA858-p5 targets several MYB protein genes (*MYB3*, *MYB4*, *MYB12* and *MYB80*), miR11334-p5 targets a phospholipase D alpha 1-like gene (*PLD1*) and a Acyl-CoA oxidase gene (*ACX1*), and a novel miRNA of PC-5p-154497_29 targets a UDP-galactose-dependent digalactosyldiacylglycerol synthase gene (*DGD2*) (Table S8). Most of above targets were down-regulated in N816S (Figure 6B).

Most negative correlations were found between the expression levels of the

target genes and their corresponding miRNAs in anthers of the sterile line N816S and its maintainer line Ning5m, implying that miRNA-mediated mRNA silencing occurred during anther development.

Discussion

Several studies have shown that miRNAs regulate anther development in plants (Yang et al., 2013; Gong et al., 2015). However, few studies on the relationships between miRNA biogenesis and CMS occurrence during anther development in pepper were very limited. MiRNAome and degradation sequencing technology provides an effective way to identify and evaluate the expression profiles of miRNAs and targets associated with CMS occurrence in plant tissues during anther development (Wei et al., 2015; Hu et al., 2016; Bai et al., 2017). To explore the roles of miRNAs during the occurrence of CMS, anthers at the early uninucleate stage were used to examine the expression profiles of the miRNAs in the sterile line N816S and its maintainer line Ning5m using high-throughput sequencing. To the best of our current knowledge, this study is the first report on identification and characterization of miRNAs, and their targets between a sterile line and its maintainer line during anther development in pepper.

In this study, a total of 76 miRNAs (59 conserved miRNAs and 17 novel miRNAs) were identified as differentially expressed between the sterile line N816S and its maintainer line Ning5m. To understand the function of the miRNAs from anthers of Pepper, targets of miRNAs were detected using degradome sequencing technology. From those targets, 80 target genes were identified to be targeted by 35 differentially expressed miRNAs (27 conserved miRNAs and 8 novel miRNAs). The qRT-PCR results indicated that most negative correlations were found between the expression levels of the target genes and their corresponding miRNAs in the anthers of the sterile line N816S and its maintainer line Ning5m. KEGG analysis showed that most gene targets of the 35 differentially expressed miRNAs were involved in pathways, including plant hormone signal transduction, starch and sucrose metabolism, oxidative phosphorylation, purine metabolism, and others.

Intriguingly, a number of genes in hormone signaling were confirmed or predicted as targets of miRNAs. Auxin regulates anther dehiscence, pollen maturation, and filament elongation in *Arabidopsis*. The miR167-guided cleavage of auxin response factor *ARF8*, miR393-guided cleavage of auxin receptor *TIR1* and *AFB2*, and miR319-guided cleavage of jasmonate acid biosynthesis related transcription factor genes *TCPs*, which were reported in previous study (Liu and Chen, 2009). In *Arabidopsis*, loss of miR167 regulation in *mARF6* and *mARF8* expression caused arrested ovule development and anther indehiscence (Wu et al., 2006). Four auxin receptor-encoding genes, *TIR1*, *AFB1*, *AFB2*, and *AFB3*, are transcribed in anthers only during late stages of development starting at the end of meiosis (Shimizu-Mitao and Kakimoto, 2014). Up-regulation of *TIR1* enhances auxin sensitivity, and causes

363 altered leave phenotype and delayed flowering in *Arabidopsis*(Chen et al., 2011).
 364 MicroRNA319-regulated TCPs can activate CO transcription and control flowering
 365 time, shaping flower structure, leaf morphology, and plant architecture in
 366 *Arabidopsis*(Chen et al., 2011). In this study, up-regulation *AFR8* was targeted by
 367 down-regulated miRNA393a/h, up-regulated *TIR1* and *AFB2* were targeted by
 368 down-regulated miR393a, and down-regulated *TCPs* (*TCP2* and *TCP4*) were targeted
 369 by up-regulated miRNA393b, indicating might modulate the hormone response to
 370 play roles in the pollen development and CMS occurrence.

371 Targets of these differentially expressed miRNAs containing important
 372 transcription factors (TFs) and functional proteins are involved in many biological
 373 processes, including signal transduction, floral organ development, and organellar
 374 gene expression(Liu and Chen, 2009). For instance, miR156 targets *SPL* transcription
 375 factor family, which is regulatory functions throughout the growth and development
 376 stages in plants. Previous report showed that miR156 regulates the timing of flower
 377 formation vis *SPL3/4/5*, activating the expression of *LEAFY*, *FRUITFULL* and
 378 *APETALA*(Jung et al., 2011). In *Arabidopsis*, multiple *SPL* genes regulate cell
 379 division, differentiation and can result in fertile flower(Zheng et al., 2019). In this
 380 study, *SPL3*, *SPL6* and *SPL9* were identified as potentially targets of miR156 miRNA
 381 family members (stu-miR156a, stu-miR156a, nta-miR156g_L+1) were identified as
 382 approximately triple up-regulated in N816S compared with Ning5M (Figure 6A),
 383 leading to disordered floral organ development in pepper, indicating that miR156 may
 384 participate in fertility regulation. Furthermore, miR858 targets *MYB* transcription
 385 factor family, which is involved in the control of plant development, determination of
 386 cell fate and identity, primary, and secondary metabolism. In rice, anther and pollen
 387 defect in floral organ development are found in the loss-of-function mutations of *MYB*.
 388 In *Arabidopsis*, *AtMYB3*, *AtMYB4*, *AtMYB7* and *AtMYB32* encode transcriptional
 389 repressors, and *AtMYB4* controls sinapate ester biosynthesis, whereas *AtMYB32*
 390 regulates pollen wall composition, and *AtMYB12* is involved in the regulation of
 391 flavonoid biosynthesis control flavonol biosynthesis(Dubos et al., 2010). The
 392 transcriptional activity of the PtMYB4 protein is positively regulated by the
 393 mitogen-activated protein kinase (MAPK) PtMAPK6 in *Pinus taeda*, which
 394 phosphorylates a Ser in the C terminal activation domain. In addition, *AtMYB80*
 395 regulated exine formation and acts downstream of *AtMYB35* to control anther
 396 development and/ or functionalities. In this present study, up-regulated miRNA858
 397 might via reducing transcript expression of *MYB3*, *MYB4*, *MYB12* and *MYB80*,
 398 leading to anther and pollen defect. In addition, MIR169j targets a heat shock protein
 399 *HSP70*, which has been widely involved in the protein peptides folding, assembly and
 400 transports, and the degradation of abnormal proteins. Studies have found that *HSP70*
 401 is associated with male sterility in plant and animal. Down regulation of *HSP70*
 402 expression level prolongs the duration of heat-induced male sterility in *Drosophila*
 403 *buzzatii*(Sarup et al., 2004). *HSP70* antisense RNA expression leads to male sterility
 404 in rice(Liu et al., 2008). In this study, up-regulated miRNA169j may through via
 405 reducing transcript expression of *HSP70*, leading to pollen abortion.

406 Many miRNA-targeted genes were involved in lipid transport and metabolism,

such as *PLD1/ACX1*, *ACA13/GIP*, *ALDH6*, *GDI1* and *DGD2*, which are cleaved by miR11334, miR159_R-3, miR319b, miRNA390b and PC-5p-154497_29, respectively. Those miRNAs may throughout their targets be involved in CMS sterile process. Phospholipase D alpha 1-like (PLD1) has been identified as cytosolic protein, which regulated cytosolic lipid droplet formation(Andersson et al., 2006). Acyl-CoA oxidase (ACX) was the first and the key step controlling enzyme involved in fatty acid β -oxidation, and mutation of *ACX1* led to petal degeneration in Chinese Cabbage(Zheng et al., 2019). In plant, *ALDH6B2* encodes a methylmalonyl semialdehyde dehydrogenase, of which involved in the degradation of valine to propionyl CoA(Brocker et al., 2013). In addition, Plant cells have multiple plasma membrane (PM)-localized calcium ATPases (ACAs) pumping calcium ions out of the cytosol. The loss of ACA13 combination with a reduction in function of other ACAs leads to seedling death at bolting, revealing the essential role of their collective function in plant growth(Yu et al., 2018). Moreover, the γ -tubulin complex (γ -TuC) Protein 3 (GCP3)-Interacting Protein 1 (GIP1) is the smallest γ -TuC component identified. In Arabidopsis, *AtGIP1* and its homologous protein *AtGIP2* mutants are impaired in establishing a fully functional mitotic spindle and exhibit severe developmental defects(Batzenschlager et al., 2013). The GDP dissociation inhibitor protein GDI1 relates to control vesicle number and transport in an amelioration of zinc toxicity, allowing yeast to survive in the presence of toxic(Ezaki and Nakakihara, 2012). Furthermore, Digalactosyl-diacylglycerol (DGD) is one of the major lipids found predominantly in the photosynthetic membrane of higher plants. *OsDGD2 β* is the sole DGDG synthase gene highly expressed in anther, and its mutation confers male sterility with pale yellow and shrunken anther, devoid of starch granules in pollen, and delayed degeneration of tapetal cells in rice(Basnet et al., 2019). All above related miRNAs are up-regulated in sterile line N816S compare to its maintainer Ning5m and those disorder miRNA-targeted genes perhaps leading to membrane-disruptive effects and cell apoptosis.

435

436 Conclusion

In the present study, small RNA libraries from anther of CMS-line N816S and its maintainer line Ning5m were generated by small RNA sequencing in pepper. A total of 76 differentially expressed miRNAs were discovered, of which 18 were further confirmed by real-time quantitative PCR (qRT-PCR). Furthermore, targets of miRNAs were identified by degradome sequencing. A total of 1292 targets that were potentially cleaved by 321 miRNAs (250 conserved miRNAs and 71 novel miRNAs) were identified. Gene Ontology (GO) and KEGG pathway analysis of target transcripts indicated that 77 target genes cleaved by 35 differentially expressed miRNAs might play roles in the regulation of CMS sterility, such as MYB, SPL, and AFR family proteins targeted by miR156, miR167, miRNA858 family. Nineteen

447 targets were selectively examined by qRT-PCR, and the results showed that there was
448 a negative correlation on the expression patterns between miRNAs and their targets.
449 These findings provide valuable information to understand the roles of miRNAs
450 during anther development and CMS occurrence in pepper.

451 **Author contributions**

452 Min Zhang designed the study. Hongyuan Zhang, Shuping Huang and Jie Tan
453 carried out the experiment, data analysis, interpretation of the results. Hongyuan
454 Zhang drafted the manuscript. Xia Chen supervised the work and revised the
455 manuscript. All authors have read and approved the final version of this submission.
456

457 **Acknowledgement**

458 This work was financially supported by the National Natural Science Foundation of
459 China (Grant No. 31701936) and the subproject of National key research and
460 development Program of China (2017YFD0101904).
461

462 **Reference**

- 463 Andersson, L., Bostrom, P., Ericson, J., Rutberg, M., Magnusson, B., Marchesan, D., Ruiz, M.,
464 Asp, L., Huang, P., Frohman, M.A., Boren, J., and Olofsson, S.O. (2006). PLD1 and
465 ERK2 regulate cytosolic lipid droplet formation. *J Cell Sci* 119, 2246-2257.
- 466 Anthony, A.M.F., G. (2005). The Arabidopsis GAMYB-Like Genes, MYB33 and MYB65 , Are
467 MicroRNA-Regulated Genes That Redundantly Facilitate Anther Development. *The*
468 *Plant Cell* 17, 705-721.
- 469 Asha, S., Sreekumar, S., and Soniya, E.V. (2016). Unravelling the complexity of
470 microRNA-mediated gene regulation in black pepper (*Piper nigrum* L.) using
471 high-throughput small RNA profiling. *Plant Cell Reports* 35, 53-63.

472 Bai, J.F., Wang, Y.K., Wang, P., Duan, W.J., Yuan, S.H., Sun, H., Yuan, G.L., Ma, J.X., Wang,
473 N., Zhang, F.T., Zhang, L.P., and Zhao, C.P. (2017). Uncovering Male Fertility
474 Transition Responsive miRNA in a Wheat Photo-Thermosensitive Genic Male Sterile
475 Line by Deep Sequencing and Degradome Analysis. *Frontiers In Plant Science* 8.

476 Barrajon-Catalan, E., Alvarez-Martinez, F.J., Borrás, F., Perez, D., Herrero, N., Ruiz, J.J., and
477 Micol, V. (2020). Metabolomic analysis of the effects of a commercial complex
478 biostimulant on pepper crops. *Food Chemistry* 310.

479 Basnet, R., Hussain, N., and Shu, Q. (2019). OsDGD2 β is the Sole Digalactosyldiacylglycerol
480 Synthase Gene Highly Expressed in Anther, and its Mutation Confers Male Sterility in
481 Rice. *Rice* 12, 66.

482 Batzenschlager, M., Masoud, K., Janski, N., Houlné, G., Herzog, E., Evrard, J.-L., Baumberger,
483 N., Ehrardt, M., Nominé, Y., Kieffer, B., Schmit, A.-C., and Chabouté, M.-E. (2013).
484 The GIP gamma-tubulin complex-associated proteins are involved in nuclear
485 architecture in Arabidopsis thaliana. *Frontiers in Plant Science* 4.

486 Bohra, A., Jha, U.C., Adhimoolam, P., Bisht, D., and Singh, N.P. (2016). Cytoplasmic male
487 sterility (CMS) in hybrid breeding in field crops. *Plant Cell Reports* 35, 967-993.

488 Brocker, C., Vasiliou, M., Carpenter, S., Carpenter, C., Zhang, Y., Wang, X., Kotchoni, S.O.,
489 Wood, A.J., Kirch, H.-H., Kopečný, D., Nebert, D.W., and Vasiliou, V. (2013).
490 Aldehyde dehydrogenase (ALDH) superfamily in plants: gene nomenclature and
491 comparative genomics. *Planta* 237, 189-210.

492 Chen, Z.H., Bao, M.L., Sun, Y.Z., Yang, Y.J., Xu, X.H., Wang, J.H., Han, N., Bian, H.W., and
493 Zhu, M.Y. (2011). Regulation of auxin response by miR393-targeted transport inhibitor

494 response protein 1 is involved in normal development in Arabidopsis. *Plant Molecular*
495 *Biology* 77, 619-629.

496 Dalmadi, A., Gyula, P., Balint, J., Szittyá, G., and Havelda, Z. (2019). AGO-unbound cytosolic
497 pool of mature miRNAs in plant cells reveals a novel regulatory step at AGO1 loading.
498 *Nucleic Acids Research* 47, 9803-9817.

499 Diaz-Manzano, F.E., Cabrera, J., Ripoll, J.J., Del Olmo, I., Andres, M.F., Silva, A.C., Barcala,
500 M., Sanchez, M., Ruiz-Ferrer, V., De Almeida-Engler, J., Yanofsky, M.F., Pineiro, M.,
501 Jarillo, J.A., Fenoll, C., and Escobar, C. (2018). A role for the gene regulatory module
502 microRNA172/TARGET OF EARLY ACTIVATION TAGGED 1/FLOWERING LOCUS
503 T (miRNA172/TOE1/FT) in the feeding sites induced by *Meloidogyne javanica* in
504 *Arabidopsis thaliana*. *New Phytologist* 217, 813-827.

505 Dubos, C., Stracke, R., Grotewold, E., Weisshaar, B., Martin, C., and Lepiniec, L. (2010). MYB
506 transcription factors in Arabidopsis. *Trends in Plant Science* 15, 573-581.

507 Ezaki, B., and Nakakihara, E. (2012). Possible involvement of GDI1 protein, a GDP
508 dissociation inhibitor related to vesicle transport, in an amelioration of zinc toxicity in
509 *Saccharomyces cerevisiae*. *Yeast* 29, 17-24.

510 Gao, S., Yang, L., Zeng, H.Q., Zhou, Z.S., Yang, Z.M., Li, H., Sun, D., Xie, F.L., and Zhang,
511 B.H. (2016). A cotton miRNA is involved in regulation of plant response to salt stress.
512 *Scientific Reports* 6.

513 German, M.A., Luo, S.J., Schroth, G., Meyers, B.C., and Green, P.J. (2009). Construction of
514 Parallel Analysis of RNA Ends (PARE) libraries for the study of cleaved miRNA targets
515 and the RNA degradome. *Nature Protocols* 4, 356-362.

516 Gong, S.M., Ding, Y.F., Huang, S.X., and Zhu, C. (2015). Identification of miRNAs and Their
517 Target Genes Associated with Sweet Corn Seed Vigor by Combined Small RNA and
518 Degradome Sequencing. *Journal Of Agricultural And Food Chemistry* 63, 5485-5491.

519 Hu, J.H., Jin, J., Qian, Q., Huang, K.K., and Ding, Y. (2016). Small RNA and degradome
520 profiling reveals miRNA regulation in the seed germination of ancient eudicot *Nelumbo*
521 *nucifera*. *Bmc Genomics* 17.

522 Huang, Q.X., Cheng, X.Y., Mao, Z.C., Wang, Y.S., Zhao, L.L., Yan, X., Ferris, V.R., Xu, R.M.,
523 and Xie, B.Y. (2010). MicroRNA Discovery and Analysis of Pinewood Nematode
524 *Bursaphelenchus xylophilus* by Deep Sequencing. *Plos One* 5.

525 Hwang, D.G., Park, J.H., Lim, J.Y., Kim, D., Choi, Y., Kim, S., Reeves, G., Yeom, S.I., Lee,
526 J.S., Park, M., Kim, S., Choi, I.Y., Choi, D., and Shin, C. (2013). The Hot Pepper
527 (*Capsicum annuum*) MicroRNA Transcriptome Reveals Novel and Conserved Targets:
528 A Foundation for Understanding MicroRNA Functional Roles in Hot Pepper. *Plos One*
529 8.

530 Jeyaraj, A., Wang, X.W., Wang, S.S., Liu, S.R., Zhang, R., Wu, and Wei, C.T. (2019).
531 Identification of Regulatory Networks of MicroRNAs and Their Targets in Response to
532 *Colletotrichum gloeosporioides* in Tea Plant (*Camellia sinensis* L.). *Frontiers In Plant*
533 *Science* 10.

534 Jha, A., and Shankar, R. (2011). Employing machine learning for reliable miRNA target
535 identification in plants. *Bmc Genomics* 12.

536 Jifon, J., Crosby, K., Patil, B., and Leskovar, D. (2019). Effects of Potassium Nutrition on Seed
537 Yield and Quality of Hybrid Chili Pepper (*Capsicum annuum* L.). *Hortscience* 54,

538 S395-S395.

539 Jovanovic, M., and Hengartner, M.O. (2006). miRNAs and apoptosis: RNAs to die for.

540 *Oncogene* 25, 6176-6187.

541 Jung, J.-H., Seo, P.J., Kang, S.K., and Park, C.-M. (2011). miR172 signals are incorporated

542 into the miR156 signaling pathway at the SPL3/4/5 genes in Arabidopsis

543 developmental transitions. *Plant Molecular Biology* 76, 35-45.

544 Kim, D.H., Kang, J.G., and Kim, B.D. (2007). Isolation and characterization of the cytoplasmic

545 male sterility-associated orf456 gene of chili pepper (*Capsicum annuum* L.). *Plant Mol*

546 *Biol* 63, 519-532.

547 Li, H.P., Peng, T., Wang, Q., Wu, Y.F., Chang, J.F., Zhang, M.B., Tang, G.L., and Li, C.H.

548 (2017). Development of Incompletely Fused Carpels in Maize Ovary Revealed by

549 miRNA, Target Gene and Phytohormone Analysis. *Frontiers In Plant Science* 8.

550 Lin, C.S., Chen, J.J.W., Huang, Y.T., Hsu, C.T., Lu, H.C., Chou, M.L., Chen, L.C., Ou, C.I.,

551 Liao, D.C., Yeh, Y.Y., Chang, S.B., Shen, S.C., Wu, F.H., Shih, M.C., and Chan, M.T.

552 (2013). Catalog of *Erycina pusilla* miRNA and categorization of reproductive

553 phase-related miRNAs and their target gene families. *Plant Molecular Biology* 82,

554 193-204.

555 Liu, L., Liu, G.Q., and Hou, N. (2008). Construction and identification of HSP70 antisense RNA

556 expression vector for genetic engineering male sterility in plant. *Anhui Agricultural*

557 *Science and Technology* 9, 84, 128.

558 Liu, Q., and Chen, Y.Q. (2009). Insights into the mechanism of plant development: Interactions

559 of miRNAs pathway with phytohormone response. *Biochemical And Biophysical*

560 *Research Communications* 384, 1-5.

561 Parra-Vega, V., Gonzalez-Garcia, B., and Segui-Simarro, J.M. (2013). Morphological markers
562 to correlate bud and anther development with microsporogenesis and
563 microgametogenesis in pepper (*Capsicum annuum* L.). *Acta Physiologiae Plantarum*
564 35, 627-633.

565 Pei, H.X., Ma, N., Chen, J.W., Zheng, Y., Tian, J., Li, J., Zhang, S., Fei, Z.J., and Gao, J.P.
566 (2013). Integrative Analysis of miRNA and mRNA Profiles in Response to Ethylene in
567 Rose Petals during Flower Opening. *Plos One* 8.

568 Reuter, J., and Mathews, D.H. (2009). RNAstructure: Software for RNA Secondary Structure
569 Prediction and Analysis. *Journal Of Biomolecular Structure & Dynamics* 26, 831-832.

570 Ru, P., Xu, L., Ma, H., and Huang, H. (2006). Plant fertility defects induced by the enhanced
571 expression of microRNA167. *Cell Research* 16, 457-465.

572 Sarup, P., Dahlgaard, J., Norup, A.M., Jorgensen, K.T., Hebsgaard, M.B., and Loschcke, V.
573 (2004). Down regulation of Hsp70 expression level prolongs the duration of
574 heat-induced male sterility in *Drosophila buzzatii*. *Functional Ecology* 18, 365-370.

575 Schuck, J., Gursinsky, T., Pantaleo, V., Burgyan, J., and Behrens, S.E. (2013).
576 AGO/RISC-mediated antiviral RNA silencing in a plant in vitro system. *Nucleic Acids*
577 *Research* 41, 5090-5103.

578 Shimizu-Mitao, Y., and Kakimoto, T. (2014). Auxin Sensitivities of All Arabidopsis Aux/IAAs for
579 Degradation in the Presence of Every TIR1/AFB. *Plant And Cell Physiology* 55,
580 1450-1459.

581 Stepień, A., Knop, K., Dolata, J., Taube, M., Bajczyk, M., Barciszewska-Pacak, M., Pacak, A.,

582 Jamolowski, A., and Szweykowska-Kulinska, Z. (2017). Posttranscriptional
583 coordination of splicing and miRNA biogenesis in plants. *Wiley Interdisciplinary*
584 *Reviews-Rna* 8.

585 Tripathi, R.K., Bregitzer, P., and Singh, J. (2018). Genome-wide analysis of the SPL/miR156
586 module and its interaction with the AP2/miR172 unit in barley. *Scientific Reports* 8.

587 Wang, Y., Stricker, H.M., Gou, D., and Liu, L. (2007). MicroRNA: past and present. *Front*
588 *Biosci* 12, 2316-2329.

589 Wei, X.C., Zhang, X.H., Yao, Q.J., Yuan, Y.X., Li, X.X., Wei, F., Zhao, Y.Y., Zhang, Q., Wang,
590 Z.Y., Jiang, W.S., and Zhang, X.W. (2015). The miRNAs and their regulatory networks
591 responsible for pollen abortion in Ogura-CMS Chinese cabbage revealed by
592 high-throughput sequencing of miRNAs, degradomes, and transcriptomes. *Frontiers*
593 *In Plant Science* 6.

594 Wu, M.F., Tian, Q., and Reed, J.W. (2006). Arabidopsis microRNA167 controls patterns of
595 ARF6 and ARF8 expression, and regulates both female and male reproduction.
596 *Development* 133, 4211-4218.

597 Xing, R.H., Lussier, Y.A., Salama, J.K., Khodarev, N.N., Huang, Y., Zhang, Q.B., Khan, S.A.,
598 Yang, X.N., Hasselle, M.D., Darga, T.E., Malik, R., Fan, H.L., Perakis, S., Filippo, M.,
599 Corbin, K., Lee, Y., Posner, M.C., Chmura, S.J., Hellman, S., and Weichselbaum, R.R.
600 (2012). MicroRNA expression characterizes oligometastasis(es). *Cancer Research*
601 72.

602 Yang, J., Liu, X., Xu, B., Zhao, N., Yang, X., and Zhang, M. (2013). Identification of miRNAs
603 and their targets using high-throughput sequencing and degradome analysis in

604 cytoplasmic male-sterile and its maintainer fertile lines of *Brassica juncea*. *BMC*
605 *Genomics* 14, 9.

606 Yeri, A., Courtright, A., Danielson, K., Hutchins, E., Alsop, E., Carlson, E., Hsieh, M., Ziegler,
607 O., Das, A., Shah, R.V., Rozowsky, J., Das, S., and Van Keuren-Jensen, K. (2018).
608 Evaluation of commercially available small RNASeq library preparation kits using low
609 input RNA. *Bmc Genomics* 19.

610 Yu, H., Yan, J., Du, X., and Hua, J. (2018). Overlapping and differential roles of plasma
611 membrane calcium ATPases in Arabidopsis growth and environmental responses. *J*
612 *Exp Bot* 69, 2693-2703.

613 Yu, S., Galvao, V.C., Zhang, Y.C., Horrer, D., Zhang, T.Q., Hao, Y.H., Feng, Y.Q., Wang, S.,
614 Schmid, M., and Wang, J.W. (2012). Gibberellin Regulates the Arabidopsis Floral
615 Transition through miR156-Targeted SQUAMOSA PROMOTER BINDING-LIKE
616 Transcription Factors. *Plant Cell* 24, 3320-3332.

617 Zhang, H.Y., Hu, J.H., Qian, Q., Chen, H., Jin, J., and Ding, Y. (2016a). Small RNA Profiles of
618 the Rice PTGMS Line Wuxiang S Reveal miRNAs Involved in Fertility Transition.
619 *Frontiers In Plant Science* 7.

620 Zhang, S.X., Liu, Y.H., and Yu, B. (2015). New insights into pri-miRNA processing and
621 accumulation in plants. *Wiley Interdisciplinary Reviews-Rna* 6, 533-545.

622 Zhang, W., Xie, Y., Xu, L., Wang, Y., Zhu, X.W., Wang, R.H., Zhang, Y., Muleke, E.M., and Liu,
623 L.W. (2016b). Identification of microRNAs and Their Target Genes Explores
624 miRNA-Mediated Regulatory Network of Cytoplasmic Male Sterility Occurrence during
625 Anther Development in Radish (*Raphanus sativus* L.). *Frontiers In Plant Science* 7.

626 Zhang, Y.-C., Yu, Y., Wang, C.-Y., Li, Z.-Y., Liu, Q., Xu, J., Liao, J.-Y., Wang, X.-J., Qu, L.-H.,
627 Chen, F., Xin, P., Yan, C., Chu, J., Li, H.-Q., and Chen, Y.-Q. (2013). Overexpression
628 of microRNA OsmiR397 improves rice yield by increasing grain size and promoting
629 panicle branching. *Nature Biotechnology* 31, 848-852.

630 Zheng, C.F., Ye, M.X., Sang, M.M., and Wu, R.L. (2019). A Regulatory Network for
631 miR156-SPL Module in Arabidopsis thaliana. *International Journal Of Molecular*
632 *Sciences* 20.

633 Zhou, X.N., Zhang, Z.H., and Liang, X.H. (2020). Regulatory Network Analysis to Reveal
634 Important miRNAs and Genes in Non-Small Cell Lung Cancer. *Cell Journal* 21,
635 459-466.

636 Zhu, Q.H., and Helliwell, C.A. (2011). Regulation of flowering time and floral patterning by
637 miR172. *Journal Of Experimental Botany* 62, 487-495.

638
639
640

Figures and Tables:

Figure 1. Frequency percent of Clean reads (A) and Unique reads (B) sequencing reads, and the length distribution of the clean reads (C) and unique reads (D) from anther of *Capsicum annuum* L.

Figure 2. Size distribution of miRNAs and characterization of converted miRNAs detected by deep sequencing. (A) Length distribution of the total miRNAs; (B) Distribution of obtained unique converted miRNAs (gp1-gp3); (C) Percentage of first nucleotide bias in the identified converted miRNAs(gp1-gp3).

655
656 Figure 3. Venn diagrams detected miRNA, the number statistics of differential
657 expression miRNAs, and qRT-PCR for verifying deep sequencing. (A) Detected
658 miRNAs in N816S and Ning5m; (B) The number statistics of up-regulated and
659 down-regulated differential expression miRNAs in conserved miRNAs and novel
660 miRNAs. (C) Detection of selected miRNA expression in N816S and Ning5m anthers
661 using qRT-PCR.

662
663

664
665 Figure 4. Typical categories of the target transcript according to the relative
666 abundance of the tags at the target mRNA sites.
667

Figure 5. (A) Differentially expressed miRNAs cluster, (B) GO functional classification of miRNA targets, (C) KEGG Pathway Classification; Notes: miRNAs cluster columns are different samples; rows are different miRNAs. Clustering with p_value, red is up-expressed miRNA, blue is down-expressed miRNA.

674

676

677 Figure 6. Quantitative real-time PCR analysis of the relative expression of
678 miRNA targets in the CMS-line N816S and its maintainer line Ning5m. (A)
679 up-regulated targets in N816S; (B) down-regulated targets in N816S.

680

681

Table 1. Overview of reads from raw data to cleaned sequences.

Category	N816S_mean		Ning5m_mean	
	Total sRNAs (%)	unique sRNAs	Total sRNAs (%)	unique sRNAs
Raw reads	14567767 (100%)	8074144 (100%)	14274100 (100%)	7568101 (100%)
		1003388		
3ADT&Length filter	1255570 (8.62%)	(12.43%)	1012608 (7.09%)	707822 (9.35%)
Junk reads	74827 (0.51%)	54461 (0.67%)	74748 (0.52%)	54565 (0.72%)
Rfam	328527 (2.26%)	10516 (0.13%)	382951 (2.68%)	11080 (0.15%)
mRNA	2437774 (16.73%)	132754 (1.64%)	2728370 (19.11%)	138920 (1.84%)
Repeats	8437 (0.06%)	306 (0.00%)	8898 (0.06%)	293 (0.00%)
	10780898	6882786	10442445	6666094
Valid reads	(74.01%)	(85.24%)	(73.16%)	(88.08%)
rRNA	254459 (1.75%)	8362 (0.10%)	323537 (2.27%)	9018 (0.12%)
tRNA	47680 (0.33%)	833 (0.01%)	32260 (0.23%)	786 (0.01%)
snoRNA	2925 (0.02%)	191 (0.00%)	2198 (0.02%)	165 (0.00%)
snRNA	2671 (0.02%)	182 (0.00%)	1554 (0.01%)	115 (0.00%)
other Rfam RNA	20793 (0.14%)	948 (0.01%)	23402 (0.16%)	997 (0.01%)

682

683

Table 2. Summary of conserved and predicted miRNA.

Groups	Pre-miRNA	Unique miRNA	miRNA type
gp1	50	77	Conserved miRNA
gp2a	94	122	Conserved miRNA
gp2b	314	261	Conserved miRNA
gp3	35	40	Conserved miRNA
gp4	421	411	Novel miRNA

684

685

Table 3. Summary of the differentially expressed conserved and novel miRNAs.

MiRNA family	MiRNA name	N816S(Norm)	Ning5m(Norm)	p_value(t_test)	up/down
miR156	nta-miR156g_L+1	3732.40	6752.02	0.003	down
	han-MIR156b-p3_1ss3CT	107.99	67.05	0.003	up
MIR159	stu-miR156a	3732.40	6752.02	0.003	down
	nta-MIR159-p5_1ss2GT	21.74	90.63	0.002	down
	nta-miR159_R-3	384.38	160.79	0.047	up
miR162 miR166	csi-miR159a-5p_1ss20AC	36.01	74.86	0.008	down
	stu-miR162a-3p	5177.75	7769.93	0.001	down
	stu-miR166a-3p_1ss21CA	1369.44	2506.50	0.041	down
	stu-miR166b_1ss20TC	52460.31	71297.86	0.029	down
	bn-miR166a_L+2	49.73	83.04	0.013	down
	aqc-MIR166a-p3_2ss1GC20AC	26.72	19.50	0.025	up
	aqc-MIR166a-p5_2ss1GC20AC	26.72	19.50	0.025	up
	nta-miR166a_R+1	62.57	97.20	0.033	down
	nta-miR166a_R+1_1ss20CT	11.51	26.44	0.045	down

miR167	stu-miR167a-5p_1ss21AT	44.40	62.92	0.010	down
	gma-miR167a_1ss21AT	44.40	62.92	0.010	down
	aly-miR167a-5p_R+1_1ss21AT	0.97	6.41	0.001	down
	nta-miR167d_R+2	17.46	30.80	0.040	down
	mdm-miR167h_1ss22AT	0.97	6.41	0.001	down
	stu-miR167a-5p	3533.81	5033.84	0.019	down
MIR169	nta-MIR169p-p3_1ss13GA	15.55	4.98	0.033	up
MIR172	nta-MIR172e-p5_2ss14CG18TA	8.43	0.64	0.002	up
	nta-MIR172e-p3_2ss15CG19TA	5.31	0.00	0.048	up
	stu-miR172a-5p_3ss2TC8AC9AT	21.28	49.93	0.010	down
	nta-MIR172f-p5_2ss14CG18TA	8.43	0.64	0.002	up
	gra-miR172b_2ss8TC9AT	21.28	49.93	0.010	down
miR319	stu-miR319-3p_L+2R-2	25.71	11.69	0.018	up
	stu-miR319b_L+1R-1	4345.23	2869.03	0.029	up
MIR390	hex-MIR390a-p5_1ss21GT	186.32	99.94	0.030	up
	hex-MIR390b-p5_2ss10TC21GA	10.01	4.27	0.001	up
	hex-MIR390b-p5_1ss21GT	186.32	99.94	0.030	up
miR391	stu-miR391-5p_L-1R+2	726.92	508.13	0.009	up
miR393	stu-miR393-5p	84.78	133.01	0.020	down
	stu-miR393-3p	0.00	41.28	0.029	down
miR396	stu-miR396-3p_L-1	1503.77	1367.94	0.038	up
miR398	stu-miR398b-3p_2ss18CG19CT	1.86	12.21	0.020	down
	stu-miR398b-3p	0.00	2.70	0.021	down
MIR399	ppe-MIR399a-p5_2ss5AT18TC	7.11	0.00	0.006	up
	fve-MIR399b-p5_2ss2GT20GA	4.89	33.61	0.005	down
	mtr-MIR399b-p5_2ss11TC20TG	22.65	18.22	0.019	up
	ppe-MIR399a-p3_2ss5AT18TC	7.11	0.00	0.006	up
	stu-miR399j-3p_1ss21GA	0.00	2.84	0.015	down
miR482	stu-miR482a-5p_1ss12AG	158.27	274.34	0.008	down
MIR858	ppe-MIR858-p5_1ss4GA	611.66	386.05	0.013	up
miR1885	bra-miR1885b	14.61	18.27	0.007	down
miR3627	stu-miR3627-5p_R-1	8.89	0.00	0.048	up
MIR10528	sly-MIR10528-p3_2ss9GA19TC	7.81	0.00	0.048	up
MIR11334	cst-MIR11334-p5_2ss9TG18TC	22.54	13.15	0.025	up
MIR5141	rgl-MIR5141-p5	16.53	40.47	0.023	down
miR6024	stu-miR6024-3p_R-1_2ss20CT21CT	20.40	9.84	0.021	up
miR6149	nta-miR6149a_L+1R-1_1ss21GC	0.00	4.06	0.034	down
miR6478	ptc-miR6478_R-1_1ss20TA	4.45	0.77	0.035	up

	ptc-miR6478_R+3_1ss21GA	3.68	11.02	0.042	down
	ptc-miR6478_R+2_1ss21GA	198.84	273.50	0.000	down
	ptc-miR6478_R+1_1ss21GA	13.40	23.60	0.013	down
MIR8005	stu-MIR8005a-p3_1ss5GA	11.39	3.39	0.036	up
MIR8021	stu-MIR8021-p3_2ss15GA22TA	9.40	13.95	0.021	down
	stu-MIR8021-p3_2ss16TC22TG	26.11	35.95	0.013	down
miR8175	ath-miR8175_L+4	0.00	3.83	0.013	down
Novel miRNAs	PC-3p-155334_29	0.00	5.53	0.001	down
	PC-3p-216061_19	0.00	6.33	0.021	down
	PC-3p-71517_65	5.84	20.24	0.050	down
	PC-3p-9747_279	24.16	51.14	0.034	down
	PC-5p-101290_47	3.77	6.09	0.020	down
	PC-5p-117714_40	0.00	13.55	0.025	down
	PC-5p-119988_39	0.00	14.12	0.040	down
	PC-5p-12980_234	62.33	116.97	0.018	down
	PC-5p-347691_9	0.00	7.12	0.040	down
	PC-5p-567932_5	0.00	5.04	0.014	down
	PC-3p-145340_32	11.33	1.54	0.014	up
	PC-3p-9335_287	78.11	53.49	0.041	up
	PC-5p-117747_40	11.26	1.96	0.023	up
	PC-5p-153350_30	14.73	2.06	0.010	up
	PC-5p-154497_29	4.86	0.00	0.038	up
	PC-5p-160111_28	9.37	1.96	0.049	up
	PC-5p-58183_77	17.38	7.49	0.030	up

686

687 Note: The miRNA name is composed of the 1st known miRNA name in a cluster, a
688 underscore, and a matching annotation: such as L-n/+n means the detected miRNA is n base
689 less or more than known rep_miRSeq in the left side; R-n/+n means the detected miRNA is n
690 base less or more than known rep_miRSeq in the right side; 2ss5TC13TA means 2
691 substitution (ss), which are T->C at position 5 and T->A at position 13.

692

693 **Supporting Information:**

694 Figure S1. Small RNAs Pearson correlation between the six libraries

695 Table S1. Primers used in this study. S1-1, Primers designed for conserved
696 miRNAs and novel miRNAs; S1-2 Primers designed for miRNA targets.

697 Table S2. Summary of five types of miRNA in this study. S2-1, gp1a type; S2-2,
698 gp2a type; S2-3, gp2b type; S2-4, gp3 type; S2-5, gp4 type.

699 Table S3. Primers used in this study. S1-1, Primers designed for conserved
700 miRNAs and novel miRNAs; S1-2, Primers designed for miRNA targets.

701 Table S4. Overview of degradome sequencing reads from raw data to mapping
702 sequences.

703 Table S5. The overview of reliable identified targets of miRNAs.

704 Table S6. Transcript annotation of targets of differential expression conserved
705 and novel miRNAs in this study.

706 Table S7. The pathways of all the miRNAs and targets involved in this study.
 707 Table S8. The transcript annotation, GO terms and KEGG pathways of
 708 differential expression miRNAs with those targets in pepper.
 709