1 Neutrophils are required during immunization with the pneumococcal conjugate vaccine
2 for protective antibody responses and host defense against infection

3 Essi Y. I. Tchalla, Elizabeth A. Wohlfert and Elsa N. Bou Ghanem^{1,2,3}

4 Author affiliations: Department of Microbiology and Immunology, University at Buffalo

5 School of Medicine, Buffalo, NY 14203, USA

6 Running title: PMNs induce antibodies post vaccination

7 Address correspondence to: Elsa N. Bou Ghanem, 955 Main Street, Buffalo, NY, 14203.

8 Telephone: 716-829-2422. Email: elsaboug@buffalo.edu

9 Word counts: Abstract 150; Manuscript 3,499

10

11

12

13

14

15

¹ The authors declare no conflict of interest

² Grant support: This work is in part supported by National Institute of Health grant R00AG051784 to ENBG.

³ Address correspondence and reprint requests to: Elsa N. Bou Ghanem, 955 Main Street, Buffalo, NY, 14203. Telephone: 716-829-2422. Email: elsaboug@buffalo.edu

Abstract

Neutrophils can shape adaptive immunity, however their role in vaccine-induced protection against infections *in vivo* remains unclear. Here, we tested their role in the clinically relevant polysaccharide conjugate vaccine against *Streptococcus pneumoniae* (pneumococcus). We antibody depleted neutrophils during vaccination, allowed them to recover, and four weeks later challenged mice with pneumococci. We found that while isotype-treated vaccinated controls were protected against an otherwise lethal infection in naïve mice, full protection was lost upon neutrophil depletion. Compared to vaccinated controls, neutrophil-depleted mice had higher lung bacterial burdens, increased incidence of bacteremia and lower survival rates. Sera from neutrophil-depleted mice had less anti-pneumococcal IgG2c and IgG3, were less efficient at inducing opsonophagocytic killing of bacteria by neutrophils *in vitro* and worse at protecting naïve mice against pneumococcal pneumonia. In summary, neutrophils are required during vaccination for optimal host protection, which has important implications for future vaccine design against pneumococci and other pathogens.

Key words: Neutrophils, antibodies, vaccines, *Streptococcus pneumoniae*

Introduction

S. pneumoniae are Gram-positive bacteria with >90 serotypes based on capsular polysaccharides [1]. These bacteria can cause pneumonia, meningitis and bacteremia [2] and remain a serious cause of mortality and morbidity worldwide, particularly in the elderly [3]. Currently, two vaccines covering common disease-causing bacterial serotypes, are available [4]. The pneumococcal polysaccharide vaccine (PPSV) consists of polysaccharides that directly cross-link B cell receptors on mature B cells leading to antibody (Ab) production independent of T cells [5]. PPSV is routinely recommended for elderly individuals >65 years old and adults with medical conditions [6]. As children <2 years old lack mature B cells, they fail to produce T-independent Abs [7]. Therefore, the pneumococcal conjugate vaccine (PCV), was introduced for use in children. PCV contains polysaccharides linked to a carrier protein that triggers a T-dependent Ab response [4]. PCV has had great efficacy in children and is currently recommended for use in immunocompromised adults and elderly individuals with underlying conditions [6]. As PCV is recommended for adults with compromised immunity including B and T cell responses [4], it is important to elucidate novel players in vaccines that could be potential targets to boost protection.

Abs against capsular polysaccharides following vaccination bind to *S. pneumoniae* and protect the host against infection [4]. The functionality of Abs is determined by their antigen affinity. Affinity to antigens is mediated via the variable regions which make up the Fab or antigen binding portions and is optimized by somatic hypermutation (SHM) [8]. The Fc or constant region of Ab, which determines their class, also shapes their function, with the different classes of Abs having distinct immune

modulating activities [8]. Abs against T-independent antigens such as bacterial polysaccharides are typically produced by marginal zone B cells in the spleen [9]. In contrast, T cell-dependent Ab production occurs in germinal centers, where a specialized subset of CD4⁺ T-follicular helper cells (TFH) [10] induce B cells to undergo class-switch recombination and SHM resulting in Abs with improved function [11]. PCV significantly boosts class switching to IgG as compared to PPSV [12] and further induces TFH cells which correlate with enhanced Ab function [13].

Polymorphonuclear leukocytes (PMNs) or neutrophils, play a crucial role in innate immunity to infections [14]. It is now appreciated that PMNs can also regulate adaptive immunity. PMNs can directly induce Ab production by B cells [11]. In the spleen, a subset of PMNs termed B helper neutrophils was described to produce APRIL, BAFF and IL-21 [9] that triggered Ab production by marginal-zone B cells [9, 15]. This was described for T-independent antigens including bacterial polysaccharides [9]. PMNs may also affect T cell dependent Ab responses [15], however, that is less established. PMNs are thought to both activate and suppress T cells [16]. PMNs produce a repertoire of chemokines that recruit T-cells and also produce cytokines that drive T-cell subset differentiation [16]. PMN-derived products can prime T cells to more efficiently respond to antigens [17]. PMNs also activate T cells via recruiting and activating antigen presenting cells [18] or acting as antigen presenting cells themselves [16, 19-23]. In contrast, PMNs produce compounds that inhibit T cell activation [24] including ROS [25], arginase-1 [26] and serine proteases [27]. Therefore, it is unclear whether upon *in vivo* vaccination, if PMNs would suppress or induce T-dependent Ab responses. Further, although there have been elegant studies characterizing mechanisms of PMN interactions

with B and T cells, most of the work has been done either *in vitro* or *in vivo* using model antigens [11, 18, 24]. Thus, studies examining the role of PMNs in clinically relevant vaccinations and how that shapes protection against *in vivo* infections are needed.

PMNs are required to control bacterial numbers following *S. pneumoniae* infection [28, 29]. PMNs also play a role in Ab responses against pneumococci. When compared to healthy controls, patients with neutropenic disorders had lower levels of Abs to some pneumococcal polysaccharides [9]. In mice, splenic PMNs localized with marginal zone B cells and were required for production of T-independent Abs during pneumococcal infection [15, 30]. However, whether PMNs shape responses to the pneumococcal conjugate vaccine and if they impair or promote Ab production remains unexplored. Here we tested the role of PMNs in response to PCV and found they were required at the time of vaccination for optimal Ab responses as well as host protection against pneumococcal infection. This study highlights the link between PMNs and Ab responses in the context of a clinically relevant immunization, which has far-reaching implications for vaccine design against *S. pneumoniae* and other pathogens.

Methods

Mice

Female C57BL/6 mice (6-8 weeks) were purchased from Jackson Laboratories (Bar Harbor, ME) and used in all experiments. Mice were housed in a specific-pathogen free facility at the University at Buffalo and all experiments were conducted in accordance with Institutional Animal Care and Use Committee (IACUC) guidelines.

Bacteria

Wild type (WT) *S. pneumoniae* TIGR4 and capsule-deletion mutant (Δcps) *S. pneumoniae* were kind gifts from Andrew Camilli. All bacteria were grown to mid-exponential phase in Todd-Hewitt broth (BD Biosciences) supplemented with Oxyrase (Oxyrase) and 0.5% yeast extract at 37°C in 5% CO₂. Aliquots were frozen at -80°C in growth media with 20% (v/v) glycerol. Prior to use, aliquots were thawed on ice, washed and diluted in PBS to the desired concentrations. Titers were confirmed by plating on Tryptic Soy Agar plates supplemented with 5% sheep blood agar (Northeast Laboratory Services).

Immunization

Mice were immunized via intramuscular (*i.m.*) injection of 50µl of the pneumococcal conjugate vaccine Prevnar-13® (Wyeth pharmaceuticals) into the caudal thigh muscle. Sera was collected from all mice prior to immunization, as well as two and four weeks post immunization and saved at -80°C for subsequent assays.

Neutrophil Depletion

Mice were treated intra-peritoneally (*i.p.*) with 50 µg of the Ly6G-depleting antibody IA8 or isotype IgG control (BioXCell) following the timeline in Fig 1A.

Adoptive Transfer of Sera

Five weeks following immunization, vaccinated, vaccinated PMN-depleted and naïve mice were euthanized and blood harvested via cardiac puncture. Sera was obtained from the blood, pooled for each group and transferred *i.p.* (250µl) into naïve recipients. Recipients were then infected one hour later [31].

Animal Infections

Mice were intra-tracheally (*i.t.*) challenged with 10⁷ colony-forming units (CFU) of WT

S. pneumoniae as previously described [31]. Following infection, one set of mice were monitored daily over one week for bacteremia as well as clinical signs of disease including weight loss, activity level, posture and breathing and blindly scored from 0 (healthy) to 21 (severely sick). Twenty-four hours post infection, another set of mice were euthanized and lung and blood were assessed for CFU.

Antibody ELISA

Sera Ab levels were measured by ELISA as previously described [31]. Nunc maxisorp® plates were coated overnight at 4°C with type 4 Pneumococcal Polysaccharide (ATCC®) at 2µg/well. Plates were washed and blocked for 2 hours. The sera were preabsorbed with a pneumococcal cell wall polysaccharide mixture (CWP-multi from Cederlane) to neutralize non-capsular Abs and then added to the plate. After a 3h incubation and washing, pneumococcal-specific Abs were detected using HRP-conjugated goat anti-mouse IgM (Invitrogen), IgG (Millipore Sigma), IgG1, IgG2b, IgG2c or, IgG3 (Southern Biotech) followed by TMB substrate (Thermo Scientific™) and readings at OD₆₅₀ using a BioTek® reader. Kinetic ELISAs were performed with readings every minute for 10 minutes. Ab units were calculated as percentages of a control hyperimmune serum included in each ELISA. Hyperimmune sera was pooled from mice that were intranasally inoculated with *S. pneumoniae* TIGR4 over four weeks as previously described [31], immunized with PCV at week 4 and injected i.p. with heat-killed bacteria at week 5.

Myeloperoxidase (MPO) ELISA

MPO levels were measured in lung homogenates using the Mouse Myeloperoxidase ELISA kit from Invitrogen™ as per manufacturer's instructions.

Isolation of PMNs

PMNs were isolated from the bone-marrow using density centrifugation with Histopaque 1119 and Histopaque 1077 (Sigma) as previously described [28]. PMNs were resuspended in Hanks' Balanced Salt Solution (HBSS without Ca^{2+} and Mg^{2+}) supplemented with 0.1% gelatin and used in subsequent experiments. Purity was confirmed by flowcytometry and the isolated cells were 85-90% Ly6G⁺.

Opsonophagocytic Killing Assay (OPH)

The ability of PMNs to kill pneumococci was assessed as previously described [28]. Briefly, 1×10^5 PMNs were infected with 1×10^3 bacteria pre-opsonized with 3% mouse sera in 100 μl reaction volumes of HBSS/0.1% gelatin (with Ca^{2+} and Mg^{2+}) and rotated at 37°C for 40 minutes. The reactions were stopped on ice and plated for CFU. The percent of bacteria killed was calculated using no PMN controls.

Flow Cytometry

One day following the last PMN depletion, mice were euthanized and blood, vaccine draining popliteal lymph nodes and spleen were harvested. Single-cell suspensions of splenocytes and lymph nodes were prepared by mashing the organs through sterile mesh screens using the plunger of a 3-ml syringe. Red blood cells were lysed with a hypotonic buffer and the cells surface stained for Ly6G (IA8 or RB6, Biolegend), CD11b (M1/70, Invitrogen), CD11c (N418, BD Bioscience), F4/80 (BM8, BD Bioscience) and Ly6C (AL-21, BD Bioscience) in the presence of Fc-block (BD Bioscience). Live cells were identified using a dead cell stain kit (Life Technologies). Fluorescence intensities were measured on a BD Fortessa and at least 20,000 events were analyzed using FlowJo.

Statistical Analysis

All statistical analysis was done using Graphad Prism version 8. Significant differences were determined by Fisher's exact test, One-way ANOVA followed by Dunnet's test or Student's t-test as appropriate. Survival analyses were performed using the log-rank (Mantel-Cox) test. p values less than 0.05 were considered significant.

Results

PMNs are required at the time of immunization with PCV for host protection against pneumococcal infection

To test if PMNs were required for protection at the time of vaccination with PCV, we used the anti-Ly6G Ab IA8 to deplete PMNs or isotype controls one day prior to and every two days throughout the first week following vaccination (timeline- Fig 1A). One day after the final treatment, we verified depletion in the blood and found ~99% reduction in the number of circulating PMNs (Fig S1A). As splenic PMNs have a role in Ab production [9, 15], we also examined PMN numbers in the spleen. We found that upon vaccination there was a ~2-fold increase in splenic PMNs (Fig S1B) and that treatment with depleting Abs resulted in ~98% depletion of those cells (Fig S1B). We also verified that Ab treatment was specific to PMNs and did not result in any changes in the number of circulating and splenic monocytes, dendritic cells or macrophages (Fig S2).

Four weeks following vaccination, we challenged mice i.t. with *S. pneumoniae* TIGR4 strain. Invasive *S. pneumoniae* infection results in pneumonia primarily, but up to 30% of patients with pneumococcal pneumonia also develop bacteremia and have a worse prognosis [32]. As *S. pneumoniae* strains can differ considerably [33], we chose the well-characterized serotype 4 isolate TIGR4, originally isolated from a bacteremic patient, as a

model of a highly invasive infection modeling pneumonia that results in bacteremia [34] and that is covered by PCV. We then monitored the disease course and as expected found that while all naïve mice rapidly succumbed to infection, 100% of vaccinated mice survived (Fig 1B). However, unlike vaccinated controls, full protection was lost in PMN-depleted mice (Fig 1). The majority of PMN-depleted mice displayed severe clinical signs of disease where 77.8% got sick as compared to only 12.5% of vaccinated controls (Fig 1C). PMN-depleted mice had between 10-100-fold higher pulmonary bacterial numbers (Fig 2A) and systemic spread (Fig 2B) into the circulation, culminating in significantly reduced survival (Fig 1B) as compared to vaccinated controls. This reduced protection in PMN-depleted mice was not due to the continued absence of PMNs at the time of challenge, as we verified that there was no difference in PMN presence in the lungs as measured by MPO levels following infection (Fig S1C). Rather, our findings suggest that PMNs are required at the time of vaccination with PCV for full protection against subsequent pneumococcal infection.

PMNs are required for optimal antibody isotype switching in response to PCV immunization

Next, we wanted to explore the mechanisms by which PMNs contributed to vaccine-induced protection. We first examined Ab production and as expected observed isotype switching to IgG by week 4 post vaccination in our control group (Fig 3). We found that PMN depletion did not alter IgM or total IgG levels against capsular polysaccharide type 4 (Fig 3A and B) or heat-killed *S. pneumoniae* (not shown). However, when we examined IgG subtypes, we found that depletion of PMNs during

vaccination resulted in slightly reduced levels of IgG2b (Fig 4A) and significantly lower levels of IgG2c (Fig 4B) and IgG3 (Fig 4C) at week 4 post vaccination. Interestingly, IgG1 levels (Fig 4D) were slightly, but not significantly elevated in PMN depleted group as compared to vaccinated controls. These data suggest that PMNs play a role in class switching to certain IgG subtypes.

PMNs are required for optimal antibody function following PCV immunization

Apart from Ab levels, Ab function is key for vaccine-efficacy [35]. We next explored if PMNs affected Ab affinity to bacterial surfaces. We tested the ability of IgG in the sera of the different mouse groups to bind the surface of *S. pneumoniae* by flow cytometry. Very little IgG bound to bacteria upon incubation with naïve sera. However, we observed a 30-fold increase in the amount of IgG bound to bacteria when sera from vaccinated mice were used (Fig 5A). As expected, in immune sera, the bound IgG was specific to capsular polysaccharides as very little IgG bound to acapsular bacteria (*Δcps* *S. pneumoniae*). Interestingly, we observed a significant decrease in the amount of IgG bound to *S. pneumoniae* opsonized with sera from PMN-depleted mice as compared sera from vaccinated controls (Fig 5A).

We next compared the opsonic capacity of Abs by comparing the ability of sera to induce opsonophagocytic (OPH) killing of *S. pneumoniae* by primary PMNs isolated from naïve mice. We found sera from vaccinated controls significantly boosted bacterial killing by PMNs where 60% of the bacterial input were killed by PMNs in the presence of immune sera as compared to ~10% with naïve sera (Fig 5B). Strikingly, sera from PMN-depleted mice failed to induce opsonophagocytic killing of *S. pneumoniae* by

PMNs where only 3% of the bacterial input was killed (Fig 5B). Abs can also activate the complement pathway and directly kill bacteria [8] but we detected no differences in the ability of sera alone from any of the mouse groups to kill pneumococci (Fig S3).

Given the difference in the *in vitro* function we observed, we finally tested the protective activity of Abs generated upon vaccination in the absence of PMNs. Naïve young mice were injected i.p. with five-week sera from either vaccinated controls, naïve mock-immunized mice or our PMN depleted vaccinated group. Mice were then challenged i.t. with *S. pneumoniae* TIGR4 one hour following sera transfer. We found that while all of the mice receiving naïve sera succumbed to infection, all of the mice receiving sera from vaccinated controls survived the challenge (Fig 5C). In contrast, only half of the mice receiving sera from the PMN depleted vaccinated group survived (Fig 5C). These data indicate that Abs produced during vaccination in the absence of PMNs are not sufficient to provide protection against subsequent pneumococcal infection.

Discussion

Traditionally, PMNs are viewed as effectors of vaccine responses where vaccination triggers Abs that bind to pathogens and promote their clearance via enhancing uptake and killing by PMNs [36]. However, the extent to which PMNs contribute to vaccine mediated protection against infections *in vivo* has not been fully elucidated. In this study, we explored the role of PMNs in immunization with the pneumococcal conjugate vaccine. We found that PMNs were needed for production of functional Abs following vaccination. Importantly, PMNs were required at the time of immunization for full protection against subsequent invasive pneumococcal infection.

Our findings highlight the *in vivo* role of PMNs as inducers of protective vaccine responses against *S. pneumoniae* infections.

The mechanisms by which PMNs mediate Ab production in response to PCV is unclear. In adults, polysaccharides can directly cross-link B cell receptors and elicit Ab production independent of T cells [5]. PCV converts this T-independent response to one that involves T cells as it consists of polysaccharides linked to the carrier protein CRM₁₉₇ [5]. This generates T cells specific to the carrier protein [37, 38]. When B cells recognize polysaccharides, they are thought to bind and internalize the polysaccharide along with its protein carrier and then display peptides derived from the carrier on MHC-II. This allows these polysaccharide-specific B cells to interact with carrier-peptide specific T cells which in turn help the B cells produce anti-polysaccharide Abs [39]. Therefore, in the context of PCV, PMNs could either be working on B cells, T cells or both. In humans, a subset of splenic PMNs directly induce Ab production by marginal zone B cells in response to T-independent antigens including bacterial polysaccharides [9]. In mice, splenic B helper PMNs were found to produce pantrexin3, which was important for IgM production following the immunization with the unconjugated pneumococcal polysaccharide vaccine [15]. Pantrexin3 was also important for T-cell independent IgM and IgG production against polysaccharides following intravenous infection with *S. pneumoniae* [15]. The role of PMNs in T-dependent responses is less clear. In mice, PMNs impaired IgA but not IgG or IgM production in response to vaccination with the adjuvant *Bacillus anthracis* edema toxin [40]. Mouse PMNs were found to directly present ovalbumin peptides to CD4⁺ T cells triggering T cell cytokine production and proliferation [20]. However, IgG2 responses to ovalbumin were not impaired in

pantrexin3^{-/-} mice [15]. In contrast, Abs against influenza PR8 and the T-dependent antigen TNP-Ficoll required pantrexin3 production by murine PMNs [15]. Similarly, human PMNs were able to present influenza hemagglutinin to CD4⁺ T cells [22]. In rhesus macaques, PMNs presented HIV-envelope glycoproteins to CD4⁺ T cells [22] and induced Ab production against SIV when co-cultured with B cells [41]. Here, in the context of immunization with PCV, PMNs are clearly required for optimal Ab responses, however whether they are acting on T cells remains to be determined.

A key finding here is that PMNs are required for the production of functional Abs. The functionality of Abs is determined by the affinity and avidity to their antigen [8]. Here, although we detected similar levels of IgM and IgG in sera from PMN-depleted and isotype-treated vaccinated mice, the ability of IgG to bind pneumococci significantly decreased when they were generated in the absence of PMNs. This suggests that Ab affinity is increased in the presence of PMNs. Ab affinity is improved by SHM of the Fab variable regions and typically occurs in germinal centers and requires T cells [42], although human splenic B helper PMNs may contribute to SHM in marginal zone B cells [9].

Ab function is also influenced by their subclass which is determined by their Fc region [8]. The Fc portion of Abs shape effector function since they determine binding to Fc receptors on PMNs as well as the ability to activate complement [8]. Here, PMNs were required for class switching to IgG2c and IgG3 but not IgG1 subtypes. The IgG subtype produced in response to PCV varies in humans based on age, with IgG2 being the predominant response in

adults [12, 43]. We found that PMNs were crucial for the ability of Abs to elicit opsonophagocytic killing of bacteria by primary immune cells. This is in line with data from humans where although the opsonophagocytic activity of IgG subtypes vary based on bacterial serotype, IgG2 was reported to have the highest activity while IgG1 had the lowest activity against serotype 4 pneumococci [44]. We also found that PMNs were required for production of Abs that protect against infection. This could be mediated by IgG2 [44] or IgG3 which was shown to be protective in mice against pneumococcal infection [45].

How PMNs are inducing class-switching to IgG2c and IgG3 but not IgG1 in response to PCV is unclear. Efficient class switching from IgM to IgG requires T cell help [39]. In adults, PCV significantly boosts class switching to IgG as compared to the unconjugated polysaccharide vaccine [12]. Cytokines produced by T cells further determine the subtype of IgG produced with IL-17 and IFN- γ enhancing switching to IgG3 and IgG2 more than IgG1 [46, 47]. As PMNs can both produce cytokines [16] and drive Th1 and Th17 cell differentiation [20], they may contribute to isotype switching by producing IFN- γ or IL-17 themselves or eliciting T cells to do so.

In summary, we demonstrate here that PMNs are required at the time of immunization with the pneumococcal conjugate vaccine for optimal protective Ab responses and host protection against subsequent *S. pneumoniae* infection. As serotype replacement by bacterial strains not covered by the current vaccines continue to emerge, novel serotype-independent vaccine formulations such as whole cell vaccines or common pneumococcal protein vaccines are being considered [48]. Therefore, future vaccine designs should take PMN responses into consideration, particularly in susceptible

populations like the elderly [2], where PMN responses are known to be dysregulated [49].

Author Contributions: EYIT conducted research, analyzed data and wrote paper. EAW provided essential reagents and expertise. ENBG designed research, wrote paper and had responsibility for final content. All authors read and approved the final manuscript.

Funding: This work was in part supported by National Institute of Health grant R00AG051784 to ENBG.

Acknowledgements: We would like to acknowledge Manmeet Bhalla for critical reading and discussion of the manuscript.

Conflict of Interest: The authors declare no conflict of interest

References:

1. Kadioglu A, Weiser JN, Paton JC, Andrew PW. The role of *Streptococcus pneumoniae* virulence factors in host respiratory colonization and disease. *Nature reviews Microbiology* **2008**; 6:288-301.
2. Chong CP, Street PR. Pneumonia in the elderly: a review of the epidemiology, pathogenesis, microbiology, and clinical features. *South Med J* **2008**; 101:1141-5; quiz 32, 79.
3. Henig O, Kaye KS. Bacterial Pneumonia in Older Adults. *Infectious disease clinics of North America* **2017**; 31:689-713.

- 360 4. Chen WH, Kozlovsky BF, Effros RB, Grubeck-Loebenstein B, Edelman R, Szein
361 MB. Vaccination in the elderly: an immunological perspective. *Trends Immunol* **2009**;
362 30:351-9.
- 363 5. Adderson EE. Antibody repertoires in infants and adults: effects of T-independent and
364 T-dependent immunizations. *Springer seminars in immunopathology* **2001**; 23:387-403.
- 365 6. Matanock A LG, Gierke R, Kobayashi M, Leidner A, Pilishvili T. Use of 13-Valent
366 Pneumococcal Conjugate Vaccine and 23-Valent Pneumococcal Polysaccharide Vaccine
367 Among Adults Aged ≥ 65 Years: Updated Recommendations of the Advisory Committee
368 on Immunization Practices. *MMWR Morb Mortal Wkly Rep* 2019;68:1069–1075. :
369 CDC, **2019**.
- 370 7. Laferriere C. The immunogenicity of pneumococcal polysaccharides in infants and
371 children: a meta-regression. *Vaccine* **2011**; 29:6838-47.
- 372 8. Vidarsson G, Dekkers G, Rispens T. IgG subclasses and allotypes: from structure to
373 effector functions. *Frontiers in immunology* **2014**; 5:520.
- 374 9. Puga I, Cols M, Barra CM, et al. B cell-helper neutrophils stimulate the diversification
375 and production of immunoglobulin in the marginal zone of the spleen. *Nature*
376 *immunology* **2011**; 13:170-80.
- 377 10. Vinuesa CG, Linterman MA, Yu D, MacLennan IC. Follicular Helper T Cells.
378 *Annual review of immunology* **2016**; 34:335-68.
- 379 11. Cerutti A, Puga I, Magri G. The B cell helper side of neutrophils. *J Leukoc Biol* **2013**;
380 94:677-82.

12. Soininen A, Seppala I, Nieminen T, Eskola J, Kayhty H. IgG subclass distribution of antibodies after vaccination of adults with pneumococcal conjugate vaccines. *Vaccine* **1999**; 17:1889-97.
13. Sterrett S, Peng BJ, Burton RL, et al. Peripheral CD4 T follicular cells induced by a conjugated pneumococcal vaccine correlate with enhanced opsonophagocytic antibody responses in younger individuals. *Vaccine* **2020**.
14. Deniset JF, Kubes P. Neutrophil heterogeneity: Bona fide subsets or polarization states? *J Leukoc Biol* **2018**; 103:829-38.
15. Chorny A, Casas-Recasens S, Sintes J, et al. The soluble pattern recognition receptor PTX3 links humoral innate and adaptive immune responses by helping marginal zone B cells. *The Journal of experimental medicine* **2016**; 213:2167-85.
16. Kalyan S, Kabelitz D. When neutrophils meet T cells: beginnings of a tumultuous relationship with underappreciated potential. *European journal of immunology* **2014**; 44:627-33.
17. Tillack K, Breiden P, Martin R, Sospedra M. T lymphocyte priming by neutrophil extracellular traps links innate and adaptive immune responses. *J Immunol* **2012**; 188:3150-9.
18. Mantovani A, Cassatella MA, Costantini C, Jaillon S. Neutrophils in the activation and regulation of innate and adaptive immunity. *Nature reviews Immunology* **2011**; 11:519-31.
19. Matsushima H, Geng S, Lu R, et al. Neutrophil differentiation into a unique hybrid population exhibiting dual phenotype and functionality of neutrophils and dendritic cells. *Blood* **2013**; 121:1677-89.

20. Abi Abdallah DS, Egan CE, Butcher BA, Denkers EY. Mouse neutrophils are professional antigen-presenting cells programmed to instruct Th1 and Th17 T-cell differentiation. *International immunology* **2011**; 23:317-26.
21. Calabro S, Tortoli M, Baudner BC, et al. Vaccine adjuvants alum and MF59 induce rapid recruitment of neutrophils and monocytes that participate in antigen transport to draining lymph nodes. *Vaccine* **2011**; 29:1812-23.
22. Vono M, Lin A, Norrby-Teglund A, Koup RA, Liang F, Lore K. Neutrophils acquire the capacity for antigen presentation to memory CD4(+) T cells in vitro and ex vivo. *Blood* **2017**; 129:1991-2001.
23. Beauvillain C, Cunin P, Doni A, et al. CCR7 is involved in the migration of neutrophils to lymph nodes. *Blood* **2011**; 117:1196-204.
24. Muller I, Munder M, Kropf P, Hansch GM. Polymorphonuclear neutrophils and T lymphocytes: strange bedfellows or brothers in arms? *Trends Immunol* **2009**; 30:522-30.
25. Schmielau J, Finn OJ. Activated granulocytes and granulocyte-derived hydrogen peroxide are the underlying mechanism of suppression of t-cell function in advanced cancer patients. *Cancer research* **2001**; 61:4756-60.
26. Munder M, Schneider H, Luckner C, et al. Suppression of T-cell functions by human granulocyte arginase. *Blood* **2006**; 108:1627-34.
27. Bank U, Reinhold D, Schneemilch C, Kunz D, Synowitz HJ, Ansorge S. Selective proteolytic cleavage of IL-2 receptor and IL-6 receptor ligand binding chains by neutrophil-derived serine proteases at foci of inflammation. *Journal of interferon & cytokine research : the official journal of the International Society for Interferon and Cytokine Research* **1999**; 19:1277-87.

28. Bou Ghanem EN, Clark S, Roggensack SE, et al. Extracellular Adenosine Protects against *Streptococcus pneumoniae* Lung Infection by Regulating Pulmonary Neutrophil Recruitment. *PLoS pathogens* **2015**; 11:e1005126.
29. Bhowmick R, Tin Maung NH, Hurley BP, et al. Systemic disease during *Streptococcus pneumoniae* acute lung infection requires 12-lipoxygenase-dependent inflammation. *J Immunol* **2013**; 191:5115-23.
30. Deniset JF, Surewaard BG, Lee WY, Kubes P. Splenic Ly6G(high) mature and Ly6G(int) immature neutrophils contribute to eradication of *S. pneumoniae*. *The Journal of experimental medicine* **2017**; 214:1333-50.
31. Bou Ghanem EN, Maung NHT, Siwapornchai N, et al. Nasopharyngeal Exposure to *Streptococcus pneumoniae* Induces Extended Age-Dependent Protection against Pulmonary Infection Mediated by Antibodies and CD138(+) Cells. *J Immunol* **2018**; 200:3739-51.
32. Lin SH, Lai CC, Tan CK, Liao WH, Hsueh PR. Outcomes of hospitalized patients with bacteraemic and non-bacteraemic community-acquired pneumonia caused by *Streptococcus pneumoniae*. *Epidemiology and infection* **2011**; 139:1307-16.
33. Kadioglu A, Andrew PW. Susceptibility and resistance to pneumococcal disease in mice. *Briefings in functional genomics & proteomics* **2005**; 4:241-7.
34. Aaberge IS, Eng J, Lemark G, Lovik M. Virulence of *Streptococcus pneumoniae* in mice: a standardized method for preparation and frozen storage of the experimental bacterial inoculum. *Microbial pathogenesis* **1995**; 18:141-52.
35. LaFon DC, Nahm MH. Measuring immune responses to pneumococcal vaccines. *Journal of immunological methods* **2018**; 461:37-43.

36. Simell B, Vuorela A, Ekstrom N, et al. Aging reduces the functionality of anti-pneumococcal antibodies and the killing of *Streptococcus pneumoniae* by neutrophil phagocytosis. *Vaccine* **2011**; 29:1929-34.
37. Lai Z, Schreiber JR. Antigen processing of glycoconjugate vaccines; the polysaccharide portion of the pneumococcal CRM(197) conjugate vaccine co-localizes with MHC II on the antigen processing cell surface. *Vaccine* **2009**; 27:3137-44.
38. Wuorimaa T, Kayhty H, Eskola J, Bloigu A, Leroy O, Surcel HM. Activation of cell-mediated immunity following immunization with pneumococcal conjugate or polysaccharide vaccine. *Scand J Immunol* **2001**; 53:422-8.
39. Rappuoli R, De Gregorio E, Costantino P. On the mechanisms of conjugate vaccines. *Proc Natl Acad Sci U S A* **2019**; 116:14-6.
40. Jee J, Bonnegarde-Bernard A, Duverger A, et al. Neutrophils negatively regulate induction of mucosal IgA responses after sublingual immunization. *Mucosal immunology* **2015**; 8:735-45.
41. Musich T, Rahman MA, Mohanram V, et al. Neutrophil Vaccination Dynamics and Their Capacity To Mediate B Cell Help in Rhesus Macaques. *J Immunol* **2018**; 201:2287-302.
42. Lau AW, Brink R. Selection in the germinal center. *Curr Opin Immunol* **2019**; 63:29-34.
43. Lottenbach KR, Mink CM, Barenkamp SJ, Anderson EL, Homan SM, Powers DC. Age-associated differences in immunoglobulin G1 (IgG1) and IgG2 subclass antibodies to pneumococcal polysaccharides following vaccination. *Infect Immun* **1999**; 67:4935-8.

44. Mikolajczyk MG, Concepcion NF, Wang T, et al. Characterization of antibodies to capsular polysaccharide antigens of *Haemophilus influenzae* type b and *Streptococcus pneumoniae* in human immune globulin intravenous preparations. *Clinical and diagnostic laboratory immunology* **2004**; 11:1158-64.
45. Briles DE, Claflin JL, Schroer K, Forman C. Mouse IgG3 antibodies are highly protective against infection with *Streptococcus pneumoniae*. *Nature* **1981**; 294:88-90.
46. Mitsdoerffer M, Lee Y, Jager A, et al. Proinflammatory T helper type 17 cells are effective B-cell helpers. *Proc Natl Acad Sci U S A* **2010**; 107:14292-7.
47. Kawano Y, Noma T, Yata J. Regulation of human IgG subclass production by cytokines. IFN-gamma and IL-6 act antagonistically in the induction of human IgG1 but additively in the induction of IgG2. *J Immunol* **1994**; 153:4948-58.
48. Pichichero ME. Pneumococcal whole-cell and protein-based vaccines: changing the paradigm. *Expert Rev Vaccines* **2017**; 16:1181-90.
49. Boe DM, Boule LA, Kovacs EJ. Innate immune responses in the ageing lung. *Clinical and experimental immunology* **2017**; 187:16-25.

Legends

Figure 1. PMNs are required at the time of vaccination for PCV-mediated protection against *S. pneumoniae* infection. C57BL/6 female mice were treated i.p. with PMN depleting antibodies (IA8) or isotype control at days -1, +1, +4 and +7 with respect to vaccination following the timeline outlined in panel A. Mice were mock treated (naïve) or administered 50µl of Prevnar-13 via intramuscular injections to the hind legs (vaccinated).

Four weeks following vaccination mice were challenged i.t. with 1×10^7 CFU *S. pneumoniae* TIGR4 and monitored for survival over time (B) and clinical signs of disease (C). (B) Data were pooled from 14 mice/group from three separate experiments and * denotes significance calculated by the log-Rank (Mantel-Cox) test. (C) Data were pooled from three separate experiments with each square representing an individual mouse. The dashed line indicates the symptomatic score threshold (above one). Fractions indicate the percent of mice that had a score above 1 and * denotes significant differences from vaccinated controls by Fisher's exact test.

Figure 2. PMNs are required at the time of PCV vaccination for subsequent control of *S. pneumoniae* burden upon pulmonary challenge. Naïve (green), Prevnar-13 immunized (black) and PMN depleted Prevnar-13 immunized mice (blue) were challenged i.t. with 1×10^7 CFU *S. pneumoniae* TIGR4 four weeks following vaccination following the timeline in Fig1A. Bacterial burden in the lungs (A) and blood (B) were also enumerated 24 hours post infection. Data were pooled from three separate experiments with each square representing an individual mouse. * denotes significant differences from vaccinated controls by One-way ANOVA followed by Dunnet's test.

Figure 3. Total levels of anti-pneumococcal IgG and IgM remain unchanged in PMN-depleted PCV immunized mice. Sera were collected from naïve (green lines), Prevnar-13 immunized (black lines) and PMN depleted Prevnar-13 immunized mice (blue lines) over time as indicated in Fig 1A. Circulating levels of IgM (A) and total IgG (B) against purified polysaccharide serotype 4 were then measured by ELISA. Antibody units were

calculated based on a hyperimmune standard (see Materials and Methods) included in each ELISA plate. p values were determined by student t-test. Asterisks ($p<0.05$) indicate significant differences with respect to vaccinated mice. Data were pooled from two separate experiments with $n=6$ mice per group and presented as means $\pm$ SD.

Figure 4. PMNs contribute to IgG2 and IgG3 production following PCV immunization. Sera were collected from naïve (green lines), Pevnar-13 immunized (black lines) and PMN depleted Pevnar-13 immunized mice (blue lines) following the timeline presented in Fig 1A. (A-D) The levels of the indicated antibodies against purified polysaccharide serotype 4 were then measured in the sera by ELISA. Antibody units were calculated based on a hyperimmune standard. p values were determined by student t-test. Asterisks ($p<0.05$) indicate significant differences with respect to vaccinated mice. Pooled data from two separate experiments with $n=6$ mice per group are presented as means $\pm$ SD.

Figure 5. PMNs are required for optimal antibody function following PCV. (A-C) Sera were collected from naïve, Pevnar-13 immunized and PMN depleted immunized mice four weeks post vaccination following the timeline indicated in Fig 1A. (A) Wild type (WT) or a capsule deletion mutant (Δcps) *S. pneumoniae* were incubated with the indicated sera for 30 minutes, washed and stained with fluorescently-labeled anti-mouse IgG. The amount (mean fluorescent intensity or MFI) of bound Abs was determined by flow cytometry. Representative data from one of three separate experiments ($n=3$ biological replicates) are shown where each condition was tested in triplicate ($n=3$

541 technical replicates) per experiment. (B) The ability of PMNs isolated from naïve mice to
 542 kill pneumococci pre-opsonized with the indicated sera was determined. Percent bacterial
 543 killing was determined with respect to a no PMN control. Data shown are pooled from
 544 three separate experiments (n=3 biological replicates) where each condition was tested in
 545 triplicate (n=3 technical replicates) per experiment. (A-B) Bar graph represent means+/-
 546 SD and asterisks indicate significant differences from vaccinated controls as calculated
 547 by One-way ANOVA followed by Dunnet's test. (C) Naïve C57BL/6 female mice were
 548 injected i.p with 200µl of pooled serum from the indicated mice then challenged i.t. 1
 549 hour later with 5x10⁵ CFU *S. pneumoniae* TIGR4. Survival was assessed over time. *,
 550 denotes significance by the log-Rank (Mantel-Cox) test. Data were pooled from 8
 551 mice/group from two separate experiments.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

A.

B.

C.