1 Fitness costs of parasites explain 2 multiple life history tradeoffs in a 3 wild mammal

4 Summary

5 Reproduction in wild animals can divert limited resources away from immune
6 defence, resulting in increased parasite burdens. A longstanding prediction of life
7 history theory states that these parasites can harm the reproductive individual,
8 reducing its subsequent fitness and producing reproduction-fitness tradeoffs. Here,
9 we examined associations among reproductive allocation, immunity, parasitism,
10 and subsequent fitness in a wild population of individually identified red deer
11 (*Cervus elaphus*). Using path analysis, we investigated whether costs of lactation
12 for downstream survival and fecundity were mediated by changes in strongyle
13 nematode count and mucosal antibody levels. Lactating females exhibited
14 increased parasite counts, which were in turn associated with substantially
15 decreased fitness in the following year in terms of overwinter survival, fecundity,
16 subsequent calf weight, and parturition date. This study offers observational
17 evidence for parasite regulation of multiple life history tradeoffs, supporting the role
18 of parasites as an important mediating factor in wild mammal populations.

19 Introduction

20 A fundamental tenet of life history theory states that reproduction should be costly
 21 for future fitness (Williams, 1966; Stearns, 1989). While evidence for such trade-
 22 offs is widespread, the mechanisms behind them remain poorly understood. One
 23 hypothesised mechanism is that reproductive costs act through parasites, where
 24 increased reproductive allocation diverts limited resources away from immune
 25 defence, resulting in increased parasite burdens, which reduce subsequent fitness
 26 (Sheldon and Verhulst, 1996; Harshman and Zera, 2007). Parasite mediation of
 27 life history tradeoffs involves two necessary components: that reproduction
 28 increases parasitism, and that these parasites cause harm or require resources to
 29 combat them, reducing subsequent fitness. There is abundant evidence for each
 30 component of this theory across a range of taxa: firstly, life history investment is
 31 often associated with weaker or altered immune allocation (Neggazi *et al.*, 2016;
 32 Rödel *et al.*, 2016; Krams *et al.*, 2017) or with increased parasitism (Festa-
 33 Bianchet, 1989; Cizauskas *et al.*, 2015; Debeffe *et al.*, 2016). Secondly, increased
 34 parasitism is often associated with decreased subsequent probability of survival
 35 (Coltman *et al.*, 1999; Leivesley *et al.*, 2019) or reproduction (Albon *et al.*, 2002;
 36 Vandegrift *et al.*, 2008; Hughes *et al.*, 2009). Despite evidence for one or other of
 37 these processes in isolation, reproduction-associated increases in parasitism have
 38 rarely been linked to downstream fitness consequences in the same study to
 39 provide full support for parasite mediation of life history tradeoffs.

40 While it is true that reproduction, immunity, and parasites all compete for host
 41 resources, mechanisms governing life history tradeoffs are hypothesised to occur
 42 in a temporal sequence rather than occurring simultaneously (Figure 1). First,
 43 reproduction diverts resources away from immunity, reducing immune allocation
 44 (Sheldon and Verhulst, 1996). Resultant weaker immunity, plus potentially
 45 increased exposure associated with altered behaviour of reproductive individuals,
 46 can then result in higher parasite burden (Knowles *et al.*, 2009; Albery *et al.*,
 47 2020). Finally, subsequent fitness is reduced by damage from parasites
 48 (Harshman and Zera, 2007; Graham *et al.*, 2011). This combination of
 49 mechanisms comprises an indirect cost of reproduction acting through parasites.
 50 Additional (direct) costs of reproduction can simultaneously act through other
 51 mechanisms such as reduced condition, hormonal or phenological regulation, or
 52 damage caused by oxidative stress (Stjernman *et al.*, 2004; Harshman and Zera,
 53 2007; Speakman, 2008; Figure 1). This causal sequence is important, because

54 parasites' observed relationship with life history traits can depend on whether a
55 preceding, contemporary, or subsequent trait is chosen to examine.

56

57

58 Figure 1: The hypothesised mechanism for parasite-dependent mediation of life history
59 tradeoffs. Blue (interior) arrows denote indirect, parasite-mediated fitness costs of
60 reproduction, while the orange (exterior) arrows denote direct costs through resource
61 allocation, hormonal regulation, or similar mechanisms.

62

63 Many studies examining reproduction-immunity-parasitism interrelationships have
64 been carried out in birds, often using experiments in which reproductive effort is
65 artificially increased by manipulating clutch sizes (Knowles *et al.*, 2009). Such
66 manipulations are not possible in many mammal species, and thus most of our
67 knowledge of these trade-offs are based on observational studies. In observational
68 contexts, or in concert with experiments, path analysis can be used to infer links
69 between parasites and their fitness consequences (Pacejka *et al.*, 1998;
70 Stjernman *et al.*, 2004; Brambilla *et al.*, 2015; Leivesley *et al.*, 2019). Notably, a
71 recent analysis in a wild population of Soay sheep used path analysis to
72 demonstrate observationally that reproduction reduced survival through increased
73 parasite count and reduced body weight (Leivesley *et al.*, 2019), but without
74 examining impacts on subsequent reproductive traits.

75 The wild red deer (*Cervus elaphus*) on the Isle of Rum provide a classic example
 76 of a life history tradeoff under natural conditions: female deer that invest in
 77 lactation have reduced future fecundity and survival probability compared to non-
 78 lactating females (Clutton-Brock *et al.*, 1989; Froy *et al.*, 2016). Females that give
 79 birth to a calf that dies within the first few months of its life have similar fitness
 80 outcomes to those that do not give birth, implying that gestation has a minimal cost
 81 relative to lactation (Clutton-Brock *et al.*, 1989). The deer are infected with several
 82 helminth parasites, with egg counts and mucosal antibody (IgA) levels measured
 83 via noninvasive collection of faecal samples (Albery *et al.*, 2018, 2020). A previous
 84 study demonstrated decreased mucosal IgA and increased parasite count in
 85 reproductive females (Albery *et al.*, 2020). Lactation (but not gestation) is
 86 associated with increased parasite counts, partially reflecting the cost of
 87 reproduction for fitness; however, investment in gestation is associated with
 88 decreased mucosal antibody levels (Albery *et al.*, 2020). These findings each
 89 demonstrate a cost of reproduction for fitness, parasitism, and immunity
 90 respectively in this population. However, we have yet to establish the degree to
 91 subsequent fitness costs of reproduction can be explained by changes in
 92 parasitism.

93 Here, we use path analysis to link reproduction-immunity-parasitism tradeoffs in
 94 the Isle of Rum red deer with survival and reproduction in the following year,
 95 investigating whether immunity and parasitism are capable of mediating life history
 96 tradeoffs, and attempting to separate immune and parasite mediation from direct
 97 effects of reproduction acting through alternative mechanisms. We expected that
 98 substantially increased parasite counts associated with lactation would be
 99 associated with decreased subsequent survival, fecundity, parturition date, and
 100 calf birth weight, so that parasites provide a mechanistic explanation for the costs
 101 of lactation seen in this system.

102

Model

Female survival

Fecundity

Calf birth weight

Calf birth date

Figure 2: The four models in the context of the red deer reproductive cycle and sampling regime, over an example two-year period. Reproduction begins in spring and summer, at the start of each deer year, and one sampling trip was undertaken each summer (August), after the calving season had finished. Mating occurs in the autumn, and the mortality season begins in winter and lasts until early spring. A second sampling trip occurred each spring (April), after mating and mortality, but before the beginning of the subsequent calving season. The fitness variables investigated were quantified at the start of the subsequent deer year: if a female survived to May 1 the following year she was counted as 1 in the survival analysis, 0 if not, and the presence, weight, and birth date of her calf in the following spring were used as response variables in the remaining three models. The sampling trips included in each model were selected according to feasibility of causal links. For example, females become pregnant in the autumn, so we did not include the spring sampling season in the reproduction model as they would already be pregnant at this point, making it unlikely that parasite counts in April have a direct effect on their probability of having a calf 1-2 months later.

113 Methods

114 Study system, sampling, and labwork

115 The study population is situated in the north block of the Isle of Rum National
 116 Nature Reserve (57°N 6°20'W). The deer are entirely wild and unmanaged, and
 117 have been monitored continuously since the 1970s (see Clutton-Brock *et al.*, 1982
 118 for an overview of the project). The life history data collected on the population
 119 provide high-resolution estimates of individuals' dates of birth and death,
 120 reproduction, and familial relationships. The "deer year" begins on May 1st, and the
 121 deer give birth ("calving") in May-June, having conceived in the previous autumn
 122 (Figure 2). Deer on Rum give birth to a single calf, and do not reproduce every
 123 year. During the calving season, we aim to capture and mark as many of the
 124 calves born as possible soon after birth, so that they can be monitored for the rest
 125 of their lives. Sex and capture weight (to the nearest 100g) are recorded. ~20% of
 126 calves die within the first few weeks of life, and giving birth to a calf that dies within
 127 this period has little cost to the mother in terms of her survival and reproduction
 128 probability the following year (Clutton-Brock *et al.*, 1989). In contrast, if a calf
 129 survives into the winter, the mother has spent ~6 months lactating to it, expending
 130 considerable resources in doing so, and this cost is associated with substantially
 131 decreased fecundity and survival probability the following year (Clutton-Brock *et al.*, 1989; Froy *et al.*, 2016).

133 During early spring (April) and late summer (August), either side of the calving
 134 season, we conducted two-week field trips to collect faecal samples from the deer
 135 noninvasively. Sampling was undertaken across five trips 2016-2018, with 701
 136 faecal samples collected in total; see Table 1 for details of datasets. We watched
 137 known individuals for defaecation, marked the spot where the droppings landed,
 138 and then collected them while minimising disturbance to the deer, generally within
 139 an hour. In the evenings, samples were processed and put into Ziploc bags for
 140 storage (Albery *et al.* 2018). A subsample was extracted by centrifugation and kept
 141 frozen for faecal antibody analysis (Watt *et al.*, 2016; Albery *et al.*, 2020). Another
 142 subsample was kept as anaerobic as possible in a Ziploc bag at 4°C to avoid egg
 143 hatching.

144 In the lab, faecal samples were counted for strongyle nematodes eggs using a salt
 145 flotation-centrifugation faecal egg count (FEC), accurate to 1 egg per gram (EPG)
 146 within 3 weeks of collection. Strongyles are ubiquitous ruminant parasites that are
 147 present at high prevalence in this population and which increase in parasitism in

lactating individuals (Albery *et al.*, 2020). Previous studies in this population have also examined the helminths *Elaphostrongylus cervi* and *Fasciola hepatica* (Albery *et al.*, 2018, 2020). We chose to examine strongyles but not *E. cervi* or *F. hepatica* for several reasons: we did not want to add too many links to the analysis for reasons of interpretability; strongyles are most expected to have strong fitness costs (Hoberg *et al.*, 2001) and exhibited the most profound reproductive tradeoff in terms of significance and magnitude (Albery *et al.*, 2020); we did not expect *E. cervi* to have strong fitness effects (Irvine *et al.*, 2006); and *F. hepatica* is present at relatively low prevalence in adult females, preventing it from being fitted easily as an explanatory variable (Albery *et al.*, 2018). We also carried out antibody detection ELISAs designed to quantify mucosal IgA in sheep (Watt *et al.*, 2016; Albery *et al.*, 2020). This protocol quantifies both total IgA levels as a measure of general immune investment, and anti-*Teladorsagia circumcincta* IgA levels (anti-Tc IgA) as a specific anti-strongyle measure. *T. circumcincta* is primarily a sheep strongyle, but the anti-Tc IgA assay shows high cross-reactivity with a range of strongyle nematodes including the mouse helminth *Heligmosomoides polygyrus* (Froy *et al.*, 2019). The deer are infected with a selection of strongyle nematodes (Irvine *et al.*, 2006), including *Teladorsagia circumcincta* (unpublished data); thus, anti-Tc IgA is used to approximate anti-strongyle immune responses in the deer (Albery *et al.*, 2020). To control for collection factors which introduce confounding variation in antibody levels we used the residuals from a model including extraction session, time to freezing, and collection day, as in previous studies (Albery *et al.*, 2019, 2020). We also assayed faecal samples collected in November (Albery *et al.*, 2018). However, females exhibited very low strongyle prevalence in the autumn compared with spring and summer, preventing our FEC data from approximating normality and providing little variation to test when fitted as an explanatory variable. Hence, autumn data were excluded from our analyses.

Statistical analysis

To investigate links among our variables we used path analysis using the D-sep method, in which a set of linear models are fitted to the data, with some variables appearing as both response and explanatory variables (Shipley, 2009). Combining the linear models in this way allows identification of potential causal links and mediating variables.

We created four Directed Acyclic Graphs (DAGs), each examining a different fitness-related trait measured in the year following measurement of parasite burden (see Figure 2). These measures included two direct fitness measures: the

184 female's overwinter survival (0/1, where 1=survived to May 1 the following year)
185 and fecundity the following year (0/1, where 1=gave birth in the following calving
186 season). We also examined two fitness-associated maternal traits: the birth weight
187 of a female's calf the following year (continuous, Gaussian distributed, based on a
188 regression of capture weight on capture age in days) and parturition date the
189 following year (continuous, Gaussian distributed, based on Julian date that year).

190 Our analyses used three immune and parasite measures, which included: Total
191 IgA level; Anti-Tc IgA level; Strongyle count per gram of faeces (continuous,
192 log(count+1)-transformed to approximate normality). We included two mutually
193 exclusive binary reproductive categories representing the reproductive cost paid
194 that year (Clutton-Brock *et al.*, 1989): Gestation (gave birth to a calf which died
195 before 1st October that year) and Gestation + Lactation (gave birth to a calf which
196 survived to 1st October; hereafter referred to as simply "Lactation", as all
197 individuals that lactated must have also undergone the cost of gestation). We also
198 included variables to control for annual, seasonal, and age-related variation: Year
199 (categorical, with three levels: 2015, 2016, 2017); Season (two levels: Summer,
200 Spring); and Age (continuous, in years).

201 Each of the four DAGs was composed of four similar models, fitted using the INLA
202 package (Rue and Martino, 2009) in R version 3.5 (R Core Team, 2018). All
203 measures included female identity as a random effect to control for
204 pseudoreplication. First, we ran a set of three "input models", where the response
205 variable was an antibody or parasite measure. The aim of these models was to
206 quantify the association between reproduction and the immune/parasite measures,
207 and to quantify links between these measures themselves.

208 The models were specified as follows for each of our analyses, with
209 immune/parasite measures in bold and reproductive traits in italics. Variables in
210 brackets were included in the models, but are not displayed in the DAGs for clarity.

- 211 1. **Total IgA** ~ *Gestation + Lactation* (+ Age + Season + Year)
- 212 2. **Anti-Tc IgA** ~ **Total IgA** + *Gestation + Lactation* (+ Age + Season + Year)
- 213 3. **Strongyles** ~ **Anti-Tc IgA** + *Gestation + Lactation* (+ Age + Season + Year)
- 214 4. Fitness-related trait ~ **Strongyles** + **Anti-Tc IgA** + **Total IgA** +
215 *Gestation + Lactation* (+ Age + Year)

216 Combining these two model sets allowed comparison of the significance and
217 magnitude of different traits' costs for fitness in the following year (Figure 1).
218 Combining the estimates from models 1-3 with the estimates from model 4 allows

219 calculation of the direct and indirect (parasite- or immune-mediated) effects of
 220 lactation and gestation on subsequent fitness traits (Figure 1). As an example, we
 221 compared the magnitude and credibility intervals of direct lactation effects (effect
 222 of lactation in the fitness model [model 4]) with indirect effects (lactation effects on
 223 strongyle count [model 3] multiplied by the effects of strongyle count on fitness
 224 [model 4]). We took 1000 posterior draws from each of the lactation-strongyle link
 225 and the strongyle-fitness link and multiplied them together, and then derived the
 226 95% credibility intervals for this link. We compared these estimates with those for
 227 the direct lactation-fitness link to investigate whether effects of lactation were likely
 228 to act independently and/or through strongyle count. The models, fitness
 229 measures, and datasets used in each analysis are described in Table 1.

230

231

Model Set	Fitness Measure	Definition	Dataset	Samples	Individuals
1	Survival	Female survival the following winter (0/1)	All females (Spring year t-1 and Summer year t)	485	134
2	Fecundity	Female reproduction the following deer year (0/1)	All females (Summer)	223	107
3	Calf birth weight	Calf weight the following deer year (Kg)	Females that reproduced the following May-June (Summer year t and Spring year t)	300	94
4	Parturition date	Date of parturition the following deer year (Days from 1 st January)		336	106

232

233

Table 1: Descriptions of path analyses and the datasets used.

Figure 3: Directed Acyclic Graphs (DAGs). Results are displayed for all four investigated fitness response variables: overwinter survival (A); fecundity (B); subsequent calf birth weight (C); subsequent parturition date (D). Link colour depends on the direction of the effect (blue=positive, red=negative); link width indicates the magnitude of the effect; and only solid, opaque links are significant (estimates did not overlap with zero). Labels denote the link-scale effect sizes (slopes) for the significant effects, derived from GLMMs (full model effects displayed in SI Figure 1).

239

240

241 Figure 4: Relationships between strongyle parasite count and fitness measures. Results are displayed
242 for all four investigated fitness response variables: overwinter survival (A); fecundity (B); subsequent
243 calf birth weight (C); subsequent parturition date (D). The lines denote the fitted slope of parasitism on
244 the response variable, with 95% credibility intervals. Credibility intervals did not overlap with zero for
245 any of the four relationships. Strongyle count was log(x+1)-transformed for analysis and plotting.

246

247

Figure 5: Comparison of direct and indirect (parasite-mediated) effects of lactation on fitness-related traits on the link scale (logistic for survival and reproduction; Gaussian for birth weight and birth date).

Points represent mean effect estimates derived from the model posterior distributions; blue corresponds to direct effects, and red corresponds to indirect effects. Parameters with asterisks were significant: i.e., their credibility intervals did not overlap with zero.

Figure 6: Higher anti-*Teladorsagia circumcincta* IgA was associated with an increased probability of reproducing the following year. Anti-Tc IgA was cube root-transformed and calculated from the residuals of a linear model including collection variables, and was then scaled to have a mean of 0 and a standard deviation of 1. The line represents the output of the reproduction probability model which includes lactation and strongyles as explanatory factors.

Results

Path analyses consistently revealed strong positive associations between lactation and parasite count, and negative associations between parasite counts and subsequent fitness in terms of all four traits (Figures 3-5). In contrast, estimates for lactation's direct association with subsequent fitness overlapped with zero for all response variables except parturition date, supporting parasite-mediated reproductive costs for fitness (Figures 3-5). Below, for each of the four fitness-related response variables, we describe the magnitude of the direct association of parasitism with fitness, the direct association of lactation with fitness, and lactation's association with parasitism multiplied by parasitism's association with fitness. The latter gives an estimate of the indirect effect of lactation on the fitness-related trait acting through strongyle count. For effect sizes we give the mean and 95% credibility intervals (CI). 1 log(EPG+1) increase corresponds to a ~3x increase in strongyle count. Full model effect sizes are displayed in the supplementary information (Figure SI1; Table SI1).

Parasite count had a strong association with subsequent survival probability despite high survival rates in the population (Figure 3A,4A). Females with the lowest counts (0 EPG, 10% of samples) had a survival probability of ~100%, while those with the highest (>25EPG, 7%

of samples) had a survival probability of <90% (Figure 4A). Lactation was associated with increased strongyle count (+0.85, CI 0.64, 0.99), so that a substantial cost of lactation for survival acted through parasitism (Figure 5). Although this effect was highly significant on the link (logistic) scale (Figure SI1, Table SI1), given the high survival rates in the population, at the mean EPG value this lactation-associated increased strongyle parasitism would correspond to only a ~2% decrease in survival probability. In contrast, estimates for the direct effect of lactation on survival overlapped widely with zero, and the point estimate was greater than zero, implying that individuals that lactate were slightly more likely to survive when the effects of parasitism were accounted for (Figures 3A, 5).

Strongyles' association with subsequent fecundity had a similar effect size to its association with overwinter survival (Figures 3B, 4B, 5; Table SI1; Figure SI1). An increase of 1 log(EPG+1) was associated with a decrease of ~15% probability of reproducing. 0 EPG (17% of samples) corresponded to a ~77% chance of reproducing the following year, and those with >20 EPG (6% of samples) had a reproduction probability of <36% (Figure 4B). The direct effect of lactation on subsequent fecundity was negligible and had very large credibility intervals, as with survival (Figure 5). In addition to the association with parasite count, individuals with higher levels of anti-Tc IgA were more likely to reproduce the following year (Figure 3B; Figure 6). An increase of 1 standard deviation of anti-Tc IgA levels corresponded to an increase of ~10% in the probability of reproducing. Individuals with the lowest anti-Tc IgA levels (less than -1 SD units) had a reproduction probability of <50%, compared to >75% for those with the highest levels (>1 SD units; Figure 6). Finally, individuals that paid the cost of gestation were much more likely to reproduce the following year, independently of the effects of antibodies and parasites (Figure 3B).

Calving traits exhibited weaker associations with parasitism than did survival and reproduction, although the results still implied an indirect cost of lactation acting through strongyle count (Figures 3-5). The DAG for calf birth weight was similar to that for survival (Figure 3C). An increase of 1 log(EPG+1) corresponded to a slight decrease in calf birth weight the following year (0.07 SD units, or about 86g; Figure 4C). Females with the highest strongyle intensities (>25 EPG) gave birth to calves which were ~400g lighter than those with the lowest intensities (0 EPG), or around 6.24 kg compared to 6.65 kg. As with survival, there was poor support for a direct association between lactation and birth weight (Figure 5). The estimates for this direct effect were close to zero, and credibility intervals overlapped substantially with zero (Figure 5). Lactation's positive effect on strongyle count once again resulted in a significant negative indirect effect of lactation on subsequent calf birth weight acting through strongyles, but the estimates were very small and nearly overlapped with zero (-0.0438kg, CI -0.111.6, -0.005.6).

In contrast to all other metrics we investigated, there was support for a positive and direct effect of lactation on parturition date the following year: that is, females whose calf survived until the winter were likely to calve later in the following year (~8.5 days later, CI: 5.9, 11.2; Figure 3D,5), regardless of parasite count. There was a much weaker association between strongyle count and parturition date: an increase of 1 log(EPG+1) produced a delay in calving of ~0.93 days (CI: 0.12, 1.75; Figure 3D,4D,5). Lactation resulted in an increase of 0.7 log(EPG+1). Combining this estimate with the effect of parasitism on birth date gives an estimate for an indirect effect of lactation acting on birth date totalling 0.58 days' delay (CI 0.06, 1.31; Figure 5). Parturition date was thus the only metric examined here for which lactation's direct effect was larger than its indirect effect acting through strongyle count (Figures 3-5).

There was a strong positive association between total IgA and anti-Tc IgA, as expected given our previous findings (Albery *et al.*, 2020; Table SI1, Figure SI1). However, lactation had no significant effect on anti-Tc IgA in our DAGs (Figure 3, Table SI1, Figure SI1).

Discussion

This study provides observational evidence for strong parasite-dependent mediation of multiple life history tradeoffs in a large wild mammal. Lactation was associated with higher parasite intensities which translated to reduced fecundity and survival probability in the subsequent year. Among individuals that did reproduce the following year, those with high strongyle counts gave birth slightly later in the year and to slightly smaller calves. These findings represent the second evidence for such mediation of reproduction-survival tradeoffs in a wild mammal (Leivesley *et al.*, 2019), and new evidence of parasites mediating reproductive tradeoffs with subsequent reproductive traits. It is likely that much of the fitness reduction associated with lactation in the Rum red deer population (Clutton-Brock *et al.*, 1989; Froy *et al.*, 2016) is caused by strongyle parasites, or that strongyle count closely corresponds to latent condition variables that are responsible for mediating fitness. This finding supports parasites' role as an important mediating factor in this system.

Lactation's negative association with fitness acted largely through strongyle count for all fitness metrics except parturition date. This may represent a parasite-mediated cost, where pathology and resource allocation associated with increased parasitism are the primary cause of increased overwinter mortality and reduced subsequent fecundity in lactating individuals (Clutton-Brock *et al.*, 1989). Allocation of resources to lactation and associated physiological changes likely reduces resources available for resistance and damage repair mechanisms, rendering lactating females more susceptible to strongyles (Sheldon and Verhulst, 1996; Speakman, 2008), while also increasing their exposure through heightened

forage intake (Albery *et al.*, 2020). High parasite counts in lactating females may cause gut pathology, interfering with nutrient absorption and thereby exacerbating the nutritional scarcity of the winter period, leading to overwinter mortality (Gulland, 1992; Pedersen and Greives, 2008; Maublanc *et al.*, 2009), as well as reducing females' ability to achieve the body condition necessary to conceive and carry a calf to term (Albon *et al.*, 1986). This reduction in body condition could likewise cause females to give birth later in the year and to a calf that is smaller. There are two time points at which strongyles may reduce fecundity: first, parasites may impact females in the resource-abundant summer and early autumn, preventing them from conceiving in the autumn mating season. In this case, strongyle-associated pathology may occur somewhat independently of overwinter nutritional scarcity. Alternatively, strongyles may cause females to lose their pregnancies over winter. This possibility may be tested in the future by investigating whether more highly parasitised females are less likely to be observed mating (demonstrating reduced conception rates), or only less likely to give birth.

Lactation exerts a substantial resource cost that results in reduced condition; therefore, it is also possible that we observed a negative relationship between parasitism and fitness because both were determined by condition, rather than because parasites were causally responsible for reducing fitness. Strongyle counts are associated with decreased body condition in shot individuals in this population, supporting this possibility (Irvine *et al.*, 2006). Similarly, an important role of condition is supported by our observation that higher anti-Tc IgA levels were associated with increased fecundity the following year, independently of any associations between 1) strongyles and fecundity and 2) anti-Tc IgA and strongyles themselves (Figure 6). It is highly likely that anti-Tc IgA is well-correlated with an unmeasured component of individual quality such as fat content (Demas *et al.*, 2003) which is linked to fitness both in the deer (Albon *et al.*, 1986) and in other systems (Milenkaya *et al.*, 2015). This possibility reflects the confounding effects of individual quality in observational studies of tradeoffs (van Noordwijk and de Jong, 1986). We were unfortunately unable to replicate previous findings of lactation costs for mucosal antibodies (Albery *et al.*, 2020), likely due to extremely reduced sample sizes (485 samples and fewer here compared to 837 samples previously), and so we were unable to link the anti-Tc IgA-fitness association as part of a reproduction-fitness tradeoff. Another potential fitness-mediating factor is body weight, which is often used to control for condition-driven versus parasite-driven fitness effects; however, analyses in Soay sheep often show that strongyles-fitness associations occur independent of, or in addition to body weight (e.g. Sparks *et al.*, 2018; Froy *et al.*, 2019; Leivesley *et al.*, 2019). In addition, although condition-parasitism relationships are well-documented, negative effects are far from ubiquitous and their slopes are relatively

shallow on average (Sánchez *et al.*, 2018). Whether or not strongyles are the effectors, our findings nevertheless support the use of these parasites as a proxy for an individual's health and as predictors of its subsequent fitness.

Strongyles will have a strong mediating effect on population dynamics, for two principal reasons: first, by reducing both survival and fecundity simultaneously, and second, by exhibiting different relationships with past and subsequent reproduction. As such, it stands to reason that their impact will prevent too many females from reproducing in a single year, potentially stabilising population fluctuations. Further years of data will reveal how parasite abundances relate to the population dynamics of the deer, and particularly whether inter-annual variation in strongyle numbers can explain population size (Wilson *et al.*, 2004). At higher population densities the deer exhibit delayed maturity and reduced fecundity (Albon *et al.*, 1983); the lactation-strongyle-fecundity tradeoff offers a potential mechanism behind this fecundity reduction, particularly as parasitism should worsen at higher densities (Altizer *et al.*, 2003; Wilson *et al.*, 2004). Local population density also influences fitness in this population (Coulson *et al.*, 1997) and parasitism demonstrates fine-scale spatial variation (Albery *et al.*, 2019), so this life history mediation could likewise occur at relatively fine spatial scales. A similar study in Soay sheep demonstrated that strongyles mediate a reproduction-survival tradeoff, but without examining similar reproduction-fecundity tradeoffs, partly because most sheep do not take years off between reproduction events (Leivesley *et al.*, 2019). The fecundity reduction seen in the deer and the strength of these parasite-mediated tradeoffs potentially contribute to the population's relatively weak population cycles, particularly compared to the strong oscillatory population dynamics of the Soay sheep (Clutton-Brock and Pemberton, 2004). As such, parasite-dependent life history mediation may be an important contributing factor determining the strength of oscillatory population dynamics.

Finally, having uncovered costs of parasitism in adult females, it would be interesting to investigate whether other age and sex categories experience similar fitness effects: e.g., do more highly parasitised males sire fewer calves, and are more highly parasitised calves less likely to survive to maturity? Do maternal costs transfer to their calves, providing another potentially important mediating mechanism (Martin and Festa-Bianchet, 2010)? Future studies in this population could elaborate on these findings by investigating how maternal and calf parasitism correlate and correspond to maternal and calf fitness, quantifying transgenerational immunity-parasitism-fitness correlations: a topic that is largely understudied and likely influences ecological and epidemiological dynamics considerably (Roth *et al.*, 2018).

References

- Albery, G. F., Kenyon, F., Morris, A., Morris, S., Nussey, D. H. and Pemberton, J. M.** (2018). Seasonality of helminth infection in wild red deer varies between individuals and between parasite taxa. *Parasitology* **145**, 1–11. doi:10.1017/S0031182018000185.
- Albery, G. F., Becker, D. J., Kenyon, F., Nussey, D. H. and Pemberton, J. M.** (2019). The fine-scale landscape of immunity and parasitism in a wild ungulate population. *Integrative and Comparative Biology* **ic2016**, 1–11. doi:10.1093/icb/ic2016.
- Albery, G. F., Watt, K. A., Keith, R., Morris, S., Morris, A., Kenyon, F., Nussey, D. H. and Pemberton, J. M.** (2020). Reproduction has different costs for immunity and parasitism in a wild mammal. *Functional Ecology* **34**, 229–239. doi:10.1101/472597.
- Albon, S. D., Mitchell, B. and Staines, B. W.** (1983). Fertility and Body Weight in Female Red Deer: A Density-Dependent Relationship. *Journal of Animal Ecology* **52**, 969–980.
- Albon, S. D., Mitchell, B., Huby, B. J. and Brown, D.** (1986). Fertility in female red deer (*Cervus elaphus*): the effects of body composition, age and reproductive status. *Journal of Zoology (London)* **209**, 447–460. doi:10.1111/j.1469-7998.1986.tb03603.x.
- Albon, S. D., Stien, A., Irvine, R. J., Langvatn, R., Ropstad, E. and Halvorsen, O.** (2002). The role of parasites in the dynamics of a reindeer population. *Proceedings of the Royal Society B: Biological Sciences* **269**, 1625–1632. doi:10.1098/rspb.2002.2064.
- Altizer, S., Nunn, C. L., Thrall, P. H., Gittleman, J. L., Antonovics, J., Cunningham, A. A., Dobson, A. P., Ezenwa, V., Jones, K. E., Pedersen, A. B., Poss, M. and Pulliam, J. R. C.** (2003). Social Organization and Parasite Risk in Mammals: Integrating Theory and Empirical Studies. *Annual Review of Ecology, Evolution, and Systematics* **34**, 517–547. doi:10.1146/annurev.ecolsys.34.030102.151725.
- Brambilla, A., Biebach, I., Bassano, B., Bogliani, G., von Hardenberg, A., Hardenberg, A. Von, Brambilla, A. and von Hardenberg, A.** (2015). Direct and indirect causal effects of heterozygosity on fitness-related traits in Alpine ibex. *Proceedings of the Royal Society B: Biological Sciences* **282**, 20141873. doi:10.1098/rspb.2014.1873.
- Cizauskas, C. A., Turner, W. C., Pitts, N. and Getz, W. M.** (2015). Seasonal patterns of hormones, macroparasites, and microparasites in wild african ungulates: The interplay among stress, reproduction, and disease. *PLoS ONE* **10**, 1–29. doi:10.1371/journal.pone.0120800.
- Clutton-Brock, T. H. and Pemberton, J. M. (Josephine M. .** (2004). *Soay sheep: population dynamics and selection on St. Kilda*. Cambridge University Press.

- 452 **Clutton-Brock, T. H., Guinness, F. E. and Albon, S. D.** (1982). *Red Deer: Behavior and*
453 *Ecology of Two Sexes*. University of Chicago Press, Chicago, IL.
- 454 **Clutton-Brock, T. H., Albon, S. D. and Guinness, F. E.** (1989). Fitness costs of gestation
455 and lactation in wild mammals. *Nature* **337**, 260–262. doi:10.1038/337260a0.
- 456 **Coltman, D. W., Pilkington, J. G., Smith, J. a, Pemberton, J. M. and Josephine, M.**
457 (1999). Parasite-Mediated Selection against Inbred Soay Sheep in a Free-Living, Island
458 Population. *Evolution* **53**, 1259–1267. doi:10.2307/2640828.
- 459 **Coulson, T., Albon, S., Guinness, F., Pemberton, J. and Clutton-Brock, T.** (1997).
460 Population Substructure, Local Density, and Calf Winter Survival in Red Deer (*Cervus*
461 *elaphus*). *Ecology* **78**, 852. doi:10.2307/2266064.
- 462 **Debeffe, L., McLoughlin, P. D., Medill, S. A., Stewart, K., Andres, D., Shury, T., Wagner,**
463 **B., Jenkins, E., Gilleard, J. S. and Poissant, J.** (2016). Negative covariance between
464 parasite load and body condition in a population of feral horses. *Parasitology* **143**, 983–
465 997. doi:10.1017/S0031182016000408.
- 466 **Demas, G. E., Drazen, D. L. and Nelson, R. J.** (2003). Reductions in total body fat
467 decrease humoral immunity. *Proceedings of the Royal Society B: Biological Sciences*
468 **270**, 905–911. doi:10.1016/j.microc.2018.08.056.
- 469 **Festa-Bianchet, M.** (1989). Individual Differences, Parasites, and the Costs of Reproduction
470 for Bighorn Ewes (*Ovis canadensis*). *Journal of Animal Ecology* **58**, 785–795.
471 doi:10.2307/5124.
- 472 **Froy, H., Walling, C. A., Pemberton, J. M., Clutton-brock, T. H. and Kruuk, L. E. B.**
473 (2016). Relative costs of offspring sex and offspring survival in a polygynous mammal.
474 *Biology Letters* **12**, 20160417. doi:10.1098/rsbl.2016.0417.
- 475 **Froy, H., Sparks, A. M., Watt, K., Sinclair, R., Bach, F., Pilkington, J. G., Pemberton, J.**
476 **M., Mcneilly, T. N. and Nussey, D. H.** (2019). Senescence in immunity against
477 helminth parasites predicts adult mortality in a wild mammal. *Science* **365**, 1296–1298.
- 478 **Graham, A. L., Shuker, D. M., Pollitt, L. C., Auld, S. K. J. R., Wilson, A. J., Little, T. J.,**
479 **Wilson, J. and Little, T. J.** (2011). Fitness consequences of immune responses:
480 Strengthening the empirical framework for ecoimmunology. *Functional Ecology* **25**, 5–
481 17. doi:10.1111/j.1365-2435.2010.01777.x.
- 482 **Gulland, F. M. D.** (1992). The role of nematode parasites in Soay sheep (*Ovis aries* L.)
483 mortality during a population crash. *Parasitology* **105** (Pt 3, 493–503.

484 doi:10.1017/S0031182000074679.

485 **Harshman, L. G. and Zera, A. J.** (2007). The cost of reproduction: the devil in the details.
 486 *Trends in Ecology and Evolution* **22**, 80–86. doi:10.1016/j.tree.2006.10.008.

487 **Hoberg, E. P., Kocan, A. A. and Rickard, L. G.** (2001). Gastrointestinal strongyles in wild
 488 ruminants. *Diseases of Wild Mammals* 193–227. doi:10.1002/9780470377000.ch8.

489 **Hughes, J., Albon, S. D., Irvine, R. J. and Woodin, S.** (2009). Is there a cost of parasites
 490 to caribou? *Parasitology* **136**, 253–65. doi:10.1017/S0031182008005246.

491 **Irvine, R. J., Corbishley, H., Pilkington, J. G. and Albon, S. D.** (2006). Low-level parasitic
 492 worm burdens may reduce body condition in free-ranging red deer (*Cervus elaphus*).
 493 *Parasitology* **133**, 465–475. doi:10.1017/S0031182006000606.

494 **Knowles, S. C. L., Nakagawa, S. and Sheldon, B. C.** (2009). Elevated reproductive effort
 495 increases blood parasitaemia and decreases immune function in birds: A meta-
 496 regression approach. *Functional Ecology* **23**, 405–415. doi:10.1111/j.1365-
 497 2435.2008.01507.x.

498 **Krams, I. A., Rumvolt, K., Saks, L., Krams, R., Elferts, D., Vrublevska, J., Rantala, M. J.,**
 499 **Kecko, S., Cīrule, D., Luoto, S. and Krama, T.** (2017). Reproduction compromises
 500 adaptive immunity in a cyprinid fish. *Ecological Research* **32**, 559–566.
 501 doi:10.1007/s11284-017-1467-y.

502 **Leivesley, J. A., Bussière, L. F., Pemberton, J. M., Pilkington, J. G., Wilson, K. and**
 503 **Hayward, A. D.** (2019). Survival costs of reproduction are mediated by parasite
 504 infection in wild Soay sheep. *Ecology Letters* ele.13275. doi:10.1111/ele.13275.

505 **Martin, J. G. A. and Festa-Bianchet, M.** (2010). Bighorn Ewes Transfer the Costs of
 506 Reproduction to Their Lambs. *The American Naturalist* **176**, 414–423.
 507 doi:10.1086/656267.

508 **Maublanc, M. L., Bideau, E., Picot, D., Rame, J. L., Dubois, M., Ferté, H., Gerard, J. F.,**
 509 **Rames, J., Dubois, M., Ferté, H. and Gerard, J. F.** (2009). Demographic crash
 510 associated with high parasite load in an experimental roe deer (*Capreolus capreolus*)
 511 population. *European Journal of Wildlife Research* **55**, 621–625. doi:10.1007/s10344-
 512 009-0298-8.

513 **Milenkaya, O., Catlin, D. H., Legge, S. and Walters, J. R.** (2015). Success but Not Survival
 514 in a Sedentary , Tropical Bird. *PLoS Neglected Tropical Diseases* **10**, 1–18.
 515 doi:10.5061/dryad.3n2j5.

- 516 **Neggazi, S. A., Noreikiene, K., Öst, M. and Jaatinen, K.** (2016). Reproductive investment
517 is connected to innate immunity in a long-lived animal. *Oecologia* **182**, 347–356.
518 doi:10.1007/s00442-016-3657-7.
- 519 **Pacejka, A. J., Gratton, C. M. and Thompson, C. F.** (1998). Do potentially virulent mites
520 affect house wren (*Troglodytes aedon*) reproductive success? *Ecology* **79**, 1797–1806.
521 doi:10.1890/0012-9658(1998)079[1797:DPVMAH]2.0.CO;2.
- 522 **Pedersen, A. B. and Greives, T. J.** (2008). The interaction of parasites and resources
523 cause crashes in a wild mouse population. *Journal of Animal Ecology* **77**, 370–377.
524 doi:10.1111/j.1365-2656.2007.01321.x.
- 525 **R Core Team** (2018). R: A language and environment for statistical computing. R
526 Foundation for Statistical Computing, Vienna, Austria.
- 527 **Rödel, H. G., Zapka, M., Stefanski, V. and von Holst, D.** (2016). Reproductive effort alters
528 immune parameters measured post-partum in European rabbits under semi-natural
529 conditions. *Functional Ecology* **30**, 1800–1809. doi:10.1111/1365-2435.12663.
- 530 **Roth, O., Beemelmanns, A., Barribeau, S. M. and Sadd, B. M.** (2018). Recent advances
531 in vertebrate and invertebrate transgenerational immunity in the light of ecology and
532 evolution. *Heredity* **121**, 225–238. doi:10.1038/s41437-018-0101-2.
- 533 **Rue, H. and Martino, S.** (2009). Approximate Bayesian inference for latent Gaussian
534 models by using integrated nested Laplace approximations. *Statistical Methodology* **71**,
535 319–392.
- 536 **Sánchez, C. A., Becker, D. J., Teitelbaum, C. S., Barriga, P., Brown, L. M., Majewska, A.**
537 **A., Hall, R. J. and Altizer, S.** (2018). On the relationship between body condition and
538 parasite infection in wildlife: a review and meta-analysis. *Ecology Letters* **21**, 1869–
539 1884. doi:10.1111/ELE.13160.
- 540 **Sheldon, B. C. and Verhulst, S.** (1996). Ecological immunology - costly parasite defenses
541 and trade- offs in evolutionary ecology. *Trends in Ecology & Evolution* **11**, 317–321.
542 doi:10.1016/0169-5347(96)10039-2.
- 543 **Shipley, B.** (2009). Confirmatory path analysis in a generalized multilevel context. *Ecology*
544 **90**, 363–368. doi:10.1890/08-1034.1.
- 545 **Sparks, A. M., Watt, K., Sinclair, R., Pilkington, J. G., Pemberton, J. M., Johnston, S. E.,**
546 **McNeilly, T. N. and Nussey, D. H.** (2018). Natural selection on antihelminth antibodies
547 in a wild mammal population. *American Naturalist* **192**, 745–760. doi:10.1086/700115.

- 548 **Speakman, J. R.** (2008). The physiological costs of reproduction in small mammals.
549 *Philosophical Transactions of the Royal Society B: Biological Sciences* **363**, 375–398.
550 doi:10.1098/rstb.2007.2145.
- 551 **Stearns, S. C.** (1989). Trade-Offs in Life-History Evolution. *Functional Ecology* **3**, 259–268.
552 doi:10.2307/2389364.
- 553 **Stjernman, M., Raberg, L. and Nilsson, J.** (2004). Survival costs of reproduction in the
554 blue tit (*Parus caeruleus*): a role for blood parasites? *Proceedings of the Royal Society*
555 *B: Biological Sciences* **271**, 2387–2394. doi:10.1098/rspb.2004.2883.
- 556 **van Noordwijk, A. J. and de Jong, G.** (1986). Acquisition and Allocation of Resources:
557 Their Influence on Variation in Life History Tactics. *The American Naturalist* **128**, 137–
558 142. doi:10.1086/284547.
- 559 **Vandegrift, K. J., Raffel, T. R. and Hudson, P. J.** (2008). Parasites prevent summer
560 breeding in white-footed mice, *Peromyscus leucopus*. *Ecology* **89**, 2251–2258.
561 doi:10.1890/07-1935.1.
- 562 **Watt, K. A., Nussey, D. H., Maclellan, R., Pilkington, J. G. and McNeilly, T. N.** (2016).
563 Fecal antibody levels as a noninvasive method for measuring immunity to
564 gastrointestinal nematodes in ecological studies. *Ecology and Evolution* **6**, 56–67.
565 doi:10.1002/ece3.1858.
- 566 **Williams, G. C.** (1966). Natural Selection, the Costs of Reproduction, and a Refinement of
567 Lack's Principle. *The American naturalist* **100**, 687–690.
- 568 **Wilson, K., Grenfell, B. T., Pilkington, J. G., Boyd, H. E. G. and Gulland, F. M. D.** (2004).
569 Parasites and their impact. In *Soay Sheep: Dynamics and Selection in an Island*
570 *Population* (ed. T. Clutton-Brock and J. Pemberton), pp. 113–165. Cambridge
571 University Press.

Supplementary Information

Figure S11: Model outputs for the component GLMMs of all four path analyses. Colours correspond to different response variables; variables on the y axis correspond to explanatory variables in each component model. Points denote mean estimates from the posterior distribution of effect sizes from the component INLA model. Error bars denote 95% credibility intervals for the distribution. Where these intervals did not overlap with 0, the effect was taken to be significant. These effects are marked with an asterisk. Notably, lactation was positively associated with strongyles (blue estimates) and strongyles were negatively associated with all fitness-related traits (purple estimates).

Survival Path Analysis				
Model	Variable	Estimate	Lower CI	Upper CI
Total IgA	Intercept	-0.3081	-0.609	-0.0075
	Gestation	-0.3182	-0.5999	-0.0367
	Lactation	-0.0912	-0.2842	0.1017
	Age	0.047	-0.0273	0.1212
	Spring	1.0171	0.8138	1.2202
	Year: 2016	0.1765	-0.0441	0.397
	Year: 2017	0.34	0.0436	0.6363
Anti-Tc IgA	Intercept	0.0571	-0.149	0.263
	Total IgA	0.767	0.7095	0.8243
	Gestation	-0.0705	-0.2829	0.1417
	Lactation	-0.1298	-0.2625	0.0027
	Age	0.0538	-0.0285	0.1362
	Spring	-0.0872	-0.2234	0.0491
	Year: 2016	0.109	-0.0265	0.2443
	Year: 2017	0.1151	-0.0715	0.3016
Strongyle Count	Intercept	-0.9499	-1.2653	-0.6347
	Gestation	0.083	-0.2426	0.4084
	Lactation	0.8496	0.6416	1.0569
	Anti-Tc IgA	-0.0785	-0.1675	0.0105
	Age	0.0192	-0.1007	0.1382
	Spring	0.4485	0.2474	0.6495
	Year: 2016	0.2491	0.0375	0.4607
	Year: 2017	0.2247	-0.0654	0.5148

Survival	Intercept	3.3842	2.1424	4.8575
	Gestation	1.6766	-0.1589	3.9344
	Lactation	0.2061	-0.7305	1.1135
	Anti-Tc IgA	0.1306	-0.4932	0.7636
	Total IgA	-0.1087	-0.7467	0.5113
	Strongyles	-0.5896	-1.0085	-0.1937
	Age	-1.3464	-1.8883	-0.8885
	Year: 2016	-0.6102	-1.6816	0.364
	Year: 2017	-0.8665	-2.1049	0.2666
Fecundity Path Analysis				
Model	Variable	Estimate	Lower CI	Upper CI
Total IgA	Intercept	-0.1121	-0.3346	0.1102
	Gestation	-0.4279	-0.8417	-0.0145
	Lactation	-0.2216	-0.471	0.0276
	Age	0.1051	-0.0108	0.2209
	Year: 2017	0.1972	-0.0388	0.4329
Anti-Tc IgA	Intercept	0.0148	-0.1561	0.1853
	Total IgA	0.755	0.6747	0.8351
	Gestation	0.1878	-0.0955	0.47
	Lactation	-0.0021	-0.1816	0.1771
	Age	0.047	-0.0583	0.1522
	Year: 2017	0.0692	-0.07	0.208
Strongyle Count	Intercept	-0.0191	-0.2778	0.2403
	Gestation	-0.5432	-0.9927	-0.0939
	Lactation	0.4618	0.1793	0.7444
	Anti-Tc	-0.1244	-0.2597	0.0106

	IgA			
	Age	-0.0228	-0.1747	0.1275
	Year: 2017	-0.2038	-0.4353	0.0286
Fecundity	Intercept	0.1424	-0.4085	0.6949
	Gestation	2.122	0.7159	3.8491
	Lactation	0.0188	-0.6041	0.6448
	Anti-Tc IgA	0.5761	0.1144	1.0549
	Total IgA	-0.2661	-0.7449	0.2034
	Strongyles	-0.5885	-0.9166	-0.2737
	Age	-0.1704	-0.4716	0.1247
	Year: 2017	-0.1285	-0.7295	0.4681
Calf Birth Weight Path Analysis				
Model	Variable	Estimate	Lower CI	Upper CI
Total IgA	Intercept	-0.0559	-0.4059	0.2937
	Gestation	-0.2193	-0.5256	0.0866
	Lactation	-0.0667	-0.3142	0.1807
	Age	0.0129	-0.0855	0.1113
	Spring	0.7025	0.4721	0.9326
	Year: 2016	0.0116	-0.2658	0.2888
	Year: 2017	-0.1004	-0.387	0.186
Anti-Tc IgA	Intercept	0.1405	-0.0992	0.3799
	Total IgA	0.7789	0.7074	0.8506
	Gestation	-0.2126	-0.4438	0.018
	Lactation	-0.1524	-0.3547	0.0498
	Age	0.0525	-0.043	0.148
	Spring	-0.1912	-0.3348	-0.0479

	Year:	0.2109	0.0333	0.388
	2016			
	Year:	0.154	-0.0338	0.3416
	2017			
Strongyle Count	Intercept	-1.0168	-1.3738	-0.6603
	Gestation	0.0489	-0.2936	0.3925
	Lactation	0.6138	0.3177	0.9118
	Anti-Tc IgA	-0.0962	-0.2046	0.0121
	Age	-0.0306	-0.162	0.1018
	Spring	0.8109	0.5991	1.0222
	Year:	0.1204	-0.15	0.3911
	2016			
	Year:	0.4678	0.1848	0.7503
	2017			
Calf Birth Weight	Intercept	0.2411	-0.0236	0.5057
	Gestation	0.2213	-0.023	0.4654
	Lactation	-0.0972	-0.3267	0.1319
	Anti-Tc IgA	0.0909	-0.0243	0.2057
	Total IgA	-0.0278	-0.1342	0.0788
	Strongyles	-0.0715	-0.1404	-0.0028
	Age	-0.1467	-0.3092	0.0156
	Year:	-0.0935	-0.2613	0.0742
	2016			
	Year:	-0.461	-0.6596	-0.2627
	2017			
Calf Birth Date Path Analysis				
Model	Variable	Estimate	Lower CI	Upper CI
Total IgA	Intercept	0.0381	-0.2974	0.3734
	Gestation	-0.1961	-0.4888	0.0963

	Lactation	-0.0889	-0.3236	0.1457
	Age	0.0342	-0.0598	0.1282
	Spring	0.6269	0.409	0.8447
	Year: 2016	-0.0365	-0.302	0.2289
	Year: 2017	-0.1125	-0.3907	0.1654
Anti-Tc IgA	Intercept	0.1135	-0.112	0.3389
	Total IgA	0.7769	0.7097	0.8441
	Gestation	-0.1909	-0.4103	0.028
	Lactation	-0.1438	-0.3309	0.0431
	Age	0.0369	-0.0546	0.1286
	Spring	-0.1948	-0.3278	-0.0619
	Year: 2016	0.1799	0.0144	0.3452
	Year: 2017	0.1978	0.0202	0.3753
Strongyle Count	Intercept	-1.0305	-1.3734	-0.6883
	Gestation	0.0273	-0.3045	0.3598
	Lactation	0.6512	0.3645	0.9407
	Anti-Tc IgA	-0.0935	-0.197	0.0098
	Age	-0.0031	-0.1375	0.1319
	Spring	0.7492	0.5533	0.9449
	Year: 2016	0.14	-0.1144	0.3951
	Year: 2017	0.5809	0.3106	0.8508
Calf Birth Date	Intercept	-0.5344	-0.7683	-0.3033
	Gestation	0.1028	-0.1294	0.3341
	Lactation	0.7067	0.4869	0.9255
	Anti-Tc IgA	0.08	-0.0294	0.1894

	Total IgA	-0.0593	-0.1619	0.0429
	Strongyles	0.0772	0.0102	0.1444
	Age	0.1201	-0.008	0.249
	Year: 2016	0.1251	-0.0301	0.2797
	Year: 2017	0.085	-0.0971	0.2668

588

589 Table SI1: effect sizes and 95% credibility intervals for the component GLMMs of each path
590 analysis

591