Spatial Distribution of Discarded Vehicle Tires and Their Influence on *Aedes albopictus* and *Culex quinquefasciatus* Populations, New Orleans, Louisiana

Mohamed F. Sallam^{*1}, Tamer Ahmed², Cynthia Sylvain-Lear³, Claudia Riegel², and Imelda K. Moise⁴

¹Navy Entomology Center of Excellence, Jacksonville, FL 32212

²New Orleans Mosquito, Termite and Rodent Control Board, 2100 Leon C. Simon
New Orleans, LA 70122

³Department of Sanitation, City of New Orleans, LA 70122

⁴Department of Geography and Regional Studies, College of Arts and Sciences, University of
Miami, 1300 Campo Sano Ave, Coral Gables, FL 33124.

* Correspondence should be addressed to mwsallam2009@aol.com,
Mohamed.e.sallam.mil@mail.mil

Abstract

Discarded vehicle tires play an important role in the colonization of container mosquito populations, particularly their geographic expansion. We assessed the spatial distribution of illegally discarded tires and their response to land use-land cover (LULC), and demographic factors using geospatial analysis and generalized regression. Multiple stepwise regressions were used to evaluate the response of the Container Index (CI) of colonized *Aedes albopictus* (Skuse), and *Culex quinquefasciatus* Say to macro- and microhabitats variables. The illegally discarded tires were distributed over 11 planning districts with clustering distribution for tires frequency and overdispersed distribution for tires number. Out of 1,137 (~37.08%) water-holding tires, 598 (~52.64%) tires at 65 (~38.46%) sites were positive for colonized mosquito populations. A total of 13 mosquito species were identified, with the highest CI of *Ae. albopictus* (44.19%) and *Cx. quinquefasciatus* (22.18%). *Aedes albopictus* colonized all 65 sample sites and *Cx. quinquefasciatus* found at 32 sites. The Container Index (CI) of colonized mosquito was clustered in seven planning districts for *Ae. albopictus* and five planning districts for *Cx. quinquefasciatus*. Microhabitat (muddy water) rather than macrohabitats variables predicted both species' colonization, especially *Ae. albopictus*. The contribution of macro- and microhabitat characteristics in predicting colonized mosquito in water-holding tires was discussed.

Keywords: arbovirus, *Aedes*, container, *Culex*, invasive, GIS, clustering

Introduction

Discarded vehicle tires play an important role in the proliferation of mosquito colonization and breeding (Shannon 1931), particularly the introduction and geographic expansion of mosquito species. The most striking example of this is the introduction of the Asian tiger mosquito, *Aedes albopictus* (Skuse) (Hawley et al. 1987) and *Ae. (Ae) japonicus* (Theobald) (Peyton et al. 1999) to the United States (US) from Asia and South Pacific. Tires are also a habitat for other mosquito species in the continental US (e.g., *Culex (Cx.) restuans*, *Cx. pipiens*, *Cx. territans*, *Cx. Salinarius* and *Cx. quinquefasciatus*) (Yee 2008). The introduction of *Ae. albopictus* to North America after World War II set the stage for the transmission of arboviruses, causing both wildlife and human disease outbreaks such as dengue, chikungunya (Paupy et al. 2009), and La Crosse encephalitis (Gerhardt et al. 2001).

Of particular concern is the durability of tires as water-holding habitats as containers for mosquitos that vector diseases. Because of their specific shape and impermeable nature, they can produce adult mosquitoes over longer periods than natural containers (e.g., leaf axils, tree holes) (Sofi 2018) or even other artificial containers such as plastic trash and food containers (Dowling et al. 2013). This has been relevant to public health because tires are most likely to be discarded near human populations and hence pose health risks (Yee 2008, Villena et al. 2017). Studies over the past two decades have provided important information on the link between the location of tires and female oviposition behavior (Kling et al. 2007, Day 2016, Allgood and Yee 2017), but few have investigated variation in the density and distributions of larvae in discarded tires (Lampman et al. 1997, Qualls and Mullen 2006).

While other studies highlighted the spatial distribution of mosquito populations in tire systems and their response to environmental variables in the U.S (Juliano and Lounibos 2005, Qualls and Mullen 2006, Bartlett-Healy et al. 2012, Yee et al. 2012) and Latin America (Manrique-Saide et al. 2008, Troyo et al. 2008, Mendonça et al. 2011, García-Rejón et al. 2012, Baak-Baak et al. 2014), they did not give detailed information on i) the frequency and density of discarded vehicle tires, ii) the general effect of the interaction between landscape and demographic subsystems on discarded tires and mosquito populations, and iii) the interplay between macro- and microhabitat characteristics and their impact on mosquito populations.

In this study, we evaluated the geospatial distribution patterns of frequency and number of discarded vehicle tires, and tires response to land use-land cover (LULC), and demographic predictors. Additionally, we assessed the response of the container index of colonized *Ae. albopictus* and *Cx. quinquefasciatus* populations to their predicting factors within macro- and microhabitat characteristics. Finally, we evaluated the interaction between macro- and microhabitat characteristics affecting the colonization of these two mosquito populations.

Materials and Methods

Study Area

The city of New Orleans Louisiana (NOLA) lies on the Mississippi River, near the Gulf of Mexico with a total land area of 438.80 km². In 2017, NOLA was inhabited by almost 343,829 people, with an average density of 892.2/km². NOLA is sub-tropical with an annual high temperature of 25°C, an annual low of 16.8°C, and average annual precipitation of 162.3 cm. The average highest precipitation occurs in July (17.9 cm).

Discarded Vehicle Tire Surveillance

We obtained 311 records of illegal discarded tire sites from the City of New Orleans through crowdsourcing data from local government and non-emergency information. The City of New Orleans Department of Sanitation reported and confirmed the calls regarding illegally discarded tires reported to the city's 311-call center or via email by residents, visitors, or businesses (www.nola.gov/311).

A purposive tire sampling based on 311 records was conducted for confirmation. Additionally, random surveillance was conducted within the reported housing blocks and neighborhood of the reported sites: i) to explore other dumping sites that have not been reported, and ii) to represent different landscape characteristics of dumping sites. The random surveillance was conducted within pre-defined sampling grids using the sampling design tool in ArcGIS ver. 10.1 (ESRI 2011) (Figure 1). Data records for discarded vehicle tires were compiled from both the 311-call database and active random surveillance during January 2015-January 2018. During this study, we did not account for legally retained or illegally dumped tires in areas not accessible such as fenced private premises.

Mosquito Sampling

All water-holding tires were sampled for mosquito colonization and container index (CI) estimates from November 2016 to January 2018. During surveillance, three types of tires were sampled : A) light truck, 8-ply non-radial, 70 cm diam x 17 cm wide; B) radial automobile, 56 x 17.5 cm, and C) tractor, 128 x 70 cm. Physical characters of water, such as pH, temperature, and water quality, were collected before sampling the colonized mosquitos. Tire water was mixed before sampling to obtain bottom-dwelling mosquito immatures. A destructive sampling method was used to collect all biological materials from water-holding tires. All water contents from the

tires were strained and transferred into Nasco Whirl-Pak bags to the New Orleans Mosquito, Termite Control Board (NOMTCB) laboratory for further sample assortment, and identification. The number of larvae and pupae was quantified at our facility. Only fourth larval instars were used for the identification of species. Early instars that were not readily identifiable were reared in plastic mosquito breeders at $27^{\circ}\text{C}\pm 2$ and 80 ± 5 RH% to the fourth instar before being identified. Pupae were similarly maintained and identified as adults. Larval and adult stages were identified by species level using the taxonomic keys of Darsie and Ward (Darsie and Ward 2005). All possible predators were quantified and recorded for each sampled tire.

Microhabitat Data Variables

Microhabitats variables were represented as seven biophysical factors: size of tires (small, large, mixed), tire status (damaged, intact), water temperature, pH value, water quality (muddy, turbid, clear), predators (midges, water fleas, midges/water fleas), and CI of colonized mosquito species (Table 1).

Macrohabitat Data Variables

Land Use-Land Cover (LULC) and Demographic Data

Because adult female mosquitoes require a blood meal for egg development (Zhu et al. 2014), we derived the recently updated (2007) LULC from the US Geological Survey (USGS) (Burkett-Cadena et al. 2010) to delineate land use and land cover patterns within the New Orleans metropolitan area, to identify areas with a higher likelihood of finding a blood meal. Urban areas were categorized into three classes to represent the degree of urbanization: (1) residential and urban, in which housing predominates in two different forms; (2) industrial and commercial services, which represents industrial settings and fewer housing structures; and (3)

other urban and built-up land with the least housing and human populations. We then used the National Land Cover Database 2011 (NLCD) to define five major classes of LULC to represent environmental predictors linked to high vector densities (Ruiz et al. 2004). These five major classes are: 1) developed high intensity, 2) developed medium intensity, 3) developed low intensity, 4) developed open space, and 5) open water areas.

Further, because vegetated vacant lots at sample sites may represent resting places and sugar meal sources for adult mosquitoes (Sallam et al. 2016) and possible disposal sites, two major classes of vegetation types were extracted from the USGS (Lewis et al. 2017). The two extracted vegetation types include non-forested wetlands and forest wetlands. Further, we included streams, canals, lakes, reservoirs, and estuaries of different sizes (>1 and ≤ 1 km) to represent permanent water bodies as possible illegal discarded tire sites (Figure 1). Demographic data (human population and housing density) at the block level was downloaded from the US Census American Community Survey (ACS) for 2012-2016.

Data Analysis

Geospatial Distribution Analysis.

We were specifically interested in evaluating the geospatial distribution of frequency (number of dumping incidences per site) and the number of illegally discarded tires. Additionally, we assessed the CI of two dominant mosquito populations in the tire systems. First, we measured the geographic distribution of illegally discarded tires within areas with an elevated risk of having a high frequency of discarded tires and reported the number using the standard deviational ellipse (SDE) statistical tools in ArcGIS. Second, we used Global Moran's I (Anselin and Getis 1992), a correlation coefficient that measures the overall spatial autocorrelation of locations (e.g., illegal

tire sites) and feature values (e.g., frequency and reported the number of tires) concurrently. Clusters of sites with a high frequency and density of illegally discarded tires were considered “high-risk areas” whereas clusters of sites with low frequency and density of illegally discarded tires (Low-Low) were considered “low-risk sites.” The value of Moran’s I is measured between -1 and 1, with the ‘Z’ score value calculated to test whether the observed clustering or dispersal is significant. Therefore, when the Z score indicates statistical significance, a positive Moran’s I indicates a tendency toward clustering while a negative value indicates a tendency toward dispersion. When the Z score value is not significantly different from zero, it is assumed that there is no spatial autocorrelation, and the pattern does not appear to be significantly different from a random distribution.

Similarly, the spatial distribution and Global Moran’s I index for the CI of *Ae. albopictus* and *Cx. quinquefasciatus* populations were evaluated. The clustering hypothesis (H_1) was tested for the frequency and density of discarded vehicle tires and the colonized mosquito populations. The null hypothesis (H_0) was that there is no spatial clustering of frequency and density of discarded vehicle tires or the CI of colonized mosquito populations in the city of New Orleans.

Statistical Analysis

To characterize the influence of the 13 LULC and demographic predictors that predict discarded tire frequency and number, two generalized regression models (GRM) with lasso tool were carried out to delineate their response to predicting factors from the sampling sites.

On the other hand, multiple stepwise linear regressions were conducted to evaluate: (1) the influence of macrohabitat characteristics on the CI of *Ae. albopictus* and *Cx. quinquefasciatus* population, (2) the influence of microhabitat characteristics on the CI of *Ae. albopictus* and *Cx.*

quinquefasciatus colonization, and (3) the interaction between macro- and microhabitat predictors and their influence on the CI of *Ae. albopictus* and *Cx. quinquefasciatus* in water-containing tires. For colonized mosquito populations, macrohabitats were denoted as a total of 13 LULC and demographic data layers, and microhabitats variables were represented as seven biophysical factors (Table 1).

The $\log_{10}(n+1)$ formula was used to transform the number of tires and area percentages of LULC to maintain the normality of their statistical distribution. All statistical analyses were conducted in SAS 14.0 (SAS 2013) using the JMP pro statistical package. Significant predictors and best prediction models were selected based on minimum corrected Akaike Information Criterion (AICc) and maximum r^2 values, with $p < 0.05$. All variables were standardized for the average and standard deviation become 1 and 0.

Results

Dumping Tires and Mosquito Sampling

Four hundred and five sites with 6,491 discarded tires were surveyed during the current study representing 13 New Orleans planning districts. A total of 236 (~58.27%) sites with 3,427 (~52.80%) tires were reported by the 311-phone service. Additionally, 3,064 (~47.20%) discarded tires were actively surveyed for mosquito immature colonization from 169 sites (~41.73%). The dumping incidences of tires reported from all sites were either one or two times, representing 64 and 341 sites, respectively.

Tire discarding sites reported by 311 services could not be surveyed for mosquito immatures because the department of sanitation of the city of New Orleans removed tires at these sites. Of

the 3,064 surveyed tires, 1,136 (~37.08%) were found to contain water, and 598 (~52.64%) of water containing tires were positive for mosquito immatures colonization.

We observed positive mosquito colonization at 65 (~38.46%) sampling sites with 2,900 mosquito immatures (larvae/pupae) in sampled tires. These tires provided suitable colonization habitats for 13 mosquito species with a new locality record of *Aedes japonicus*³⁵ (Table 2).

Colonized *Ae. Albopictus* populations were the highest (CI = ~44.19), followed by *Cx.*

quinquefasciatus (CI = ~22.18). *Aedes albopictus* was recorded from 502 tires at 65 sites.

Whereas, *Cx. quinquefasciatus* was collected from 252 tires at 32 sites. The dominance of *Ae. albopictus* population also was demonstrated on spatial scale, as it was collected from seven districts (~54%). In the meantime, *Cx. quinquefasciatus* was reported in five districts (~39%).

Both species co-existed in 21 sampling sites distributed in four planning districts : Bywater, Gentilly, Lower Ninth Ward, and New Orleans East.

Geospatial Distribution Analysis

We used the Moran's I to measure global clustering patterns of illegal discarded tire frequency.

We found spatial clustering of dumping frequency (Z-score = 1.76, $p < 0.10$). Most of these discarded tires were in areas of the 13 New Orleans planning districts (Figure 2). In contrast, the number of discarded tires showed a random spatial distribution (Z-score value = -0.17, $p > 0.05$).

However, the concentration centers of illegally discarded tire frequency and reported numbers were in Mid-city. We also found that certain areas of New Orleans have an elevated risk for discarded tires in both frequency and numbers. This area covers a total area of 24.8 Km² (5.64%) and 23.3 Km² (5.31%) of the total land area of New Orleans and is mostly located in 11 planning districts.

Similarly, spatial autocorrelation and geographic distribution patterns for the CI of colonized *Ae. albopictus* and *Cx. quinquefasciatus* populations were evaluated to test the clustering hypothesis (H_1). Only the CI of *Ae. albopictus* showed the significant clustered distribution and the center of concentration was found to be in Bywater.

Meanwhile, the CI of *Cx. quinquefasciatus* demonstrated a random distribution pattern, and the concentration center of its distribution also was found to be in Bywater. The distance between mean centers of concentration for both species was 0.47 Km.

The clustered and random distribution patterns were confirmed by the significant positive Z-score values (*Ae. albopictus*= 3.61, *Cx. quinquefasciatus*= 1.62, $p < 0.05$). Accordingly, the clustering hypothesis (H_1) for the CI of *Ae. albopictus* colonized in discarded tires was accepted.

The geospatial distribution demonstrated by SDE polygons showed areas under high risk of increased CI of colonized *Ae. albopictus* and *Cx. quinquefasciatus* covered 55.19 Km² (~12.58%) and 41.88 Km² (~9.54%), respectively, of the total area of the city. Areas under risk of both mosquito populations that colonized discarded tires were found to represent 39.95 Km² (9.10%) of the studied sites. These areas were found to represent seven planning districts:

Bywater, Gentilly, Lower Ninth Ward, Mid-City, New Orleans East, Algiers, and Venetian Isles (Figures 3 and 4). The SDEs of both mosquito populations were found to overlap in an area of 36.06 Km² (~8.22%). This finding reflects the co-existence of both species within the same sampling sites and tire habitats.

Statistical Regression Models

Spatial Characteristics of Illegal Discarded Tires

Two factors were significantly associated with illegally discarded tire frequency ($r^2=4.58\%$, $AICc=321.63$, $p<0.05$). Industrial areas were positively associated with discarded tire frequency (Table 3). None of LULC variables showed any significant association with discarded tire frequency. Meanwhile, five variables were significantly associated with the increased number of dumping tires ($r^2=3.52\%$, $AICc=1237.24$, $p<0.05$). All variables showed a negative association with an increased number of tires dumping with the exception of industrial areas (Table 3).

Spatial Characteristics of Mosquito Populations

The scale of regression models, whether macro- or microhabitats, influenced the number and type of predicting variables for the colonization of each mosquito species. For *Ae. albopictus*, two macrohabitat variables have a significant maximum influence in predicting the CI. These variables were developed medium intensity ($\beta= 7.16E-6$, $r^2=5.23\%$, $AICc=1674$, $p<0.05$), and developed low intensity ($\beta= -1.75E-6$, $r^2=6.64\%$, $AICc=1673$, $p<0.05$) as indicated by r^2 and $AICc$ values. Developed low-intensity areas were found to be negatively associated with the increased CI of colonized *Ae. albopictus* population (Table 4).

On the other hand, the microhabitat variables demonstrated an increased prediction effect on the CI of colonized *Ae. albopictus* ($r^2= 23.23\%$, $AICc=1647$, $p<0.05$). Five predicting variables were found to increase colonized *Ae. albopictus* populations (Table 4). The water quality demonstrated the maximum prediction power with the highest r^2 and minimum $AICc$ values ($\beta= 5.5$, $r^2=23.23\%$, $AICc=1647$, $p<0.05$). Additionally, the presence of other colonized mosquito populations in water-containing tires used tire status and water alkaline pH value shared reduced prediction gain in increasing CI of colonized *Ae. albopictus*.

The combined regression model approach highlighted how the interaction between macro- and microhabitat variables could change the number and type of influential factors that affect mosquito density. The combined regression model showed the best prediction power in characterizing suitable habitats for the colonization of *Ae. albopictus* ($r^2 = 25.40\%$, $AICc = 1640$, $p < 0.01$). The combined model demonstrated the contribution of three microhabitats and one macrohabitat variables in affecting the CI (Table 4). Accordingly, the combined regression approach was selected as the best prediction model because of the recorded minimum $AICc$ value for the most predictive variable.

For *Cx. quinquefasciatus*, the developed medium intensity areas were the only macrohabitat variable that significantly predict increased CI values ($\beta = 1.69$, $r^2 = 6.0\%$, $AICc = 1673$, $p < 0.05$). Four microhabitat variables were found to be the most influential factors in predicting increased CI values ($r^2 = 18.68\%$, $AICc = 1567$, $p < 0.05$). Muddy and clear water quality demonstrated the highest prediction gain in affecting the colonization of *Cx. quinquefasciatus* ($\beta = 6.46$, $r^2 = 18.68\%$, $AICc = 1567$, $p < 0.05$). Additionally, intact tire status (undamaged), large tire size (tractor, 128 x 70 cm), and CI values of colonized *Ae. albopictus* were positively correlated with increased CI values of *Cx. quinquefasciatus* (Table 5). The combined regression model showed that the four microhabitat variables superseded the developed medium intensity areas in predicting increased CI values. The combined regression approach was selected as the best prediction model because of the recorded minimum $AICc$ value.

Discussion

To our knowledge, this study is the first in New Orleans to investigate geographic distribution patterns, habitat characteristics and ecological suitability of illegally discarded vehicle tires and

their associated mosquito populations. Previous work in New Orleans highlighted natural *Ae. albopictus* population in tires (Marten 1990b, a, Comiskey et al. 1999) and container-breeding *Ae. aegypti* (Focks et al. 1981). Focks et al. (1981) highlighted the significance of tires as compared to other available water-holding containers as potential breeding sites for *Ae. aegypti* and *Cx. quinquefasciatus*. Other local reports conducted by the City of New Orleans and department of health concluded that tires are strong breeding sites for mosquitoes as they are easily filled with water and collect leaf litter, providing a prime spot for mosquito vectors that can transmit West Nile, Zika and dengue (NOLA 2019). However, the previous studies either focused on biological control or the biology of these mosquito populations in tire systems. Our preliminary findings characterized areas under risk of increased frequency and number of discarded tires and two mosquito vectors populations. We investigated the number and type of influential factors that predict the container index (CI) of colonized *Ae. albopictus* and *Cx. quinquefasciatus* populations within their macro- and microhabitats. We also highlighted the significance of evaluating the interaction between macro- and microhabitat variables in predicting increased CI values of the colonized mosquito populations.

The conflicting findings between the clustered patterns of the dumping frequency and their number are possible because of the impact of landscape and demographic factors such as housing density or vegetation. Adding the number of tires to the algorithm of spatial autocorrelation reflects the magnitude of influence of certain landscape and demographic factors. For example, average housing density, other built-up urban and residential/urban were negatively associated with increased dumping frequency and number of discarded tires. A possible explanation for this is that highly populated areas were either not suitable as dumping sites or people have the

tendency to discard tires in areas with low population density. This finding would improve our understanding of their influence in predicting the likelihood of an increased number of dumped tires. This would also shed light on the significant negative association of other urban built-up areas and average housing density with the dumping problem in certain areas. We recommend prioritizing neighborhoods with similar LULC types for surveillance and control efforts.

The current study found that industrial areas were positively associated with both discarded tire frequency and number. No correlations were found between the frequency or number of dumped tires and population density. Additionally, the negative associations between the number of dumped tires and housing density, or residential and urban settings reflect that the number of dumping reports were not correlated with populated areas. Previous studies in other areas showed that the dumping frequency and number of tires were positively associated with grown vegetation in vacant lots and/or community perceptions, as people would store a variety of containers in their yard (Andreadis 1988, Mazine et al. 1996). For example, discarded tire frequency and density increased in neighborhoods with lower housing density. This finding is confirmed by the significant negative association between developed medium areas with increased discarded tire frequency and number. These patterns of discarded tire frequency and number in New Orleans have ecological consequences that affect both ecosystem services and human health (Lewis et al. 2017). However, further studies are needed to include the influence of other socioeconomic factors such as community perception, income rates, and education level. This not only will help in highlighting high risk areas of dumping frequency but also will shed the light on community engagement in the proliferation of illegally discarded tires.

Out of 65 positive mosquito sites, both *Ae. albopictus* and *Cx. quinquefasciatus* co-existed in 21 sampling sites representing 214 tires (35.79%) from four planning districts: Bywater, Gentilly, Lower Ninth Ward, and New Orleans East. The clustering pattern of CI for *Ae. albopictus* demonstrates their association with certain biophysical characters that predict their distribution. Meanwhile, the random distribution pattern of *Cx. quinquefasciatus* generated by standard deviational ellipsoid may indicate their response to microhabitat characteristics or the contribution of other potential habitats that predict their colonization, which should be addressed in a separate investigation.

The consistent interactions between macro- and microhabitat variables were demonstrated by the variation in the number and type of variables that predict the CI of mosquito populations. For example, our regression models showed that the distribution range of *Ae. albopictus* was predicted by a combination of macro- and microhabitat variables such as muddy water quality ($r^2=25.40$) and developed medium intensity areas ($r^2=22.44$) (Sallam et al. 2017b). These findings were consistent with other researches, which found a link between *Ae. albopictus* population density and water quality (e.g., organic muddy, dark turbid) (Higa et al. 2010, Nasir et al. 2017), developed medium intensity (Comiskey et al. 1999), and pH (Honório et al. 2006). Our findings were confirmed by the maximum r^2 values generated by combined regression models (Tables 4 and 5). This finding broadly supports the work of other studies in this area linking microhabitat suitability with *Ae. albopictus* breeding in discarded tires (Comiskey et al. 1999, Vezzani et al. 2005, Maciel-de-Freitas et al. 2007, Vezzani and Albiocco 2009, Dória et al. 2010, Murrell et al. 2011). This mosquito vector was found to be affected by detritus and tires with high nutrient contents (Merritt et al. 1992, Mazine et al. 1996).

In contrast, the microhabitat factors predicted the presence of *Cx. quinquefasciatus*. In particular, although macrohabitat factors demonstrated high prediction gain ($r^2=6.0$, AICc=1673), the combination of macro- and microhabitat variables improved the prediction gain, and the combined regression model was selected as the best model in delineating habitat factors for *Cx. quinquefasciatus* ($r^2=18.67$, AICc=1566). This model indicated an intact/undamaged tire status and muddy/clear water quality as the best predictors for *Cx. quinquefasciatus*. *Culex. quinquefasciatus* also occurred at much lower densities and exhibited a much smaller geographic range that was indicated by the standard deviational ellipsoid. This finding supports evidence from the previous observations (Merritt et al. 1992, Qualls and Mullen 2006). However, the predicting variables of *Cx. quinquefasciatus* in water-containing habitats seemed to be context dependent in terms of interspecific competition with other mosquito populations, host preference, type of breeding sites and season (Sallam et al. 2017a). This mosquito species showed resilience to ecological requirements for their presence and abundance and demonstrated biological fitness to breed in a wide range of breeding sites. Another study need to be conducted to quantify all potential breeding sites of *Cx. quinquefasciatus* in the City of New Orelans to understand their predicting factors in their breeding sites.

Generally, the presence of both *Ae. albopictus* and *Cx. quinquefasciatus* in discarded tires was influenced much more by microhabitat factors than macrohabitat factors. This is important because mosquito dietetic necessities are met through the intake of both dead and living organic material (Merritt et al. 1992, Mazine et al. 1996, Yee et al. 2004). This finding warrants the need to not only reinforce discarded tire pick-up initiatives in New Orleans but also control of mosquito breeding in this urban landscape by exploring water quality conditions of *Aedes* and

Culex larval habitats that proliferate mosquitoes. Our study is a preliminary step for a comprehensive model that integrates crowd-sourcing data with active surveillance to more completely understand Spatio-temporal distribution of container breeding mosquitos in their potential colonization sites.

Conclusion

Our results indicate that although the problem of discarded vehicle tires is widespread (in both frequency and number) in the 13 planning districts of New Orleans, their center of concentration was found to be in Mid-city. We identified 13 mosquito species from 65 sites (38.46%) representing 1,136 (~37.08%) sampled tires in seven of the planning districts. The two most dominant mosquito species, *Ae. albopictus* and *Cx. quinquefasciatus*, were found to co-exist in 21 sampling sites (~32.31%) representing 214 tires (35.79%) from four planning districts: Bywater, Gentilly, Lower Ninth Ward, and New Orleans East. The clustered and random spatial distribution polygons of *Ae. albopictus* and *Cx. quinquefasciatus* were found to be in seven planning districts. Both species were significantly predicted by microhabitat characteristics rather than macrohabitats, especially *Ae. albopictus*. Given the diverse mosquito fauna of Louisiana, we recommend rigorous ecological studies be conducted to identify ecological and socioeconomic factors that may influence species composition within and among sites and their temporal distribution pattern in New Orleans. This is a critical step towards understanding the contribution of macro- and microhabitat characteristics in predicting colonized mosquito populations in water-holding tires. Also, this study adds to the growing body of research that indicates that different factors found in the tire environments influence oviposition and/or larval performance.

Acknowledgments

We thank Matt Torri and the employees of the City of New Orleans Sanitation Department, the City of New Orleans 311 Call Center, and City of New Orleans Mosquito and Termite Control Board staff who helped to locate tire sites and assist with mosquito identification. A special thank you to all the property owners for allowing us access to study sites. This work was partially supported by Grant ZIKA-CDC-RFA-CK14-140199HF, building domestic surveillance, laboratory, vector control, and pregnancy registry capacity to respond to Zika virus. The views expressed in this article are those of the author and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U. S. Government.

Data Availability

The authors declare that all the relevant data supporting the study findings are available from the City of New Orleans upon request.

Authors' contributions

TA, CSL, and CR carried out the experimental design and data sampling. MFS: carried out geo-database and model building, data analysis, and writing the first draft of the manuscript. IKM helped in validating models and editing manuscript. All authors approved the final version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Ethics approval and consent to participate

Not applicable

References

- Allgood, D. W., and D. A. Yee. 2017. Oviposition preference and offspring performance in container breeding mosquitoes: evaluating the effects of organic compounds and laboratory colonisation. *Ecological Entomology* 42: 506-516.
- Andreadis, T. G. 1988. A survey of mosquitoes breeding in used tire stockpiles in Connecticut. *J Am Mosq Cont. Assoc.* 4: 256-260.
- Anselin, L., and A. Getis. 1992. Spatial statistical analysis and geographic information systems. *The Annals of Regional Science* 26: 19-33.
- Baak-Baak, C. M., R. Arana-Guardia, N. Cigarroa-Toledo, M. Puc-Tinal, C. Coba-Tún, V. Rivero-Orsorno, D. Lavallo-Kantun, M. A. Loroño-Pino, C. Machain-Williams, G. C. Reyes-Solis, B. J. Beaty, L. Eisen, and J. E. García-Rejón. 2014. Urban mosquito fauna in Mérida City, México: Immatures collected from containers and storm-water drains/catch basins. *Southwestern Entomologist* 39: 291-306, 216.
- Bartlett-Healy, K., I. Unlu, P. Obenauer, T. Hughes, S. Healy, T. Crepeau, A. Farajollahi, B. Kesavaraju, D. Fonseca, G. Schoeler, R. Gaugler, and D. Strickman. 2012. Larval mosquito habitat utilization and community dynamics of *Aedes albopictus* and *Aedes japonicus* (Diptera: Culicidae). *Journal of Medical Entomology* 49: 813-824.
- Burkett-Cadena, N. D., R. A. Ligon, M. Liu, H. K. Hassan, G. E. Hill, M. D. Eubanks, and T. R. Unnasch. 2010. Vector-host interactions in avian nests: do mosquitoes prefer nestlings over adults? *The American journal of tropical medicine and hygiene* 83: 395-399.
- Comiskey, N. M., R. C. Lowrie, Jr., and D. M. Wesson. 1999. Role of habitat components on the dynamics of *Aedes albopictus* (Diptera: Culicidae) from New Orleans. *Journal of Medical Entomology* 36: 313-320.
- Darsie, R. F., and R. A. Ward. 2005. Identification and geographical distribution of the mosquitos of North America, North of Mexico, University Press of Florida.
- Day, J. F. 2016. Mosquito oviposition behavior and vector control. *Insects* 7: 65.
- Dória, G. A. A., W. J. Silva, G. A. Carvalho, P. B. Alves, and S. C. H. Cavalcanti. 2010. A study of the larvicidal activity of two Croton species from northeastern Brazil against *Aedes aegypti*. *Pharmaceutical Biology* 48: 615-620.
- Dowling, Z., P. Armbruster, S. L. LaDeau, M. DeCotiis, J. Mottley, and P. T. Leisnham. 2013. Linking mosquito infestation to resident socioeconomic status, knowledge, and source reduction practices in suburban Washington, DC. *EcoHealth* 10: 36-47.
- ESRI. 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.
- Focks, D. A., S. R. Sackett, D. L. Bailey, and D. A. Dame. 1981. Observations on container-breeding mosquitoes in New Orleans, Louisiana, with an estimate of the population density of *Aedes Aegypti* (L.). *The American journal of tropical medicine and hygiene* 30: 1329-1335.
- García-Rejón, J. E., M. P. López-Urbe, M. A. Loroño-Pino, R. Arana-Guardia, M. Puc-Tinal, G. M. López-Urbe, C. Coba-Tún, C. M. Baak-Baak, C. Machain-Williams, G. C. Reyes-Solis, S. Lozano-Fuentes, K. Saavedra-Rodriguez, W. C. t. Black, B. J. Beaty, and L. Eisen. 2012. *Aedes* (Stegomyia) *aegypti* and *Aedes* (Howardina) *cozumelensis* in Yucatán State, México, with a summary of published collection records for *Ae. cozumelensis*. *Journal of vector ecology : journal of the Society for Vector Ecology* 37: 365-372.
- Gerhardt, R. R., K. L. Gottfried, C. S. Apperson, B. S. Davis, P. C. Erwin, A. B. Smith, N. A. Panella, E. E. Powell, and R. S. Nasci. 2001. First isolation of La Crosse virus from naturally infected *Aedes albopictus*. *Emerging Infectious Diseases* 7: 807-811.
- Hawley, W. A., P. Reiter, R. S. Copeland, C. B. Pumpuni, and G. B. Craig. 1987. *Aedes albopictus* in North America: probable introduction in used tires from northern Asia. *Science* 236: 1114-1116.

- Higa, Y., N. T. Yen, H. Kawada, T. H. Son, N. T. Hoa, and M. Takagi. 2010. Geographic distribution of *Aedes aegypti* and *Aedes albopictus* collected from used tires in Vietnam. *Journal of the American Mosquito Control Association* 26: 1-9, 9.
- Honório, N. A., P. H. Cabello, C. T. Codeço, and R. Lourenço-de-Oliveira. 2006. Preliminary data on the performance of *Aedes aegypti* and *Aedes albopictus* immatures developing in water-filled tires in Rio de Janeiro. *Memórias do Instituto Oswaldo Cruz* 101: 225-228.
- Juliano, S. A., and L. P. Lounibos. 2005. Ecology of invasive mosquitoes: effects on resident species and on human health. *Ecology Letters* 8: 558-574.
- Kling, L. J., S. A. Juliano, and D. A. Yee. 2007. Larval mosquito communities in discarded vehicle tires in a forested and unforested site: detritus type, amount, and water nutrient differences. *Journal of vector ecology : journal of the Society for Vector Ecology* 32: 207-217.
- Lampman, R., S. Hanson, and R. Novak. 1997. Seasonal abundance and distribution of mosquitoes at a rural waste tire site in Illinois. *Journal of the American Mosquito Control Association* 13: 193-200.
- Lewis, J. A., W. C. Zipperer, H. Ernstson, B. Bernik, R. Hazen, T. Elmqvist, and M. J. Blum. 2017. Socioecological disparities in New Orleans following Hurricane Katrina. *Ecosphere* 8: e01922.
- Maciel-de-Freitas, R., W. A. Marques, R. C. Peres, S. P. Cunha, and R. Lourenço-de-Oliveira. 2007. Variation in *Aedes aegypti* (Diptera: Culicidae) container productivity in a slum and a suburban district of Rio de Janeiro during dry and wet seasons. *Memórias do Instituto Oswaldo Cruz* 102: 489-496.
- Manrique-Saide, P., C. R. Davies, P. G. Coleman, E. Rebollar-Tellez, A. Che-Medoza, F. Dzul-Manzanilla, and A. Zapata-Peniche. 2008. Pupal Surveys for *Aedes aegypti*: Surveillance and potential targeted control in residential areas of Mérida, México. *Journal of the American Mosquito Control Association* 24: 289-298, 210.
- Marten, G. G. 1990a. Evaluation of cyclopoid copepods for *Aedes albopictus* control in tires. *J Am Mosq Cont. Assoc.* 6: 681-688.
- Marten, G. G. 1990b. Elimination of *Aedes albopictus* from tire piles by introducing *Macrocyclus albidus* (Copepoda, Cyclopidae). *J Am Mosq Cont. Assoc.* 6: 689-693.
- Mazine, C. A. B., M. L. G. Macoris, M. T. M. Andrighetti, S. Yasumaro, M. E. Silva, M. J. Nelson, and P. J. Winch. 1996. Disposable containers as larval habitats for *Aedes aegypti* in a city with regular refuse collection: a study in Marília, Sao Paulo State, Brazil. *Acta Tropica* 62: 1-13.
- Mendonça, H. F. M. S. d., A. L. Ferreira, C. B. d. Santos, H. R. Rezende, G. E. M. Ferreira, G. R. Leite, and A. Falqueto. 2011. Breeding sites of *Aedes aegypti* in metropolitan vacant lots in Greater Vitória, State of Espírito Santo, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical* 44: 243-246.
- Merritt, R. W., R. H. Dadd, and E. D. Walker. 1992. Feeding behavior, natural food, and nutritional relationships of larval mosquitoes. *Annual Review of Entomology* 37: 349-374.
- Murrell, E. G., K. Damal, L. P. Lounibos, and S. A. Juliano. 2011. Distributions of competing container mosquitoes depend on detritus types, nutrient ratios, and food availability. *Annals of the Entomological Society of America* 104: 688-698.
- Nasir, S., F. Jabeen, S. Abbas, I. Nasir, and M. Debboun. 2017. Effect of climatic conditions and water bodies on population dynamics of the dengue vector, *Aedes aegypti* (Diptera: Culicidae). *Journal of Arthropod-Borne Diseases* 11: 50-59.
- NOLA. 2019. Recent rains provide breeding grounds for mosquitoes. <https://midcitymessenger.com/2019/07/15/recent-rains-provide-breeding-grounds-for-mosquitoes/>.
- Paupy, C., H. Delatte, L. Bagny, V. Corbel, and D. Fontenille. 2009. *Aedes albopictus*, an arbovirus vector: From the darkness to the light. *Microbes and Infection* 11: 1177-1185.
- Peyton, E. L., S. R. Campbell, T. M. Candeletti, M. Romanowski, and W. J. Crans. 1999. *Aedes* (Finlaya) *japonicus japonicus* (Theobald), a new introduction into the United States. *Journal of American Mosquito Association* 15: 238-241.

- Qualls, W. A., and G. R. Mullen. 2006. Larval survey of tire-breeding mosquitoes in Alabama. *Journal of the American Mosquito Control Association* 22: 601-608, 608.
- Ruiz, M. O., C. Tedesco, T. J. McTighe, C. Austin, and U. Kitron. 2004. Environmental and social determinants of human risk during a West Nile virus outbreak in the greater Chicago area, 2002. *International Journal of Health Geographics* 3: 8-8.
- Sallam, M., S. Michaels, C. Riegel, R. Pereira, W. Zipperer, B. Lockaby, and P. Koehler. 2017a. Spatio-Temporal Distribution of Vector-Host Contact (VHC) Ratios and Ecological Niche Modeling of the West Nile Virus Mosquito Vector, *Culex quinquefasciatus*, in the City of New Orleans, LA, USA. *International Journal of Environmental Research and Public Health* 14: 892.
- Sallam, M. F., R.-D. Xue, R. M. Pereira, and P. G. Koehler. 2016. Ecological niche modeling of mosquito vectors of West Nile virus in St. John's County, Florida, USA. *Parasites & Vectors* 9: 371.
- Sallam, M. F., C. Fizer, A. N. Pilant, and P.-Y. Whung. 2017b. Systematic Review: Land cover, meteorological, and socioeconomic determinants of *Aedes* mosquito habitat for risk mapping. *Int. J. Environ. Res. Public Health* 14.
- SAS. 2013. SAS Institute Inc., Cary, NC, USA.
- Shannon, R. C. 1931. The classification of Brazilian Culicidae with special reference to those capable of harboring the yellow fever virus. *Proc. Ent. Soc. Washington* 33: 150-152.
- Sofi, A. 2018. Effect of waste tyre rubber on mechanical and durability properties of concrete – A review. *Ain Shams Engineering Journal* 9: 2691-2700.
- Troyo, A., D. O. Fuller, O. Calderón-Arguedas, and J. C. Beier. 2008. A geographical sampling method for surveys of mosquito larvae in an urban area using high-resolution satellite imagery. *Journal of vector ecology : journal of the Society for Vector Ecology* 33: 1-7.
- Vezzani, D., and A. P. Albiocco. 2009. The effect of shade on the container index and pupal productivity of the mosquitoes *Aedes aegypti* and *Culex pipiens* breeding in artificial containers. *Medical and Veterinary Entomology* 23: 78-84.
- Vezzani, D., A. Rubio, S. M. Velázquez, N. Schweigmann, and T. Wiegand. 2005. Detailed assessment of microhabitat suitability for *Aedes aegypti* (Diptera: Culicidae) in Buenos Aires, Argentina. *Acta Tropica* 95: 123-131.
- Villena, O. C., I. Terry, K. Iwata, E. R. Landa, S. L. LaDeau, and P. T. Leisnham. 2017. Effects of tire leachate on the invasive mosquito *Aedes albopictus* and the native congener *Aedes triseriatus*. *PeerJ* 5: e3756-e3756.
- Yee, D. A. 2008. Tires as habitats for mosquitoes: A review of studies within the Eastern United States. *Journal of Medical Entomology* 45: 581-593.
- Yee, D. A., B. Kesavaraju, and S. A. Juliano. 2004. Larval feeding behavior of three co-occurring species of container mosquitoes. *Journal of vector ecology : journal of the Society for Vector Ecology* 29: 315-322.
- Yee, D. A., D. Allgood, J. M. Kneitel, and K. A. Kuehn. 2012. Constitutive differences between natural and artificial container mosquito habitats: Vector communities, resources, microorganisms, and habitat parameters. *Journal of Medical Entomology* 49: 482-491, 410.
- Zhu, G., W. Ju, J. M. Chen, and Y. Liu. 2014. A Novel Moisture Adjusted Vegetation Index (MAVI) to reduce background reflectance and topographical effects on LAI retrieval. *PLoS ONE* 9: e102560.

Figures

Figure 1. Map of the city of New Orleans showing different land use-land cover and random sampling points generated for surveillance.

Figure 2. Standard Deviational Ellipse (SDE) for dumping frequency and number of tires.

Figure 3. Spatial distribution of colonized mosquito populations in the city of New Orleans.

Figure 4. Standard Deviational Ellipse (SDE) of Container Index of colonized mosquito populations in the city of New Orleans.

Table 1. Proposed macro- and microhabitat variables used in regression models.

Habitat Type	Variable name	Data Source	Unit
Macrohabitat	Residential and urban settings	USGS ¹	Integer values
	Industrial and commercial services	USGS ¹	Integer values
	Other urban and build-up land	NLCD ²	Integer values
	Developed high intensity	NLCD ²	Integer values
	Developed medium intensity	NLCD ²	Integer values
	Developed low intensity	NLCD ²	Integer values
	Developed open space	NLCD ²	Integer values
	Open water areas	NLCD ²	Integer values
	Non-forested wetland	USGS ¹	Integer values
	Forest wetland	USGS ¹	Integer values
	Streams, canals, lakes, reservoirs and estuaries	USGS ¹	Integer values
	Human density	NOLA ³	No. household/block
	Housing density	NOLA ³	No. houses/area
Microhabitat	Container index of other mosquito species	Surveillance	Integer values
	pH value	Surveillance	Integer values
	Predators (midges, water fleas, midges/water fleas)	Surveillance	Categorical values
	Size of tires (small, large, mixed)	Surveillance	Categorical values
	Tire status (damaged, intact)	Surveillance	Categorical values
	Water quality (muddy, turbid, clear)	Surveillance	Categorical values
	Water temperature	Surveillance	Integer values

¹ USGS available at <http://water.usgs.gov/GIS/dsdl/ds240/> (accessed 3 March 2019), ² Multi-Resolution Land Characteristics Consortium (U.S.). "National Land Cover Dataset (NLCD)" [Research Triangle Park, NC]:[Multi-Resolution Land Characteristics Consortium], ³ data.nola.gov (accessed 7 March 2019).

Table 2. Container Index (CI), number of larvae and pupae sampled from discarded tires in the City of New Orleans, LA

Mosquito Species	CI	No of larvae	No. of pupae
<i>Aedes albopictus</i>	44.19	1370	1340
<i>Culex quinquefasciatus</i>	22.18	850	630
<i>Cx. nigripalpus</i>	18.49	252	504
<i>Ae. aegypti</i>	17.65	241	301
<i>Cx. salinarius</i>	9.63	131	229
<i>Cx. restuans</i>	1.75	24	48
<i>Anopheles quadrimaculatus</i>	0.67	10	19
<i>Cx. territans</i>	0.59	9	15
<i>Ae. japonicus</i> *	0.1	5	9
<i>Anopheles crucians</i>	0.08	2	5
<i>Orthopodomyia signifera</i>	0.07	2	4
<i>An. punctipennis</i>	0.05	2	3
<i>Toxorhynchites rutilius</i>	0.02	2	3

* New locality record

Table 3. The response of frequency and number of increased dumping tires to landscape and demographic variables.

Model Type	Influential Factor	β	Wald χ^2
------------	--------------------	---------	---------------

Frequency of dumping tires	Industrial areas	1.72E-7	6.32**
	Average housing density	-0.01	6.45**
Number of dumping tires	Other urban built-up areas	-7.4E-6	22.15*
	Average housing density	-0.06	7.47*
	Industrial areas	6.66E-7	5.11**
	Residential and urban settings	-9.42E-7	5.01**
	Non-forested areas	-2.62E-7	3.91**

* $p < 0.01$, ** $p < 0.05$

Table 4. The response of colonized *Ae. albopictus* population to micro- and macrohabitat variables.

Model Type	Influential Factor	β	r^2	AICc
Microhabitat	Water quality (muddy)	5.5**	23.23	1647
	CI of other mosquito species	0.24**	21.53	1649
	Tire status (used)	4.84**	19.48	1651
	pH (6)	22.61**	16.48	1654
	CI of <i>Cx. quinquefasciatus</i>	0.35*	11.52	1663
Macrohabitat	Developed low intensity	-1.75E-6**	6.64	1673
	Developed medium intensity	7.16E-6*	5.23	1674
Combined model [‡]	Water quality (muddy)	6.24*	25.4	1640
	Developed medium intensity	8.19E-06*	22.44	1645
	pH (6)	22.91*	16.98	1654
	CI of <i>Cx. quinquefasciatus</i>	0.35*	11.52	1663

[‡] Best model, * $p < 0.01$, ** $p < 0.05$

Table 5. The response of colonized *Cx. quinquefasciatus* population to micro- and macrohabitat variables.

Model Type	Influential Factor	β	r^2	AICc
Microhabitat	Water quality (muddy and clear)	6.46**	18.67	1567
	Tire status (intact)	5.42**	17.01	1568
	Tire size (large)	6.68**	14.21	1572
	CI of <i>Ae. albopictus</i>	0.25*	11.52	1575
Macrohabitat	Developed medium intensity	1.69**	6.00	1673
Combined model [†]	Water quality (muddy and clear)	4.42**	18.67	1566
	Tire status (intact)	5.14**	17.01	1568
	Tire size (large)	7.35**	14.21	1572
	CI of <i>Ae. albopictus</i>	0.26*	11.52	1575

[†] Best model, * $p < 0.01$, ** $p < 0.05$