1 Exploring spatiotemporal changes in ecosystem 2 service values and hotspots in Southwest of China, 3 Mekong Region

4 Zhuoya Zhang^{1,2}, Junjie Yang², Xin Yang^{3,*}, Fuming Xie⁴

5 ¹ Faculty of Environmental Science and Engineering, Kunming University of Science
6 and Technology, Yunnan, China

7 ² Faculty of Geography and Ecotourism, Southwest Forestry University, Yunnan,
8 China

9 ³ Communist Youth League Committee, Southwest Forestry University, Yunnan,
10 China

11 ⁴ Institute of International River and Eco-security, Yunnan University, Yunnan, China

12 * Corresponding author:

13 Xin Yang, Communist Youth League Committee, Southwest Forestry University, No.
14 300, Bailong Road, Kunming, Yunnan, People's Republic of China.

15 Email: yangxin@swfu.edu.cn

16

17 **Abstract:** Xishuangbanna in the southwestern border of China is located in the upper
18 reaches of the Mekong River between the Indo-China Peninsula and the East Asian
19 continent. It is the largest tropical rain forest and monsoon forest in China. They have
20 an irreplaceable ecological service function and are an important barrier to
21 maintaining the ecological security of the Mekong River transboundary basin of the
22 Lancang River. Based on three sets of remote sensing data (1996, 2003, 2010 and
23 2016), this study has made an exploration on spatio-temporal changes of

24 Xishuangbanna, and also made an assessment on different ecosystem values of six
25 types of ecosystem in Xishuangbanna based on related theories of ecological
26 economics. Results showed that the values of Xishuangbanna ecosystem services in
27 1996, 2003, 2010, and 2016 were 70.4014billion yuan, 70.2115 billion yuan, 68.5129
28 billion yuan and 63.6098 billion yuan, respectively. The total value shows a
29 continuous decreasing trend, reflecting the continuous decline in the ability of the
30 Xishuangbanna ecosystem to provide services to humans.If divided by ecosystem
31 types, forest and rubber were two types that have the greatest proportion of service
32 values. In the perspective of service types, soil formation and protection account for
33 the largest proportion, followed by gas regulation and biodiversity
34 protection. Studying Xishuangbanna's ecosystem service values is helpful to explore
35 the sustainable development of resources and economy.

36 **Keywords:** Ecosystem service values, hotspots, spatiotemporal changes, Southwest
37 of China, Mekong Region

38 **Highlight:**

39 This paper used coefficient of sensitivity to determine the dependence of ESV on the
40 change of ecosystem value coefficient over time.

41 The total amount of ESV shows a continuous downward trend, indicating that the
42 ecological environment of Xishuangbanna is still severe.

43 The temporal-spatial change of ESV was analyzed for two decades.

44

45 **Introduction**

46 Ecosystem services (ES) refer to the various benefits that humans receive from
47 ecosystems, including supply services, regulation services, support services and
48 cultural services (Alcamo 2003, MEA 2005) . The intensification of human activities

49 has greatly accelerated the climate, environment and ecosystems of the planet. Urban
50 ecosystems are increasingly threatened by urban population growth, urban land use
51 expansion(Storkey, Döring et al. 2015, Leverkus and Castro 2017, Yang, Guan et al.
52 2018, Kuriqi, Pinheiro et al. 2019), and socio-economic activities(Mensah, Veldtman
53 et al. 2017, Song and Deng 2017), which affects the value of ecosystem services and
54 ultimately the sustainable development of human society(Costanza, d'Arge et al. 1997,
55 Sun, Liu et al. 2016) . Accurate assessment of ecosystem services value (ESV) is
56 essential for urban construction planning and the improvement and restoration of
57 urban ecosystems(Cui, Xiao et al. 2017), so it has received increasing attention from
58 the research community(Costanza, Chichakly et al. 2014, Costanza, De Groot et al.
59 2014, Leverkus and Castro 2017, Ossola and Hopton 2017).

60 As Costanza et al proposed the value of ecosystem services, especially after the
61 Millennium Ecosystem Assessment, the impact of human activities on ecosystems has
62 become an important research direction(Costanza, d'Arge et al. 1997, Robert,
63 Costanza et al. 2005). Many scholars have emphasized that human activities are an
64 important driving force for changing ecosystem services(Danz, Niemi et al. 2007,
65 Robards, Schoon et al. 2011). The true economic value of ecosystem services depends
66 on the interaction between the supply of ecosystems and the needs of society(Braat
67 and de Groot 2012) (Robert, Costanza et al. 2005). Monetary valuation of the
68 importance of ecosystem services to society can serve as a powerful and essential
69 communication tool to provide a basis for better and more balanced decisions(De
70 Groot, Brander et al. 2012). Scholars have found that basic income transfers can be a
71 convenient way to determine ESVs globally and nationally, assuming a constant unit
72 value per hectare for a given ecosystem type multiplied by the area of each type to
73 arrive at a total (Costanza, De Groot et al. 2014).

The value of ecosystem services not only reflects the functions of ecosystem services, but also reflects the significance of human ecological environment and the demand for ecosystem services. Similarly, many scholars have analyzed the spatio-temporal changes of ESV, and their responses to land-use changes(Fengqin, Yan et al. 2016, Kindu, Schneider et al. 2016, Fei, Shuwen et al. 2018) or other human activities(Camacho-Valdez, Ruiz-Luna et al. 2014). Although it is important to have a better understanding of the temporal changes of ESVs, there is an increasing focus on determining the spatial changes of ESVs by identifying "hot spots" on ecosystem services(Li, Fang et al. 2016). These spatial studies can provide a range of useful tools that can effectively integrate ecosystem services into planned or current conservation plans(Naidoo, Balmford et al. 2008)], assess the effectiveness of implementing ecological policies, and prioritize the management of ecosystem services field(Egoh, Reyers et al. 2011, Li, Fang et al. 2016). This information is particularly important for modifying current ecological protection plans and policies in a more beneficial and targeted manner. For example, by linking the spatial changes in wetland area with the provision of ecosystem services and economic value, some scholars have analyzed the spatiotemporal changes in the service value of coastal landscapes in Southern Sinaloa (Mexico) (Camacho-Valdez, Ruiz-Luna et al. 2014). Bottalico [28] evaluated the spatial distribution of Molise ESV by developing a spatial explicit method(Bottalico, Pesola et al. 2016). Li et al. verified the spatiotemporal changes of ESV and its hot and cold issues in China(Li, Fang et al. 2016).However, ESVs are less well-characterized by hotspots that change over a specific spatial range, especially globally and nationally.

With the change of policies in Xishuangbanna, economic development and the advancement of urbanization, the transformation and occupation of land resources has

intensified, resulting in a periodical sharp change in land use types and areas. This change has a great impact on ESV in Xishuangbanna. In this paper, we investigated the spatiotemporal changes of ESV and identified hotspots and hotspots of ESV changes. The analysis was based on Xishuangbanna four-phase remote sensing image data from 1996 to 2016. In this context, studying the impact of ecosystem services in Xishuangbanna from the perspective of land use change in Xishuangbanna can clarify the state of the ecosystem under the "stress-state-response" in Xishuangbanna. This is of great significance for adjusting and optimizing the land use pattern in Xishuangbanna, promoting the coordination of sustainable development in Xishuangbanna, and protecting the stability of cross-border ecological security.

1. Methods

1.1 Study area

Xishuangbanna is located at 21 ° 10'-22 ° 40 ' north latitude and 99 ° 55'-101 ° 50 east longitude. It is on the tropical edge south of the Tropic of Cancer. The land area is 19,124.50 square kilometers. It is bordered by Puer City in the northeast and northwest, Laos in the southeast, and Myanmar in the southwest. The border line is 966.3 kilometers long. The highest altitude is 2429 meters and the lowest altitude is 477 meters. The whole state has jurisdiction over one city and two counties, Jinghong City, Menghai County, and Mengla County. The climate of Xishuangbanna is warm and moist all year round. There are no four seasons except the dry and wet seasons. The dry season runs from November to April of the year and the wet season runs from May to October.

1.2 Data sources and processing

The social and economic data of Xishuangbanna include the publicly released relevant data such as the website of the National Bureau of Statistics of the People's

Republic of China and the statistical yearbook of Xishuangbanna. The remote sensing image data include: availability images using vegetation-free cloud-free datasets, which consist of four Landsat TM / TM + / OLI remote sensing images from March 1996 to April 1996, 2003, 2010, and April 2016, respectively.

Based on field surveys, combined with remote sensing data spectral information, we referenced the second-level survey data of forest resources in 2006 and 2016 to construct a remote sensing interpretation mark for land use in Xishuangbanna. Support Vector Machine (SVM) supervised classification method was used for The land use type data were interpreted and the land use types in the fourth phase of the study area were obtained. After the four-phase image classification, the total accuracy Kappa coefficients are 85.9%, 86.7%, 89.9%, and 93.5%, which meet the research needs.

1.3 Calculation of ESV in Xishuangbanna

1.3.1 Ecosystem classification

According to China's latest classification standard for land use status (GB / T21010-2007), and combined with the actual local conditions in Xishuangbanna, the ecosystem in Xishuangbanna is divided into forest ecosystem, rubber ecosystem, tea garden ecosystem, farmland ecosystem, watershed ecosystem, and construction land ecosystem.

1.3.2 Evaluation method of ESV in Xishuangbanna

(1) Modification of Xishuangbanna'ESV Equivalent Factor

In this study, the tea garden ecosystem was taken as the average value of woodland and grassland(Xiao-qing, Ze-xian et al.). The forest ecosystem equivalent factor in Xishuangbanna was revised to 1.96 times the national average(Xiaosai, Yongming et al. 2015), and the rubber ecosystem equivalent factor was revised to

1.60 times the national average(Yuanfan, Qingzhong et al. 2010). Based on the above amendments, the Xishuangbanna ESV Equivalent Scale was obtained.

(2) Functional value of food production per unit area of cultivated land ecosystem in Xishuangbanna

In order to eliminate the influence of crop price fluctuations on the total value in each year and region, the basic data were selected from the sown area, total output and average price of three main crops (rice, upland rice, and corn) in 1997 in Xishuangbanna. According to Xie Gaodi's research, the economic value of a standard unit ecosystem service value equivalent factor is equal to 1/7 of the average market value of a single grain(Gaodi, Chunxia et al. 2003, Xiaosai, Yongming et al. 2015). The calculation is as follows:

$$E_n = \frac{1}{7} \sum_{i=1}^n \frac{m_i q_i p_i}{M} (n = 1, 2, 3)$$

Among them, E_n is the economic value (yuan / hm^2) of providing food production and service functions for a unit of cultivated land ecosystem. i is the type of crop. p_i is the price of i crops (yuan / kg). q_i is the yield of i food crops (kg / hm^2). m_i is the area of i food crops (hm^2). M is the total area of n food crops (hm^2).

(3) Calculation method of Xishuangbanna 'ESV

We used the ESV formula of Costanza et al. to calculate the ESV of Xishuangbanna (Robert, Costanza et al.). In order to facilitate the analysis of the changes in the value of ecosystem services in Xishuangbanna and to compare the value of ecosystem services in different regions, this study adds two indicators to the calculation method of ecosystem service values, the Ecosystem Services Contribution

Rate (ESCR) and Ecosystem Services Value per unit (UESV) (Tianhai, field et al. 2018). The value of the ecosystem per unit area is to reduce the difference in the size of different ecosystems and to facilitate comparison. The calculation is as follows:

$$ESV = \sum V_i \times A_i$$

$$ESV_k = \sum V_{ki} \times A_{ki}$$

$$ESCR = \frac{\Delta ESV_k}{ESV}$$

$$UESV = \frac{ESV}{A}$$

Among them, ESV is the total value of ecosystem services in the study area. V_i is the ecosystem service value coefficient of ecosystem type i per unit area. A_i is the area of ecosystem type in the study area. ESV_k is the value of the k -th service of the ecosystem. V_{ki} is the k -th ecosystem service value coefficient of the i -th ecosystem type. $ESCR$ is the change in the value of a single ecosystem service ΔESV_k to the total value of ecosystem services. $UESV$ is the ratio of total ecosystem service value ESV to total area A .

(4) Calculation method of Coefficients of Sensitivity(CS)

This study applied CS to measure the representativeness of individual ecosystem services to each ecosystem type and the accuracy of the value of ecosystem services per unit area (Yao, Rusong et al. 2012). CS is the change in the ecosystem service value coefficient V_i of the ecosystem type i per unit area caused by a 1% change in the ESV of the total ecosystem service value in the study area. $CS > 1$, which indicates that ESV is elastic to V_i and the accuracy of V_i is low. $CS < 1$, it indicates that ESV is inelastic to V_i , which indicates that V_i is more accurate and ESV estimation is

accurate. The value of the ecosystem service value of each ecosystem type is increased or decreased by 50%, and the change of the elasticity coefficient of the total value of ecosystem services is analyzed (Xiao-qing, Ze-xian et al. , Yao, Rusong et al. 2012). The calculation is as follows:

$$CS = \left| \frac{(ESV_j - ESV_i)/ESV_i}{(VC_{jk} - VC_{ik})/VC_{ik}} \right|$$

Among them, ESV is the total value of ecosystem services in the study area. VC is the coefficient of ecosystem service value per unit area. A_i is the area of ecosystem type in the study area. k is the k th service of the ecosystem. i and j are the adjusted values of initial value and ecological value coefficient.

2 Results

2.1 Changes in total values of ecosystem services in Xishuangbanna from 1996 to 2016

2.1.1 ESV of different ecosystem types in Xishuangbanna from 1996 to 2016

According to calculations, the values of different ecosystem services from 1996 to 2016 are shown in Table 1.

Table 1 Ecosystem service values of each ecosystem type in Xishuangbanna from 1996 to 2016

	1996		2003		2010		2016	
	Value		Value		Value		Value	
Ecosystem types	(100	%	(100	%	(100	%	(100	%
	million		million		million		million	
	yuan)		yuan)		yuan)		yuan)	
Forest Ecosystem	583.2746	82.23	561.6264	79.99	460.6084	67.23	427.0692	66.77

Tea garden Ecosystem	4.2859	0.60	10.4002	1.48	8.584	1.25	22.522	3.52
Cultivated Land Ecosystem	6.1635	0.87	4.5839	0.65	5.6547	0.83	9.6756	1.51
Watershed Ecosystem	3.1857	0.45	2.8703	0.41	2.5494	0.37	3.7308	0.58
Rubber Ecosystem	114.3118	16.12	124.7308	17.77	209.0921	30.52	178.7769	27.95
Construction Ecosystem	-1.8814	-0.27	-2.0961	-0.30	-1.3601	-0.20	-2.1648	-0.34
Total	709.3401	100.00%	702.1154	100.00%	685.1285	100.00%	639.6097	100.00%

213

214 Among the ESV of different ecosystem types in Xishuangbanna, the ecological
215 value coefficients of forest ecosystems and rubber ecosystems are high. Changes in
216 the area of such ecosystems will greatly affect the changes in ecosystem service
217 values in Xishuangbanna.

218 From Table 1, it can be concluded that the ESV of Xishuangbanna in 1996, 2003,
219 2010, and 2016 showed a continuous decrease, which reflects that the ability of
220 Xishuangbanna ecosystem to provide services to human beings has been continuously
221 decreasing. The values of Xishuangbanna ecosystem services in 1996, 2003, 2010,
222 and 2016 were 70.4014billion yuan, 70.2115 billion yuan, 68.5129 billion yuan and
223 63.6098 billion yuan, respectively. The total value shows a continuous decreasing
224 trend, reflecting the continuous decline in the ability of the Xishuangbanna ecosystem
225 to provide services to humans.

226 Among them, in terms of the proportion of ecosystem service value, the forest
227 land ecosystem has the highest proportion, which is the most important component of
228 the ESV in Xishuangbanna (66.77% -82.23%), followed by rubber ecosystem (16.12%
229 -30.52%). Xishuangbanna is an area with high forest coverage, and the ecological
230 environment is generally well preserved. However, with the development of human

society and economy, the ecosystem services in Xishuangbanna show a continuous decline. In the case where the forest land and rubber ecosystem in Xishuangbanna account for a large proportion, the ESV of the forest land ecosystem and rubber ecosystem accounts for 94.71% -98.35% of the total value. It can also be seen that woodland and rubber ecosystems dominate the ecosystem services in Xishuangbanna.

The construction land ecosystem is the lowest and has a negative value, and the service value of the construction land ecosystem continues to show negative growth, from -6.6847 billion yuan in 1996 to -7.6916 billion yuan, which is also in line with the reality of the increasing area of construction land.

According to research, the ESV of rubber in Xishuangbanna is far less than that of tropical rain forests and secondary vegetation (67.149 million yuan / (hm².a)) , the proportion of which is about 56.74%. The ecological service value of evergreen coniferous forest (13,315 yuan / (hm².a)) is large(Gaodi, Chunxia et al. 2003). The service value of the rubber forest ecosystem in Xishuangbanna is far greater than the average value of tropical forests in China (16.056 million yuan / (hm².a)) (Yuanzhao 2003). The ecological and economic value of rubber forests is only about half of the value of tropical rain forests and secondary vegetation. Cutting down tropical rain forests and planting rubber forests is not desirable from the total regional ecological and economic value, which is not conducive to regional natural asset appreciation and local sustainable development(Tiyuan, Jiayong et al. 2009).

The direct economic output of rubber forest is much larger than that of tropical rain forest. This is the source of huge economic pressure on tropical rain forest protection. It is recommended to adopt environmental and economic policies such as ecological compensation to comprehensively protect tropical rain forest(Tiyuan, Jiayong et al. 2009).

2.1.2 Changes in ESV of different ecosystem types in Xishuangbanna from 1996 to 2016

The total reduction of ESV from 2010 to 2016 (16.17153 billion yuan) was 630.19% of the total reduction of ESV from 1996 to 2003, which is the time period when ESV changes the most. During 2010-2016, the rapid development of urbanization, the continuous decline in rubber prices, the rise in the prices of tea and crops, and the continuous advancement of the country's policy of “returning rubber to forests” have made the value of the Xishuangbanna ecosystem the most significant. Total reduction in ESV from 2003 to 2010 (6.03301 billion yuan). The period from 1996 to 2003 was the smallest total reduction in ESV (2.556614 billion yuan).

The ESV of forest ecosystems continues to decline. The change of forest ecosystem ESV is the most dramatic change among all ecosystem types, from 82.23% of the total ESV in 1996 to 66.77% of the total ESV in 2016. The period when the ESV of forestland ecosystem changed the most was from 2003 to 2010. During this period, especially from 2003 to 2007, the continuous increase of rubber prices in the world has caused a large number of forest land ecosystems to be replaced by rubber ecosystems. The rubber ecosystem ESV has increased sharply and the forest land ecosystem ESV has decreased sharply. The total reduction in ESV of forest land ecosystem from 2003 to 2010 (35.87842 billion yuan) is 64.67% of the total reduction from 1996 to 2016 (55.49324 billion yuan).

The ESV of the tea garden ecosystem decreased first, then increased sharply, and increased overall. From 2003 to 2010, the ESV of the tea garden ecosystem decreased. During this period, the tea price in Xishuangbanna was low, and rubber was the main source of the local economy. From 2010 to 2016, the tea market in Xishuangbanna was opened. The prices of tea, especially those of several famous tea mountains, rose

sharply, and the ESV (4.995 billion yuan) of the tea garden ecosystem increased sharply.

The ESV of cultivated land ecosystem showed a trend of first decline, then slow increase, and overall growth. The ESV of cultivated land ecosystem decreased by 56106 million yuan from 1996 to 2003, and continued to increase from 2003 to 2010 and from 2010 to 2016. Among them, the biggest change was from 2010 to 2016. The total increase in ESV of forest land ecosystem (1.42819 billion yuan) from 2010 to 2016 was 114.49% of the total increase of ESV of forest land ecosystem (1.224746 billion yuan) from 1996 to 2016.

The ESV of water ecosystems are decreasing first, then increasing, and generally increasing slowly.

2.2 Changes of total values in single ecosystem service in Xishuangbanna from 1996 to 2016

2.2.1 ESV in the individual ecosystem services in Xishuangbanna from 1996 to 2016

It can be seen from Table 2 that among the various ESVs in Xishuangbanna, soil formation and protection account for the largest proportion (17.86% -17.93%), followed by air regulation (accounting for 15.80% -15.92%) and biodiversity protection (accounting for 14.86% -14.91%), water conservation (accounting for 14.48% -14.57%), climate regulation (accounting for 12.34% -12.35%), raw material production (accounting for 11.61% -11.78%), and food production accounting for the smallest proportion (accounting for Than 0.56% -0.70%).

Compared with urban areas with low forest coverage, this ranking result is quite different. For example, in the study of urban ecosystem service value, hydrological regulation accounts for the largest proportion in various ESVs(Hui, Wenwu et al.

2017, Tianhai, field et al. 2018). The second largest proportion is waste treatment and biodiversity maintenance (Hui, Wenwu et al. 2017, Tianhai, field et al. 2018).

308

309 Table 2 Evaluation of ecosystem service of different types in Xishuangbanna from
310 1996 to 2016

	1996		2003		2010		2016	
Individual ecosystem services	Value (100 million yuan)	%	Value (100 million yuan)	%	Value (100 million yuan)	%	Value (100 million yuan)	%
Food production	4.1512	0.59	3.9549	0.56	4.0058	0.59	4.4911	0.70
Raw material	83.4826	11.77	82.6823	11.78	80.5464	11.76	74.2826	11.61
Air regulation	112.8175	15.90	111.8080	15.92	108.9442	15.90	101.0697	15.80
Climate regulation	87.5521	12.34	86.7146	12.35	84.5643	12.34	78.9300	12.34
Water conservation	103.2227	14.55	101.9738	14.52	99.8082	14.57	92.6174	14.48
Waste treatment	44.4557	6.27	43.7798	6.24	42.9431	6.27	41.5826	6.50
Soil formation and protection	126.6771	17.86	125.5667	17.88	122.4488	17.87	114.7029	17.93
Biodiversity conservation	105.6038	14.89	104.6699	14.91	101.9862	14.89	95.0425	14.86
Recreation and culture	41.3773	5.83	40.9656	5.83	39.8816	5.82	36.8908	5.77
Total	709.3401	100.00	702.1154	100.00	685.1285	100.00	639.6097	100.00

311

312 The essence of ESV change is the area change of different ecosystems and the
313 difference of equivalent factors in different ecosystems. Therefore, the trend of the

value of individual ecosystem services in Xishuangbanna from 1996 to 2016 is consistent (Figure 1)

Figure 1 Structure of ecosystem service value in Xsihuangbanna, 1996-2016.

2.2.2 ESV changes in the individual ecosystem services in Xishuangbanna from 1996 to 2016

According to Table 3, from the contribution of each component to the reduction of the total ESV, the contribution of ESV change in water conservation from 1996 to 2003 was the largest (17.29%). The contribution of ESV change to soil formation and protection from 2003 to 2010 was the largest (18.35%). The largest contribution of gas-regulated ESV changes during the period 2010-2016 (17.30%). The largest contribution of changes in soil formation and protection of ESV during the period 1996-2016 (17.17%).

Table 3 Ecosystem services values change and its contribution rate

Fisrt classification	Second classification	Value billion/（亿元）				Contribution rate of value change/%			
		1996	2003	2010	2016	1996-2003	2003-2010	2010-2016	1996-2016

Supply services	Food production	4.1512	3.9549	4.0058	4.4911	2.72	-0.30	-1.07	-0.49
Supply services	Raw material	83.4826	82.6823	80.5464	74.2826	11.08	12.57	13.76	13.19
Regulation services	Air regulation	112.8175	111.8080	108.9442	101.0697	13.97	16.86	17.30	16.85
Regulation services	Climate regulation	87.5521	86.7146	84.5643	78.9300	11.59	12.66	12.38	12.36
Regulation services	Water conservation	103.2227	101.9738	99.8082	92.6174	17.29	12.75	15.80	15.21
Regulation services	Waste treatment	44.4557	43.7798	42.9431	41.5826	9.36	4.93	2.99	4.12
Support services	Soil formation and protection	126.6771	125.5667	122.4488	114.7029	15.37	18.36	17.02	17.17
Support services	Biodiversity conservation	105.6038	104.6699	101.9862	95.0425	12.93	15.80	15.25	15.15
Cultural services	Recreation and culture	41.3773	40.9656	39.8816	36.8908	5.70	6.38	6.57	6.43
Total		709.3401	702.1154	685.1285	639.6097	100.00	100.00	100.00	100.00

329

330 2.2.3 Changes in the value of individual ecosystem services in different
331 administrative regions of Xishuangbanna from 1996 to 2016

332 An analysis of the ecosystem service value of the three counties (Figure 2) shows
333 that Xishuangbanna has higher ESV values in Erhai County and Mengla County.
334 Xishuangbanna's urban construction is dominated by Jinghong City (the seat of the
335 state government), and tourism in Jinghong City is developing rapidly. Erhai County
336 is not suitable for planting rubber, and the proportion of the tea garden ecosystem is

relatively large. It is the main tea production place in Xishuangbanna.

Figure 2 Changes in the value of individual ecosystem services in different administrative regions of Xishuangbanna from 1996 to 2016

2.3 Spatiotemporal changes of ESV in Xishuangbanna

As can be seen from the figure 3, there are obvious spatial differences and temporal changes in ESV in Xishuangbanna. In 1996, land with an ESV in Xishuangbanna of more than 3,000 yuan / ha accounted for more than 90% of the total area. The ESV value was the highest in the year. ESV dropped in 2003, mainly at 1000-2000 yuan / ha. In 2010, ESV rose, but the area was larger than 3,000 yuan / ha in 1996, and the change was mainly concentrated in Erhai County. The area of

ESV 1000-2000 yuan / ha in Erhai County increased. In 2016, ESV became more complicated and fragmented. High ESV areas were concentrated in Jinghong City and Mengla County, while Erhai County had fewer high ESV areas.

Figure3. Spatiotemporal changes of ESV in Xishuangbanna from 1996 to 2016

2.4 Ecosystem services sensitivity analyses

By calculation, the results of CS of all ecosystem types except forest in 1996, 2003, 2010 and 2016 were all lower than 1. The equivalence factors used in the study appear credible and accurate. This shows that the elasticity of the value coefficient of ESV is good, and the estimation results of this study are credible.

2.5 Cold and hot spot distribution of ESV in Xishuangbanna

Based on the Getis-Ord Gi * tool in the spatial analysis module, cold and hot

spot identification was performed for the four ESVs of Xishuangbanna in 1996, 2003, 2010, and 2016, and the spatial distribution of the high and low ESV agglomerations in the fourth phase of the study area was obtained. Compared with 1996, the area of high-value ESV accumulation in Xishuangbanna in 2003, 2010 and 2016 has been significantly reduced. The hot spots of ESV in 1996 were mainly distributed in Jinghong City and Mengla County. Cold spots are distributed in Erhai County. Since 2003, 99% of cold spots have increased and hot spots have disappeared.

Figure4. Cold and hot spot distribution ofESV in Xishuangbanna from 1996 to 2016

3. Discussions and conclusions

(1) When evaluating regional ESV, most scholars directly used the Xie Gaudi and other Chinese terrestrial ecosystems per unit area ecological service value equivalent scale(Gaodi, Chunxia et al. 2003, Xiaosai, Yongming et al. 2015), and used national scale equivalent factors to evaluate ESV in counties and cities, without reflecting regional differences.

(2) During 1996-2016, the ESV in Xishuangbanna continued to decrease. In Xishuangbanna, due to the gradual decrease in the area of forest ecosystems, they have gradually been replaced by rubber, construction land, tea gardens, etc. in areas with low ecosystem service values. The overall value of ecosystem services in Xishuangbanna has continued to decline. As the forest land is the most important part of the ESV in Xishuangbanna, in the future development of Xishuangbanna, a corresponding ecological compensation system should be introduced to strengthen the protection of forest land, especially natural forests, and continue to promote the policy of “returning rubber to forests” to avoid excessive development and natural forest cover. The replacement of economic forests leads to ecological imbalances and ecological risks, and eventually triggers international ecological security issues.

(3) In this study, the ecological service value equivalent factor calculation method is used. Compared with other research results using the market value method, the calculation results are consistent, but the total value is smaller than the market value method (Zhaopeng and Youxin 2012) (Heli, Anyi et al. 2014).

(4) There are obvious spatial differences and temporal changes in ESV and hot and cold spots in Xishuangbanna. In 1996, land with an ESV in Xishuangbanna of more than 3,000 yuan / ha accounted for more than 90% of the total area. ESV dropped in 2003, mainly at 1000-2000 yuan / ha. In 2010, ESV rose, and changes were mainly concentrated in Erhai County. The area of ESV 1000-2000 yuan / ha in Erhai County increased. In 2016, ESV became more complicated and fragmented. High ESV areas were concentrated in Jinghong City and Mengla County, while Erhai County had fewer high ESV areas. Compared with 1996, the area of high-value ESV accumulation in Xishuangbanna in 2003, 2010 and 2016 has been significantly reduced. The hot spots of ESV in 1996 were mainly distributed in Jinghong City and

Mengla County. Cold spots are distributed in Erhai County. Since 2003, 99% of cold spots have increased and hot spots have disappeared.

(5) The ESV assessment in this study is only for the first-class classification system, and there is no further subdivision of paddy fields and dry fields in cultivated land ecosystems, and there are woodlands and shrubs in woodland ecosystems.

Implications for conservation

Acknowledgements

Our special thanks go to Prof Yang Yuming and Prof Ye Wen for providing constructive suggestions.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interests with respect to the research, authorship, and/or publication of this article.

Funding

This work was supported by the 2017 Yunnan Provincial Department of Education Science Research Foundation Project (Grant 2017ZZX210).

References

Alcamo, J. (2003). Ecosystems and human well-being: a framework for assessment,

Island Press.

Bottalico, F., L. Pesola, M. Vizzarri, L. Antonello, A. Barbati, G. Chirici, P. Corona, S.

Cullotta, V. Garfi and V. Giannico (2016). "Modeling the influence of alternative forest

management scenarios on wood production and carbon storage: A case study in the

- 430 Mediterranean region." Environmental research **144**: 72-87.
- 431 Braat, L. C. and R. de Groot (2012). "The ecosystem services agenda:bridging the
432 worlds of natural science and economics, conservation and development, and public
433 and private policy." Ecosystem Services **1**(1): 4-15.
- 434 Camacho-Valdez, V., A. Ruiz-Luna, A. Ghermandi, C. A. Berlanga-Robles and P. A. L. D.
435 Nunes (2014). "Effects of Land Use Changes on the Ecosystem Service Values of
436 Coastal Wetlands." Environmental Management **54**(4): 852-864.
- 437 Costanza, R., K. Chichakly, V. Dale, S. Farber, D. Finnigan, K. Grigg, S. Heckbert, I.
438 Kubiszewski, H. Lee and S. Liu (2014). "Simulation games that integrate research,
439 entertainment, and learning around ecosystem services." Ecosystem Services **10**:
440 195-201.
- 441 Costanza, R., R. d'Arge, R. De Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, S.
442 Naeem, R. V. O'Neill and J. Paruelo (1997). "The value of the world's ecosystem services
443 and natural capital." nature **387**(6630): 253-260.
- 444 Costanza, R., R. De Groot, P. Sutton, S. Van der Ploeg, S. J. Anderson, I. Kubiszewski, S.
445 Farber and R. K. Turner (2014). "Changes in the global value of ecosystem services."
446 Global environmental change **26**: 152-158.
- 447 Cui, Y., X. Xiao, Y. Zhang, J. Dong, Y. Qin, R. B. Doughty, G. Zhang, J. Wang, X. Wu and
448 Y. Qin (2017). "Temporal consistency between gross primary production and
449 solar-induced chlorophyll fluorescence in the ten most populous megacity areas over

450 years." *7*(1): 14963.

451 Danz, N. P., G. J. Niemi, R. R. Regal, T. Hollenhorst, L. B. Johnson, J. M. Hanowski, R. P.

452 Axler, J. J. H. Ciborowski, T. Hrabik and V. J. Brady (2007). "Integrated Measures of

453 Anthropogenic Stress in the U.S. Great Lakes Basin." *Environmental Management* **39**(5):

454 631-647.

455 De Groot, R., L. Brander, S. van der Ploeg, R. Costanza, F. Bernard, L. Braat, M. Christie,

456 N. Crossman, A. Ghermandi and L. Hein (2012). "Global estimates of the value of

457 ecosystems and their services in monetary units." *Ecosystem Services* **1**(1): 50-61.

458 Egoh, B. N., B. Reyers, M. Rouget and D. M. Richardson (2011). "Identifying priority

459 areas for ecosystem service management in South African grasslands." *Journal of*

460 *Environmental Management* **92**(6): 1642-1650.

461 Fei, L., Z. Shuwen, Y. Jiuchun, C. Liping, Y. Haijuan and B. Kun (2018). "Effects of land

462 use change on ecosystem services value in West Jilin since the reform and opening of

463 China." *Ecosystem Services* **31**: 12-20.

464 Fengqin, Yan, Shuwen, Zhang, Xingtuo, Liu, Dan, Chen, Jing and Chen (2016). "The

465 Effects of Spatiotemporal Changes in Land Degradation on Ecosystem Services Values

466 in Sanjiang Plain, China."

467 Gaodi, X., L. Chunxia, L. Yunfa, Z. Du and L. Shuangcheng (2003). "Ecological assets

468 valuation of the Tibetan Plateau." *Journal of Natural Resources* **18**(2): 189-196.

469 Heli, L., N. Anyi, Z. Jincai, L. Xiaowan and L. Xiaojing (2014). "Evaluation of ecosystem

470 services in Xishuangbanna based on land use change." Jiangsu Agricultural Science

471 **42**(5): 278-281.

472 Hui, W., Z. Wenwu, Z. Xiao and W. Xinzhi (2017). "Regional ecosystem service value

473 evaluation based on land use changes: A case study in Dezhou , Shandong Province ,

474 China." Acta Ecologica Sinica **37**(11): 3830-3839.

475 Kindu, M., T. Schneider, M. D?llerer, D. Teketay and T. Knoke (2016). "Scenario

476 modelling of land use/land cover changes in Munessa-Shashemene landscape of the

477 Ethiopian highlands." Science of the Total Environment **622-623**: 534-546.

478 Kuriqi, A., A. N. Pinheiro, A. Sordo-Ward and L. Garrote (2019). "Influence of

479 hydrologically based environmental flow methods on flow alteration and energy

480 production in a run-of-river hydropower plant." Journal of Cleaner Production **232**:

481 1028-1042.

482 Leverkus, A. B. and J. Castro (2017). "An ecosystem services approach to the ecological

483 effects of salvage logging: Valuation of seed dispersal." Ecological Applications.

484 Li, G., C. Fang and S. Wang (2016). "Exploring spatiotemporal changes in

485 ecosystem-service values and hotspots in China." Science of the Total Environment

486 **545**: 609-620.

487 MEA, M. E. A. (2005). "Ecosystems and human well-being: current state and trends."

488 Millennium Ecosystem Assessment, Global Assessment Reports.

489 Mensah, S., R. Veldtman, A. E. Assogbadjo, C. Ham, R. Glèlè Kaka? and T. Seifert (2017).

490 "Ecosystem service importance and use vary with socio-environmental factors: A study
491 from household-surveys in local communities of South Africa." Ecosystem Services **23**:
492 1-8.

493 Naidoo, R., A. Balmford, R. Costanza, B. Fisher, R. E. Green, B. Lehner, T. Malcolm and T.
494 H. Ricketts (2008). "Global mapping of ecosystem services and conservation priorities."
495 Proceedings of the National Academy of Sciences **105**(28): 9495-9500.

496 Ossola, A. and M. Hopton (2017). "Measuring tree loss dynamics across residential
497 landscapes." **612**: 940-949.

498 Robards, M. D., M. L. Schoon, C. L. Meek and N. L. Engle (2011). "The importance of
499 social drivers in the resilient provision of ecosystem services." Global Environmental
500 Change **21**(2): 0-529.

501 Robert, Costanza, and, Ralph, d'Arge, and, Rudolf, de, Groot and and "The value of the
502 world's ecosystem services and natural capital."

503 Robert, Costanza, and, Ralph, d'Arge, and, Rudolf, de, Groot and and (2005). "The
504 value of the world's ecosystem services and natural capital."

505 Song, W. and X. Deng (2017). "Land-use/land-cover change and ecosystem service
506 provision in China." Science of the Total Environment **576**: 705-719.

507 Storkey, J., T. Döring, J. Baddeley, R. Collins, S. Roderick, H. Jones and C. Watson (2015).
508 "Engineering a plant community to deliver multiple ecosystem services." Ecological
509 Applications **25**(4).

- 510 Sun, Z., Z. Liu, H. E. Chunyang, J. Wu and Technology (2016). "Multi-scale analysis of
511 ecosystem service trade-offs in urbanizing drylands of China:A case study in the
512 Hohhot-Baotou-Ordos-Yulin region." Acta Ecologica Sinica.
- 513 Tianhai, Z., field, X. Shu, T. Lina and G. Wei (2018). "The evolvement of land use
514 patterns in coastal cities and its influence on ecosystem service values." Acta Ecologica
515 Sinica **38**(21): 55-64.
- 516 Ti yuan, X., W. Jiayong, D. Changqun and D. Liangjun (2009). "Ecological economic
517 valuation of rubber plantation in Xishuangbanna." Journal of East China Normal
518 University (Natural Science)(2): 27-34.
- 519 Xiao-qing, Z., G. Ze-xian and G. Xiang-yu "LAND USE AND LAND-COVER CHANGE
520 AND IT' S IMPACT ON ECOSYSTEM SERVICES VALUES IN A REGION WITH
521 LARGE-AREA ARTIFICIAL GARDENS." Resources and Environment in the Yangtze Basin
522 **v.25**(01): 91-100.
- 523 Xiao-qing, Z., G. Ze-xian and G. Xiang-yu "LAND USE AND LAND-COVER CHANGE
524 AND IT' S IMPACT ON ECOSYSTEM SERVICES VALUES IN A REGION WITH
525 LARGE-AREA ARTIFICIAL GARDENS." Resources and Environment in the Yangtze Basin
526 **v.25**(1): 91-100.
- 527 Xiaosai, L., Z. Yongming, Z. Li, T. Jingjing and L. Jing (2015). "Ecosystem services value
528 change in Qinglong County from dynamically adjusted value coefficients." Chinese
529 Journal of Eco-Agriculture **23**(3): 373-381.

- 530 Yang, J., Y. Guan, J. C. Xia, C. Jin and X. Li (2018). "Spatiotemporal variation
- 531 characteristics of green space ecosystem service value at urban fringes: A case study
- 532 on Ganjingzi District in Dalian, China." Science of the Total Environment **639**:
- 533 1453-1461.
- 534 Yao, S., W. Rusong, W. K. House and Y. Wenrui (2012). "An analysis of the spatial and
- 535 temporal changes in Chinese terrestrial ecosystem service functions." Chinese Science
- 536 Bulletin(9): 42-53.
- 537 Yuanfan, Z., W. Qingzhong, T. Jing, H. Chaolang, S. Jingxin and Y. Dong (2010).
- 538 "Evaluation of the values for ecological service function of tropical natural forest in
- 539 Xishuangbanna." Forest Inventory and Planning **35**(1): 1-6.
- 540 Yuanzhao, H. (2003). "Distribution, Types and Characteristics of China's Tropical
- 541 Forest." World Forestry Research **16**(3): 47-51.
- 542 Zhaopeng, J. and M. Youxin (2012). "Dynamic evaluation on ecosystem service values
- 543 of Xishuangbanna, Yunnan, China." Journal of Central South University of Forestry
- 544 &Technology **9**(32): 87-93.
- 545