GROUPS OF SPECIES RESPONDING TO SOIL AND LIGHT IN THE CERRADO

Glaucia Soares Tolentino¹, Cristina Máguas^{2*}, Luiz Fernando Silva Magnago^{1,3}, Tillmann Buttschardt⁴,
João Augusto Alves Meira-Neto^{1,2*}

¹ Laboratory of Ecology and Evolution of Plants - LEEP, Universidade Federal de Viçosa, Campus UFV
s/n, Viçosa, State of Minas Gerais, Brazil, 36570-000.

² Centre for Ecology, Evolution and Environmental Changes – CE3C, Faculdade de Ciências da
Universidade de Lisboa, Edifício C2, Lisboa, Portugal, 1749-016.

³ Centro de Formação em Ciências e Tecnologias Agroflorestais, Universidade Federal do Sul da Bahia,
Rodovia BR-415, Parque Verde, Itabuna, Bahia State, Brazil, 45604811

⁴ Institute of Landscape Ecology - ILÖK, University of Münster, D-48149 Münster, Germany.

* cmhanson@fc.ul.pt, j.meira@ufv.br
JAAMN-ORCID 0000-0001-5953-3942

Abstract

Woody plants in the Cerrado have been understood to comprise two groups: calcicole species responding to soil Al-Ca gradient (CAL) and leguminous species responding to inorganic soil N gradient (LEG). However, the bulk of Cerrado woody flora was not inside these groups. Our main question was whether CAL and LEG are distinct from each other as well as from the bulk of Cerrado species that occur rather in savannas than in quasi-forests (SAV) while responding only to soil variables and light. We performed the study among five soil categories, and measured canopy openness. We evaluated the functional traits of the 34 most relevant species of the studied Cerrado. We measured specific leaf area, bark thickness, leaf nitrogen, leaf carbon, leaf C/N, maximum height, maximum, stem elongation, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$. We determined wood density by consulting a database. RLQ and Fourth-corner analyses confirmed CAL and LEG as consistent groups as well as found a third group, the SAV. CAL and LEG were found to dominate plots with low canopy openness; only CAL were negatively associated with aluminium saturation; only LEG were negatively associated with pH; only SAV were negatively associated with inorganic nitrogen and CEC. LEG presented the lowest leaf C/N and the highest leaf N meanwhile SAV presented the highest C/N, the lowest leaf N and the lowest stem elongation. Higher values of $\delta^{13}\text{C}$ in leaves suggest higher water use efficiency for CAL and LEG than for SAV.

Keywords: Aluminium, nitrogen, calcicoles, light, $\delta^{13}\text{C}$, water use efficiency.

Introduction

A biodiversity hotspot, the Cerrado is one among tropical savannas that occur on acidic soils as some Australian and African savannas (Matarechera et al. 1998; Hutley et al. 2000). In a biodiversity hotspot as the Cerrado, disentangle plant responses to environmental factors is time consuming because the high plant diversity, the soil complexity and a myriad of other environmental variables usually blur the plant-soil patterns and challenge researchers that seek a general interpretation for vegetation responding to soil gradients (de Assis et al. 2011; Neri et al. 2012; Meira-Neto et al. 2017). Grouping species into functional groups with unique ecological attributes (Díaz and Cabido, 2001; Lavorel and Garnier, 2002) frames plant diversity into a few environmentally responsive groups, facilitates the understanding of diverse communities and allows increase complexity step by step. Grouping species of the Cerrado has demanded a long and continuous effort. As far as we know, the first grouping separated the typical woody species that give phytogeographical unity to the Cerrado from species that occur in the Cerrado but occur mainly in other vegetation types, noticeably Atlantic and Amazon forests (Rizzini 1963). Concerning the soil influence on Cerrado plants, plant functional groups have been understood to comprise woody plants responding to the Al-Ca gradient (i.e., pH gradient) and to the inorganic nitrogen gradient (Ratter *et al.*, 1977; Durigan and Ratter, 2006; Meira-Neto *et al.* 2017). For that, isolating soil from other explanatory environmental variables is important in order to avoid blurring the plant-soil patterns.

Light variation may be included in such studies of tropical savannas as an explanatory variable since it is influenced by soil. Cerrado has high irradiance and typical Cerrado species demand high levels of light (Ribeiro and Walter, 1998). Thus, these species are expected to be displaced in shaded environments by shade-tolerant species (Pinheiro *et al.*, 2016) more adapted to the Cerrado quasi-forests physiognomies (cerradões) with more soil resources. Soil influences Cerrado physiognomies (e.g., (Goodland 1971; Oliveira-Filho and Ratter 2002) and the physiognomic variation promotes a light gradient with species responses (Dalmagro et al. 2014); therefore, canopy cover (i.e., light regime proxy) should be considered as a main ecological factor (Goodland 1971; Goodland and Pollard 1973; Felfili and Fagg 2007; Simon et al. 2009; Dantas et al. 2013) influencing plant growth and soil humidity in Cerrado and in vegetation in general (Valiente-Banuet and Verdú 2007; Meira-Neto et al. 2018). Fire opens Cerrado's physiognomies decreasing coverage of woody species (e.g., (Coutinho 1990; Hoffmann 2002; Simon et al. 2009), influencing light regime. However, in the absence of fire in the Cerrado, soil properties are expected to be the main generators of physiognomic variations as well as the main drivers of light regime.

The ubiquitous Al-Ca gradient is a soil pH gradient in tropical terrestrial ecosystems that explains much of the physiognomic variation in the Cerrado, from the most oligotrophic and acidic soils (i.e., highest Al^{3+} content) to the relatively fertile mesotrophic soils (Ratter *et al.*, 1977, 2003). The cation exchange capacity may be taken as a synthesis of soil fertility and is related to Al-Ca gradient is positively related to pH and has been used to explain physiognomic variation (Neri *et al.*, 2013). The levels of exchangeable aluminium are also related to the Al-Ca gradient (negatively related to pH) and were found as a main source of stress for plants in tropical savannas on acidic soils (Furley and Ratter 1988; Motta et al. 2002; Sugihara et al. 2014). As a result, open physiognomies are associated with dystrophic soils with toxic levels of Al^{3+} (Goodland and Pollard, 1973; Ruggiero *et al.*, 2002; Neri *et al.*, 2012), thus Al-Ca gradient could determine biomass and abundances in Cerrado (Silva *et al.*, 2013), especially calcicole species. Calcicoles are recognized as functional groups of plants worldwide (Ström 1997; Wohlgemuth and Gigon 2003) and Cerrado is a well-studied tropical savanna concerning calcicoles. The first functional group related to the Cerrado soil was the group of calcicole species (CAL), which respond readily to the Al-Ca gradient (Ratter et al. 1977). These species occur throughout all Cerrado physiognomies, but are dominant in cerradões on mesotrophic soils (mesotrophic cerradões) (Ratter *et al.* 1977; Durigan and Ratter, 2006) and are commonly found as dominant species in tropical dry forests on limestone hills (Oliveira-Filho *et al.*, 2006; Meira-Neto *et al.*, 2017).

Another functional group, the nitrogen-fixing leguminous trees (LEG) occur throughout Cerrado physiognomies, but dominate the cerradões on acidic dystrophic soils (dystrophic cerradões) (Meira-Neto *et al.*, 2017). Therefore, dystrophic cerradões are quasi-forest physiognomies with high abundance of trees, with much higher biomass than open savannas, also occur on dystrophic and acidic soils with high levels of Al^{3+} and with high levels of inorganic nitrogen dominated by leguminous nitrogen-fixers (Goodland 1971; Oliveira-Filho and Ratter 2002; Meira-Neto et al. 2017).

These two groups, CAL and LEG, were found to respond in opposite ways to the Al-Ca gradient, with CAL biomass and abundances positively associated with pH and negatively with inorganic nitrogen at the uppermost soil layers, and LEG biomass and abundances negatively associated with pH and positively with inorganic nitrogen at the uppermost soil layers (Meira-Neto et al. 2017). However, this CAL-LEG model only considers the gradient between mesotrophic soils with higher pH and dystrophic soils with higher contents of inorganic nitrogen. Additionally, open savanna physiognomies dominated by the bulk of Cerrado's woody species were still unknown in this CAL-LEG model.

Climate, altitude, topography, water table and fire are important rulers of the Cerrado functioning and physiognomic variation (Rossatto et al. 2012; Cianciaruso et al. 2012; Dantas et al. 2013; Franco et al. 2014). Without variation of climate, of altitude, of topography, of water table depth and of disturbances (mainly fire), the woody species of Cerrado respond essentially to soil and light. In such a situation, the growth of woody species should respond to light, to soil humidity, to N, to P, to K and to Mg soil concentrations as well as to the soil Al-Ca gradient (Goodland and Pollard 1973; de Assis et al. 2011; Neri et al. 2012). The CAL and the LEG are in different situations on the gradient of inorganic soil nitrogen as nitrogen is a key factor for plant growth and development (Sprent 2007, 2009), especially in oligotrophic soils (Bustamante et al. 2004). Therefore, the inorganic soil nitrogen can vary independently from the most of soil nutrients along the Al-Ca gradient. Moreover, the other species that are not CAL or LEG comprise a species-rich set of woody plants in the Cerrado and their responses to environmental factors of the Al-Ca gradient working along with the inorganic-N gradient are still unknown. This work aims to answer i) whether without influence of other main environmental variables, CAL and LEG are distinct from each other as well as from the set of other species considering their responses to soil factors and to light regime in the analyses. Another question is ii) whether the other Cerrado species that are not CAL and LEG can be encompassed as typical Cerrado savanna-species (SAV) *sensu* Rizzini (1963). A final question is iii) whether these analyses reveal additional axes that explain the environmental gradient of the Cerrado (and possibly of tropical savanna on acidic soils). In order to test the soil and light effects, avoiding other environmental biases, we sampled a Cerrado with plots in five different soil types with minimum variation in topography, in altitude, within the same climate, within the same fragment (i.e., the same landscape) and without fire disturbance for more than 50 years. For that, we analysed and modelled soil factors and canopy openness for all plots as well as we analysed and modelled leaf contents of nitrogen, of carbon, leaf $\delta^{13}\text{C}$, leaf $\delta^{15}\text{N}$, maximum plant height, maximum stem diameter, bark thickness, wood density and specific leaf area for all plots and for all sampled species of woody plants.

Material and methods

Study area

The study was carried out in the 200-ha Paraopeba Reserve in the state of Minas Gerais (19°20'S, 44°20'W), Brazil. The climate of the region is classified as Aw (tropical humid) by the Köppen system with a rainy summer from October to April and a dry season from May to September. The mean annual temperature and mean annual rainfall are 20.9°C and 1328 mm, respectively. The vegetation of the reserve

is the product of regeneration after clear-cutting in 1952. There are records of fires in 1960 and 1963, since which the area as a whole has been protected from fire (Neri et al. 2012).

We carried out the study on a wide range of soils spread over a short distance, where differences in vegetation are ascribed to soil attributes since variation in climate and topography was negligible (Neri et al. 2012).

The soils are classified as i) dystrophic Haplic Cambisol, ii) Yellow Latosol, iii) Red-Yellow Latosol, iv) dystrophic Red Latosol and v) mesotrophic Red Latosol (EMBRAPA 2006; Neri et al. 2012). Along this gradient the darker the soil the greater the biomass and density of woody plants. Colluvial materials from weathered limestone or *in situ* alteration of such calcareous rocks influence both eutrophic and dystrophic red soils. On the other hand, Cerrado savannas are related to either shallow or deep yellowish soils (Cambisols/Latosols) developed from slate, a pelitic Al-rich nutrient-poor metamorphic rock. Since there is a wide range of soil characteristics within a small area, this fragment of Cerrado is a suitable system for increasing knowledge regarding plant functional groups with different strategies for resource use and stress tolerance without climatic, topographical and dispersion limitation biases.

Vegetation, soil and light

The structure of the woody layer was evaluated by equally distributing 24 plots (20-m x 20-m) along the aforementioned soil gradient. We sampled three plots of Inceptisol (Cambisol), three plots of mesotrophic Red Latosol, three plots of dystrophic Red Latosol, six plots of Yellow Latosol and nine plots for Red-Yellow Latosol (Yellow Latosol and Red-Yellow Latosol are intermediary soils with more plots to better represent the intermediate portion of the gradients). All plots were randomly selected inside soil types areas. The soil characterization is in previous publications (Neri et al. 2012, 2013). From July 2011 to September 2012 we recorded all individual woody plants with stem circumferences equal to or greater than 10 cm at ground level. The classification of species into families followed APG IV (The Angiosperm Phylogeny Group 2016), with the nomenclature of species and abbreviations being in agreement with Brazilian Flora Checklist (www.floradobrasil.jbr.gov.br).

Soil samples were collected from all 20-m x 20-m plots, each one comprising 10 subsamples from a depth of 0–10 cm. Soil samples taken for chemical analysis were air-dried and sieved. Since Cerrado soils generally possess low nutrient status and fertility, which have been shown to influence its vegetation, we used cation exchange capacity (CEC), inorganic N and pH as proxies for soil fertility. Furthermore, since aluminium saturation determines the fraction of Al^{3+} in CEC, it was considered a main soil factor. The pH values were obtained from aqueous solution while inorganic N was gathered from the sum of nitrate (NO_3^-

) and ammonium (NH₃) content. Colorimetric analyses were used for measuring NO₃⁻ and NH₃. All analyses were performed in the Soil Laboratory at the Universidade Federal de Viçosa, following methods for Brazilian tropical soils (EMBRAPA, 1997).

Canopy openness was measured as a proxy for light availability by taking hemispherical photographs (Beaudet and Messier 2002). The photos were taken using a Nikon Coolpix 5700 digital camera, with a Nikon UR-E12 extender and a Nikon FC-E9 fisheye lens, supported at 1.5 m above the ground by a tripod with the zenith centred and a north orientation. The photos were taken at dusk or during perfectly overcast days (Breshears and Ludwig, 2010) during the middle and the end of the rainy season (February 2012) when the canopy cover was maximum and light could act as a limiting factor for plants. Each plot was divided into four 10-m x 10-m subplots and photos were taken at the center of each subplot. Photos were analyzed with Gap Light Analyzer (GLA) software (Frazer et al. 1999). Canopy openness values were obtained for each plot by averaging canopy openness of four 10-m x 10-m subplots.

In order to choose the dominant species that totalize ~75% of abundance, we used data from a previous survey on 3 hectares with 15 plots of 20-m x 20-m on each soil type, 75 plots in total. We found 14,671 individuals of 174 woody species of 51 families in the woody communities of the studied vegetation. Of these, 10,866 individuals belonged to the 34 species that made up 74.06% of the total abundance of the community. We used the 75 plots only to choose that 34 species with ~75% of abundance. All other analyses used the 24 chosen plots. Thus, those 34 species were sampled and analysed in the 24 chosen plots previously described.

Functional traits

Despite the high floristic richness of the Cerrado, the abundance of the woody community are usually dominated by a few species. As 75% of dominance is the lower limit for highest level of dominance for sampling vegetation methods as Braun-Blanquet and Domin-Krajina (Mueller-Dombois and Ellenberg 2003; Maarel 2009), we considered the species that collectively comprise ~ 75% of the local abundance as rulers of the Cerrado functioning. Thus, we evaluated functional traits for 34 selected species, which accounted for 74.06% of the cumulative relative abundance in the Cerrado of Paraopeba Reserve. All individuals of selected species were assessed for measuring traits.

Trait-based analysis requires selection of critical traits to the community processes of interest (Kraft et al. 2007). Six key traits (Table 1) related to competitive ability, resources use and tolerance were used (Cianciaruso et al. 2012; Pérez-Harguindeguy et al. 2013). Wood density was determined by the

Global Wood Density Database, subsection Tropical South America (Chave et al. 2009). Because data on wood density was obtained from the online database, it had no variability. Other traits were measured by widely accepted methods (Cornelissen et al. 2003; Pérez-Harguindeguy et al. 2013). The sampling was plot-specific, which allowed us to assess environmental factors (Cianciaruso et al. 2009). Thus, we used functional traits measured on individuals of a plot as a proxy of functional traits of that plot.

Carbon, nitrogen and stable isotopes analyses ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$)

We used stable isotopes analyses in order to understand the conditions of functional groups and set of species in each plot under different soil and light conditions. Three leaves healthy and fully developed from all individuals sampled in each plot were taken during the rainy season (March of 2013). The leaves were sampled at a similar distance from the ground and at the north side of the canopy. The leaves were dried to constant weight (65°C) without petioles and midribs and ground to a powder using a ball mill (Retsch, Haan, Germany) for measuring carbon, nitrogen, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. Nitrogen and carbon concentration, and $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were analysed using an elemental analyzer (HEKAtech, Weinberg, Germany) with a continuous-flow stable isotope ratio mass spectrometer (ISOPRIME, GV, Manchester, UK), and measured against an ammonium sulphate standard (IAEA.N2). C and N isotope ratios were presented in δ notation:

$$\delta^{13}\text{C sample} = R_{\text{sample}} - R_{\text{standard}} / R_{\text{standard}}$$

where R_{standard} is the $^{13}\text{C}/^{12}\text{C}$ of Pee Dee Belemnite (PDB) and R_{sample} is the $^{13}\text{C}/^{12}\text{C}$ ratio of the sampled leaves.

$$\delta^{15}\text{N sample} = R_{\text{sample}} - R_{\text{standard}} / R_{\text{standard}}$$

where R_{standard} is the $^{15}\text{N}/^{14}\text{N}$ ratio of atmospheric N_2 and R_{sample} is the $^{15}\text{N}/^{14}\text{N}$ ratio of the sampled leaves. The repeated measurement precision was 0.2 ‰.

Species groups and data analysis

Different groups of woody species were considered based on the dispersion of species along the ordination axis of a multidimensional scaling – MDS using two functional traits as variables for species:

mean leaf C/N and mean aluminium saturation. Mean leaf C/N was calculated as arithmetic mean of leaf C/N of all individuals of each species. Mean aluminium saturation was calculated giving the aluminium saturation of soil of each plot to each individual occurring in that plot; then, was calculated the arithmetic mean of aluminium saturation among individuals of each species. The CAL was considered the calcicoles species occurring in the Cerrado with lowest aluminium saturation means. LEG are the leguminous woody species that occur in the Cerrado, most of them belonging to nitrogen-fixing genera (Sprent 2009) as well as other species with the lowest leaf C/N. SAV were the woody species with the highest leaf C/N. and are not nitrogen-fixing.

In order to investigate relationships between soil and light effects and abundance of the found species groups, we created generalized linear models (GLMs) applying abundance as a response variable of soil cation exchange capacity (CEC), of soil pH, of soil inorganic N, of soil aluminium saturation and of canopy openness.

We constructed a numerical matrix, with averaged values of SLA, ratio of maximum height to maximum diameter (Hmax/Dmax) as stem elongation, leaf C/N, bark thickness and woody density for the 34 species and species-weighted averaged values for CAL, LEG and SAV in plots.

Statistical analysis

RLQ analysis was carried out using the functions ‘dudi.coa’, ‘dudi.hillsmith’, ‘dudi.pca’ and ‘rlq’ of the ade4 package (Dray et al. 2018) for canonical ordination of species abundances, functional traits (SLA, Hmax/Dmax, %C and %N), plots and three soil variables (CEC, aluminium saturation and inorganic nitrogen, all square-rooted transformed). The Fourth-corner analysis was carried out using the ‘fourthcorner’ function of the Ade4 package performing 50 000 permutations with FDR adjustment of P values for multiple comparisons between the soil variables and the functional traits of the RLQ analysis (Peres-Neto et al. 2017; Braak et al. 2018). All analyses were performed within the R statistical environment (R Development Core Team, 2015). The canonical ordination was the basis on which the groups CAL, LEG and SAV were proposed.

ANOSIM test was performed for testing the 34 species divided into the three groups by means of the SLA, Hmax/Dmax, %C and %N for calculation of Euclidean Distance dissimilarity using the functions ‘dist’ and ‘anosim’ from the ‘Vegan’ package in R statistical environment (R Development Core Team 2015).

GLMs were carried out using the ‘glm’ function of the MASS package with Poisson error distribution for count data (i.e., abundance) to test the response of CAL, LEG and SAV abundances to soil pH, to soil

inorganic N, to soil aluminium saturation and to canopy openness. All explaining variables were scaled. We used the dredge function of the MuMIn package (Bartón 2018) to select the models that explain our data among all possible combinations of two predictive variables that are not correlated ($r < 0.6$, https://github.com/rojaff/dredge_mc) or one predictive variable (i.e., overfitting avoidance). The Akaike Information Criterion (AIC) was used to select the best models (Symonds and Moussalli 2011), and all models with ΔAIC less than 2 was considered equally parsimonious. From all selected models, we calculated average parameters and unconditioned standard errors weighted by the likelihood ratio using the 'model.avg' function of the MuMIn package. The function performs significance tests for each predictor estimated in this average model. The final model will be validated by plotting residual x adjusted values, residual versus predictors and observing the distribution of residues. All analyses were performed within the R statistical environment (R Development Core Team 2015).

For each one of the 34 species was calculated a mean value for each functional trait. Then were calculated mean values for each functional group based on the values of each species. GLMs with gaussian error distribution were performed using the glm function of the MASS package to test differences between means of groups for leaf C/N, leaf nitrogen content, specific leaf area, woody density, bark thickness and Hmax/Dmax (stem elongation). The difference between groups for each functional trait was additional tested using dunnTest function of the dunn.test package in cases that the differences between groups were dubious by GLMs. All analyses were performed within the R statistical environment (R Development Core Team 2015).

Results

The RLQ analysis showed that three groups of species were related mainly to stem elongation (H/DSH), specific leaf area (SLA), leaf nitrogen contents (%N) and leaf carbon contents (%C). The 34 woody species were grouped into three groups proposed based on the canonical ordination of RLQ: Calicole species (CAL) mainly associated to high H/DSH and high SLA, leguminous species (LEG) mainly associated to high %N and high %C, and savanna species (SAV) mainly associated to low H/DSH, low SLA, low %N and low %C (Figure 1, Appendix S1). The ANOSIM analysis testing the consistency of the three proposed groups with 34 species responding to SLA, Hmax/Dmax, %C and %N was significant ($p=0.01796$).

The fourth-corner analysis showed that the H/DSH and SLA are positively related to soil CEC and negatively related to soil aluminium saturation (m). Leaf nitrogen contents (%N) and leaf carbon contents (%C) are positively related to soil inorganic nitrogen (Figure 2).

The best GLM full models (i.e., two uncorrelated variables for each species group [$r < 0.6$], delta AIC ≤ 2 , Appendix S2) for CAL species showed aluminium saturation, and canopy openness as explanatory variables of abundance (Figure 3). LEG species, the best full models (i.e. two uncorrelated variables [$r < 0.6$], delta AIC ≤ 2 , Appendix S2) showed pH, and canopy openness as explanatory variables negatively associated with abundance (Figure 4). The best full models for SAV species (i.e. two uncorrelated variables [$r < 0.6$], delta AIC ≤ 2 , Appendix S2) showed inorganic nitrogen, and CEC as explanatory variables of abundance (Figure 5, Table 1). All selected models were negative relations between abundance of species groups and environmental variables. CAL were severely limited by aluminium toxicity and excessive light indicating that excessive light and high aluminium are a damaging combination. The CAL limitation by aluminium saturation is clearly a stress limitation since aluminium is a known and ubiquitous stress in plants (Kochian 1995; Jansen et al. 2002). The CAL limitation by canopy openness is likely light stress caused by photon flux density higher than used by photosynthesis. This light stress would be caused by excessive light or by photosynthesis limitation because of indirect stressors as aluminium (i.e., the other variable of the global model) that limit root growth and nutrient uptake (Demmig-Adams and Adams 1992; Akaya and Takenaka 2001). LEG were strongly limited by increasing pH and to increasing light, in a deleterious combination of high pH and excessive light stresses. Possibly in the LEG case, interconnected stresses are acting since the light stress is caused combined with pH increasing, which is possibly an effect of photosynthesis limitation caused by a low phosphorus uptake in soils with higher pH (Riley and Barber 1971; Hocking 2001; Thomas et al. 2006). SAV species are not benefitted from the increasing resources of soil inorganic nitrogen and soil CEC. Thus, the SAV limitation not caused by stresses, but the lack of ability to use efficiently soil resources (i.e., inorganic nitrogen and CEC) to grow.

Among the 34 species (fourteen CAL, six LEG, fourteen SAV) CAL presented greater specific leaf area (SLA) than SAV, LEG presented the highest leaf nitrogen content and SAV the lowest, SAV presented the highest C/N and LEG the lowest (Figure 6). CAL presented the higher wood density than SAV and CAL and LEG presented higher stem elongation (Hmax/Dmax) than SAV. Bark thickness was not different between groups, but CAL presented low variance and SAV presented high variance (Figure 7).

The C/N did not differ among groups, but was negatively related to leaf $\delta^{13}\text{C}$ (Figure 8). Leaf $\delta^{15}\text{N}$ neither differed significantly nor was related to any other functional trait (data and results not shown).

Discussion

CAL, LEG and SAV species can be used as different groups of species responding to environmental variables and with different functional profiles. CAL, LEG and SAV abundances responded differently to the explanatory variables of soil and to canopy openness of (i.e., light). CAL, LEG and SAV differed from each other in leaf traits (i.e., nitrogen content, C/N and SLA) and in stem traits (wood density and Hmax/Dmax) indicating distinctive functioning. Therefore, the CAL, LEG and SAV presented distinctive abundance responses to soil/light variables as well as distinctive set of functional traits. The proposed species grouping was significantly consistent with the traits SLA, Hmax/Dmax, %C and %N as well as the soil variables.

CAL abundance was the only group negatively related to aluminium saturation. CAL was negatively related to canopy openness as well. CAL presented greater stem elongation and greater SLA compared to SAV, indicating that CAL species are associated with more fertile soils of higher productivity than SAV, congruently to their high SLA (Reich et al. 1997; Westoby 1998). The deciduousness (i.e., short leaf lifespan) of the most of CAL species (Oliveira-Filho et al. 2006) suggests also high productivity (Reich *et al.*, 1997) in plots dominated by CAL species. Therefore, functional traits and responses to environmental variables that distinguish CAL species are related to competition ability for light, to long lifespan (i.e., high Hmax/Dmax, low canopy openness and high wood density), to habitats with high productivity (i.e., high SLA) (Westoby 1998; Moles et al. 2009; Pérez-Harguindeguy et al. 2013; Kunstler et al. 2016) as well as to intolerance to soil stress.

The abundance of LEG was negatively related to pH, and to canopy openness. Their low leaf C/N and highest leaf nitrogen content are congruent with the nitrogen-fixing function and with low canopy openness (Bustamante et al. 2004, 2006). Functional traits and environmental variables that distinguish LEG species are related to competition for light, to long lifespan (i.e., medium to high wood density and low canopy openness), to high nitrogen status (i.e., low C/N and high leaf nitrogen), to soil nutritional stress-tolerance (i.e., low pH).

The SAV abundance is negatively related with soil inorganic nitrogen and with CEC. The negative relation of SAV abundance with soil resources are congruent with their low SLA, suggesting suitability to habitats with low productivity (Dantas et al. 2013; Pérez-Harguindeguy et al. 2013). Functional traits

suggest SAV species as bad competitors for light with short lifespan (i.e., low stem elongation and low wood density), tolerant to soil-nutritional stress of habitats with low productivity (i.e., high C/N, low leaf nitrogen content, low SLA, low soil CEC and low inorganic soil N).

Within CAL species, there are six species that would be grouped separately forming another intermediate group or even be placed into another group. However, as the objective of the grouping is to decrease diversity in order to find the main soil and light influences in the Cerrado, we chose to use three groups. Therefore, our choice was to include *Alibertia edulis*, *Erythroxylum daphnitis*, *Lithraea molleoides*, *Myrsine umbellata*, *Pera glabrata* and *Terminalia argentea* inside CAL group.

From that first grouping effort that listed ‘Cerrado typical species’ and separated them from species that ‘mainly occur in other vegetation types’ (Rizzini 1963), the CAL (calcicoles) were in the species list that mainly occur in other vegetation types (Ratter et al. 1977; Oliveira-Filho et al. 2006; Durigan and Ratter 2006) and the LEG had species in both lists (Meira-Neto et al. 2017). The Cerrado typical species *sensu* Rizzini (1963) are the bulk of SAV, but some SAV were listed as species that ‘mainly occur in other vegetation types’ by the author. It is the case of *Callistene major*, *Curatella americana*, *Miconia albicans*, *Styrax camporum*, *Tapirira guianensis* and *Vochysia tucanorum*. Therefore, the Rizzini’s groups may be a good hint, but they are not consistent functionally.

The negative relation between C/N and $\delta^{13}\text{C}$ among all groups show that the higher the N nutritional status, the more the ^{13}C proportion in leaves. A possible explanation of low $\delta^{13}\text{C}$ in some of these species may be that deep roots can access water during the dry season on deep soils in Cerrado (Rawitscher et al. 1943; Ferri 1944; Rachid 1947; Rossatto et al. 2012). Therefore, low ^{13}C proportion might be because the higher stomatal conductance and high CO_2 assimilation rates lower the $\delta^{13}\text{C}$ as Rubisco freely discriminates and prefers ^{12}C to ^{13}C (Kohn, 2010). Thus, the high nitrogen status associated with high $\delta^{13}\text{C}$ may enhance drought tolerance (Wright et al. 2001, 2003) and should be further investigated in LEG and CAL that presented lower C/N than SAV. The high $\delta^{13}\text{C}$ in leaves suggest stomatal closure as well as higher water use efficiency (Goldstein et al. 1989; Scartazza et al. 1998). The $\delta^{13}\text{C}$ of the cerradões dominating groups (CAL+LEG) is higher than the $\delta^{13}\text{C}$ of SAV ($p=0.0317$, GLM, gaussian distribution, results not shown). Therefore, our results suggest that ecosystems dominated by CAL and LEG (cerradões) are on soils with higher nutritional status, with less canopy openness and rather tolerate drought stress more than open Cerrado plots dominated by SAV. Considering that $\delta^{13}\text{C}$ enrichment is associated to stomatal closure as a response to drought stress, ^{13}C enriched functional groups can tolerate longer periods of water

stress (Werner and Máguas 2010; Máguas et al. 2011) than SAV (i.e., ^{13}C depleted functional group). Moreover, decreased leaf C/N of LEG and CAL may to improve plant growth and canopy coverage. Thus, higher $\delta^{13}\text{C}$ values suggest longer periods of biomass incorporation during droughts and different phenological developments (Werner and Máguas 2010). Therefore, the results show that as leaf C/N decreases, leaf $\delta^{13}\text{C}$ increases, drought tolerance increases and water use efficiency increases, similarly to found results of other studies (Goldstein et al. 1989; Scartazza et al. 1998; Coletta et al. 2009). As CAL and LEG are ^{13}C enriched and dominate cerradões, the relation between high nitrogen status and high water use efficiency must be further assessed in CAL and LEG.

Despite the reported evidence of nitrogen fixation by nodulated genera in the Cerrado (Sprent *et al.* 1996), our results of leaf $\delta^{15}\text{N}$ were not significant. However, the proportion of this stable isotope should be further studied in both soil and plants to determine whether the sources of N for the different functional groups are the same (Marshall et al. 2007; Craine et al. 2015), and whether there are ecological processes involved in nitrogen uptake by plants (e.g., mycorrhizal and nitrogen-fixing bacteria associations). In addition to $\delta^{15}\text{N}$, the existence of variation in leaf N/P among groups of species should be determined (Bustamante *et al.*, 2004, 2006; Townsend *et al.*, 2007). *Copaifera* was the only leguminous genera that is not considered nitrogen fixing among LEG (Sprent 2009) and deserves attention in future studies concerning nitrogen fixing species.

Based on the GLMs results and fourth-corner analysis, we propose a model with three main gradients and two functional traits to explain the responses of CAL, LEG and SAV: canopy openness, soil aluminium saturation, pH, CEC, and soil inorganic nitrogen. The model presented in Figure 9 shows the main gradient for each functional group. CAL, associated with high SLA, dominate soils/plots with low aluminium saturation, and low canopy openness. LEG, associated to low C/N, dominate soils/plots with high pH and with low canopy openness. The SAV, associated to low stem elongation, dominate soils/plots with low inorganic nitrogen low CEC and with low inorganic nitrogen. That model was proposed based on the figures 1 to 8 and respective statistical tests. The model is hypothetical and should be tested in any tropical savannas on acidic soils.

The proposed model may generate hypotheses and questions for different situations throughout the Cerrado and other tropical savannas. The first question might be how general is the model. The hypotheses derived from each side of the triangle would deepen the understanding of particular situations in the Cerrado, as demonstrated by the side represented between the mesotrophic and dystrophic cerradões gradient (CAL-LEG) (Meira-Neto et al. 2017) which would be interpreted as the merged red and green

gradients of the model. Thus, are the mesotrophic cerradões-savanna gradient (CAL-SAV) as well as dystrophic cerradões-savanna gradient (LEG-SAV) the resulting merged gradients of the other triangle sides? What would be the responses of the groups where one of them is underrepresented or even missing?

Acknowledgements

The authors thank CNPq (proc. 446698/2014-8), FAPEMIG (PPM 0584-16), PIRSES (GA-2010-269206), CAPES and PPGBot/UFV for scholarships, grants and financial support. The authors also thank the Paraopeba Reserve staff for providing help and facilities. JAAMN has a productivity fellowship from CNPq (307591/2016-6). The research complies with the current Brazilian law.

408 Literature

- 409
- 410 Akaya M, Takenaka C (2001) Effects of aluminum stress on photosynthesis of *Quercus glauca* Thumb.
411 Plant Soil 237:137–146
- 412 Bartón K (2018) Package ‘MuMIn, 1.40.4
- 413 Beaudet M, Messier C (2002) Variation in canopy openness and light transmission following selection
414 cutting in northern hardwood stands: an assessment based on hemispherical photographs. Agric
415 For Meteorol 110:217–228. [https://doi.org/10.1016/S0168-1923\(01\)00289-1](https://doi.org/10.1016/S0168-1923(01)00289-1)
- 416 Braak CJF ter, Peres-Neto PR, Dray S (2018) Simple parametric tests for trait–environment association. J
417 Veg Sci 29:801–811. <https://doi.org/10.1111/jvs.12666>
- 418 Breshears DD, Ludwig JA (2010) Near-ground solar radiation along the grassland–forest continuum:
419 Tall-tree canopy architecture imposes only muted trends and heterogeneity. Austral Ecol 35:31–
420 40. <https://doi.org/10.1111/j.1442-9993.2009.02009.x>
- 421 Bustamante MMC, Martinelli LA, Silva DA, et al (2004) 15N natural abundance in woody plants and
422 soils of central brazilian savannas (cerrado). Ecol Appl 14:200–213. <https://doi.org/10.1890/01-6013>
- 424 Bustamante MMC, Medina E, Asner GP, et al (2006) Nitrogen cycling in tropical and temperate
425 savannas. Biogeochemistry 79:209–237. <https://doi.org/10.1007/s10533-006-9006-x>
- 426 Chave J, Coomes D, Jansen S, et al (2009) Towards a worldwide wood economics spectrum. Ecol Lett
427 12:351–366. <https://doi.org/10.1111/j.1461-0248.2009.01285.x>
- 428 Cianciaruso MV, Silva IA, Batalha MA, et al (2012) The influence of fire on phylogenetic and functional
429 structure of woody savannas: Moving from species to individuals. Perspect Plant Ecol Evol Syst
430 14:205–216. <https://doi.org/10.1016/j.ppees.2011.11.004>
- 431 Cianciaruso MV, Silva IA, Batalha MA (2009) Diversidades filogenética e funcional: novas abordagens
432 para a Ecologia de comunidades. Biota Neotropica 9:93–103. <https://doi.org/10.1590/S1676-06032009000300008>
- 434 Coletta L, Nardoto GB, Latansio S (2009) Isotopic view of vegetation and carbon and nitrogen cycles in a
435 Cerrado ecosystem, southeastern Brazil. Sci Agric 66:467–475
- 436 Cornelissen JHC, Lavorel S, Garnier E, et al (2003) A handbook of protocols for standardised and easy
437 measurement of plant functional traits worldwide. Aust J Bot 51:335–380
- 438 Coutinho LM (1990) Fire in the Ecology of the Brazilian Cerrado. In: Goldammer DJG (ed) Fire in the
439 Tropical Biota. Springer Berlin Heidelberg, pp 82–105
- 440 Craine JM, Brookshire ENJ, Cramer MD, et al (2015) Ecological interpretations of nitrogen isotope ratios
441 of terrestrial plants and soils. Plant Soil 396:1–26. <https://doi.org/10.1007/s11104-015-2542-1>
- 442 Dalmagro HJ, Lobo F de A, Vourlitis GL, et al (2014) The physiological light response of two tree
443 species across a hydrologic gradient in Brazilian savanna (Cerrado). Photosynthetica 52:22–35.
444 <https://doi.org/10.1007/s11099-014-0001-5>
- 445 Dantas VL, Batalha MA, Pausas JG (2013) Fire drives functional thresholds on the savanna–forest
446 transition. Ecology 94:2454–2463. <https://doi.org/10.1890/12-1629.1>
- 447 de Assis ACC de, Coelho RM, da Silva Pinheiro E da S, Durigan G (2011) Water availability determines
448 physiognomic gradient in an area of low-fertility soils under Cerrado vegetation. Plant Ecol
449 212:1135–1147. <https://doi.org/10.1007/s11258-010-9893-8>

- 450 Demmig-Adams B, Adams WW (1992) Photoprotection and Other Responses of Plants to High Light
451 Stress. *Annu Rev Plant Physiol Plant Mol Biol* 43:599–626.
452 <https://doi.org/10.1146/annurev.pp.43.060192.003123>
- 453 Díaz S, Cabido M (2001) Vive la différence: plant functional diversity matters to ecosystem processes.
454 *Trends Ecol Evol* 16:646–655. [https://doi.org/10.1016/S0169-5347\(01\)02283-2](https://doi.org/10.1016/S0169-5347(01)02283-2)
- 455 Dray S, Dufour A-B, Thioulouse J, et al (2018) ade4: Analysis of Ecological Data: Exploratory and
456 Euclidean Methods in Environmental Sciences. Version 1.7-11URL [https://CRAN.R-](https://CRAN.R-project.org/package=ade4)
457 [project.org/package=ade4](https://CRAN.R-project.org/package=ade4)
- 458 Durigan G, Ratter JA (2006) Successional changes in Cerrado and Cerrado/forest ecotonal vegetation in
459 western São Paulo State, Brazil. *Edinb J Bot* 63:119–130.
460 <https://doi.org/10.1017/S0960428606000357>
- 461 EMBRAPA (2006) Sistema Brasileiro de Classificação de Solos. Centro Nacional de Pesquisa de Solos,
462 Rio de Janeiro
- 463 EMBRAPA (1997) Manual de métodos de análise de solo, 2nd edn. EMBRAPA - Centro Nacional de
464 Pesquisa de Solos, Rio de Janeiro, Brazil
- 465 Felfili JM, Fagg CW (2007) Floristic composition, diversity and structure of the “cerrado” sensu stricto
466 on rocky soils in northern Goiás and southern Tocantins, Brazil. *Braz J Bot* 30:375–385.
467 <https://doi.org/10.1590/S0100-84042007000300004>
- 468 Ferri MG (1944) Transpiração de plantas permanentes dos cerrados. *Bol Fac Filos Ciênc E Let USP -*
469 *Botânica* 4:159–224
- 470 Franco AC, Rossatto DR, de Carvalho Ramos Silva L, da Silva Ferreira C (2014) Cerrado vegetation and
471 global change: the role of functional types, resource availability and disturbance in regulating
472 plant community responses to rising CO₂ levels and climate warming. *Theor Exp Plant Physiol*
473 26:19–38. <https://doi.org/10.1007/s40626-014-0002-6>
- 474 Frazer GW, Canham CD, Lertzman KP (1999) Gap Light Analyzer (GLA): Imaging software to extract
475 canopy structure and gap light transmission indices from true-colour fisheye photographs. Users
476 Manual and Program Documentation, Version 2.0. Simon Fraser University and Institute of
477 Ecosystem Studies, Burnaby, British Columbia
- 478 Furley PA, Ratter JA (1988) Soil Resources and Plant Communities of the Central Brazilian Cerrado and
479 Their Development. *J Biogeogr* 15:97–108. <https://doi.org/10.2307/2845050>
- 480 Goldstein G, Rada F, Sternberg L, et al (1989) Gas exchange and water balance of a mistletoe species and
481 its mangrove hosts. *Oecologia* 78:176–183. <https://doi.org/10.1007/BF00377153>
- 482 Goodland R (1971) A physiognomic analysis of the ‘Cerrado’ vegetation of central Brasil. *J Ecol* 59:411–
483 419. <https://doi.org/10.2307/2258321>
- 484 Goodland R, Pollard R (1973) The Brazilian Cerrado vegetation: a fertility gradient. *J Ecol* 61:219–224.
485 <https://doi.org/10.2307/2258929>
- 486 Hocking PJ (2001) Organic acids exuded from roots in phosphorus uptake and aluminum tolerance of
487 plants in acid soils. In: *Advances in Agronomy*. Academic Press, pp 63–97
- 488 Hoffmann WA (2002) Direct and Indirect Effects of Fire on Radial Growth of Cerrado Savanna Trees. *J*
489 *Trop Ecol* 18:137–142
- 490 Hutley LB, O’Grady AP, Eamus D (2000) Evapotranspiration from Eucalypt open-forest savanna of
491 Northern Australia. *Funct Ecol* 14:183–194. <https://doi.org/10.1046/j.1365-2435.2000.00416.x>

- 492 Jansen S, Broadley MR, Robbrecht E, Smets E (2002) Aluminum hyperaccumulation in angiosperms: a
493 review of its phylogenetic significance. *Bot Rev* 68:235–269. <https://doi.org/10.1663/0006->
494 8101(2002)068[0235:AHIAAR]2.0.CO;2
- 495 Kochian LV (1995) Cellular mechanisms of aluminum toxicity and resistance in plants. *Annu Rev Plant*
496 *Physiol Plant Mol Biol* 46:237–260. <https://doi.org/10.1146/annurev.pp.46.060195.001321>
- 497 Kohn MJ (2010) Carbon isotope compositions of terrestrial C3 plants as indicators of (paleo)ecology and
498 (paleo)climate. *Proc Natl Acad Sci U S A* 107:19691–19695.
499 <https://doi.org/10.1073/pnas.1004933107>
- 500 Kraft NJB, Cornwell WK, Webb CO, Ackerly DD (2007) Trait evolution, community assembly, and the
501 phylogenetic structure of ecological communities. *Am Nat* 170:271–283.
502 <https://doi.org/10.1086/519400>
- 503 Kunstler G, Falster D, Coomes DA, et al (2016) Plant functional traits have globally consistent effects on
504 competition. *Nature* 529:204–207. <https://doi.org/10.1038/nature16476>
- 505 Lavorel S, Garnier E (2002) Predicting changes in community composition and ecosystem functioning
506 from plant traits: revisiting the Holy Grail. *Funct Ecol* 16:545–556.
507 <https://doi.org/10.1046/j.1365-2435.2002.00664.x>
- 508 Maarel E van der (2009) *Vegetation Ecology*. John Wiley & Sons
- 509 Máguas C, Rascher KG, Martins-Loução A, et al (2011) Responses of woody species to spatial and
510 temporal ground water changes in coastal sand dune systems. *Biogeosciences* 8:3823–3832.
511 <https://doi.org/10.5194/bg-8-3823-2011>
- 512 Marshall J, Brooks R, Lajtha K (2007) Sources of variation in the stable isotopic composition of plants.
513 In: Michener R, Lajtha K (eds) *Stable Isotopes in Ecology and Environmental Science*. Wiley -
514 Blackwell, Malden, pp 22–60
- 515 Materechera S, Mandiringana O, Mbokodi P, Nyamapfene K (1998) Organic matter, pH and nutrient
516 distribution in soil layers of a savanna Thornveld sUbjected to different burning frequencies at
517 Alice in the Eastern Cape. *South Afr J Plant Soil* 15:109–115
- 518 Meira-Neto JAA, da Silva MCNA, Tolentino GS, et al (2018) Shading, nitrogen and soil texture rule a
519 sandy savanna: Does facilitation rule its patchy physiognomy as well? *Flora* 239:45–51.
520 <https://doi.org/10.1016/j.flora.2017.11.007>
- 521 Meira-Neto JAA, Tolentino GS, da Silva MCNA da, et al (2017) Functional antagonism between
522 nitrogen-fixing leguminous trees and calcicole-drought-tolerant trees in the Cerrado. *Acta Bot*
523 *Bras* 31:11–18. <https://doi.org/10.1590/0102-33062016abb0380>
- 524 Moles AT, Warton DI, Warman L, et al (2009) Global patterns in plant height. *J Ecol* 97:923–932.
525 <https://doi.org/10.1111/j.1365-2745.2009.01526.x>
- 526 Motta P, Curi N, Franzmeier D (2002) Relation of Soils and Geomorphic Surfaces in the Brazilian
527 Cerrado: Ecology and Natural History of a Neotropical Savanna. In: *The Cerrados of Brazil:*
528 *ecology and natural history of a neotropical savanna*, OLIVEIRA, P.S. & MARQUIS, R.J.
529 Columbia University Press, Chicago
- 530 Mueller-Dombois D, Ellenberg H (2003) *Aims and Methods of Vegetation Ecology*. Blackburn Press
- 531 Neri AV, Schaefer CEGR, Silva AF, et al (2012) The influence of soils on the floristic composition and
532 community structure of an area of Brazilian Cerrado vegetation. *Edinb J Bot* 69:1–27.
533 <https://doi.org/10.1017/S0960428611000382>
- 534 Neri AV, Schaefer CEGR, Souza AL, et al (2013) Pedology and plant physiognomies in the cerrado,
535 Brazil. *An Acad Bras Ciênc* 85:87–102. <https://doi.org/10.1590/S0001-37652013000100007>

- 536 Oliveira-Filho A, Ratter JA (2002) Vegetation physiognomies and woody flora of the Cerrado Biome. In:
537 The Cerrados of Brazil: ecology and natural history of a Neotropical savanna. Columbia
538 University Press, New York, pp 91–120
- 539 Oliveira-Filho AT, Jarenkow JA, Rodal M (2006) Floristic relationships of seasonally dry forests of
540 eastern South America based on tree species distribution patterns. In: Neotropical savannas and
541 seasonally dry forests: plant diversity, biogeography, and conservation, R . Toby Pennington ,
542 Gwilym P . Lewis , and James A . Ratter eds. CRC Press, pp 159–192
- 543 Peres-Neto PR, Dray S, Braak CJF ter (2017) Linking trait variation to the environment: critical issues
544 with community-weighted mean correlation resolved by the fourth-corner approach. *Ecography*
545 40:806–816. <https://doi.org/10.1111/ecog.02302>
- 546 Pérez-Harguindeguy N, Díaz S, Garnier E, et al (2013) New handbook for standardised measurement of
547 plant functional traits worldwide. *Aust J Bot* 61:167–234
- 548 Pinheiro LFS, Kolb RM, Rossatto DR (2016) Changes in irradiance and soil properties explain why
549 typical non-arboreal savanna species disappear under tree encroachment. *Aust J Bot* 64:333–341.
550 <https://doi.org/10.1071/BT15283>
- 551 R Development Core Team (2015) R: A language and environment for statistical computing.
552 <http://www.R-project.org/>. Accessed 19 Feb 2015
- 553 Rachid M (1947) Transpiração e sistemas subterrâneos da vegetação de verão dos campos cerrados de
554 Emas. *Bol Fac Filos Ciênc E Let USP - Botânica* 5:1–140
- 555 Ratter JA, Bridgewater S, Ribeiro JF (2003) Analysis of the floristic composition of the Brazilian Cerrado
556 vegetation III: comparison of the woody vegetation of 376 areas. *Edinb J Bot* 60:57–109.
557 <https://doi.org/10.1017/S0960428603000064>
- 558 Ratter JA, Montgomery R, Askew D, Gifford DR (1977) Observações adicionais sobre o cerradão de
559 solos mesotróficos no Brasil Central. In: IV Simpósio Sobre o Cerrado. Itatiaia, São Paulo, pp
560 303–316
- 561 Rawitscher F, Ferri MG, Rachid M (1943) Profundidade dos solos e vegetacao em campos cerrados do
562 Brasil meridional. *An Acad Bras Ciênc* 15:267–294
- 563 Reich PB, Walters MB, Ellsworth DS (1997) From tropics to tundra: Global convergence in plant
564 functioning. *Proc Natl Acad Sci* 94:13730–13734
- 565 Ribeiro JF, Walter BMT (1998) Fitofisionomias do Bioma Cerrado. In: Sano SM, Almeida SP (eds)
566 Cerrado: ambiente e flora, S.M. Sano & S.P. Almeida. Planaltina, pp 89–166
- 567 Riley D, Barber SA (1971) Effect of Ammonium and Nitrate Fertilization on Phosphorus Uptake as
568 Related to Root-Induced pH Changes at the Root-Soil Interface. *Soil Sci Soc Am J* 35:301.
569 <https://doi.org/10.2136/sssaj1971.03615995003500020035x>
- 570 Rizzini CT (1963) A flora do Cerrado: análise florística das savanas centrais. In: Simpósio sobre o
571 Cerrado. Editora da Universidade de São Paulo, São Paulo, pp 125–177
- 572 Rossatto DR, de Carvalho Ramos Silva L, Villalobos-Vega R, et al (2012) Depth of water uptake in
573 woody plants relates to groundwater level and vegetation structure along a topographic gradient
574 in a neotropical savanna. *Environ Exp Bot* 77:259–266.
575 <https://doi.org/10.1016/j.envexpbot.2011.11.025>
- 576 Ruggiero PGC, Batalha MA, Pivello VR, Meirelles ST (2002) Soil-vegetation relationships in cerrado
577 (Brazilian savanna) and semideciduous forest, Southeastern Brazil. *Plant Ecol* 160:1–16.
578 <https://doi.org/10.1023/A:1015819219386>

579 Scartazza A, Lauteri M, Guido MC, Brugnoli E (1998) Carbon isotope discrimination in leaf and stem
580 sugars, water-use efficiency and mesophyll conductance during different developmental stages
581 in rice subjected to drought. *Funct Plant Biol* 25:489–498

582 Silva LCR, Hoffmann WA, Rossatto DR, et al (2013) Can savannas become forests? A coupled analysis
583 of nutrient stocks and fire thresholds in central Brazil. *Plant Soil* 373:829–842.
584 <https://doi.org/10.1007/s11104-013-1822-x>

585 Simon MF, Grether R, de Queiroz LP, et al (2009) Recent assembly of the Cerrado, a neotropical plant
586 diversity hotspot, by in situ evolution of adaptations to fire. *Proc Natl Acad Sci* 106:20359–
587 20364. <https://doi.org/10.1073/pnas.0903410106>

588 Sprent JI (2009) Legume nodulation: a global perspective. Wiley-Blackwell, Chichester, UK

589 Sprent JI (2007) Evolving ideas of legume evolution and diversity: a taxonomic perspective on the
590 occurrence of nodulation. *New Phytol* 174:11–25

591 Sprent JI, Geoghegan IE, Whitty PW, James EK (1996) Natural abundance of ¹⁵N and ¹³C in nodulated
592 legumes and other plants in the cerrado and neighbouring regions of Brazil. *Oecologia* 105:440–
593 446. <https://doi.org/10.1007/BF00330006>

594 Ström L (1997) Root Exudation of Organic Acids: Importance to Nutrient Availability and the Calcifuge
595 and Calcicole Behaviour of Plants. *Oikos* 80:459–466. <https://doi.org/10.2307/3546618>

596 Sugihara S, Shibata M, Ze ADM, et al (2014) Effect of vegetation on soil C, N, P and other minerals in
597 Oxisols at the forest-savanna transition zone of central Africa. *Soil Sci Plant Nutr* 60:45–59.
598 <https://doi.org/10.1080/00380768.2013.866523>

599 Symonds MRE, Moussalli A (2011) A brief guide to model selection, multimodel inference and model
600 averaging in behavioural ecology using Akaike’s information criterion. *Behav Ecol Sociobiol*
601 65:13–21. <https://doi.org/10.1007/s00265-010-1037-6>

602 The Angiosperm Phylogeny Group (2016) An update of the Angiosperm Phylogeny Group classification
603 for the orders and families of flowering plants: APG IV. *Bot J Linn Soc* 181:1–20.
604 <https://doi.org/10.1111/boj.12385>

605 Thomas DS, Montagu KD, Conroy JP (2006) Leaf inorganic phosphorus as a potential indicator of
606 phosphorus status, photosynthesis and growth of *Eucalyptus grandis* seedlings. *For Ecol Manag*
607 223:267–274. <https://doi.org/10.1016/j.foreco.2005.11.006>

608 Townsend AR, Cleveland CC, Asner GP, Bustamante MMC (2007) Controls over foliar N:P ratios in
609 tropical rain forests. *Ecology* 88:107–118. [https://doi.org/10.1890/0012-9658\(2007\)88\[107:COFNRI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2007)88[107:COFNRI]2.0.CO;2)

611 Valiente-Banuet A, Verdú M (2007) Facilitation can increase the phylogenetic diversity of plant
612 communities. *Ecol Lett* 10:1029–1036. <https://doi.org/10.1111/j.1461-0248.2007.01100.x>

613 Werner C, Máguas C (2010) Carbon isotope discrimination as a tracer of functional traits in a
614 mediterranean macchia plant community. *Funct Plant Biol* 37:467–477.
615 <https://doi.org/10.1071/FP09081>

616 Westoby M (1998) A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant Soil* 199:213–227.
617 <https://doi.org/10.1023/A:1004327224729>

618 Wohlgemuth T, Gigon A (2003) Calcicole plant diversity in Switzerland may reflect a variety of habitat
619 templates. *Folia Geobot* 38:443–452. <https://doi.org/10.1007/BF02803251>

620 Wright IJ, Reich PB, Westoby M (2001) Strategy shifts in leaf physiology, structure and nutrient content
621 between species of high- and low-rainfall and high- and low-nutrient habitats. *Funct Ecol*
622 15:423–434. <https://doi.org/10.1046/j.0269-8463.2001.00542.x>

623 Wright IJ, Reich PB, Westoby M (2003) Least-Cost Input Mixtures of Water and Nitrogen for
624 Photosynthesis. *Am Nat* 161:98–111. <https://doi.org/10.1086/344920>
625

Table 1. Plant traits used to analyze functional groups of a Cerrado community at Paraopeba Reserve, Minas Gerais. (*) indicates a trait that was not directly used in the data analysis, but which is part of a rate. SLA = specific leaf area, Hmax = maximum height, Dmax = maximum diameter, C/N = carbon to nitrogen ratio of leaf.

Trait	Unit	Functional significance
1. SLA	mm ² .mg ⁻¹	Correlated with aspects of physiology related to the efficiency of resource uptake and use
2. Hmax*	m	Reflects competitive vigour and ability to persist after disturbances
3. Dmax*	cm	
4. Hmax/Dmax	-	Stem elongation considering vertical/horizontal growth
5. C/N	-	Indicates nitrogen limitation, proxy of nutritional status
6. N%	-	Indicates leaf nitrogen content, proxy of nutritional status
6. C%	-	Indicates leaf carbon content, proxy of polysaccharides and photoassimilates
6. Bark thickness	mm	Indicates if a species is adapted to poor soils and to high disturbance regimes: the thicker the bark, the poorer the soil and the higher disturbance resistance
7. Wood density	mg.mm ⁻³	Key trait for plant growth

Figure 1- Canonical RLQ ordination of 34 woody species and environmental variables; red dots are calcicole species (CAL), green dots are leguminous species (LEG), purple dots are savanna species (SAV). SLA – specific leaf area, elongation – maximum height to maximum diameter ratio, % nitrogen – leaf nitrogen content (%), % carbon – leaf carbon content (%).

631

	CEC	Aluminium saturation	Inorganic nitrogen
SLA			
Hmax/Dmax			
%N			
%C			

632

633 **Figure 2.** Results of the fourth-corner analysis. Adjusted significances of positive relations: between Cation
634 Exchange Capacity (CEC) and specific leaf area (SLA), P value adjusted = 0.044; between CEC and stem
635 elongation (Hmax/Dmax), P value adjusted = 0.044; between inorganic nitrogen and leaf nitrogen
636 percentage (%N), P value adjusted = 0.040; between inorganic nitrogen and leaf carbon percentage (%C),
637 P value adjusted = 0.044. Adjusted significances of negative relations: between aluminium saturation (m)
638 and specific leaf area (SLA), P value adjusted = 0.044; between aluminium saturation and Hmax/Dmax, P
639 value adjusted = 0.024.

640

Figure 3. The best GLM full models for Calcicole species (Ni=abundance) in relation to soil and canopy openness in the Cerrado of Paraopeba Reserve. Model: $\text{glm}(\text{CAL} \sim \text{Al saturation} + \text{Inorganic nitrogen} + \text{CEC} + \text{Canopy openness} + \text{pH}, \text{family}=\text{poisson})$, $\Delta \text{AIC} < 2$. Dots are 20-m x 20-m plots. Soil: Inc - Cambisol (Inceptisol); RLd – dystrophic Red Latosol; RLM – mesotrophic Red Latosol; RYL – Red-Yellow Latosol; YL – Yellow Latosol.

Figure 4. The best GLM full models for Leguminous species (Ni=abundance) in relation to soil and canopy openness in the Cerrado of Paraopeba Reserve. Model: $\text{glm}(\text{CAL} \sim \text{Al saturation} + \text{Inorganic nitrogen} + \text{CEC} + \text{Canopy openness} + \text{pH}, \text{family}=\text{poisson})$, Delta AIC < 2. Dots are 20-m x 20-m plots. Soil: Inc - Cambisol (Inceptisol); RLd – dystrophic Red Latosol; RLm – mesotrophic Red Latosol; RYL – Red-Yellow Latosol; YL – Yellow Latosol.

658
659 **Figure 5.** The best GLM full models for Savanna species (Ni=abundance) in relation to
660 soil and canopy openness in the Cerrado of Paraopeba Reserve. Model: glm (CAL ~ Al
661 saturation + Inorganic nitrogen + CEC + Canopy openness + pH, family=poisson), Delta
662 AIC < 2. Dots are 20-m x 20-m plots. Soil: Inc - Cambisol (Inceptisol); RLd – dystrophic
663 Red Latosol; RLM – mesotrophic Red Latosol; RYL – Red-Yellow Latosol; YL – Yellow
664 Latosol.
665

666 **Figure 6.** Box and whiskers plots of the GLMs for functional groups by functional traits.

667 Leaf nitrogen content (Leaf N%), Leaf carbon to nitrogen ratio (C/N) and Specific leaf

668 area (SLA) by groups. Different lowercases letters indicate significant differences

669 ($p < 0.005$).

670

671 **Figure 7.** Box and whiskers plots of the GLMs for functional groups by functional traits.
672 Wood density, bark thickness and ratio of maximum height to maximum diameter (i.e.,
673 stem elongation, Hmax/Dmax) by groups. Different lowercases letters indicate significant
674 differences ($p < 0.005$).
675

676
677 **Figure 8.** GLM of the best model for leaf C/N in response to functional traits (SLA,
678 Hmax/Dmax, Bark thickness, leaf $\delta^{13}\text{C}$ and leaf $\delta^{15}\text{N}$) of the 34 Cerrado plant species of
679 the three functional groups CAL and LEG and SAV. The best model is between leaf C/N
680 and leaf $\delta^{13}\text{C}$ (d13C) values ($p=0.00148$).

681

682
683 **Figure 9.** Model for the main gradients determining the distribution of the Calcicole
684 species (CAL); the Leguminous species (LEG), the Savanna species (SAV); Inorganic N
685 = inorganic nitrogen; C/N = leaf carbon-to-nitrogen ratio; pH = hydrogenionic potential;
686 CEC = cation exchange capacity. Arrows indicate the gradients where, at one extremity,
687 one group preferentially occurs, on the other extremity, two other groups preferentially
688 occur.