“Prevalence of of ABO and Rhesus (Rh) Blood Groups in Eastern UP Population”

Authors:

Pradeep Kumar

Vandana Rai

Affiliation:

Department of Biotechnology

VBS Purvanchal University

Jaunpur (UP) 222001

Corresponding Author:

Dr. Pradeep Kumar

Human Molecular Genetics Laboratory

Department of Biotechnology

VBS Purvanchal University

Jaunpur (U P)222001

India

Pradipk14@yahoo.co.in

Running Title:

Prevalence of ABO and Rh Blood Groups

Abstract:

Approximately 300 different types of blood groups are identified so far, the ABO and Rh antigens are still the clinically most significant and genetically most polymorphic of all human blood group systems to date. A total of 200 unrelated individuals from Uttar Pradesh were studied for the phenotype and allele frequency distribution of ABO and Rh (D) blood groups. In total 200 samples analyzed, phenotype B blood type has the highest frequency 36.5% (n=73), followed by O (34.5%; n=69), A (20.5%; n=41) and AB (8.5%; n=17). The O, A and B frequencies were 0.5849, 0.1571 and 0.2580 respectively. The overall phenotypic frequencies of ABO blood groups were B>O>A>AB. The variation in phenotypic frequencies between male and female might be due to small sample size of male sample. The allelic frequency of Rh-negative was 0.2.

Keywords:

ABO blood groups, Antigen, Allele frequency, Phenotype

Introduction:

Blood groups are antigenic determinants on the surface of red cells, platelets and granulocytes, but the use of the term is often restricted to antigens on red blood cells. Blood groups are defined by antibodies, usually alloantibodies produced by individuals who lack the corresponding antigen. Some blood group antibodies, such as anti-A and anti-B, are present in the plasma of a person, whose red cells lack the corresponding antigen, but most blood group antibodies are formed only in response to antigen positive red cells as the result of transfusion or pregnancy such as Rh (Daniels, 2001). Nearly 300 blood grouping antigens have been reported, the ABO and Rh are recognized as the major and clinically significant blood group antigens (Bauer, 1982). Rhesus blood group system was the 4th system discovered and yet it is second most important blood group from the point of view of transfusion (Molison, 1979).

The ABO locus is located on chromosome 9 at 9p34.1-q34.2 and encodes glycosyltransferases. And ABO gene has three alleles namely *IA*, *IB* and *i* determine blood groups. *IA* produces A antigen, *IB* produces B antigen whereas *i* produces neither. *IA* and *IB*

are mutant alleles and show codominance with each other but both are dominant over the wild type allele i . The concept of wild type allele is based on the fact that the allele more frequent in a population is a wild type allele (Gardner et al, 2001). Rh antigens are determined by three pairs of closely linked allelic genes located on chromosome 1.

The ABO and Rh blood groups have been studied extensively for their involvement in incompatibility selection and their distribution are studied in almost all the races and populations of the world like-Nigeria (Enosolease and Bazuaye, 2008), Kenya (Lyko et al., 1992), Palestine (Alishtayeh et al., 1988), Iraq (Mohamad and Jaff, 2010), Sudan (Hassan,2010); Pakistan (Hameed et al., 2002; Anees and Mirza,2005), Bangladesh (Talukdar and Das,2010), Saudi Arab (Adullah,2010), Jordan (Hanania et al,2007), Iran (Boskabady et al, 2005), Nepal (Pramanik and Pramanik,2000), and India (Chakraborty, 2010; Kumar et al.,2010; Rai and Kumar, 2010; Deepa et al, 2011;Kumar and Rai,2012) .

The genotype frequencies for a particular gene(s) in a population depend on the gene frequency and the proportions of different alleles of a gene in a Mendelian (panmictic) population are known as gene frequency. Estimates of gene's frequency provide very valuable information on the genetic similarity of different populations and to some extent on their ancestral genetic linkage, despite the cultural and religious differences of the two populations. Keeping this in view, the present study was designed to see frequency of ABO and Rh blood group antigens in UP population.

Materials and method:

The study was done in Uttar Pradesh (UP). Over three months period (December 2010 to February 2011), a total of 200 unrelated individuals of both genders from Jaunpur and Varanasi, Azamgarh and Mau districts of UP were selected. Blood samples were collected from each subject and relevant data were also collected after taking written consent. Confidentially, of the data were maintained. Each subject, who accepted to participate in the study, received two sheets, (consent form, and questionnaire). The first sheet was a declaration form for each participant that she/he understood the project/study well, the second sheet was, a questionnaire that included profile/demographic data of the participants. The blood samples were collected by finger prick with sterile lancet and cleaning the puncture site with 70% ethyl alcohol. A drop of monoclonal anti-A, anti-B, monoclonal/polyclonal anti-D (Span) was added to a drop of finger prick blood on clean

slide and mixed well. Results of agglutination were recorded immediately for ABO blood groups and after 2 minutes in Rh(D) (Bhasin and Chahal, 1996). The gene frequencies for this system were calculated according to the method of Mourant et al. (1976).

Results and Discussion:

In total 200 samples analyzed, phenotype B blood type has the highest frequency 36.5% (n=73, followed by O (34.5%; n=69), A (20.5%; n=41) and AB (8.5%; n=17) (Table 1). The O, A and B frequencies were 0.5849, 0.1571 and 0.2580 respectively (Table 1). The overall phenotypic frequencies of ABO blood groups were B>O>A>AB. Total numbers of samples were categorized by gender-wise, 68 samples were of females and 132 of males (Table 2). In female samples 20 individuals have O blood group, 17 individuals have A blood group, 25 individuals have B blood group and 6 individuals have AB group. In 132 male samples, O, A, B and AB blood groups were found in 49, 24, 48 and 11 individuals respectively. In female samples the phenotypic frequencies were B> O> A>AB, whereas in male samples overall phenotypic frequencies were O> B> A>AB (Table 2). The variation in phenotypic frequencies between male and female might be due to small sample size of male sample. Out of total 200 samples, 192 (96%) samples were Rh-positive and 8 (4 %) were Rh-negative. The allelic frequency of Rh-negative was 0.2 (Table 3).

In India ABO and Rh frequency is studied in other states very well-Karnataka (Prabhakar et al, 2005; Dore Raj and Reddy, 2010), Andhra Pradesh (Naidu et al,2002; Rao et al,2003; Reddy and Reddy,2003, 2005; Reddy and Sudha,2009), Maharastra (Mukherjee et al, 1997; Warghat et al, 2011), Dadar and Nagar Haveli (Meitei and Kshatriya,2010), Rajasthan (Thukral and Bhasin, 1990),Haryana (Kushwaha et al,1990), Punjab (Sidhu, 2003), Uttarakhand (Patni and Yadav, 2003; Guniyal, 2006; Pattanayak,2006), Jammu and Kashmir (Calcutti et al, 2003), Himanchal Pradesh (Mukhopadhyaya and Kshatriys, 2004), Assam (Chakraborty, 2010), Manipur (Devi and Singh, 2008), Orissa (Mohanty and Das, 2010) , Uttar Pradesh (Mandal ,1992 ; Ara et al 2008; Kumar et al.,2009a,b,c,2010; Rai and Kumar, 2010,2011, Rai,2011; Rai et al., 2009a,b,c, 2010,)), but the UP population is very less explored.

We compared our result with other studies carried out in different parts of the world like-Britain, Guinea, Nigeria, Yamen, Iran, Bahrain, Turkey, Palestine, Saudi Arabia, Pakistan, Nepal and Bangladesh (Mwangin et al,1999; Pramanik and pramanik,2000; Al-Arrayed et

al, 2001; Bashwari et al, 2001; Hameed et al,2002; Bahaj ,2003; Boskbady et al, 2005; Dilek et al, 2006; Skaik and El-Zyan,2006; Loua et al, 2007; Firkin et al, 2008; Talukdar and Das,2010). A comparison of frequency of blood group with this study to other is shown in the table 4. Except turkey, Palestine, Nepal and India (present study), frequency of O blood group is highest in Britain (47%), Guinea (48.9%), Nigeria (54.2%), Yamen (51.5%), Iran (34.7%), Bahrain (49.6%), Saudi Arabia (51%), Pakistan (49.6%) and Bangladesh (40.6%) and there is no marked difference in incidence of O blood group in these countries (table 4). Difference is marked in case of A blood group between Turkey and Bangladesh (43.8% vs 26.6%) and is highest phenotype in Turkey (43.8%), Palestine (40%) and Nepal (34%). In Britain B blood group is exceptionally low i.e. 8%, whereas it is highest in our study (36.5%). Marked difference of incidence of AB group is observed between Nigeria and Bangladesh (2.8% vs 9.6%). Rh-negative frequency is lowest in Palestine (2.7%) and highest in Iran (11.3%). Rh-negative frequency in present study is comparable with other countries-Guinea, Nepal and Bangladesh (between 3 to 4%) but is exceptionally high in Britain (17%).

Acknowledgement:

We are grateful to the individuals/subjects who participated in this study and without their cooperation this study could not be completed.

References:

- Abdullah S M . (2010).Frequency of ABO and Rh blood groups in the Jazan region of Saudi Arabia .Pak J Med Sci ,26(4): 818-821.
- Al-Arrayed S, Shome D K, Hafadh N, Amin S, Al Mukhareq H, Al Mulla M et al. (2001). ABO blood group and RhD phenotypes in Bahrain: Results of screening school children and blood donors. Baharain Med Bull, 75: 147-153.
- Alishtayeh MS, Hamlin AH, Fendy YR. (1988). Distribution of ABO blood groups and Rh factor in Palestinians living in the northern part of the West Bank. Najah Res J, 2(1): 35-41.
- Anees M and Mirza M S. (2005). Distribution of ABO and Rh Blood Group Alleles in Gujrat Region of Punjab. Pakistan. Proc. Pakistan Acad.. Sci. 4 2(4):233-238.
- Ara G, Siddique Yh, Beg T, Afzal M. (2008). Gene diversity among some muslim population of Western Uttar Pradesh. Anthropologist, 2008 ;10(1):5-9.

Bahaj AA. (2003). ABO and Rhesus blood groups distribution in Hadhramont population. Hadhramont for Studies and researches, 2003;4:52-58.

Bashwari LA, Al-Mulhim A A, Ahmad M S, Ahmed M A.(2001). Frequency of ABO blood groups in eastern region of Saudi Arabia. Saudi Med J, 22:1008-1012.

Bauer JD. (1982) Clinical laboratory methods. 9th.Edition. USA: Mosby Company MI, .p. 353–76.

Bhasin MK, Chahal SMS. (1996). A Laboratory Manual for Human Blood Analysis. Delhi: Kamla-Raj Enterprise.

Boskabady MH, Shademan A, Ghamami G, Mazloom R. (2005). Distribution of blood groups among population of the city of Mashhad (North East of Iran). Pak J Med Sci Q, 21(2):194–198.

Calcutti RA, Lone MK, basher SA, Shah A , Jan N (2003). Blood groups in the Kashmir valley. JK Science: Journal of Medical Education & Research, 5(3).

Chakraborty S.(2010). Genetic Analysis on Frequency of Alleles for Rh and ABO Blood Group Systems in the Barak Valley Populations of Assam. Not Sci Biol 2 (2) 2010, 31-34.

Daniels,G.(2009). The molecular genetics of blood group polymorphism.Hum Genet,126:729–742.

Deepa, Alwar V A, Rameshkumar K and Ross C .(2011). ABO blood groups and malaria related clinical outcome. *J Vector Borne Dis* 48: 7–11.

Devi KR , Singh TS (2008). Incidence of ABO and Rh(D) incompatibility among the Meities of Kwatha village Manipur, India. Anthropologist, 10(1):65-69.

Dilek I, Demir C, Bay A, Akdeniz H, Oner AF. (2006). ABO and Rh blood groups frequency in men and women living in eastern Turkey. International Journal of Hematology and Oncology, 16:23-26.

Dore Raj M L and Rajasekhara Reddy K .(2010).The Study of ABO and Rh (D) Blood Groups among the Vishwakarma Population of Mysore District in Karnataka, India. Anthropologist, 12(3): 227-228.

Enosolease ME, Bazuaye GN. (2008). Distribution of ABO and Rh-D blood groups in the Benin area of Niger-Delta: implication for regional blood transfusion. *Asian J Trans Sci*. 2008;2(1):3–5.

Firkin F, Chesterman C, Penington D, and Rush B (2008) De Gruchy's Clinical haematology in medical Practice. Blackwell Science Publisher, 5th Edition. pp 475.

Gardener, E. J., M. J. Simmons, and D. P. Snustad, (2001). Principles of genetics. 8th edition John Wiley and Sons (Asia) Pvt Ltd.

Guniyal M. (2006). The study of blood group A1A2BO and Rh among the Brahmins of Mussoorie, Uttaranchal. Anthropologist, 8(2): 53-54.

Hanania S, Hassawi D, Irshaid N. (2007). Allele frequency and molecular genotypes of ABO blood group system in Jordanian population. J Med Sci ,7(1):51-8.

Hameed A, Hussain W, Ahmad J, Rabbi F, Qureshi JA. (2002). Prevalence of phenotypes and genes of ABO and Rhesus (Rh) blood groups in Faisalabad, Pakistan. Pak J Biol Sci, 5:722-724.

Hassan F M .(2010). .Frequency of ABO, Subgroup ABO and Rh(D) Blood Groups in Major Sudanese Ethnic Groups. Pak J Med Res, 49: 21-24.

Kumar P, Yadav M, Rai V. (2008) ABO and Rh(D) blood groups among Yadav of district Jaunpur. Pb. Univ. Res. J (Sci.),58:79-81.

Kumar P, Singh VK, Rai V (2009a) Study of ABO and Rh(D) Blood Groups in Kshatriya (Rajput) of Jaunpur District, Uttar Pradesh. Anthropologist, 11(4): 303-304.

Kumar P, Maurya S, Rai V. (2009b) Distribution of ABO and Rh(D) Blood Groups among Koari (Backward Caste) Population of Jaunpur District. Anthropologist, 11(4), 309-310.

Kumar P, Maurya AK, Rai V. (2009c) ABO and Rh(D) blood groups among Maurya (Backward caste) population of Jaunpur district (UP), India. Asian J Microbiol. Biotech. Env. Sci., 11(4) : 723-726.

Kumar P, Saima, Rai V. (2010) Study of ABO and Rh(D) blood groups in Sunni Muslims of Jaunpur district. Anthropologist, 2010; 12(3) 225-226.

Kumar P, Rai V. (2012) Red Blood Cell Antigen Distribution in Scheduled Caste Population of Uttar Pradesh. Asian J Microbiol. Biotech. Env. Sci. 14(2):217-222.

Kushwaha KPS, Gaur JSR, Sangwan SK, Yadav AS.(1990). ABO and Rh (D) blood groups among 19 caste populations of Haryana. Bionature, 10: 73-75.

Loua A, Lamah MR, Haba NY et al (2007). Frequency of blood groups ABO and Rhesus D in the Guinean population. Transfus Clin Biol. 14(5):435-9.

Lyko J, Gaertner H, Kaviti JN, Karithi MW, Akoto B. (1992). The blood groups antigens ABO and Rh in Kenyans. *Hamdard Medicus*, 35(3):59-67.

Mandal PK. (1992). The ABO and Rh(D) blood groups among five Caste population of Uttar Pradesh. *J Hum Ecol*, 3: 57-58.

Meitei K S and Kshatriya G K. (2010). The Study of Rhesus System among the Rajputs and Varlis of Dadra and Nagar Haveli. *Anthropologist*, 12(1): 63-64.

Mohamad S Jaff (2010). ABO and rhesus blood group distribution in Kurds. *Journal of Blood Medicine*, 1: 143-146.

Mohanty R, Das PK (2010). A search for operation of natural selection in ABO blood groups: Evidence from four ethnic groups of Orissa. *Anthropologist*, 12(1):1-11.

Molison PL. (1979). *Blood transfusion in clinical medicine* 6th Edition. Oxford UK: Blackwell Scientific Publication, p. 239-66.

Mourant AE, Kopec ADA, Domaniewska-Sobezek K. (1976). *The ABO Blood Groups-Comprehensive Tables and Maps of World Distribution*. Oxford, London: Blackwell Scientific Publication.

Mukhopadhyay R, Kshatriya GK. (2004). Distribution of blood groups among Brahmins and Rajputs of Himanchal Pradesh. *Anthropologist*, 6(4): 293-294.

Mwangi J. (1999). Blood group distribution in an urban population of patient targeted blood donors. *East Afr Med J*. 76(11):615-8.

Patni S, Yadav A. (2003). Blood groups among the Bhoksa of Vikasnagar block of Dehradun, Uttaranchal. *Anthropologist*, (2):137-138

Pattanayak I. (2006). Distribution of A1A2BO and Rh blood group among the Rajputs of Uttaranchal. *Anthropologist*, 8 (2): 139-140.

Prabhakar SCJ, Gangadhar MR, Reddy KR. (2005). ABO and Rh (D) blood groups among the Vishwakarmas of Mysore district, Karnataka. *Anthropologist*, 7(1):71-72.

Pramanik T, Pramanik S. (2000) Distribution of ABO and Rh blood groups in Nepalese medical students: a report. *East Mediterr Health J*, Jan(1):156-8.

Rai V, Patel RP, Kumar P. (2009a) Study of ABO and Rh(D) blood groups in Scheduled Caste of Jaunpur district. *Anthropologist*, 11(2): 151-152.

Rai V, Maurya SK, Kumar P. (2009b) ABO and Rh (D) blood groups among Khatik (Scheduled caste) population of Jaunpur district (UP). *PURJ (Science)*. 59:107-109.

Rai V, Verma AK, Kumar P. (2009c) A study of ABO and Rh(D) blood groups among Kurmi (Backward caste) of Jaunpur district. *Anthropologist*, 11(4), 305-306.

Rai V, Kumar P.(2010) The incidence of ABO blood group in Muslim population of Uttar Pradesh,India. *J Appl Biosci.*, 36(2):191-195.

Rai V, Jahan S, Kumar P. (2010) Demographic study of ABO and Rh (D) Blood Groups among Muslim Population of Jaunpur District (U P). *Asian Jr. Microbiol. Biotech. Env. Sc.*,12(2):429-432.

Rai V. Genetic Analysis of ABO and Rh Blood Groups in Brahmin Population of Uttar Pradesh, India”. *Nature Precedings*, 1-1.

Rai V, Kumar P. (2011) Genetic Analysis of ABO and Rh Blood Groups in Backward Caste Population of Uttar Pradesh, India. *Not Sci Biol*, 2011, 3(3):07-14.

Reddy BKC, Reddy CS. (2005). ABO and Rh(D) blood group distribution among Voddas, a backward caste population of Andhra Pradesh. *Anthropologist*, 7(3): 235-236.

Sidhu S. (2003). Distribution of the ABO blood groups and Rh(D) factors among the scheduled caste population of Punjab. *Anthropologist*, 5 (3): 203-204.

Skaik YA, EL-Zyan N. (2006). Spectrum of ABO and Rh (D) blood groups amongst the Palestinian students at Al-Azhar University-Gaza. *Pak J Med Sci*, 22:333-335.

Talukder S I and Das R K. (2010). Distribution of ABO and Rh Blood Groups among Blood Donors of Dinajpur District of Bangladesh. *Dinajpur Med Col J* , 3 (2):55-58.

Thukral R., Bhasin, MK (1990). Blood groups of Meghwal and Salvi, the Scheduled Caste of Rajasthan. *Hum Eco*, 11: 183-184.

Tewari SC and Bhasin MK (1968). The blood groups of the Brahmins and the Rajputs of Garhwal. *Hum Biol*, 40:386-395.

Warghat NE, Sharma NR, Baig MM (2001). ABO and Rh Blood Group distribution among Kunbis (Maratha) population of Amravati District, Maharashtra-India *Asiatic Journal of Biotechnology Resources*, 2(04)479-483

Table 1. Distribution of the ABO blood group and their allele frequencies among eastern UP Population

Phenotype	Observed Number	Percentage	Expected Number	Allele frequency
O	69	34.5%	68.4	O= 0.5849
A	41	20.5%	41.7	A=0.1571
B	73	36.5%	73.7	B=0.2580
AB	17	8.5%	16.2	

Table 2. Total number of samples classified according to gender

Sections	No. of O phenotype	No. of A phenotype	No. of B phenotype	No. of AB phenotype	Total
Females	20	17	25	6	68
Males	49	24	48	11	132
Total	69	41	73	17	200

Table 3. Rh blood group among eastern UP population

Phenotypes	Observed		Allele frequencies
	Number	Percentile	
Rh(Anti D)+	192	96%	0.8
Rh(Anti D)-	8	4%	0.2
Total	200		

Table 4. Comparison of percentage frequencies of ABO and Rh blood groups in different studies carried out in different countries

Country	Study	O	A	B	AB	Rh-positive	Rh-negative
Britain	Firkin et al,2008	47	42	8	3	83	17
Guinea	Loua et al,2007	48.9	22.5	23.7	4.7	95.9	4
Nigeria	Mwahgin et al,1999	54.2	21.6	21.4	2.8	95.2	4.8
Yamen	Bahaj,2003	51.5	29.6	16.5	2.3	92.9	7.1
Iran	Boskabady et al,2005	34.7	33.1	23.3	8.9	88.7	11.3
Bahrain	Al-Arrayed et al,2001	49.6	21.5	24.4	4.5	94.5	4.5
Turkey	Dilek et al,2006	30.8	43.8	16.2	9.2	86	14
Palestine	Skaik and El-Zyan,2006	32	40	22	6	97.3	2.7
Saudi	Bashwari et al,2001	51	26	18	4	92	8

Arabia							
Pakistan	Hameed et al,2002	49.6	23.8	38	10	89.1	10.9
Nepal	Pramanik and Pramanik,2000	32.5	34	29	4	96.7	3.3
Bangladesh	Talukdar and Das,2010	40.6	26.6	23.2	9.6	96.8	3.2
India	Present study	34.5	20.5	36.5	8.5	96	4