1 **Perturbation of GABA Biosynthesis Links Cell Cycle to** 2 **Control *Arabidopsis thaliana* Leaf Development**

3 **Yaxin Gong, Han Yue, Yu Xiang, Guanghui Yu***

4 Hubei Provincial Key Laboratory for Protection and Application of Special Plants in
5 Wuling Area of China, Engineering Research Centre for the Protection and Utilization
6 of Bioresource in Ethnic Area of Southern China, College of Life Sciences,
7 South-Central University for Nationalities, Wuhan 430074, China

8

9 *** Correspondence:**

10 Guanghui Yu

11 yuguanghui@scuec.edu.com

12

13 **Keywords:** *Arabidopsis thaliana*; CDKA;1; endoreplication; GABA; Reactive
14 Oxygen Species

15 **Running title:** Endoreplication response to GABA deficit

16 **Abstract**

17 To investigate the molecular mechanism underlying increasing leaf area in
18 γ -Aminobutyric acid (GABA) biosynthetic mutants, the first pair of true leaves of
19 GABA biosynthetic mutants was measured. The results showed that the leaf blade
20 area in GABA biosynthetic mutants was larger than that of the wild type to different
21 extents, and the area of the leaf epidermal cells in mutants was larger than that of the
22 wild type. DNA polyploid analysis showed that polyploid cells in GABA biosynthetic
23 mutants were appearing earlier and more abundant than in the wild type. To check the
24 correlation between cell size and endoreplication, the expression of factors involving
25 endocycles, including D-type cyclin gene (*CYCD3;1*, *CYCD3;2*, *CYCD3;3*, and
26 *CYCD4;1*) and kinase *CKDA;1*, were analysed by qRT-PCR. The results showed that
27 *CKDA;1* in GABA biosynthetic mutants was downregulated, and four types of
28 *CYCDs* showed different expression patterns in different GABA biosynthetic mutants.
29 Inconsistent with this result, for *CCS52A* (*CELL CYCLE SWITCH 52A*) (controlling
30 the endocycle entry) in *gad2* and *gad1/gad2* mutants, the expression of *CCS52A2* was
31 significantly higher than that in the wild type. The expression of *SIM* (*SIAMESE*) and
32 *SMR* (*SIAMESE-RELATED*), which inhibit kinase activity, were also upregulated
33 compared with the control. To further study the possible potential relationship
34 between GABA metabolism and endoreplication, we analysed the reactive oxygen
35 species (ROS) levels in guard cells using ROS fluorescent probes. ROS levels were
36 significantly higher in GABA biosynthetic mutants than the control. All results
37 indicated that cyclin, the cyclin-dependent kinase, and its inhibitory protein were

Endoreplication response to GABA deficit

38 coordinated to participate in endoreplication control at the transcription level in the
39 leaves of GABA biosynthetic mutant *Arabidopsis*.

40

Endoreplication response to GABA deficit

41 Contribution to the field statement

42 γ -Aminobutyric acid (GABA) metabolic pathway plays a dual role in plant
43 development. This research investigated the perturbation of GABA biosynthesis on
44 *Arabidopsis* leave endoreplication for the first time. In the GABA biosynthetic
45 mutants, many genes, participating in cell division regulation, are coordinately
46 transcriptionally expressed to trigger the onset and maintenance of endoreplication,
47 and this led to the cell expansion and the increase leaf blade area. However, this
48 initiation of endoreplication links with the decrease of endogenous GABA level and
49 the increase Reactive oxygen species (ROS). This may be a compensation mechanism
50 to adapt to abnormal GABA level in plant leaf development. Present evidence
51 provided hypothesized that the normal GABA level in plant leaf development plays a
52 brake to inhibit the immature cell expansion and differentiation, and this negative
53 regulation functions a guarantee mechanism to watchdog the normal leaf development.
54 In all, this contribution provides an updated perspective on the role of GABA in plant
55 development.

56

57 1 Introduction

58 γ -Aminobutyric acid (GABA) is a four-C, non-protein component amino acid
59 commonly found in organisms and is prevalent in bacteria, plants, and vertebrates
60 (Seifikalhor et al., 2019). The precursor of GABA synthesis is L-glutamic acid (Glu),
61 which is catalysed by glutamate decarboxylase (GAD) in cytoplasm. The *Arabidopsis*
62 genome has five genes encoding GAD named GAD1–5. The oxidative metabolism of
63 GABA occurs in mitochondria and entry is mediated by a GABA permease (Michaeli
64 et al., 2011). In mitochondria, using α -ketoglutarate or pyruvate as an amino acceptor,
65 GABA is catalysed by GABA transaminase (GABA-T) to produce glutamic acid (Glu)
66 or Alanine (Ala) and succinic semialdehyde (SSA), respectively. SSA is oxidized by
67 SSA dehydrogenase (SSADH) to succinic acid (SucA), and then SucA enters the
68 tricarboxylic acid (TCA) cycle for further metabolism. This metabolic pathway is
69 referred to as the GABA shunt. Under hypoxic or high-light conditions, SSA can be
70 reduced to gamma-hydroxybutyrate (GHB) by SSA reductase (SSR, also known as
71 GHB dehydrogenase) in the cytoplasm, mitochondria, and chloroplasts (Allan et al.,
72 2008). Previous studies have reported that unique cytosolic and plastid glyoxylate
73 reductase isoforms in *Arabidopsis* are known as GLYR1 and atGLYR2, respectively,
74 and they catalyse the conversion of SSA to GHB and glyoxylic acid to glycolic acid
75 via an NADPH-dependent reaction (Brikis et al., 2017). The balance of the redox state
76 was maintained by the accumulation of GHB and the reduction of SSA via the GABA
77 shunt.

78 GABA biosynthesis and accumulation from the glutamic acid pathway and its
79 metabolism are at the junction of N and C metabolism, providing a useful metabolic
80 substrate for the TCA cycle, electron transport chain, and C skeleton, which are

Endoreplication response to GABA deficit

involved in the balancing of C:N metabolism (Jacoby et al., 2011). It is generally believed that GABA metabolism is considered to be involved in metabolic signalling, which plays a dual role in plant development, including both metabolic and signal regulation (Häusler et al., 2014; Ramesh et al., 2017; Podlešáková et al., 2019). In addition, aldehyde chemical groups (i.e. H-C=O) produced by GABA biosynthesis and metabolic pathways in plants have high molecular activity, and aldehydes accumulated under stressed conditions are highly toxic and can react with DNA, lipids of oxidative membranes, and modified proteins, or affect the transcription of stress-related genes, leading to cellular and ontogenetic problems in plants. Previous studies have reported that GABA metabolism regulates leaf pattern morphogenesis. The SSADH gene is involved in the formation of the paraxial–abaxial (upper–lower) leaf model in *Arabidopsis thaliana* (Toyokura et al., 2011; 2012). Mutation of *enfl* (*enlarged fil expression domain1*) alters the expression pattern of the *FIL* (*YABBY1*, *FILAMENTOUS FLOWER*) gene, which is characteristic of meristem and organs in *A. thaliana* (Sawa et al., 1999), on the abaxial surface of leaf primordium (Sawa et al., 1999; Siegfried et al., 1999). However, here is a dearth of research on the role of GABA biosynthesis in the leaf development of *A. thaliana*. In the present study, GABA biosynthetic mutants, *gad1*, *gad2*, and *gad1/gad2* were examined to explore the molecular mechanism underlying the GABA negative feedback that regulates leaf cell endoreplication during leaf development. Our findings will provide evidence for further understanding the role of GABA in plant development.

2 Materials and methods

2.1 Experimental Materials

A. thaliana Col wild-type seeds, Col ecotype *gad1* mutant, *gad2* mutant, and *gad1/gad2* double mutant were provided by Prof. Barry Shelp (University of Guelph, Canada). The seeds of the wild type and mutants were sterilized for ~2 h in a sealed container with chlorine gas, and then inoculated on MS solid medium and synchronized for 3 d at 4 °C. After synchronization, the plate was taken out and placed in greenhouse under a photoperiod of 16/8 h light/dark and incubated at 22 °C.

2.2 Measurement of leaf blade area

From the 4th day after *A. thaliana* seedlings being transferred to the greenhouse, their growth condition was photographed every 24 h with a Brinno time-lapse camera (TLC100) at a fixed distance to record the leaf blade area.

2.3 Microscopic observation and measurement of cell area

From the 7th day after *A. thaliana* seedlings being transferred to the greenhouse, the first pair of true leaves was collected daily, the leaf abaxial epidermis located 25% and 75% from the distance between the tip and the base of the leaf blade was photocopied with nail polish, and then photographed using an Olympus DP80 microscope, and the area was calculated with the software that came with the microscope. Under the microscope, a certain area of the epidermis was confined to count the number of cells, and then the average cell area was calculated.

2.4 DNA ploidy analysis

Endoreplication response to GABA deficit

Approximately two or three leaves at the young stage or one or two leaves at the middle and late leaf developmental stages were chopped with a razor blade in nuclei extraction buffer (CyStain® UV Precise P, Sysmex Partec), and then transferred to the staining buffer (CyStain® UV Precise P, Sysmex Partec) according to the manufacturer's instructions. The ploidy level of DNA in leaf cells was determined using a CyFlow Ploidy Analyser (Sysmex Partec).

2.5 qRT-PCR analysis

Total RNA from *A. thaliana* leaves was extracted by AxyPrep Multisource Total RNA Miniprep Kit (Axygen Science, Inc). Reverse transcription was performed by Goldenstar RT6 Gene Synthesis Kit (Tingke Biotechnology Co., Ltd.) with a reverse transcription reaction system of 20 µL. qRT-PCR analysis was performed by MyGo Pro qPCR System (IT-IS Life Science Ltd.). The gene *isoprenyl diphosphate delta isomerase II* (*IPP2*, AT3G02780) was selected as the internal reference gene according to previous research (Fung-Uceda et al., 2018). The primers of the genes involved in cell cycle regulation, endocycle initiation, progression, and exit are listed in the supporting materials (Table S1).

2.6 In vivo reactive oxygen species imaging

For GC reactive oxygen species (ROS) staining, H2DCF-DA stock solution (10 mmol/L stock in Dimethyl Sulfoxide) was diluted in deionized water to yield a final concentration of 6.25 µmol/L with a final dimethyl sulfoxide concentration of 0.0125% (v/v) (Watkins et al., 2017). Epidermal strips were peeled with adhesive tape and directly stained for 30 min in the above solution. After rinsing with deionized water for 5 min, images were taken by Olympus DP80 fluorescent microscopy. The intensity of coloration was quantified using ImageJ software (National Institutes of Health, USA).

2.7 Statistical analyses

Data are expressed as averages ± standard error (SE). Experiments were conducted with two or three independent replicates. One-way ANOVA was employed using SPSS 20.0.

3 Results

3.1 The leaf blade area of the GABA biosynthetic mutant was larger than that of the wild type

At the early stage of leaf development (4th day after stratification), there was no significant difference in the area of the first pair of true leaves between the wild-type and GABA biosynthetic mutants (Figure 1). As time progressed, the area of the mutant leaves increased more rapidly than that of the wild type. Around the 8th day, the leaf area of the first pair of true leaves in the wild type was about 1.8 mm², and the first pair of true leaves in *gad1* and *gad2* mutants was about 2.4 mm², at 1.3-fold that of the wild type, and that of the *gad1/gad2* mutants was about 3.2 mm² with 1.8-fold that of the wild type (Figure 1). The leaf area between the GABA biosynthetic mutants and wild type was significantly different ($P < 0.05$).

Endoreplication response to GABA deficit

3.2 Cell size in GABA biosynthetic mutants was larger than those of the wild type

To determine the reasons for the leaf area in GABA biosynthetic mutants being larger than that in the wild type, we compared the area of leaf epidermal cells. The average cell area in the epidermis of the first pair of true leaves of the wild type was about 2700 μm^2 at the 8th day after stratification (Figure 2). The average cell area of the epidermis of the first pair of leaves of *gad1* and *gad2* mutants was ~3200 μm^2 , 1.2-fold that of the wild type ($P < 0.05$). The average cell area of the epidermis of the first pair of true leaves in *gad1/gad2* mutants was ~3800 μm^2 , and was 1.4-fold that of the wild type with significant differences ($P < 0.05$) (Figure 2).

3.3 The polyploidy in GABA biosynthetic mutant leaf occurred earlier and higher than that of the wild type

The correlation between cell size and DNA ploidy level (Matsunaga et al., 2013) prompted us to explore whether the cell size of GABA biosynthetic mutants is related to DNA polyploidy level. On the 7th day of leaf development after stratification, the cells in the first pair of true leaves in wild-type *A. thaliana* were mostly diploid and tetraploid, indicating that leaf cells mainly divided at this stage (Figure 3A). However, leaf cells in *gad1*, *gad2*, and *gad1/gad2* mutants appeared to be in different proportions of 8-ploid cells except the diploid and tetraploid cells (Figure 3B). The appearance of 8-ploid cells in leaves is a marker of endoreplication. On the 8th day of leaf development, 8-ploid cells were not found in wild-type *A. thaliana*, and the proportion of octoploid cells in three kind of GABA-biosynthetic mutants increased further (Figure 3B).

To further confirm the relationship between cell enlargement and cell ploidy level, we compared the cell ploidy level in the middle and late leaf development stages (21st day after stratification). The proportion of octoploid cells in the wild type was about 18% (Figure 3C). However, the proportion of the octoploid cells in *gad1*, *gad2*, and *gad1/gad2* mutants was about 30–40%, which was significantly higher than that in the wild type (Figure 3C).

Overall, at the early stage of leaf development (7–8th day after stratification), the 8-ploid cells were only detected in the GABA biosynthetic mutants (8-ploid cells in *gad1* and *gad2* mutants was 3–5%, and that in *gad1/gad2* double mutants was about 5–9%) (Figure 3D). At the late stage of leaf development (21st day after stratification), the proportion of 8-ploid cells in wild-type leaves was also detected; however, its level was significantly lower than that in *gad* mutants (Figure 3D).

3.4 Type-D cyclin genes involved in endocycle regulation in GABA biosynthetic mutants

Considering that endoreplication is a special form of cell division (De Veylder et al., 2011) and is the reason for cell polyploidy (Matsunaga et al., 2013), type-D relative gene participation in cell cycle regulation was analysed by qRT-PCR.

CYCD3;1 (At4g34160) is a key component to balance cell proliferation/division and endoreplication (Dewitte et al., 2007). Downregulation of *CYCD3;1* can lead to

Endoreplication response to GABA deficit

endoreplication, and its overexpression can cause excessive cell proliferation and inhibit cell differentiation (Dewitte et al., 2003). qRT-PCR confirmed that the expression of *CYCD3;1* was significantly downregulated in GABA biosynthetic mutants (Figure 4). In *gad1* and *gad2* single mutants, the relative expression of *CYCD3;1* was about 0.2-fold that of the wild type ($P < 0.01$), and in *gad1/gad2* double mutants, the relative expression of *CYCD3;1* was about 0.1-fold that of the wild type ($P < 0.01$) (Figure 4), indicating that blocking GABA synthesis inhibited the expression of *CYCD3;1*.

CYCD3;2 (At5g67260) and *CYCD3;3* (At3g50070) participate in the symmetrical division of guard mother cells and guard cells (GCs) in the late stomatal lineage development (Yang et al., 2014). In the *gad1* mutant, the expression of *CYCD3;2* was significantly downregulated ($P < 0.05$); in the *gad2* and *gad1/gad2* mutants, the relative expression of *CYCD3;2* was about 0.6-fold that of the wild type with significant differences ($P < 0.01$) (Figure 4). Similarly, the expression of *CYCD3;3* in the *gad1* mutant was about 0.3-fold that of the wild type ($P < 0.01$), and in the *gad2* and *gad1/gad2* mutants, the expression of *CYCD3;3* was 0.7 and 0.5-fold that of the wild type, respectively ($P < 0.05$) (Figure 4). *CYCD4;1* (At5g65420) could activate the cell cycle of root apical meristem (Masubelele et al., 2005), and the expression of *CYCD4;1* in the *gad2* mutant was significantly downregulated ($P < 0.05$), and in the *gad1* and *gad1/gad2* mutants the relative expression of *CYCD4;1* decreased to become the most significantly different to that of the wild type ($P < 0.01$) (Figure 4).

3.5 The expression of *CDKA;1* in GABA biosynthetic mutants was significantly downregulated

CDKA;1 (At3g48750) is one of the core components of cell cycle regulation, mainly controlling the transition of the G1/S and G2/M mitotic phase (Nowack et al., 2012). Inhibition of its activity can block mitosis to initiate leaf cell endoreplication (Verkest et al., 2005). The expression of *CDKA;1* in GABA biosynthetic mutants was significantly lower than that in the wild type ($P < 0.05$). These results indicated that blocking GABA synthesis is linked with the decreased expression of *CDKA;1* (Figure 5).

The transcription factor *E2Fa* (At2g36010) stimulates cell proliferation and delayed differentiation (Boudolf et al., 2004). In the *gad1* and *gad2* mutants, the expression of *E2Fa* was significantly downregulated ($P < 0.05$). In *gad1/gad2*, the level of *E2Fa* was 0.2-fold that in the wild type ($P < 0.01$) (Figure 5).

3.6 Different expression patterns of *CCS52A2* and *CDC6* were observed in GABA synthetic mutants

CCS52A2 (*CELL CYCLE SWITCH52*, At4g11920) encodes a ubiquitin ligase regulating cell cycle division phase (M) and a substrate-specific activator of anaphase promotion complex/cyclosome (Fülöp et al., 2005). Inhibition of kinase activity after *CCS52A2* expression and enhancement of its activity are prerequisites for endoreplication (Heyman et al., 2017; Umeda et al., 2019). Mutations in the *CCS52* gene resulted in delayed endoreplication, and its overexpression resulted in increased DNA ploidy levels (Heyman et al., 2017). In the *gad1* mutant, the relative expression of *CCS52A2* was not different from that of the wild type; however, in the *gad2* and

Endoreplication response to GABA deficit

250 *gad1/gad2* mutants, the expression of *CCS52A2* was significantly higher than that of
251 the wild type ($P < 0.05$) (Figure 6).

252 *CDC6* (At2g29680) encodes a homolog of cell division regulatory protein 6
253 (Castellano et al., 2001), which is a license gene for DNA replication (Fung-Uceda et
254 al., 2018). *CDC6* participates in the initiation of DNA replication and is an important
255 factor for maintaining endoreplication. The ectopic expression of *CDC6* can increase
256 the ploidy level caused by endoreplication (Castellano et al., 2001; 2004). In the *gad1*
257 mutant, the relative expression of *CDC6* was 1.7-fold as much as that in the wild type
258 ($P < 0.05$) (Figure 6). The expression of *CDC6* in the *gad2* mutant was about 3-fold
259 that in the wild type ($P < 0.01$), and the expression of *CDC6* in the *gad1/gad2* double
260 mutant was about 8-fold that in the wild type ($P < 0.01$) (Figure 6).

261 3.7 The expression of *SIAMESE* and *SIAMESE-RELATED* in GABA 262 biosynthetic mutants was significantly up-regulated

263 *SIAMESE* (*SIM*, At5g04470) is an inhibitory protein of cyclin-dependent kinase and
264 a regulator of mitotic inhibition and endoreplication (Churchman et al., 2006).
265 Overexpression of *SIM* results in dwarfing of plants, serrated leaves, and cells with
266 higher nuclear DNA content (Churchman et al., 2006). The relative expression of the
267 *SIM* in *gad1* single mutant is about twice that of the wild type ($P < 0.05$), in the *gad2*
268 single mutant is about 4-fold that of the wild type ($P < 0.05$), and in the *gad1/gad2*
269 double mutant is about 7-fold that of the wild type ($P < 0.01$) (Figure 7).

270 Most components of the *SIAMESE-RELATED* (*SMR*) gene family function in mitosis
271 inhibition and endocycle promotion (Yi et al., 2014; Kumar et al., 2015; Dubois et al.,
272 2018). In the chosen genes, the expression of *SMR1* (At3g10525) in *gad1* and *gad2*
273 mutants was not different to that of the wild type. In the *gad1/gad2* mutants, its
274 expression is 1.5-fold that of the wild type ($P < 0.05$) (Figure 7). However, the
275 expression of other components, *SMR2* (At1g08180), *SMR5* (At1g07500), and *SMR8*,
276 was markedly different to that of the wild type (Figure 7). The expression of *SMR2* in
277 *gad1*, *gad2*, and *gad1/2* reached 5-, 6-, and 11-fold of that in the wild type,
278 respectively ($P < 0.01$). Similarly, the expression of *SMR5* in *gad1*, *gad2*, and *gad1/2*
279 reached 3-, 5-, and 17-fold that in the wild type ($P < 0.01$), respectively, and
280 the expression of *SMR8* in *gad1*, *gad2*, and *gad1/2* reached 4-, 4-, and
281 6-fold that in the wild type ($P < 0.01$), respectively.

282 3.8 The level of ROS in leaves of GABA biosynthetic mutant was higher than 283 that of the wild type

284 The fact that redox regulates cell proliferation and the cell cycle (Schippers et al.,
285 2016), and GABA could scavenge ROS (Liu et al., 2011; Seifikalhor et al., 2019), led
286 us to postulate that ROS levels in GABA biosynthetic mutants could be increased.
287 Imaging of H2DCFDA (DCFH-DA; 2', 7'-Dichlorodihydrofluorescein diacetate) with
288 the fluorescent probe of ROS showed that the average ROS level intensity in GCs of
289 *gad* mutants was higher than that in the wild type ($P < 0.05$) (Figure 8). These results
290 suggest that GABA synthetic mutants are indeed correlated with ROS accumulation.

291 4 Discussion

Endoreplication response to GABA deficit

The GABA metabolic pathway plays a dual role during plant development, including metabolic and signal regulation (Bown & Shelp, 2016; Seifikalhor et al., 2019). Abnormal GABA biosynthesis and metabolism have important effects on plant development (Baum et al., 1996; Palanivelu et al., 2003; Renault et al., 2013). The expression of the *Petunia* GAD gene, which lacked the calmodulin-binding domain in transgenic tobacco, resulted in abnormal plant development, which was shorter and more branched than that of normal plants (Baum et al., 1996). In the Arabidopsis *GABA-T* (*pop2-1*) mutant, the growth of pollen tubes could not accurately target the ovule sac in the pistils (Palanivelu et al., 2003). The molecular mechanism was related to the destruction of the gradient distribution of GABA in the stigma of the pistils (Palanivelu et al., 2003). Furthermore, the growth retardation in hypocotyl epidermal cells and root cortex cells is related to the limitation of cell elongation (Renault et al., 2011), and inadequate expression of *GABA-T* may lead to developmental defects in roots and hypocotyls and composition change in cell walls (Renault et al., 2013).

In the present study, the molecular mechanism underlying leaf-area enlargement in GABA biosynthetic mutants was investigated. The growth of plant leaves is largely limited by the development of the epidermis. Polyploidy in pavement cells is strongly correlated with cell size ($P < 0.01$) (Melaragno et al., 1993). We confirmed that the larger leaf area and cell size in GABA biosynthetic mutants (Figures 1, 2) was related to much higher percentages of polyploid cells in GABA biosynthetic mutant leaves, which occurred earlier and in higher abundance than that of the wild-type leaf cells (Figure 3).

Endoreplication is a special kind of cell division (De Veylder et al., 2011), and the essence of endoreplication is to maintain CDK activity below the threshold that triggers mitosis (De Veylder et al., 2011). During leaf development of *A. thaliana*, the decrease in transcription levels of mitotic CDK and cyclin genes resulted in endoreplication (Beemster et al., 2006). Consistent with these results, the qRT-PCR analysis confirmed that the type-D cyclin genes (*CYCD3;1*, *CYCD3;2*, *CYCD3;3*, and *CYCD4;1*) were differently downregulated in GABA biosynthetic mutants, and the expression of *CDKA;1* exhibited a decreasing expression trend (Figures 4, 5).

Other factors interrelated with endoreplication regulation, from endoreplication initiation, progression, and maintenance, and exit (Breuer et al., 2014), synergistically regulate the occurrence of endoreplication in the leaves of GABA biosynthetic mutants. *CCS52A* (*CELL CYCLE SWITCH 52A*) plays an important role in cell cycle exit and endoreplication entry (Lammens et al., 2009; Vlieghe et al., 2005). The concentration of *CCS52A* in mitotic cells remains below a critical threshold to prevent immaturely initiating endoreplication (Lammens et al., 2009; Vlieghe et al., 2005). In the *gad2* and *gad1/gad2* mutants, the relative expression of *CCS52A2* was significantly higher than that of the wild type (Figure 6). The expression of cyclin-dependent kinase inhibitor *SIM* (*SIAMESE*) and its *SMR* (*SIAMESE-RELATED*) was significantly higher than that of the control (Figure 7). These results demonstrated that the endoreplication regulators, including cyclin, cyclin-dependent kinase, and kinase inhibitor, at the transcriptional level, all harmoniously contributed to the regulation of endoreplication in mutants.

It is worth noting that *SMR5*, a member of *SMR* (*SIAMESE-RELATED*), is an ROS-induced gene, which is more highly expressed in GABA biosynthetic mutants

Endoreplication response to GABA deficit

(3–17-fold that of the control) (Figure 7). ROS is an important signalling molecule regulating leaf development and is involved in triggering endoreplication (Schippers et al., 2016). Previous evidence proved that exogenous GABA could scavenge ROS (Liu et al., 2011), and the GABA level in mutants was much lower than that in the wild type (Figure S1); thus, we speculated that the perturbation of GABA biosynthesis may be intrinsically linked with the ROS level. Detection using an ROS fluorescence probe confirmed that the level of ROS in GCs of GABA biosynthesis mutants was significantly higher than that in the control (Figure 8). Therefore, we speculated that blocking the GABA metabolic shunt pathway will lead to the accumulation of ROS intermediates (Bouché et al., 2003; Fait et al., 2005), which may trigger endoreplication (Figure 9). In this hypothesis, normal GABA metabolism functioned as a signal and antioxidant to effectively inhibit the production of ROS. In the GABA biosynthetic mutants, this inhibition was relieved. The production of ROS promotes the expression of *SIAMESE* (*SIM*) and *SIAMESE-RELATED* (*SMR*) expression, of which, their encoding products inhibited kinase activity to initiate endoreplication (Figure 9). In contrast, normal GABA metabolism postponed endoreplication, and this negative regulation functions to prevent premature cell differentiation and to ensure normal leaf development. However, this conclusion must be made cautiously because, based on current data, it is difficult to demonstrate a causal relationship between GABA level and ROS in terms of triggering endoreplication during the development of Arabidopsis leaves.

Overall, the results of the present study demonstrate that the increase in polyploidy level in GABA biosynthetic mutant leaves is achieved through endoreplication, which is reflected in the increase in average cell size (Breuer et al., 2010). In this regulation, multiple regulators are involved in the initiation, maintenance, and exit of endoreplication. However, the premature termination of mitosis and the immature occurrence of endoreplication are perhaps a compensatory mechanism for leaf immature development, which resulted in increased cell ploidy. This mechanism may be mediated by ROS signalling, but the complex regulatory mechanism requires further research.

5 Conclusion

Using GABA biosynthetic mutants, the present study focused on the perturbation of GABA biosynthesis on endoreplication in Arabidopsis leaf development. The endoreplication cells that occurred in mutants were earlier and of higher abundance than those in the wild type. This is the reason to lead to the increase in cell size and leaf blade area. For transcription-level regulation, qRT-PCR confirmed that many genes involved in cell cycle regulation were synergistically participating in the initiation, progression, and maintenance of endoreplication. Among the regulators, *SMR5*, encoding protein inhibited CDK activity, was markedly upregulated in GABA biosynthetic mutants. Owing to the perturbation of GABA biosynthesis, the content of ROS increased in GABA biosynthetic mutants, which is a potential signal to trigger endoreplication in mutant leaves. Present evidence indicates that normal GABA metabolism inhibited endoreplication to prevent immature cell differentiation in leaf development. This research provides a deeper understanding of the role of GABA in plant development.

6 Acknowledgments

Endoreplication response to GABA deficit

We apologize if we inadvertently omitted citations of major contributions to this area. We thank Prof. Barry Shelp (University of Guelph, Canada) for generously providing the mutants seeds. We would like to thank Editage (www.editage.cn) for English language editing.

7 Author Contributions

YXG studied the root phenotype in relation to GABA metabolism and wrote the draft of the manuscript. HY and YX studied the role of ROS in cell-cycle regulation. GHY studied the role of small molecules in plant development and revised the manuscript.

8 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

9 Funding

This work was supported by the Special Fund for Basic Scientific Research of Central Colleges, South-Central University for Nationalities (CZP17051), National Natural Science Foundation of China (31270361), studies on the regulation mechanisms of agronomic traits in important crops and *A. thaliana* supported by State Administration of Foreign Experts Affairs in Ministry of Science and Technology of the People's Republic of China (P193009007) and Fund for Key Laboratory Construction of Hubei Province (Grant No.2018BFC360).

10 Data Availability Statement

No datasets were generated for this study.

11 References

- Allan WL, Simpson JP, Clark SM, Shelp BJ. Gamma-hydroxybutyrate accumulation in Arabidopsis and tobacco plants is a general response to abiotic stress: putative regulation by redox balance and glyoxylate reductase isoforms. J Exp Bot 2008; 59: 2555–64.
- Baum, G., S Lev-Yadun, S., Fridmann, Y., Arazi, T., Katsnelson, H., Zik, M., and Fromm, H. (1996). Calmodulin binding to glutamate decarboxylase is required for regulation of glutamate and GABA metabolism and normal development in plants. The EMBO J. 15, 2988-2996
- Beemster GT, Vercruysse S, De VL, et al. The Arabidopsis leaf as a model system for investigating the role of cell cycle regulation in organ growth. Journal of Plant Research, 2006, 119(1):43-50.

Endoreplication response to GABA deficit

- 417 Bouché N, Fait A, Bouchez D, Møller SG, Fromm H. Mitochondrial
418 succinic-semialdehyde dehydrogenase of the gamma-aminobutyrate shunt is
419 required to restrict levels of reactive oxygen intermediates in plants. Proc Natl
420 Acad Sci USA 2003; 100:6843-8.
- 421 Boudolf V, Vlieghe K, Beemster GT, Magyar Z, Torres Acosta JA, Maes S, Van Der
422 Schueren E, Inzé D, De Veylder L. (2004). The plant-specific cyclin-dependent
423 kinase CDKB1;1 and transcription factor E2Fa-DPa control the balance of
424 mitotically dividing and endoreduplicating cells in Arabidopsis. Plant Cell.
425 16(10):2683-2692.
- 426 Bown AW, Shelp BJ. Plant GABA: Not Just a Metabolite. Trends Plant Sci. 2016,
427 21(10):811-813.
- 428 Breuer C, Braidwood L, Sugimoto K. Endocycling in the path of plant development.
429 Curr Opin Plant Biol. 2014, 17:78-85.
- 430 Breuer C, Ishida T, Sugimoto K. Developmental control of endocycles and cell
431 growth in plants. Current Opinion in Plant Biology, 2010, 13(6):654-660.
- 432 Brikis CJ, Zarei A, Trobacher CP, DeEll JR, Akama K, Mullen RT, Bozzo GG, Shelp
433 BJ. Ancient Plant Glyoxylate/Succinic Semialdehyde Reductases: GLYR1s Are
434 Cytosolic, Whereas GLYR2s Are Localized to Both Mitochondria and Plastids.
435 Front Plant Sci. 2017; 8: 601.
- 436 Castellano MM, Boniotti MB, Caro E, et al. DNA replication licensing affects cell
437 proliferation or endoreplication in a cell type-specific manner. The Plant cell,
438 2004, 16(9): 2380-93.
- 439 Castellano MM, del Pozo JC, Ramirez-Parra E, Brown S, Gutierrez C. Expression and
440 stability of Arabidopsis CDC6 are associated with endoreplication. Plant Cell.
441 2001, 13(12):2671-86.
- 442 Churchman ML, Brown ML, Kato N, Kirik V, Hülskamp M, Inzé D, De Veylder L,
443 Walker JD, Zheng Z, Oppenheimer DG, Gwin T, Churchman J, Larkin JC.
444 SIAMESE, a plant-specific cell cycle regulator, controls endoreplication onset in
445 *Arabidopsis thaliana*. Plant Cell. 2006, 18(11):3145-57.

Endoreplication response to GABA deficit

- 446 Coelho, C., Dante, R. Sabelli, P., Sun, Y., Dilkes, B., Gordon-Kamm, W., and Larkins,
447 B. (2005). Cyclin-dependent kinase inhibitors in maize endosperm and their
448 potential role in endoreduplication. *Plant Physiol.* 138, 2323-2336.
- 449 Coleman ST, Fang TK, Rovinsky SA, Turano FJ, Moye-Rowley WS. Expression of a
450 glutamate decarboxylase homologue is required for normal oxidative stress
451 tolerance in *Saccharomyces cerevisiae*. *J Biol Chem.* 2001, 276(1):244-50.
- 452 De Veylder L, Larkin JC, Schnittger A. Molecular control and function of
453 endoreplication in development and physiology. *Trends Plant*
454 *Sci.* 2011,16(11):624-34.
- 455 De Veylder L., Beeckman T., Beemster G.T.S., Krols L., Terras F., Landrieu I., van
456 der Schueren E., Maes S., Naudts M., Inze D. (2001). Functional analysis of
457 cyclin-dependent kinase inhibitors of *Arabidopsis*. *Plant Cell* 13:1653-1667
- 458 Dewitte W, Riou-Khamlichi C, Scofield S, Healy JM, Jacquemard A, Kilby NJ, Murray
459 JA. Altered cell cycle distribution, hyperplasia, and inhibited differentiation in
460 *Arabidopsis* caused by the D-type cyclin CYCD3. *Plant Cell.* 2003, 15(1):79-92.
- 461 Dewitte W, Scofield S, Alcasabas AA, Maughan SC, Menges M, Braun N, Collins C,
462 Nieuwland J, Prinsen E, Sundaresan V, Murray JA. *Arabidopsis* CYCD3 D-type
463 cyclins link cell proliferation and endocycles and are rate-limiting for cytokinin
464 responses. *Proc Natl Acad Sci USA.* 2007, 104(36):14537-42.
- 465 Dubois M, Selden K, Bediee A, *et al.* SIAMESE-RELATED1 Is Regulated
466 Posttranslationally and Participates in Repression of Leaf Growth under
467 Moderate Drought. *Plant physiology*, 2018, 176(4): 2834-50.
- 468 Fait A, Yellin A, Fromm H. GABA shunt deficiencies and accumulation of reactive
469 oxygen intermediates: insight from *Arabidopsis* mutants. *FEBS Lett.* 2005,
470 579(2):415-20.
- 471 Fülöp K, Tarayre S, Kelemen Z, Horváth G, Kevei Z, Nikovics K, Bakó L, Brown S,
472 Kondorosi A, Kondorosi E. *Arabidopsis* anaphase-promoting complexes:
473 multiple activators and wide range of substrates might keep APC perpetually
474 busy. *Cell Cycle.* 2005, 4(8):1084-92.

Endoreplication response to GABA deficit

- 475 Fung-Uceda J, Lee K, Seo PJ, Polyn S, De Veylder L, Mas P. The Circadian Clock
476 Sets the Time of DNA Replication Licensing to Regulate Growth in Arabidopsis.
477 Dev Cell. 2018, 45 (1):101-113.
- 478 Häusler RE, Ludewig F, Krueger S. Amino acids--a life between metabolism and
479 signaling. Plant Sci. 2014, 229:225-237.
- 480 Heyman J, Polyn S, Eekhout T, De Veylder L. Tissue-Specific Control of the
481 Endocycle by the Anaphase Promoting Complex/Cyclosome Inhibitors UVI4
482 and DEL1. Plant Physiol. 2017, 175(1):303-313.
- 483 Jacoby RP, Taylor NL, Millar AH. The role of mitochondrial respiration in salinity
484 tolerance. Trends Plant Sci 2011; 16:614–23.
- 485 Jun SE, Okushima Y, Nam J, Umeda M, Kim GT. Kip-related protein 3 is required
486 for control of endoreduplication in the shoot apical meristem and leaves of
487 *Arabidopsis*. Mol Cells. 2013, 35(1):47-53.
- 488 Kumar N, Harashima H, Kalve S, Bramsiepe J, Wang K, Sizani BL, Bertrand LL,
489 Johnson MC, Faulk C, Dale R, Simmons LA, Churchman ML, Sugimoto K,
490 Kato N, Dasanayake M, Beemster G, Schnittger A, Larkin JC. Functional
491 Conservation in the SIAMESE-RELATED Family of Cyclin-Dependent Kinase
492 Inhibitors in Land Plants. Plant Cell. 2015, 27(11):3065-80.
- 493 Lammens T, Li J, Leone G, et al. Atypical E2Fs: new players in the E2F transcription
494 factor family. Trends in Cell Biology, 2009, 19(3):111-118.
- 495 Liu CL, Zhao L, Yu GH. (2011). The dominant glutamic acid metabolic influx to
496 produce γ -amino butyric acid over proline in *Nicotiana tabacum* leaves under
497 water stress relates to its significant role of antioxidant activity. Integr. Plant Biol.
498 53:608-618.
- 499 Masubelele NH, Dewitte W, Menges M, Maughan S, Collins C, Huntley R,
500 Nieuwland J, Scofield S, Murray JA. D-type cyclins activate division in the root
501 apex to promote seed germination in Arabidopsis. Proc Natl Acad Sci USA.
502 2005, 25; 102(43):15694-9.

Endoreplication response to GABA deficit

- 503 Matsunaga, S., Katagiri, Y., Nagashima, Y., Sugiyama, T., Hasegawa, J., Hayashi, K.
504 and Sakamoto, T. (2013). New insights into the dynamics of plant cell nuclei and
505 chromosomes. *Int. Rev. Cell Mol. Biol.* 305, 253-301.
- 506 Melaragno J E, Mehrotra B, Coleman A W. Relationship between Endopolyploidy
507 and Cell Size in Epidermal Tissue of Arabidopsis. *Plant Cell*, 1993,
508 5(11):1661-1668
- 509 Michaeli S, Fait A, Lagor K, Nunes-Nesi A, Grillich N, Yellin A, et al. A
510 mitochondrial GABA permease connects the GABA shunt and the TCA cycle,
511 and is essential for normal carbon metabolism. *Plant J* 2011; 67:485–498.
- 512 Nowack MK, Harashima H, Dissmeyer N, *et al.* Genetic framework of
513 cyclin-dependent kinase function in *Arabidopsis*. *Developmental cell*, 2012,
514 22(5): 1030-40.
- 515 Palanivelu R, Brass L, Edlund AF, Preuss D. Pollen tube growth and guidance is
516 regulated by POP2, an Arabidopsis gene that controls GABA levels. *Cell*. 2003,
517 114(1):47-59.
- 518 Podlešáková K, Ugena L, Spíchal L, Doležal K, De Diego N. Phytohormones and
519 polyamines regulate plant stress responses by altering GABA pathway. *New*
520 *Biotechnol.* 2019, 48:53-65.
- 521 Ramesh SA, Tyerman SD, Gilliam M, Xu B. γ -Aminobutyric acid (GABA)
522 signalling in plants. *Cell Mol Life Sci.* 2017, 74(9):1577-1603.
- 523 Renault H, El Amrani A, Berger A, Mouille G, Soubigou-Taconnat L, Bouchereau A,
524 et al. γ -Aminobutyric acid transaminase deficiency impairs central carbon
525 metabolism and leads to cell wall defects during salt stress in Arabidopsis roots.
526 *Plant Cell Environ* 2013; 36:1009–18.
- 527 Renault H, El Amrani A, Palanivelu R, Updegraff EP, Yu A, Renou JP, Preuss D,
528 Bouchereau A, Deleu C. GABA accumulation causes cell elongation defects and
529 a decrease in expression of genes encoding secreted and cell wall-related proteins
530 in Arabidopsis thaliana. *Plant Cell Physiol.* 2011, 52(5):894-908.

Endoreplication response to GABA deficit

- 531 Sawa S, Watanabe K, Goto K, Liu YG, Shibata D, Kanaya E, Morita EH, Okada K.
532 FILAMENTOUS FLOWER, a meristem and organ identity gene of Arabidopsis,
533 encodes a protein with a zinc finger and HMG-related domains. *Genes Dev.* 1999,
534 13(9):1079-88.
- 535 Schippers JH, Foyer CH, Van Dongen JT. Redox regulation in shoot growth, SAM
536 maintenance and flowering. *Current Opinion in Plant Biology*, 2016,
537 29:121-128.
- 538 Seifikalhor M, Aliniaefard S, Hassani B, Niknam V & Lastochkina O. (2019).
539 Diverse role of γ -aminobutyric acid in dynamic plant cell responses. *Plant Cell*
540 *Reports*. 38(8):847-867.
- 541 Siegfried KR, Eshed Y, Baum SF, Otsuga D, Drews GN, Bowman JL. Members of
542 the YABBY gene family specify abaxial cell fate in *Arabidopsis*. *Development*.
543 1999, 126(18):4117-28.
- 544 Toyokura K, Hayashi M, Nishimura M, and Okada K. Adaxial-abaxial patterning: a
545 novel function of the GABA shunt. *Plant Signal Behav.* 2012, 7 (7):705-7.
- 546 Toyokura K, Watanabe K, Oiwa A, Kusano M, Tameshige T, Tatematsu K,
547 Matsumoto N, Tsugeki R, Saito K, Okada K. Succinic semialdehyde
548 dehydrogenase is involved in the robust patterning of Arabidopsis leaves along
549 the adaxial-abaxial axis. *Plant Cell Physiol.* 2011, 52(8):1340-53.
- 550 Umeda M, Aki SS, Takahashi N. Gap 2 phase: making the fundamental decision to
551 divide or not. *Curr Opin Plant Biol.* 2019, 51:1-6.
- 552 Verkest A, Manes CL, Vercruysse S, Maes S, Van Der Schueren E, Beeckman T,
553 Genschik P, Kuiper M, Inzé D, De Veylder L (2005). The cyclin-dependent
554 kinase inhibitor KRP2 controls the onset of the endoreduplication cycle during
555 Arabidopsis leaf development through inhibition of mitotic CDKA; 1 kinase
556 complexes. *Plant Cell.* 2005, 17(6):1723-1736.
- 557 Vlieghe K, Boudolf V, Beemster G T, et al. (2005). The DP-E2F-like Gene DEL1,
558 Controls the Endocycle in Arabidopsis thaliana. *Current Biology*, 15(1):59-63.

Endoreplication response to GABA deficit

Watkins JM, Chapman JM, Muday GK (2017). Absciscic Acid-Induced Reactive Oxygen Species Are Modulated by Flavonols to Control Stomata Aperture. *Plant Physiol.* 175(4):1807-1825.

Yang K, Wang H, Xue S, *et al.* Requirement for A-type cyclin-dependent kinase and cyclins for the terminal division in the stomatal lineage of *Arabidopsis*. *J Exp Bot*, 2014, 65(9): 2449-2461.

Yi D, Alvim Kamei CL, Cools T, Vanderauwera S, Takahashi N, Okushima Y, Eekhout T, Yoshiyama KO, Larkin J, Van den Daele H, Conklin P, Britt A, Umeda M, De Veylder L. The *Arabidopsis* SIAMESE-RELATED cyclin-dependent kinase inhibitors SMR5 and SMR7 regulate the DNA damage checkpoint in response to reactive oxygen species. *Plant Cell*. 2014, 26(1):296-309.

Figure Legends

Figure 1. Leaf area of γ -Aminobutyric acid (GABA) biosynthetic mutants was larger than that of wild-type *Arabidopsis*.

Top view of leaves at 8 d after stratification of A. wild type; B. *gad1* mutant, C. *gad2* mutant and D. *gad1/gad2* mutants; E. Leaf growth curve of GABA mutants and wild type of *Arabidopsis* in the early stage (from 4–8 d after stratification). Results are presented as averages $\pm$ SE of three separate experiments (n = 15). Bar in A–D = 860 μ m.

Figure 2. Epidermal cell size in the first true leaves of γ -Aminobutyric acid (GABA) biosynthetic mutants was larger than that of wild-type *Arabidopsis* at the early growth stage.

Epidermal cell of A. wild type; B. *gad1* mutant, C. *gad2* mutant and D. *gad1/gad2* mutants; E. Cell size comparison among GABA mutants and wild type of *Arabidopsis* in the early stage (7–9 d after stratification). Results are presented as averages $\pm$ SE of 60 cells calculated across three separate experimental replicates. Asterisks represent significant differences between the mutants and the wild type ($P < 0.05$) and were determined using one-way ANOVA. Bar in A–D = 55 μ m.

Figure 3. Octoploid (8n) cells occurred earlier in biosynthetic mutants than that in the wild type. Top panel: DNA ploidy analysis in leaf early growth stage (7–8 days after stratification) between γ -Aminobutyric acid (GABA) biosynthetic mutants and the wild type. Left bottom panel: DNA ploidy analysis in leaf middle-late stage (21 days after stratification) between GABA mutants and the control. Right bottom panel: 8C percentage comparison among mutants and the wild type during the long growth period.

Endoreplication response to GABA deficit

Figure 4. Expression of *Cyclin D* components in biosynthetic mutants was lower than that in the wild type at the 8th day after stratification. Results are presented as averages $\pm$ SE of triplicate experiments. Asterisks represent significant differences between the mutants and the wild type (* P < 0.05; ** P < 0.01) and were determined using one-way ANOVA.

Figure 5. Expression of *CDA;I* and *E2Fa* in biosynthetic mutants was lower than that in the wild type at the 8th day after stratification. Results are presented as averages $\pm$ SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (* P < 0.05; ** P < 0.01) and were determined using one-way ANOVA.

Figure 6. Expression of *CCS52A* and *CDC6* in biosynthetic mutants was higher than that in the wild type at the 8th day after stratification. Results are presented as averages $\pm$ SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (* P < 0.05; ** P < 0.01) and were determined using one-way ANOVA.

Figure 7. Expression of *SIAMESE (SIM)* and *SIAMESE-RELATED (SMR)* components in biosynthetic mutants was higher than that in the wild type at the 8th day after stratification. Results are presented as averages $\pm$ SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (* P < 0.05; ** P < 0.01). Statistics were determined using one-way ANOVA.

Figure 8. Reactive oxygen species (ROS) level in guard cells (GCs) of γ -Aminobutyric acid (GABA) mutants are higher than that of control A–D. DCF fluorescence (green) in epidermal GCs of GABA biosynthetic mutants and wild type Arabidopsis. E. DCF fluorescence was quantified in the mutants and control. Results are presented as averages $\pm$ SE of three separate experiments (n = 16). Asterisks represent significant differences between the mutant and the wild type (P < 0.05) and were determined using one-way ANOVA. Bar in A–D = 55 μ m.

Figure 9. Working hypothesis of γ -Aminobutyric acid (GABA) negatively control the endoreplication of Arabidopsis leaves.

GABA, as a normal metabolic signal and antioxidant, effectively inhibits reactive oxygen species (ROS) production. In the GABA biosynthetic mutants, owing to biosynthetic perturbation, the inhibition of GABA on ROS was relieved. The production of ROS promotes the expression of *SIAMESE (SIM)* and *SIAMESE-RELATED (SMR)*. Both components are inhibitors of Cyclin-dependent Kinase complexes (e.g. CYC3;1-CDKA;1), and low levels of kinases initiate endoreplication in Arabidopsis. Other components, such as CDC6, synergistically respond to low levels of GABA to prime endoreplication. In the wild type, the normal metabolism of GABA delays endoreplication and prevents premature endoreplication.

Supporting Materials

Table S1. The primers of the genes involved in cell cycle regulation, endocycle initiation, progression, and exit.

Figure S1. The γ -Aminobutyric acid (GABA) analysis was performed on whole-shoot samples. Results are presented as averages $\pm$ standard error (SE) of three independent

Endoreplication response to GABA deficit

639 replicates. Significant differences from the control are indicated with asterisks: * P
640 < 0.05 , ** $P < 0.01$, by one-way ANOVA.

641

Endoreplication response to GABA deficit

Figure 1. Leaf area of γ -Aminobutyric acid (GABA) biosynthetic mutants was larger than that of wild-type Arabidopsis.

Top view of leaves at 8 d after stratification of A. wild type; B. *gad1* mutant, C. *gad2* mutant and D. *gad1/gad2* mutants; E. Leaf growth curve of GABA mutants and wild type of Arabidopsis in the early stage (from 4–8 d after stratification). Results are presented as averages $\pm$ SE of three separate experiments (n = 15). Bar in A–D = 860 μ m.

Endoreplication response to GABA deficit

Figure 2. Epidermal cell size in the first true leaves of γ -Aminobutyric acid (GABA) biosynthetic mutants was larger than that of wild-type Arabidopsis at the early growth stage.

Epidermal cell of A. wild type; B. *gad1* mutant, C. *gad2* mutant and D. *gad1/gad2* mutants; E. Cell size comparison among GABA mutants and wild type of Arabidopsis in the early stage (7–9 d after stratification). Results are presented as averages $\pm$ SE of 60 cells calculated across three separate experimental replicates. Asterisks represent

Endoreplication response to GABA deficit

significant differences between the mutants and the wild type ($P < 0.05$) and were determined using one-way ANOVA. Bar in A–D = 55 μm .

Figure 3. Octoploid (8n) cells occurred earlier in biosynthetic mutants than that in the wild type. Top panel: DNA ploidy analysis in leaf early growth stage (7–8 days after stratification) between γ -Aminobutyric acid (GABA) biosynthetic mutants and the wild type. Left bottom panel: DNA ploidy analysis in leaf middle-late stage (21 days after stratification) between GABA mutants and the control. Right bottom panel: 8C percentage comparison among mutants and the wild type during the long growth period.

Endoreplication response to GABA deficit

678

679

680 **Figure 4.** Expression of *Cyclin D* components in biosynthetic mutants was lower than
681 that in the wild type at the 8th day after stratification. Results are presented as averages
682 $\pm$ SE of triplicate experiments. Asterisks represent significant differences between the
683 mutants and the wild type (* P < 0.05; ** P < 0.01) and were determined using
684 one-way ANOVA.

685

Endoreplication response to GABA deficit

Figure 5. Expression of *CDKA;1* and *E2Fa* in biosynthetic mutants was lower than that in the wild type at the 8th day after stratification. Results are presented as averages $\pm$ SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (* $P < 0.05$; ** $P < 0.01$) and were determined using one-way ANOVA.

Endoreplication response to GABA deficit

Figure 6. Expression of *CCS52A* and *CDC6* in biosynthetic mutants was higher than that in the wild type at the 8th day after stratification. Results are presented as averages $\pm$ SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (* P < 0.05; ** P < 0.01) and were determined using one-way ANOVA.

Endoreplication response to GABA deficit

Figure 7. Relative expression of *SIM* gene

Figure 7. Expression of *SIAMESE* (*SIM*) and *SIAMESE-RELATED* (*SMR*) components in biosynthetic mutants was higher than that in the wild type at the 8th day after stratification. Results are presented as averages $\pm$ SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (* P < 0.05; ** P < 0.01). Statistics were determined using one-way ANOVA.

Endoreplication response to GABA deficit

Figure 8. Reactive oxygen species (ROS) level in guard cells (GCs) of γ -Aminobutyric acid (GABA) mutants are higher than that of control A–D. DCF fluorescence (green) in epidermal GCs of GABA biosynthetic mutants and wild type Arabidopsis. E. DCF fluorescence was quantified in the mutants and control. Results are presented as averages $\pm$ SE of three separate experiments (n = 16). Asterisks

Endoreplication response to GABA deficit

723 represent significant differences between the mutant and the wild type ($P < 0.05$) and
724 were determined using one-way ANOVA. Bar in A–D = 55 μm .

725

Endoreplication response to GABA deficit

Figure 9. Working hypothesis of γ -Aminobutyric acid (GABA) negatively control the endoreplication of Arabidopsis leaves.

GABA, as a normal metabolic signal and antioxidant, effectively inhibits reactive oxygen species (ROS) production. In the GABA biosynthetic mutants, owing to biosynthetic perturbation, the inhibition of GABA on ROS was relieved. The production of ROS promotes the expression of *SIAMESE* (*SIM*) and *SIAMESE-RELATED* (*SMR*). Both components are inhibitors of Cyclin-dependent Kinase complexes (e.g. CYC3;1-CDKA;1), and low levels of kinases initiate endoreplication in Arabidopsis. Other components, such as CDC6, synergistically respond to low levels of GABA to prime endoreplication. In the wild type, the normal metabolism of GABA delays endoreplication and prevents premature endoreplication.

Endoreplication response to GABA deficit

Supporting Materials

Table S1: The primers of the genes involved in cell cycle regulation, endocycle initiation, progression and exit.

Locus tag	Gene name	Primer	Primer sequences
AT3G02780	<i>IPP2</i>	IPP2_EXP_F	CATGCGACACACCAACACCA
		IPP2_EXP_R	TGAGGCGAATCAATGGGAGA
AT4G34160	<i>CYCD3;1</i>	CYCD3;1_EXP_F	CCTCTCTGTAATCTCCGATTC
		CYCD3;1_EXP_R	AAGGACACCGAGGAGATTAG
AT5G67260	<i>CYCD3;2</i>	CYCD3;2_EXP_F	TCTCAGCTTGTTGCTGTGGCTTC
		CYCD3;2_EXP_R	TCTTGCTTCTTCCACTTGGAGGTC
AT3G50070	<i>CYCD3;3</i>	CYCD3;3_EXP_F	TCCGATCGGTGTGTTTGATGCG
		CYCD3;3_EXP_R	GCAGACACAACCCACGACTCATTC
AT5G65420	<i>CYCD4;1</i>	CYCD4;1_EXP_F	GAAGGAGAAGCAGCATTTGCCAAG
		CYCD4;1_EXP_R	ACTGGTGTACTTCACAAGCCTTCC
AT4G11920	<i>CCS52A2</i>	CCS52A2_EXP_F	CGTAGATACCAACAGCCAGGTGTG
		CCS52A2_EXP_R	CGTGTGTGCTCACAAGCTCATTC
AT2G29680	<i>CDC6</i>	CDC6_EXP_F	AGGCTCTATGTGTCTGCAGGAG
		CDC6_EXP_R	ACCACTTGACACTCTGGAAGTGG
AT2G31270	<i>CDT1a</i>	CDT1a_EXP_F	AATCGCTCTTCGGAAAGTGTTCG
		CDT1a_EXP_R	CCTCTGGAACCTTCATCACCCCTGAG
AT3G48750	<i>CDKA;1</i>	CDKA;1_EXP_F	ACTGGCCAGAGCATTCGGTATC
		CDKA;1_EXP_R	TCGGTACCAGAGAGTAACAACCTC
AT2G36010	<i>E2Fa</i>	E2Fa_EXP_F	TAGATCGGGAGGAAGATGCTGTGCG

Endoreplication response to GABA deficit

		E2Fa_EXP_R	TTGTCGCCTTCTCTTTTCGTGAAG
AT2G23430	<i>KRP1</i>	KRP1_EXP_F	ACGGAGCCGGAGAATTGTTTATG
		KRP1_EXP_R	CGAAACTCCATTATCACCGACGAC
AT3G50630	<i>KRP2</i>	KRP2_EXP_F	TAGGAGATTATGGCGGCGGTTAGG
		KRP2_EXP_R	TTTCACCGTCGTCGTCGTAAGTC
AT2G32710	<i>KRP4</i>	KRP4_EXP_F	AAGCTTCAACAGGACCACAAGGG
		KRP4_EXP_R	GGGTTGTCATGATTTCAGGCCTTC
AT1G49620	<i>KRP7</i>	KRP7_EXP_F	GAGGCTCATGAAATCTCCGAAACC
		KRP7_EXP_R	CCGAGTCCATTTCTGCTGTTTCTC
AT5G04470	<i>SIM</i>	SIM_EXP_F	AGCCATCAAGATCCGAGCCAAC
		SIM_EXP_R	TTGTGGTCGGAAGAAGTGGGAGTG
AT3G10525	<i>SMR1</i>	SMR1_EXP_F	CAAAGAAGGACGAAGGTGATGACG
		SMR1_EXP_R	TGTTCTTGGGATGTGGGTGTGC
AT1G60783	<i>SMR2</i>	SMR2_EXP_F	TCACAAGATTCCGGAGGTGGAGAC
		SMR2_EXP_R	ATCTCACGCGGTGCTTTCTTG
AT1G07500	<i>SMR5</i>	SMR5_EXP_F	ACGCCTACACGTGATGATTGCC
		SMR5_EXP_R	TATCCCTTCTTCGGTGGTTCCC
AT1G10690	<i>SMR8</i>	SMR8_EXP_F	GCGGTTTCCGTCAGAATTCCAAG
		SMR8_EXP_R	GCACTTCAACGACGGTTTACGC

746

747

Endoreplication response to GABA deficit

Figure S1: GABA analysis was performed on whole shoot samples. Averages $\pm$ standard error (SE) of three independent replicates. Significant differences from the control are indicated with asterisks: * $P < 0.05$, ** $P < 0.01$, by one-way ANOVA.

GABA levels in the leaves of the mutants were determined essentially as described earlier (Allan and Shelp 2006).

Allan WL, Simpson JP, Clark SM, Shelp BJ. Gamma-hydroxybutyrate accumulation in Arabidopsis and tobacco plants is a general response to abiotic stress: putative regulation by redox balance and glyoxylate reductase isoforms. J Exp Bot 2008; 59: 2555–64.