Immunization of mice with chimeric antigens displaying selected epitopes confers protection against intestinal colonization and renal damage caused by Shiga toxin-producing *Escherichia coli*

David A. Montero^{1,2}, Felipe Del Canto¹, Juan C. Salazar¹, Sandra Cespedes¹, Leandro Cádiz¹, Mauricio Arenas-Salinas³, José Reyes⁴, Ángel Oñate⁴, Roberto M. Vidal^{1,5 *}

1. Programa de Microbiología y Micología, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile
2. Programa Disciplinario de Inmunología, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile
3. Centro de Bioinformática y Simulación Molecular, Facultad de Ingeniería, Universidad de Talca, Talca, Chile
4. Departamento de Microbiología, Facultad de Ciencias Biológicas, Universidad de Concepción, Concepción, Chile
5. Instituto Milenio de Inmunología e Inmunoterapia, Facultad de Medicina, Universidad de Chile, Santiago, Chile

* Corresponding author. Correspondence should be addressed to R.V. (email: rvidal@uchile.cl)

Abstract

Shiga toxin-producing *Escherichia coli* (STEC) cause diarrhea and dysentery, which may progress to hemolytic uremic syndrome (HUS). Vaccination has been proposed as a preventive approach against STEC infection; however, there is no vaccine for humans and those used in animals reduce but do not eliminate the intestinal colonization of STEC. The OmpT, Cah and Hes proteins are widely distributed among clinical STEC strains and are recognized by serum IgG and IgA in patients with HUS. Here, we develop a vaccine formulation based on two chimeric antigens containing epitopes of OmpT, Cah and Hes proteins against STEC strains. Intramuscular and intranasal immunization of mice with these chimeric antigens elicited systemic and local long-lasting humoral responses. However, the class of antibodies generated was dependent on the adjuvant and the route of administration. Moreover, while intramuscular immunization with the combination of the chimeric antigens conferred protection against colonization by STEC O157:H7 and the intranasal conferred protection against renal damage caused by STEC O91:H21. This pre-clinical study supports the potential use of this formulation based on recombinant chimeric proteins as a preventive strategy against STEC infections.

Keywords: Shiga toxin-producing *Escherichia coli*, STEC, chimeric antigen-based vaccine, Hemolytic uremic syndrome.

73 Introduction

74 Shiga toxin-producing *Escherichia coli* (STEC) are a group of food-borne pathogens
 75 causing acute and bloody diarrhea, which may progress to life-threatening
 76 complications such as hemolytic uremic syndrome (HUS).¹ To date there is no specific
 77 treatment for STEC infection and antibiotic use is contraindicated due to increased risk
 78 of HUS development.² However, some drugs have been specifically designed to
 79 protect against the effects of the presence of Shiga toxins and are in different stages of
 80 clinical trials.^{3,4} While STEC O157:H7 is the serotype most frequently associated with
 81 diarrhea outbreaks and HUS cases worldwide, there are other serotypes, the incidence
 82 and impact of which on public health and the food industry have increased.^{5,6}

83 STEC colonizes the human colon and produces Shiga toxins (Stx) that can enter the
 84 blood stream and disseminate to organs such as the kidneys and central nervous
 85 system. Once Stx reach the target organs and enter the cells, the toxins inhibit protein
 86 synthesis, leading to autophagy and apoptosis and ultimately tissue damage, which
 87 may lead to HUS.⁷

88 To colonize the human colon, STEC requires several virulence factors like those
 89 encoded in the locus of enterocyte effacement (LEE) pathogenicity island (PAI). LEE-
 90 mediated adherence causes the formation of the "attaching and effacing" lesion and
 91 loss of microvilli of the intestinal epithelial cells.⁸ In addition, STEC strains lacking LEE
 92 (LEE-negative STEC) harbor other PAIs like the Locus of Adhesion and
 93 Autoaggregation (LAA), which encodes virulence factors involved in intestinal
 94 colonization.^{9,10} In fact, the presence of two or more PAIs in single isolates of clinically
 95 relevant STEC serotypes is common, suggesting that the cumulative acquisition of
 96 mobile genetic elements encoding virulence factors may contribute additively or
 97 synergistically to pathogenicity.^{10,11}

98 Vaccination of the infant population, which is the highest-risk group for STEC
 99 infections, and animal reservoirs have been proposed as a preventive approach that
 100 could reduce their incidence and prevalence. However, there is no approved STEC
 101 vaccine for humans, and commercial vaccines used in cattle reduce but do not
 102 eliminate colonization and shedding of these bacteria.¹² Therefore, the development of
 103 an effective STEC vaccine is still underway. STEC proteins involved in attachment to
 104 host tissues are eligible targets for vaccine development, as they determine initial steps
 105 during infection; however, the selection of antigens that may provide a broadly and
 106 protective immune response among their diverse adhesion and colonization
 107 mechanisms is a pivotal point to consider.¹³ An additional difficulty for the development

of an effective STEC vaccine has been the lack of an animal model of infection that can reproduce the pathologies caused in humans.¹⁴ Despite these limitations, several STEC vaccine candidates have been evaluated in laboratory animals (mice, rats and rabbits) and in cattle, with promising results. They include Stx subunit-based vaccines,^{15–17} protein and peptide-based vaccines,^{17–21} attenuated bacteria-based vaccines,²² bacterial ghost-based vaccines,²³ DNA-based vaccines,^{24,25} and more recently nanoparticle-based vaccines.²⁶ While most of these vaccine candidates are based on LEE-encoded antigens and Stx subunits, there are several antigens encoded outside LEE that are expressed *in vivo* during human infection that could be suitable targets for vaccine development.^{13,27}

For instance, Outer membrane protease I (OmpT) and Calcium binding antigen 43 homologue (Cah) proteins have been shown to be recognized by IgG and IgA antibodies present in sera from patients who develop HUS (hereinafter referred to as HUS sera). Notably, the *ompT* gene has been identified in almost all clinical STEC strains and, in the case of the *cah* gene, its detection frequency is higher than 70%.¹³ Another promising antigen is the Hemagglutinin from STEC (Hes), which is recognized by IgG present in HUS sera.¹³ In addition, the *hes* gene, which is carried by the LAA PAI, is identified in about 40% and 46% of LEE-negative STEC strains isolated from humans and cattle, respectively.^{9,10,28} Thus, these three antigens are widespread among clinical STEC strains, but it is also important to note that they are mostly absent in commensal *E. coli* strains,^{9,10,13} which could diminish the probability of cross-reactivity with commensal microbiota. Nevertheless, the production and purification of outer membrane proteins (OMPs) such as OmpT, Cah and Hes poses a challenge due to their partially hydrophobic surfaces, flexibility and lack of stability that affect their solubility and efficient purification. In addition, strong detergents are used in the purification of this class of proteins and therefore the loss of conformational epitopes may impair their antigenicity and efficiency as immunogens.²⁹

To circumvent these issues and to develop a STEC vaccine targeting the OmpT, Cah and Hes proteins, we implemented a vaccine development approach based on the identification of linear B-cell epitopes for the design of chimeric antigens that include them. Here we demonstrate that the immunization of mice with chimeric antigens displaying selected epitopes of OmpT, Cah and Hes proteins induces immune responses that reduce intestinal colonization and prevent renal damage caused by STEC. Overall, this study revealed the feasibility of using such type of formulation based on recombinant chimeric proteins against STEC colonization and more relevantly to protect against kidney damage by Shiga toxin-producing *Escherichia coli*.

144 We anticipate that our comprehensive experimental approach will contribute to the
145 development and evaluation of future chimeric antigen-based vaccines, and while our
146 candidate was initially intended to protect humans against colonization/infection by
147 STEC, we believe that it could also have an effect on STEC elimination in the animal
148 reservoir (bovine and pig) and realistically such a trial is most likely to be conducted
149 first.

150

151 Results

152 **The OmpT, Cah and Hes proteins have several linear B-cell epitopes that are**
153 **recognized by IgG and IgA from HUS sera but not from control sera.**

154

155 An overview of our vaccine development approach is shown in Figure 1. We carried out
156 a high-throughput screening of linear B-cell epitopes in the OmpT, Cah and Hes
157 proteins by using a peptide microarray assay (see Methods). A total of 6, 6 and 5
158 epitope-like spot patterns were identified in the peptide slides of OmpT, Cah and Hes
159 proteins (**Fig. 2a-2c**), respectively. As a complementary approach, we used a number
160 of immunoinformatics tools and found that 13 out of 17 of the experimentally identified
161 B-cell epitopes were predicted *in silico* by one or more algorithms (**Table 1**). This result
162 highlights the advances in the accuracy of these bioinformatics tools. On the other
163 hand, it is known that MHC class II (MHC-II) epitopes included in peptide vaccines
164 enhance T-cell-dependent antibody responses³⁰. Thus, we also performed an *in silico*
165 analysis for the prediction of MHC-II binding peptides and found two putative T-cell
166 epitopes in the OmpT and Cah proteins (**Table 2**).

167

168 Because the peptide microarray assay was performed with a mix of three HUS sera,
169 we sought to confirm the reactivity of these epitopes using a larger number of sera. We
170 also tested the reactivity of the epitopes against three sera obtained from children with
171 no medical record of STEC-related disease (hereinafter referred to as control sera). For
172 this, short peptides ranging from 15 to 24 amino acids (aa) which include each epitope
173 were chemically synthesized, and their reactivity to IgG and IgA was assessed by
174 ELISA. In general, the peptides were recognized at higher levels by IgG than by IgA
175 from HUS sera (**Fig. 2d-i**). Furthermore, the peptides derived from the same antigen
176 were recognized by similar levels of IgG or IgA, with the exception of Cah_5 and
177 Hes_5, which showed a higher level of reactivity to IgG compared to Cah_4 and
178 Hes_4, respectively (**Fig. 2f and 2h**). The frequency of reactivity of the peptides to IgG

179 ranged from 60% (Hes_4) to 100% (Cah_5, Cah_6 and Hes_5), while for IgA it was
180 between 29% (Cah_2) and 100% (Hes_5) (**Table 1**). As expected, a polyclonal
181 antibody response was observed within individuals, evidenced by the variable
182 concentration of IgG and IgA antibodies that recognize each tested peptide (**Fig. 2j-k**).
183 Importantly, none of the peptides was recognized by IgG or IgA from control sera
184 (**Table 1, Fig. 2j-k**). Taken together, these results indicate that the identified linear B-
185 cell epitopes are broadly recognized and immunodominant during immune responses
186 against STEC.

187

Chimeric proteins displaying linear B-cell epitopes of OmpT, Cah and Hes proteins are recognized by IgG and IgA from HUS sera but not from control sera.

190

191 We consider that the best epitopes for vaccine development are those that are
192 conserved, broadly distributed among clinical isolates, with higher levels of
193 immunoreactivity and surface exposition in the native antigen. As a result, the Hes_4
194 (lower reactivity and detection frequency) and OmpT_5 (limited surface exposition)
195 epitopes were discarded and not used in further assays. For the *in silico* design of
196 proteins we implemented two different approaches. Firstly, we noticed that the linear B-
197 cell epitopes of OmpT and Hes are overlapped or consecutively arranged along the
198 protein (**Fig. 2a and 2c**), suggesting that they form antigenic domains (AD). We took
199 advantage of this and designed a chimeric protein containing these AD through the
200 fusion of 135 AA and 127 AA from OmpT and Hes proteins, respectively (**Fig. 3a**). We
201 named this protein Chimera 1 (Chi1; 262 AA and 29 kDa), which includes a total of
202 eight B-cell epitopes (OmpT_1, OmpT_2, OmpT_3, OmpT_4, Hes_1, Hes_2, Hes_3
203 and Hes_5) and two predicted T-cell epitopes (**Table 2**). In the second approach, we
204 used the passenger domain of Cah (α Cah) as a carrier (keeping its epitopes) and
205 incorporated several B-cell epitopes of OmpT and Hes (**Table 2, Fig. 3d**). Thus, this
206 second protein named Chimera 2 (Chi2; 559 AA and 56 kDa) includes a total of eleven
207 B-cell epitopes (OmpT_1, OmpT_6, Hes_1, Hes_2 and Hes_5, Cah_1, Cah_2, Cah_3,
208 Cah_4, Cah_5, Cah_6) and two predicted T-cell epitopes (**Table 2**).

209

210 We predicted the 3D structures for both chimeric proteins, which were refined and
211 validated (see Methods) (**Fig. 3a and 3d**). A Ramachandran plot analysis revealed that
212 94.6% and 86.6% of amino acid residues from the Chi1 and Chi2 modeled structures
213 were in favored regions, respectively (**Fig. 3b and 3e**). In addition, the Z-score of the
214 Chi1 and Chi2 modeled structures were -2.52 and -7.01 respectively, which are within
215 the range of scores found for native proteins of similar size (**Fig. 3c and 3f**). The

216 predicted solubility, *in vivo* half-life and instability index of both chimeric proteins
 217 suggested that their expression and purification could be feasible (**Table 2**). Consistent
 218 with the above, the production of these recombinant proteins in *E. coli* showed that
 219 they are stable, water-soluble and have the predicted molecular weight (**Fig. 3g**).
 220 Further, we confirm that both the Chi1 and Chi2 proteins are recognized by IgG and
 221 IgA of HUS sera (**Fig. 3h and 3i**). We also found that the reactivity of Chi2 to IgG of
 222 HUS sera was significantly higher than α Cah (**Fig. 3h**), indicating that the incorporation
 223 of B-cell epitopes of OmpT and Hes increased the seroreactivity. However, this
 224 difference was not observed in the reactivity to IgA of HUS sera (**Fig. 3i**). Importantly,
 225 none of the proteins was seroreactive to IgG and IgA of control sera (not shown).

226

227 **Chi1 and Chi2 antigens, administered alone or in combination, trigger long-** 228 **lasting systemic and local humoral responses in mice.**

229

230 Having established that the Chi1 and Chi2 proteins are seroreactive to HUS sera, we
 231 sought to evaluate them as immunogens. For this, BALB/c mice were immunized
 232 following the scheme described in Figure 1d. Immunity achieved by vaccination is
 233 influenced to a large extent by the administration route and the type of adjuvant^{31–33}.
 234 Therefore, we also compared systemic and local immune responses triggered by the
 235 chimeric antigens when administered with Imject Alum or Sigma adjuvants by
 236 intramuscular or intranasal route, respectively.

237

238 The measurement of Chi1 and Chi2-specific IgG antibodies in serum showed that mice
 239 immunized with Chi2 or Chi1 plus Chi2 by either intramuscular or intranasal route
 240 elicited significantly higher levels of IgG on days 28 and 42 than the PBS control group
 241 (**Fig. 4a**). In general, intramuscular immunization induced higher levels of specific IgG
 242 antibodies than the intranasal route, this difference being significant in mice immunized
 243 with Chi1 plus Chi2. Similarly, specific IgA antibodies in serum were significantly higher
 244 in mice immunized with Chi2 or Chi1 plus Chi2 by both administration routes on days
 245 14, 28 and 42 than in the PBS control group (**Fig. 4b**). It was also observed that
 246 intramuscular immunization with Chi1 plus Chi2 induced higher levels of specific IgA
 247 antibodies than the intranasal route, this difference being significant on day 42. In
 248 contrast, there were no differences on the levels of specific IgG and IgA antibodies in
 249 serum between mice immunized with Chi1 and the PBS control group (**Fig. 4a-b**).
 250 Regarding specific IgM antibodies in serum, all vaccine formulations and administration
 251 routes induced significant higher antibody levels on days 14, 28 and 42 than the PBS
 252 control group (**Fig. 4c**).

To evaluate the induction of mucosal responses, specific secretory IgA (sIgA) antibodies were determined in feces. Mice immunized with Chi1 and Chi1 plus Chi2 by intranasal route elicited significantly higher levels of specific sIgA on days 14 and 28 than the PBS control group (**Fig. 4d**). However, the sIgA levels for both experimental groups were similar on day 42 compared to the PBS control group. On day 42, only mice immunized with Chi2 by intranasal route elicited significantly higher levels of specific sIgA than the PBS control group. Taken together, these results indicate that immunization with Chi1, Chi2 or Chi1 plus Chi2 induces a systemic and local humoral response influenced by the type of adjuvant and the administration route as long as 42 days after the third immunization.

263

Chi1 and Chi2 antigens administered in combination by intramuscular route reduce intestinal colonization and fecal shedding of STEC O157:H7

266

We next evaluated whether immune responses elicited by vaccination with the chimeric antigens may confer protection against intestinal colonization by STEC. For this, two weeks after the last booster immunization, mice were treated with streptomycin and then orally challenged with STEC O157:H7 str. 86-24 (**Fig. 1d**; see Methods). Notably, mice immunized with Chi1 plus Chi2 by intramuscular route showed a significantly lower fecal shedding of the STEC 86-24 strain from day 8 post-infection to the end of the experiment (day 12) compared to the PBS control group (**Fig. 5a**). In mice immunized by the intranasal route, only the Chi2 group showed a slight but significant decrease in fecal shedding of STEC 86-24 strain on days 7 and 8 compared to the PBS control group. However, on day 9 and later, this difference was not observed (**Fig. 5b**).

On day 12 post-infection, mice were euthanized and the level of colonization of the challenge strain in the cecum was determined. Consistent with the above results, recovery of the STEC 86-24 strain was significantly lower in mice immunized with Chi1 plus Chi2 by intramuscular route than in the PBS control group, which corresponded to 2.1 log of protection (**Table 3**). The other experimental groups presented levels of colonization similar to the PBS control group. These data demonstrate that immunization of mice with Chi1 plus Chi2 by intramuscular route induces protective immune responses against intestinal colonization of STEC O157:H7 in this model.

286

Chi1 and Chi2 antigens administered alone or in combination by intranasal route avoid renal damage caused by STEC O91:H21

288

Kidney damage is one of the most severe clinical outcomes that can occur during STEC infection due to the action of Shiga toxins. In streptomycin-treated mice, Stx2d-producing *E. coli* strains have been shown to affect renal function leading to death.^{34,35} For instance, in a previous study we showed that the Stx2d-producing *E. coli* O91:H21 str. V07-4-4 is lethal to mice when they are orally inoculated with a dose of 10^9 CFU.¹⁰ However, with a dose lower than 10^5 CFU of the STEC V07-4-4 strain, mice develop renal pathologies but survive for at least 12 days (unpublished results). Therefore, we conducted further work to investigate whether immunization with the chimeric antigens confers protection against renal damage caused by the STEC V07-4-4 strain. For this, two weeks after the last booster immunization, mice were treated with streptomycin and orally challenged with 10^5 CFU of the STEC V07-4-4 strain. Our results showed that on days 7 and 12 post-infection, mice intranasally immunized with Chi1, Chi2 or Chi1 plus Chi2 had creatinine levels in urine similar to uninfected mice (**Fig. 5c and 5d**). This was particularly evident in mice immunized with Chi1 plus Chi2. In contrast, mice immunized by intramuscular route and the PBS control group showed significantly higher levels of creatinine in urine than uninfected mice. Moreover, the histopathological analysis of kidney tissue obtained on day 12 post-infection showed that mice immunized by intranasal route had mild or no evident tissue injuries (**Fig 5e**). Conversely, mice immunized by intramuscular route and the PBS control group showed moderate and severe tissue injuries, respectively (**Fig 5e**). Urinary clinical markers such as the number of leukocytes and urobilinogen levels on day 12 post-infection also evidenced the protection achieved by intranasal immunizations (**Table 4**). Together, these results indicate that immunization with the chimeric antigens by intranasal route protects against kidney damage caused by the STEC V07-4-4 strain.

Discussion

Sera obtained from Chilean-hospitalized pediatric patients diagnosed with HUS, after STEC primoinfection, recognizes antigens such as OmpT, Cah and Hes, as result of a primary immune response to an initial STEC antigen exposure with the development of immunological memory (sera obtained from the convalescent phase with IgG and IgA that recognize STEC antigens).¹³ Therefore, the key to designing a new vaccine also considers the bacterial target selected for this process. Since STEC infection recurrence is an uncommon process,^{36,37} it is likely that STEC primoinfection can trigger a successful immune memory directed against key bacterial antigens and that remains over time.

Here we demonstrated that immunization of mice with chimeric antigens displaying selected epitopes of OMPs confers protection against intestinal colonization and renal damage caused by STEC. It is well established that a suitable vaccine must be composed of different antigens to boost the immune response with a wide range of protection. Interestingly, the antigens selected for our vaccine design are different from those used in most trials;^{62–65} however, they are widely distributed in STEC and involved in several of its pathogenicity mechanisms.^{10,13} While Cah and Hes are related to the bacterial-host interaction,^{9,66} OmpT participates in the biogenesis of bacterial outer membrane vesicles (OMVs)⁵⁸ and the degradation of antimicrobial peptides like LL-37.^{61,67} To our knowledge, this is the first formulation based on recombinant chimeric proteins that includes a virulence factor exclusive of LEE-negative STEC strains in a vaccine design.

In general, subunit and protein-based vaccines have proven to be safe and to have a defined and homogeneous composition between production batches. This latter aspect is an important advantage over other types of vaccines, which may present complex manufacturing processes, with subsequent regulatory and safety issues.^{38–40} In Gram-negative bacteria, OMPs are primary components interacting with host cells; therefore, vaccines targeting these proteins may be effective by blocking key pathogenic mechanisms.^{13,41} However, as previously mentioned, the purification and efficient production of OMPs pose a major challenge. Thus, construction of water-soluble and stable chimeric antigens displaying epitopes from OMPs is an approach worth exploring.

Our vaccine development approach takes advantage of immunoinformatics tools and *in vitro* assays (epitope mapping, ELISA) to predict and identify linear B-cell epitopes, respectively (**Fig. 1**). Previous studies have also demonstrated the usefulness of immunoinformatics tools for the development of vaccine candidates against STEC and other pathogens.^{24,42} Both, *in silico* and *in vitro* assays, allowed us to select the best epitopes (**Fig. 2, Table 1, Table 2**).

Importantly, our results provide proof-in-principle that incorporating selected linear B-cell epitopes into a carrier protein, such as α Cah, may result in an increase in antigenicity (**Fig. 3h**). Many autotransporter (AT) proteins and especially their passenger domains are potential vaccine targets.^{43,44} In fact, the pertactin AT from *Bordetella pertussis* is a component of licensed pertussis vaccines.⁴⁵ Thus, other passenger domains from AT proteins could be used as carriers to design chimeric antigens.

359 The immunization studies showed that the Chi1 and Chi2 antigens, administered alone
 360 or in combination, induce humoral responses which remain active until day 42
 361 postimmunization (**Fig 4**). However, the class of antibodies generated was in general
 362 dependent on the adjuvant and the administration route. This dependency was most
 363 obvious in mice immunized with Chi1 plus Chi2, which elicited significantly higher
 364 levels of specific IgG and IgA antibodies in serum when intramuscularly immunized
 365 than when intranasally immunized (**Fig. 4a-b**). Further, the sIgA detection in feces after
 366 systemic immunization is interesting; however, while this seems to be a controversial
 367 issue, there is literature describing this type of results suggesting that vaccines
 368 systemically administered can trigger mucosal immune responses.⁴⁶ Many questions
 369 await an answer, and one of them is how the immune response moves to the mucosa
 370 after systemic immunization, which for some researchers, in addition to producing a
 371 paradigm shift, may also mean a modification in vaccine design and delivery.⁴⁷

372 Secretory IgA is the most abundant immunoglobulin of the mammalian mucosa, playing
 373 a fundamental role in the immunity of the gastrointestinal tract.⁴⁸ Therefore, the
 374 development of vaccines against intestinal pathogens has traditionally given priority to
 375 immunogens that induce significant levels of sIgA. However, since some
 376 gastrointestinal infections can be eliminated in the absence of sIgA; other classes of
 377 antibodies such as IgM and IgG may also play an important role in the intestinal
 378 immunity.⁴⁹ Unfortunately, the presence and effector functions of IgG in the intestinal
 379 mucosa have been largely ignored in the literature.⁵⁰ A recent study by Kamada et al.,
 380 2015,⁵¹ revealed that IgG in the murine intestine leads to the selective elimination of a
 381 virulent *Citrobacter rodentium* subpopulation by luminal neutrophils. The protective role
 382 of IgG against other enteropathogens such as rotavirus has also been demonstrated.⁵²

383 Our challenge experiments using the STEC O157:H7 strain showed that only
 384 immunization with Chi1 plus Chi2 by intramuscular route confers protection against
 385 intestinal colonization (**Fig. 5a-b, Table 3**). In the murine model of infection, the
 386 permanent addition of streptomycin throughout the challenge, promotes STEC
 387 O157:H7 colonization by preventing the interference of the microbiota. This situation
 388 might explain the decrease in the final part of the protection test (days 8-12), only
 389 associated to immune response against STEC O157:H7. Probably, if we had removed
 390 the treatment with streptomycin, the interfering activity of the microbiota added to the
 391 immune response would have affected early colonization by STEC O157:H7 (before 8
 392 days).⁵³ Since intramuscular immunization with Chi1 plus Chi2 did not lead to
 393 significant production of specific fecal sIgA antibodies (**Fig. 4d**), it is possible to
 394 correlate the protection achieved with other classes of immunoglobulins and more likely

395 with the IgG. On the other hand, although Chi2 includes epitopes of the other two
 396 antigens in addition to Cah, it was observed that the set of intranasally immunized
 397 mice, yielded a mild immune response against O157:H7 on days 7 and 8, but this was
 398 neither sufficiently protective nor maintained over time. Other vaccine candidates that
 399 generated significant levels of specific IgG antibodies in serum but not fecal sIgA
 400 antibodies have also conferred protection to mice against colonization by STEC
 401 O157:H7.⁵⁴ Consequently, our results and those reported by others support the idea of
 402 a pivotal role of the IgG antibodies in the defense against enteropathogens. This is a
 403 major finding that will be relevant to the development of vaccines against these
 404 pathogens by avoiding biases in the selection of the "best" immunogens based mainly
 405 on the ability to induce sIgA antibody responses.

406 Because the challenge studies were performed only to day 12 post-inoculation, it was
 407 not possible to determine whether the immune responses triggered by the
 408 intramuscular immunization with Chi1 and Chi2 may lead to complete clearance of
 409 STEC O157:H7. Long-term protection experiments in mice and other animal models
 410 immunized will complement the evaluation of this formulation based on recombinant
 411 chimeric proteins.

412 An effective STEC vaccine may also confer protection against the action of the Stx.
 413 STEC export the Shiga toxins along with a number of OMPs and cytoplasmic proteins
 414 via outer membrane vesicles (OMVs).⁵⁵⁻⁵⁷ These OMVs may be endocytosed in a
 415 dynamin-dependent manner by intestinal epithelial cells, and then OMV-associated
 416 virulence factors are differentially separated from vesicles during intracellular
 417 trafficking.^{57,58} Recently, it was reported that in the case of Stx2 but not Stx1, once the
 418 toxin is internalized, it can be released from eukaryote cells in microvesicles that have
 419 exosome markers.⁵⁹ Therefore, immunity against Stx may be mediated by neutralizing
 420 Stx-specific antibodies or by immune mechanisms that prevent the toxin from entering
 421 the eukaryotic cell.

422 Our chimeric antigens did not display Stx-associated epitopes. Consequently, the
 423 protection conferred by intranasal immunizations against renal damage caused by
 424 STEC O91:H21 (**Fig. 5c-e, Table 4**) could be explained by two different mechanisms.
 425 The first, the generated immune response might reduce the intestinal colonization of
 426 the STEC O91:H21 strain, which could be correlated with a lower release and number
 427 of OMVs carrying Stx; however, the intestinal colonization by STEC O91:H21 was not
 428 measured. As a result, we cannot conclude that the decrease in colonization correlates
 429 with protection against renal damage. The second possible explanation is that sIgA

antibodies generated by intranasal immunizations could prevent the release and/or endocytosis of OMVs via immune exclusion. The latter explanation is supported by the fact that OmpT and Ag43 (a Cah homologue protein) are transported in OMVs.^{58,60}

In some vaccines, it has been shown that the combination of administration routes, for example mucosal priming followed by systemic boosting or systemic priming followed by mucosal boosting, leads to robust humoral and cellular responses that improve their efficacy.^{68,69} In future studies we will investigate whether the combination of systematic and mucosal immunizations with the Chi1 and Chi2 antigens leads to a more robust and complete immune response characterized by the production of both systemic and secretory antibodies. Also, we will endeavor to reveal the mechanism by which intranasal immunization with Chi1 and Chi2 confers protection against renal damage caused by STEC O91:H21.

Our main focus has always been to protect human health. This is based on the fact that we have seen that the selected antigens are present in a wide range of STEC serotypes previously associated to human illness. However, there are a number of studies that also link these virulence factors to interaction mechanisms between STEC and intestinal epithelial cells in cattle and pigs. In this context, we speculate that our candidate might also be used as a vaccine in animals to prevent STEC colonization, another way to protect the human health.

In conclusion, we developed a promising formulation based on recombinant chimeric proteins that confers protection against STEC intestinal colonization and more relevantly against renal damage caused by Stx. Our study presents interesting results that support the potential use of recombinant chimeras containing epitopes of different antigens of STEC as a preventive strategy.

Methods

Bacterial strains and growth conditions

Spontaneously derived streptomycin resistant (Str^r) mutants of STEC O157:H7 86-24 and STEC O91:H21 V07-4-4 strains were used in this study. Bacterial cultures were routinely grown at 37 °C in Luria-Bertani (LB) broth.

Human sera

Sera were obtained from 20 pediatric patients in the convalescent phase who presented diarrhea within two weeks prior to HUS diagnosis (HUS sera). Control sera

were obtained from two patients with no history of STEC-associated diarrhea. These sera were collected from 1990 to 1993 and from 1999 to 2003 in various healthcare centers in Santiago, Chile, with the written consent of the parents or legal guardians. All procedures were approved by the Ethics Committee of the Facultad de Medicina, Universidad de Chile.

Peptide Microarray

Epitope mapping assays were performed by PEPperPRINT (Heidelberg, Germany). Briefly, the *ompT*, *cah* and *hes* sequences were translated into 15, 12 and 10 amino acid peptides with peptide-peptide overlaps of 14, 11 and 9 amino acids. The microarray contained peptides printed in duplicate framed by HA (YPYDVPDYAG) control peptides. Peptide slides were incubated with a mix of three HUS sera at a dilution of 1:100 followed by secondary antibody Goat anti-human IgA (DyLight800) at a dilution of 1:1000, in the presence of the monoclonal anti-HA (12CA5)-DyLight680 control antibody at a dilution of 1:2000. The read-out was performed with a LI-COR Odyssey Imaging System and the image analysis with the PepSlide® Analyzer.

In silico prediction of B-cell and T-cell epitopes

Prediction of B-cell epitopes was done using several tools available at the IEDB server⁷⁰, including BepiPred 2.0⁷¹, Kolaskar and Tongaonker antigenicity method⁷² and Ellipro⁷³. Peptides binding to MHC-II molecules were also predicted on the IEDB server⁷⁰.

Validation of epitopes by ELISA assay

Seventeen short peptides from 15 to 24 amino acids (**Table 1**) containing linear B-cell epitopes were chemically synthesized at Genic Bio Ltd (Shanghai, China). These short peptides were evaluated for their reactivity to IgG and IgA of individual HUS sera by ELISA. Briefly, 96-well ELISA plates (Nunc Maxisorp or Nunc Immobilizer Amino Plates, ThermoFisher, USA) were incubated with 1.2 µg of each peptide diluted in 100 µl of phosphate-buffered saline (PBS; pH 7.2) overnight at 4 °C. Standard curves were obtained by dilutions in PBS of purified human IgG (Cat. 02-7102, Invitrogen, USA) or IgA (Cat. 3860-1AD-6, Mabtech, USA) ranging from 1.2 µg/ml to 0.0047 µg/ml. Plates were washed three times with PBS containing 0.05% Tween 20 (TPBS) and then incubated with blocking solution (TPBS + 0.5% bovine serum albumin) for 15 min at room temperature (RT). The HUS and control sera were diluted 1:25 (dilution determined from serum titration experiments in a range of 1:10 to 1:100) in blocking solution (100 µl / well) and incubated for 60 min at 37 °C. After six washes with T-PBS,

501 goat anti-human IgG (H + L), peroxidase-labeled (Cat. 04-10-06, KPL, USA) or goat
502 anti-human IgA alpha chain (alkaline phosphatase) (Cat. Ab97212, Abcam, UK),
503 diluted 1:1000 in blocking solution, were added and plates were incubated for 60 min at
504 37 °C. After six washes with Tris-buffered saline (TBS; pH 7.5) containing 0.05%
505 Tween 20, ABTS® peroxidase substrate (Cat. 50-66-18, KPL, USA) or pNPP substrate
506 (Cat. N2600-10, USBiological, USA) were added and plates were incubated for 12 or
507 30 min at RT, respectively. The reaction was stopped with 5% sodium dodecyl sulfate
508 or 3 M sodium hydroxide dissolved in distilled water. The absorbance was determined
509 at 405 nm (A_{405}) using a Synergy HT microplate reader (Biotek Instruments, USA).
510 Each sample was determined twice in duplicate. The relation between absorbance
511 values and the IgG or IgA concentration of each well was calculated from standard
512 curves using a four-parameter logistic regression in GraphPad Prism 8 software.

513

514 ***In silico* modeling and design of chimeric proteins**

515 Predicted three-dimensional structure of Chimera 1 was constructed based on the
516 crystal structures of the Opa60 (PDB_ID: 2MLH)⁷⁴ and OmpT (PDB_ID: 1I78)⁷⁵
517 proteins. The template for Chimera 2 structure was the crystal structure of the Ag43
518 protein (PDB_ID: 4KH3).⁷⁶ Comparative modeling of chimeric proteins was performed
519 in Modeller v9 software⁷⁷, using default parameters. The modeled structures were
520 solvated and embedded in a water box using ions (Na⁺, Cl⁻) to neutralize the system
521 with the TCL script using VMD software.⁷⁸ The models were optimized with cycles of
522 energy minimization and dynamics using the NAMD 2.12 software.⁷⁹ A molecular
523 dynamics simulation was performed under periodic bordering conditions and isobaric-
524 isothermal set (NPT). The entire system was relaxed by molecular dynamics (MD)
525 simulations using NAMD 2.12 software for 10 ns and subsequently balanced for 30 ns,
526 using the force field CHARMM v2.7.⁸⁰ Quality evaluation and validation of the models
527 were carried out by Ramachandran plot analysis on the RAMPAGE server
528 (<http://mordred.bioc.cam.ac.uk/~rapper/rampage.php>) and ProSA-web.⁸¹ Chemical
529 and physical properties of the chimeric proteins were predicted by Protein-Sol⁸² and
530 ProtParam⁸³ tools. The modelled structures were visualized with UCSF Chimera
531 1.10.2.⁸⁴

532

533 **Purification of proteins**

534 Synthetic genes and production of Chimera 1, Chimera 2 and αCah proteins were
535 ordered to GenScript (USA). Synthetic genes were optimized for *E. coli* expression and
536 cloned into vector pET30a with N-terminal 6xHis-tag. *E. coli* strain BL21(DE3) was then
537 transformed with recombinant plasmids containing synthetic genes. For purification of

recombinant proteins, transformant BL21(DE3) strains were grown in Terrific Broth containing kanamycin (50 µg/ml) at 37° C. When the culture reached an optical density at 600 nm of ~1.2, it was supplemented with IPTG for 4 h. Later, cells were harvested by centrifugation, resuspended with lysis buffer followed by sonication. The sediment obtained after centrifugation was dissolved using urea. Denatured protein was obtained by one-step purification using a Ni-column. The target protein was refolded and sterilized by 0.22 µm filter before being stored in aliquots. The concentration was determined by Bradford protein assay with BSA as standard. The protein purity and molecular weight were determined by standard SDS-PAGE along with Western blot confirmation (Supplementary Figure 1). Target proteins were obtained with purity >85% and endotoxin level <2 EU/mg (LAL Endotoxin Assay Kit, GenScript, Cat. No. L00350). In addition, reactivity of Chimeric and αCah proteins to IgG and IgA of individual HUS was assessed by ELISA assay as described above.

551

552 Immunization studies

All animal experiments were performed at the Universidad de Concepción, Concepción, Chile, following protocols and guidelines approved by the Bioethics Committee of the Faculty of Biological Sciences. Female BALB/c mice (5 to 6 weeks old; purchased from the Instituto de Salud Pública, Santiago, Chile) were randomly distributed into seven experimental groups (each group n=20) and housed in conventional animal facilities with water and food *ad libitum*. Mice were anesthetized with 10 mg/ml of ketamine and 250 µg/ml of acepromazine and immunized by intramuscular (i.m.) or intranasal (i.n.) route with the corresponding protein formulation along with 50 µl of Imject™ Alum Adjuvant (ThermoFisher Scientific, USA) or 20 µl of Sigma Adjuvant System® oil (Sigma-Aldrich, USA), respectively (**Fig. 1d**). The intramuscular immunization was performed in the hamstring muscle and the other group of animals immunized intranasally were anesthetized with a ketamine/xylazine mixture and the corresponding volume of vaccine was administered through the nose. Experimental groups 1, 3 and 5 were i.m. immunized with either 20 µg of Chi1, 20 µg of Chi2 or 10 µg of Chi1 plus 10 µg of Chi2, respectively. Experimental groups 2, 4 and 6 were i.n. immunized with either 20 µg of Chi1, 20 µg of Chi2 or 10 µg of Chi1 plus 10 µg of Chi2, respectively. The control group were injected with PBS plus adjuvants. Two booster immunizations were performed on days 15 and 30 using similar amounts of protein formulations and adjuvants. In the BALB/c model, STEC O157:H7 does not cause morbidity or mortality but does allow us to evaluate intestinal colonization.⁸⁵ In contrast, STEC O91:H21 and its mucus-activated Shiga toxin variant 2d (Stx2d) allowed us to measure kidney damage in infected BALB/c mice.^{86,87}

575

576 **Sera and feces collection**

577 Sera were obtained from five mice per group by tail vein bleeding on days -2, 14 and
578 28 before each immunization and at day 42 (two weeks after the last booster
579 immunization), according to conventional techniques. Briefly, blood samples were left
580 at 37 °C for 30 min and then centrifuged at 1,000 x g for 10 min. Supernatant was
581 collected, the complement was inactivated at 56 °C for 30 min and aliquots were stored
582 at -80 °C until IgG, IgA and IgM determinations by ELISA. For secretory IgA (sIgA)
583 measurement, feces were collected on days -2, 14, 28 and 42. Feces were weight,
584 homogenized and diluted to 0.1 g/ml with PBS containing 0.1% sodium azide and 1
585 mM of phenylmethylsulfonyl fluoride (PMSF). Fecal suspension was centrifuged at
586 15,000 x g for 5 min at 4 °C, the supernatant fluid recovered and again centrifuged at
587 15,000 x g for 15 min at 4 °C and stored at -80 °C until use.

588

589 **Measurement of humoral response**

590 Chimeric proteins were diluted to 1 µg/ml in carbonate buffer (pH 9.6) and used to coat
591 polystyrene 96-well high-binding ELISA plates (100 µl/well; Nunc-Immuno plate with
592 MaxiSorp surface). After overnight incubation at 4 °C, plates were washed with
593 washing buffer (Tris-buffered saline [pH 7.4] with 0.05% Tween 20) and blocked with
594 0.8% gelatin in TPBS for 1 h at 37 °C and then incubated with either sera or
595 supernatant from fecal suspensions, at a dilution of 1:100, for 2.5 h at room
596 temperature and washed four times. Next, isotype-specific goat anti-mouse HRP
597 conjugates (BioLegend, USA) were added (100 µl/well) at a dilution of 1:1000 and
598 incubated for 1 h min at room temperature followed by washing with TPBS. Then, 200
599 µl/well of OPD Peroxidase Substrate (Cat. P9187-5SET, Sigma-Aldric, USA) was
600 added for 30 min. The reaction was stopped with 50 µl/well of 2 N H₂SO₄ and the
601 absorbance at 450 nm was measured on a microplate reader.

602

603 **Challenge studies**

604 Two weeks after the last booster immunization (day 45), the infection experiments were
605 performed in the streptomycin-treated mouse model of STEC infection as described
606 elsewhere^{87,88} with minor modifications. Briefly, mice were given water *ad libitum*
607 containing streptomycin (5 g/l) 24–48 h prior to inoculation and for the duration of the
608 experiment. Feces were documented to be free of streptomycin-resistant *E. coli* at the
609 time of inoculation. STEC O157:H7 86-24 and STEC O91:H21 V07-4-4 strains were
610 grown overnight in agitated LB broth containing 50 µg/ml streptomycin at 37 °C.
611 Cultures were centrifuged, washed once with PBS and resuspended in a 20% sucrose

(w/v) and 10% NaHCO₃ (w/v) solution in sterile water to 1×10^{10} CFU/ml (STEC 86-24 strain) or 1×10^6 CFU/ml (STEC V07-4-4 strain). Prior to inoculation, mice were starved of food and water overnight (12 h). The next morning mice were orally infected by pipette feeding with 100 μ l of bacterial suspension containing 10^9 CFU or 10^5 CFU of STEC 86-24 or STEC V07-4-4 strains, respectively. After challenge, food and water were reintroduced and provided *ad libitum*. The fecal shedding of the 86-24 strain was recorded daily for 12 days. For this, feces were collected, weighed, homogenized, suspended in 1 ml PBS and, after serial dilutions, plated on MacConkey agar plates supplemented with streptomycin (50 μ g/ml) for bacterial counts. For determination of intestinal colonization of the STEC 86-24 strain, 12 days after challenge (day 57), mice were euthanized, and cecum was collected under aseptic conditions, homogenized and diluted in PBS. Suspensions were serially diluted and plated on MacConkey supplemented with streptomycin (50 μ g/ml) for bacterial counts. Log₁₀ units of protection were obtained by subtracting the mean Log₁₀ CFU for each experimental group from the mean Log₁₀ CFU of the PBS control group. Mice inoculated with the STEC V07-4-4 strain were used to measure kidney damage (*vide infra*).

Urinalysis

Urine samples were collected as previously described⁸⁹, on days 45, 51 and 57 (0, 7 and 12 post-infection, respectively) from mice infected with the STEC V07-4-4 strain. Biochemical estimation of urine creatinine concentration was assessed using the Creatinine Kit (BioSystems, Spain) according to the manufacturer's instructions. Other clinical urine markers were measured by using Combur10 Test®M semiquantitative test strips (Roche Diagnostics GmbH, Germany). Each test strip consists of colorimetric reaction spots for 10 markers: specific gravity (1.000 to 1.030), pH (5.0 to 9.0), leukocytes (range, negative to 500 cells/ μ L), nitrites (negative or positive), proteins (negative to 500 mg/dl), glucose (negative to 55 mmol/l), ketones (negative to 15 mmol/L), urobilinogen (normal to 200 μ mol/L), bilirubin (negative to +3) and blood (negative; trace of non-hemolyzed; or hemolyzed, 10 to 250 cells/ μ L). Each square was wet with a drop of urine and the marker value was determined through comparison with a colorimetric standard.

Histopathological analysis of kidney tissue

For the histological analysis of kidney tissue, mice infected with the STEC V07-4-4 strain were euthanized at day 57, the kidneys collected, fixed in 10% formaldehyde (pH 6.9), embedded in paraffin wax for sectioning at 5 μ m and stained with hematoxylin and eosin (H/E). Pathological evaluation of H/E-stained tissue sections was carried out by a

649 pathologist blinded to the experimental design. Histopathological changes were
650 evaluated by the degree of perivascular edema, leukocyte infiltration, vascular
651 congestion, mesangial cell expansion and injury of the glomerular filtration barrier
652 (glomerular hypertrophy or glomerular hypoperfusion). Each sample was quantitated by
653 ten randomly selected fields with the following criteria: 0, no damage; 1, <25%; 2, 25–
654 50%; 3, >50%; 4, >75% of damage. Differences between experimental groups were
655 evaluated by a one-way ANOVA followed by Tukey's multiple comparisons test.

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683 **Data availability**

684 The data that support the findings of this study are available on request from the
685 corresponding author R.V. Amino acid sequences of Chimeric proteins and identified
686 epitopes are not publicly available due to legal restrictions and an ongoing international
687 patent application.

688

689 **Acknowledgments**

690 This study was supported by FONDEF ID16I10140 and FONDECYT 1161161 grants.
691 We thank Dr. Helen Lowry for the careful revision and edition of the manuscript. We
692 thank Dr. Mauricio Farfán for sharing the STEC O157:H7 str. 86-24.

693

694 **Competing Interests**

695 Currently, an application for an international patent had been presented for the
696 chimeric antigens developed and uses thereof (PCT/IB2019/054554). The authors
697 declare that the research was conducted in the absence of any commercial or financial
698 relationships that could be construed as a potential conflict of interest.

699

700 **Author contributions**

701 Conceptualization and experimental design: Roberto Vidal, David A. Montero, Felipe
702 Del Canto, Juan C. Salazar and Angel Oñate. Data acquisition: Sandra Céspedes,
703 Leandro Cádiz, José Reyes, Mauricio Arenas and David A. Montero. Data analysis and
704 interpretation: Roberto Vidal, David A. Montero, Felipe Del Canto, Juan C. Salazar and
705 Angel Oñate. Writing – original draft: David A. Montero; Review and edition: Roberto
706 Vidal, Felipe Del Canto, Juan C. Salazar, Mauricio Arenas and Angel Oñate.

707

708

709

710

711

712

713

714

715

716

717

718 **References**

- 719 1 Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev* 1998;
720 **11**: 142–201.
- 721 2 Freedman SB, Xie J, Neufeld MS, Hamilton WL, Hartling L, Tarr PI. Shiga toxin–
722 producing *Escherichia coli* infection, antibiotics, and risk of developing Hemolytic
723 Uremic Syndrome: A meta-analysis. *Clin Infect Dis* 2016; **62**: 1251–1258.
- 724 3 Thallion Pharmaceuticals. Study of chimeric monoclonal antibodies to Shiga
725 Toxins 1 and 2. ClinicalTrials.gov Identifier: NCT01252199.
726 2013.<https://clinicaltrials.gov/ct2/show/study/NCT01252199>.
- 727 4 Immunova S.A. Anti-shiga toxin hyperimmune equine immunoglobulin F(ab')₂
728 fragment (INM004) in healthy volunteers. ClinicalTrials.gov Identifier:
729 NCT03388216. 2019. <https://clinicaltrials.gov/ct2/show/study/NCT03388216>.
- 730 5 Majowicz SE, Scallan E, Jones-Bitton A, Sargeant JM, Stapleton J, Angulo FJ *et*
731 *al.* Global incidence of human Shiga toxin–producing *Escherichia coli* infections
732 and deaths: A systematic review and knowledge synthesis. *Foodborne Pathog*
733 *Dis* 2014; **11**: 447–455.
- 734 6 Hazard identification and characterization: Criteria for categorizing Shiga Toxin–
735 producing *Escherichia coli* on a risk basis. *J Food Prot* 2019; **82**: 7–21.
- 736 7 Lee MS, Tesh VL. Roles of shiga toxins in immunopathology. *Toxins (Basel)*
737 2019; **11**: 1–26.
- 738 8 Farfán MJ, Torres AG. Molecular mechanisms that mediate colonization of Shiga
739 toxin-producing *Escherichia coli* strains. *Infect Immun* 2012; **80**: 903–913.
- 740 9 Montero DA, Velasco J, Del Canto F, Puente JL, Padola NL, Rasko DA *et al.*
741 Locus of Adhesion and Autoaggregation (LAA), a pathogenicity island present in
742 emerging Shiga Toxin-producing *Escherichia coli* strains. *Sci Rep* 2017; **7**: 7011.
- 743 10 Montero DA, Canto F Del, Velasco J, Colello R, Padola NL, Salazar JC *et al.*
744 Cumulative acquisition of pathogenicity islands has shaped virulence potential
745 and contributed to the emergence of LEE-negative Shiga toxin-producing
746 *Escherichia coli* strains. *Emerg Microbes Infect* 2019; **8**: 486–502.
- 747 11 Coombes BK, Wickham ME, Mascarenhas M, Gruenheid S, Finlay BB, Karmali
748 M a. Molecular analysis as an aid to assess the public health risk of non-O157
749 shiga toxin-producing *Escherichia coli* strains. *Appl Environ Microbiol* 2008; **74**:

750 2153–2160.

751 12 O’Ryan M, Vidal R, del Canto F, Carlos Salazar J, Montero D. Vaccines for viral
752 and bacterial pathogens causing acute gastroenteritis: Part II: Vaccines for
753 *Shigella*, *Salmonella*, enterotoxigenic *E. coli* (ETEC), enterohemorrhagic *E. coli*
754 (EHEC) and *Campylobacter jejuni*. *Hum Vaccin Immunother* 2015; **11**: 601–619.

755 13 Montero D, Orellana P, Gutiérrez D, Araya D, Salazar JC, Prado V *et al*.
756 Immunoproteomic analysis to identify Shiga toxin-producing *Escherichia coli*
757 outer membrane proteins expressed during human infection. *Infect Immun* 2014;
758 **82**: 4767–4777.

759 14 Garcia-Angulo V a, Kalita A, Torres AG. Advances in the development of
760 enterohemorrhagic *Escherichia coli* vaccines using murine models of infection.
761 *Vaccine* 2013; **31**: 3229–3235.

762 15 Cai K, Gao X, Li T, Wang Q, Hou X, Tu W *et al*. Enhanced immunogenicity of a
763 novel Stx2Am-Stx1B fusion protein in a mice model of enterohemorrhagic
764 *Escherichia coli* O157:H7 infection. *Vaccine* 2011; **29**: 946–952.

765 16 Mejias MP, Gherzi G, Craig PO, Panek CA, Bentancor L V, Baschkier A *et al*.
766 Immunization with a chimera consisting of the B subunit of Shiga toxin type 2
767 and brucella lumazine synthase confers total protection against Shiga toxins in
768 mice. *J Immunol* 2013; **191**: 2403–2411.

769 17 Martorelli L, Garimano N, Fiorentino GA, Vilte DA, Garbaccio SG, Barth SA *et al*.
770 Efficacy of a recombinant Intimin, EspB and Shiga toxin 2B vaccine in calves
771 experimentally challenged with *Escherichia coli* O157:H7. *Vaccine* 2018; **36**:
772 3949–3959.

773 18 Gu J, Liu Y, Yu S, Wang H, Wang Q, Yi Y *et al*. Enterohemorrhagic *Escherichia*
774 *coli* trivalent recombinant vaccine containing EspA, intimin and Stx2 induces
775 strong humoral immune response and confers protection in mice. *Microbes*
776 *Infect* 2009; **11**: 835–841.

777 19 Cheng Y, Feng Y, Luo P, Gu J, Yu S, Zhang W *et al*. Fusion expression and
778 immunogenicity of EHEC EspA-Stx2AI protein: Implications for the vaccine
779 development. *J Microbiol* 2009; **47**: 498–505.

780 20 Gao X, Cai K, Li T, Wang Q, Hou X, Tian R *et al*. Novel fusion protein protects
781 against adherence and toxicity of enterohemorrhagic *Escherichia coli* O157:H7
782 in mice. *Vaccine* 2011; **29**: 6656–6663.

- 783 21 Wan C, Zhou Y, Yu Y, Peng LJ, Zhao W, Zheng X. B-cell epitope KT-12 of
784 enterohemorrhagic *Escherichia coli* O157:H7: A novel peptide vaccine
785 candidate. *Microbiol Immunol* 2011; **55**: 247–253.
- 786 22 Liu J, Sun Y, Feng S, Zhu L, Guo X, Qi C. Towards an attenuated
787 enterohemorrhagic *Escherichia coli* O157:H7 vaccine characterized by a deleted
788 ler gene and containing apathogenic Shiga toxins. *Vaccine* 2009; **27**: 5929–
789 5935.
- 790 23 Cai K, Tu W, Liu Y, Li T, Wang H. Novel fusion antigen displayed-bacterial
791 ghosts vaccine candidate against infection of *Escherichia coli* O157:H7. *Sci Rep*
792 2015; **5**: 17479.
- 793 24 García-Angulo VA, Kalita A, Kalita M, Lozano L, Torres AG. Comparative
794 genomics and immunoinformatics approach for the identification of vaccine
795 candidates for enterohemorrhagic *Escherichia coli* O157:H7. *Infect Immun* 2014;
796 **82**: 2016–2026.
- 797 25 Riquelme-Neira R, Rivera A, Sáez D, Fernández P, Osorio G, del Canto F *et al.*
798 Vaccination with DNA encoding truncated enterohemorrhagic *Escherichia coli*
799 (EHEC) Factor for Adherence-1 gene (efa-1') confers protective immunity to
800 mice infected with *E. coli* O157:H7. *Front Cell Infect Microbiol* 2016; **5**: 1–9.
- 801 26 Sanchez-Villamil JI, Tapia D, Torres AG. Development of a gold nanoparticle
802 vaccine against Enterohemorrhagic *Escherichia coli* O157:H7. *mBio* 2019; **10**:
803 pii: e01869-19.
- 804 27 John M, Kudva IT, Griffin RW, Dodson AW, McManus B, Krastins B *et al.* Use of
805 *in vivo*-induced antigen technology for identification of *Escherichia coli* O157:H7
806 proteins expressed during human infection. *Infect Immun* 2005; **73**: 2665–2679.
- 807 28 Colello R, Vélez MV, González J, Montero DA, Bustamante A V, Canto F Del *et al.*
808 First report of the distribution of Locus of Adhesion and Autoaggregation
809 (LAA) pathogenicity island in LEE-negative Shiga toxin-producing *Escherichia*
810 *coli* isolates from Argentina. *Microb Pathog* 2018; **123**: 259–263.
- 811 29 Carpenter EP, Beis K, Cameron AD, Iwata S. Overcoming the challenges of
812 membrane protein crystallography. *Curr Opin Struct Biol* 2008; **18**: 581–586.
- 813 30 Kaumaya PTP, Kobs-Conrad S, Seo YH, Lee H, Vanbuskirk AM, Feng N *et al.*
814 Peptide vaccines incorporating a 'promiscuous' T-cell epitope bypass certain
815 haplotype restricted immune responses and provide broad spectrum

816 immunogenicity. *J Mol Recognit* 1993; **6**: 81–94.

817 31 Reed SG, Orr MT, Fox CB. Key roles of adjuvants in modern vaccines. *Nat Med*
818 2013; **19**: 1597–1608.

819 32 Mohanan D, Slütter B, Henriksen-Lacey M, Jiskoot W, Bouwstra JA, Perrie Y *et*
820 *al.* Administration routes affect the quality of immune responses: A cross-
821 sectional evaluation of particulate antigen-delivery systems. *J Control Release*
822 2010; **147**: 342–349.

823 33 Belyakov IM, Ahlers JD. What role does the route of immunization play in the
824 generation of protective immunity against mucosal pathogens? *J Immunol* 2009;
825 **183**: 6883–6892.

826 34 Melton-Celsa AR, O'Brien AD, Feng PCH. Virulence potential of activatable
827 Shiga Toxin 2d-producing *Escherichia coli* isolates from fresh produce. *J Food*
828 *Prot* 2015; **78**: 2085–2088.

829 35 Melton-Celsa AR, Darnell SC, O'Brien AD. Activation of Shiga-like toxins by
830 mouse and human intestinal mucus correlates with virulence of
831 enterohemorrhagic *Escherichia coli* O91:H21 isolates in orally infected,
832 streptomycin-treated mice. *Infect Immun* 1996; **64**: 1569–76.

833 36 Commereuc M, Weill F-X, Loukiadis E, Gouali M, Gleizal A, Kormann R *et al.*
834 Recurrent hemolytic and uremic syndrome induced by *Escherichia coli*. *Medicine*
835 *(Baltimore)* 2016; **95**: e2050.

836 37 Siegler RL, Griffin PM, Barrett TJ, Strockbine NA. Recurrent hemolytic uremic
837 syndrome secondary to *Escherichia coli* O157:H7 infection. *Pediatrics* 1993; **91**:
838 666–668.

839 38 Cox MMJ. Recombinant protein vaccines produced in insect cells. *Vaccine* 2012;
840 **30**: 1759–1766.

841 39 Hollingshead S, Jongerius I, Exley RM, Johnson S, Lea SM, Tang CM.
842 Structure-based design of chimeric antigens for multivalent protein vaccines. *Nat*
843 *Commun* 2018; **9**: 1–10.

844 40 Li W, Joshi M, Singhania S, Ramsey K, Murthy A. Peptide Vaccine: Progress
845 and challenges. *Vaccines* 2014; **2**: 515–536.

846 41 Chang YF, Chen CS, Palaniappan RUM, He H, McDonough SP, Barr SC *et al.*
847 Immunogenicity of the recombinant leptospiral putative outer membrane proteins

848 as vaccine candidates. *Vaccine* 2007; **25**: 8190–8197.

849 42 Tapia D, Ross BN, Kalita A, Kalita M, Hatcher CL, Muruato LA *et al*. From *in*
850 *silico* protein epitope density prediction to testing *Escherichia coli* O157:H7
851 vaccine candidates in a murine model of colonization. *Front Cell Infect Microbiol*
852 2016; **6**: 1–8.

853 43 Wells TJ, Tree JJ, Ulett GC, Schembri M a. Autotransporter proteins: novel
854 targets at the bacterial cell surface. *FEMS Microbiol Lett* 2007; **274**: 163–172.

855 44 Harris J a., Roy K, Woo-Rasberry V, Hamilton DJ, Kansal R, Qadri F *et al*.
856 Directed evaluation of Enterotoxigenic *Escherichia coli* autotransporter proteins
857 as putative vaccine candidates. *PLoS Negl Trop Dis* 2011; **5**: e1428.

858 45 Klein NP. Licensed pertussis vaccines in the United States. *Hum Vaccin*
859 *Immunother* 2014; **10**: 2684–2690.

860 46 Su F, Patel GB, Hu S, Chen W. Induction of mucosal immunity through systemic
861 immunization: Phantom or reality? *Hum Vaccines Immunother* 2016; **12**: 1070–
862 1079.

863 47 Clements JD, Freytag LC. Parenteral vaccination can be an effective means of
864 inducing protective mucosal responses. *Clin Vaccine Immunol* 2016; **23**: 438–
865 441.

866 48 Brandtzaeg P. Mucosal immunity: Induction, dissemination, and effector
867 functions. *Scand J Immunol* 2009; **70**: 505–515.

868 49 Eckmann L, Stappenbeck TS. IgG ‘detoxes’ the intestinal mucosa. *Cell Host*
869 *Microbe* 2015; **17**: 538–539.

870 50 Horton RE, Vidarsson G. Antibodies and their receptors: Different potential roles
871 in mucosal defense. *Front Immunol* 2013; **4**.

872 51 Kamada N, Sakamoto K, Seo S-U, Zeng MY, Kim Y-G, Cascalho M *et al*.
873 Humoral immunity in the gut selectively targets phenotypically virulent attaching-
874 and-effacing bacteria for intraluminal elimination. *Cell Host Microbe* 2015; **17**:
875 617–627.

876 52 Westerman LE, McClure HM, Jiang B, Almond JW, Glass RI. Serum IgG
877 mediates mucosal immunity against rotavirus infection. *Proc Natl Acad Sci* 2005;
878 **102**: 7268–7273.

879 53 Saito K, Suzuki R, Koyanagi Y, Isogai H, Yoneyama H, Isogai E. Inhibition of

880 enterohemorrhagic *Escherichia coli* O157:H7 infection in a gnotobiotic mouse
881 model with pre-colonization by bacteroides strains. *Biomed Reports* 2019; **10**:
882 175–182.

883 54 Amani J, Salmanian AH, Rafati S, Mousavi SL. Immunogenic properties of
884 chimeric protein from *espA*, *eae* and *tir* genes of *Escherichia coli* O157:H7.
885 *Vaccine* 2010; **28**: 6923–6929.

886 55 Kolling GL, Matthews KR. Export of virulence genes and Shiga toxin by
887 membrane vesicles of *Escherichia coli* O157:H7. *Appl Environ Microbiol* 1999;
888 **65**: 1843–1848.

889 56 Kim SH, Lee SR, Kim KS, Ko A, Kim E, Kim YH *et al*. Shiga toxin A subunit
890 mutant of *Escherichia coli* O157:H7 releases outer membrane vesicles
891 containing the B-pentameric complex. *FEMS Immunol Med Microbiol* 2010; **58**:
892 412–420.

893 57 Kunsmann L, Rüter C, Bauwens A, Greune L, Glüder M, Kemper B *et al*.
894 Virulence from vesicles: Novel mechanisms of host cell injury by *Escherichia coli*
895 O104:H4 outbreak strain. *Sci Rep* 2015; **5**:13252.

896 58 Bielaszewska M, Rü Ter 2 C, Bauwens A, Greune L, Jarosch K-A, Steil D *et al*.
897 Host cell interactions of outer membrane vesicle-associated virulence factors of
898 enterohemorrhagic *Escherichia coli* O157: Intracellular delivery, trafficking and
899 mechanisms of cell injury. *PLoS Pathog* 2017; **13**: e1006159.

900 59 Watanabe-Takahashi M, Yamasaki S, Murata M, Kano F, Motoyama J, Yamate
901 J *et al*. Exosome-associated Shiga toxin 2 is released from cells and causes
902 severe toxicity in mice. *Sci Rep* 2018; **8**: 10776.

903 60 Boysen A, Borch J, Krogh J, Hjernø K, Møller-Jensen J. SILAC-based
904 comparative analysis of pathogenic *Escherichia coli* secretomes. *J Microbiol*
905 *Methods* 2015; **116**: 66–79.

906 61 Premjani V, Tilley D, Gruenheid S, Le Moual H, Samis JA. Enterohemorrhagic
907 *Escherichia coli* OmpT regulates outer membrane vesicle biogenesis. *FEMS*
908 *Microbiol Lett* 2014; **355**: 185-192.

909 62 Babiuk S, Asper DJ, Rogan D, Mutwiri GK, Potter A a. Subcutaneous and
910 intranasal immunization with type III secreted proteins can prevent colonization
911 and shedding of *Escherichia coli* O157:H7 in mice. *Microb Pathog* 2008; **45**: 7–
912 11.

- 913 63 Gu J, Liu Y, Yu S, Wang H, Wang Q, Yi Y *et al.* Enterohemorrhagic *Escherichia*
914 *coli* trivalent recombinant vaccine containing EspA, intimin and Stx2 induces
915 strong humoral immune response and confers protection in mice. *Microbes*
916 *Infect* 2009; **11**: 835–841.
- 917 64 Zhang X-H, He K-W, Zhang S-X, Lu W-C, Zhao P-D, Luan X-T *et al.*
918 Subcutaneous and intranasal immunization with Stx2B-Tir-Stx1B-Zot reduces
919 colonization and shedding of *Escherichia coli* O157:H7 in mice. *Vaccine* 2011;
920 **29**: 3923–3929.
- 921 65 Iannino F, Herrmann CK, Roset MS, Briones G. Development of a dual vaccine
922 for prevention of *Brucella abortus* infection and *Escherichia coli* O157:H7
923 intestinal colonization. *Vaccine* 2015; **33**: 2248–2253.
- 924 66 Torres AG, Perna NT, Burland V, Ruknudin A, Blattner FR, Kaper JB.
925 Characterization of Cah, a calcium-binding and heat-extractable autotransporter
926 protein of enterohaemorrhagic *Escherichia coli*. *Mol Microbiol* 2002; **45**: 951–
927 966.
- 928 67 Urashima A, Sanou A, Yen H, Tobe T. Enterohaemorrhagic *Escherichia coli*
929 produces outer membrane vesicles as an active defence system against
930 antimicrobial peptide LL-37. *Cell Microbiol* 2017; **19**: 1–11.
- 931 68 Ralli-Jain P, Tifrea D, Cheng C, Pal S, de la Maza LM. Enhancement of the
932 protective efficacy of a *Chlamydia trachomatis* recombinant vaccine by
933 combining systemic and mucosal routes for immunization. *Vaccine* 2010; **28**:
934 7659–7666.
- 935 69 Vajdy M, Singh M, Ugozzoli M, Briones M, Soenawan E, Cuadra L *et al.*
936 Enhanced mucosal and systemic immune responses to *Helicobacter pylori*
937 antigens through mucosal priming followed by systemic boosting immunizations.
938 *Immunology* 2003; **110**: 86–94.
- 939 70 Kim Y, Ponomarenko J, Zhu Z, Tamang D, Wang P, Greenbaum J *et al.* Immune
940 epitope database analysis resource. *Nucleic Acids Res* 2012; **40**: 525–530.
- 941 71 Jespersen MC, Peters B, Nielsen M, Marcatili P. BepiPred-2.0: Improving
942 sequence-based B-cell epitope prediction using conformational epitopes. *Nucleic*
943 *Acids Res* 2017; **45**: W24–W29.
- 944 72 Kolaskar a S, Tongaonkar PC. A semi-empirical method for prediction of
945 antigenic determinants on protein antigens. *FEBS Lett* 1990; **276**: 172–174.

946 73 Ponomarenko J, Bui H-H, Li W, Fusseder N, Bourne PE, Sette A *et al.* ElliPro: a
947 new structure-based tool for the prediction of antibody epitopes. *BMC*
948 *Bioinformatics* 2008; **9**: 514.

949 74 Horst R, Stanczak P, Wüthrich K. NMR Polypeptide backbone conformation of
950 the *E. coli* outer membrane protein W. *Structure* 2014; **22**: 1204–1209.

951 75 Vandeputte-Rutten L, Kramer RA, Kroon J, Dekker N, Egmond MR, Gros P.
952 Crystal structure of the outer membrane protease OmpT from *Escherichia coli*
953 suggests a novel catalytic site. *EMBO J* 2001; **20**: 5033–5039.

954 76 Heras B, Totsika M, Peters KM, Paxman JJ, Gee CL, Jarrott RJ. The antigen 43
955 structure reveals a molecular Velcro-like mechanism of autotransporter-
956 mediated bacterial clumping. *Proc Natl Acad Sci U S A* 2014; **111**: 457–462.

957 77 Sánchez R, Šali A. Evaluation of comparative protein structure modeling by
958 MODELLER-3. *Proteins Struct Funct Genet* 1997; **29**: 50–58.

959 78 Ricker EW. Stoch and recruitrnt. *J Fish Res Board Canada* 1954; **11**: 559–
960 623.

961 79 Biller JR, Elajaili H, Meyer V, Rosen GM, Eaton SS, Eaton GR. Electron spin–
962 lattice relaxation mechanisms of rapidly-tumbling nitroxide radicals. *J Magn*
963 *Reson* 2013; **236**: 47–56.

964 80 Foloppe N, Mackerell AD. All-atom empirical force field for nucleic acids: I.
965 Parameter optimization based on small molecule and condensed phase
966 macromolecular target data. *J Comput Chem* 2000; **21**: 86–104.

967 81 Wiederstein M, Sippl MJ. ProSA-web: Interactive web service for the recognition
968 of errors in three-dimensional structures of proteins. *Nucleic Acids Res* 2007; **35**:
969 407–410.

970 82 Hebditch M, Carballo-Amador MA, Charonis S, Curtis R, Warwicker J. Protein-
971 Sol: A web tool for predicting protein solubility from sequence. *Bioinformatics*
972 2017; **33**: 3098–3100.

973 83 Wilkins MR, Gasteiger E, Bairoch A, Sanchez JC, Williams KL, Appel RD *et al.*
974 Protein identification and analysis tools in the ExPASy server. *Methods Mol Biol*
975 1999; **112**: 531–552.

976 84 Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC *et*
977 *al.* UCSF chimera - A visualization system for exploratory research and analysis.

978 *J Comput Chem* 2004; **25**: 1605–1612.

979 85 Mohawk KL, O'Brien AD. Mouse models of *Escherichia coli* O157:H7 infection
980 and shiga toxin injection. *J Biomed Biotechnol* 2011; **2011**: 258185.

981 86 Melton-Celsa AR, Rogers JE, Schmitt CK, Darnell SC, O'Brien AD. Virulence of
982 Shiga toxin-producing *Escherichia coli* (STEC) in orally-infected mice correlates
983 with the type of toxin produced by the infecting strain. *Jpn J Med Sci Biol* 1998;
984 **51 Suppl**: S108-14.

985 87 Lindgren SW, Melton AR, O'Brien AD. Virulence of enterohemorrhagic
986 *Escherichia coli* O91:H21 clinical isolates in an orally infected mouse model.
987 *Infect Immun* 1993; **61**: 3832–3842.

988 88 Wadolkowski E a., Burris J a., O'Brien a. D. Mouse model for colonization and
989 disease caused by enterohemorrhagic *Escherichia coli* O157:H7. *Infect Immun*
990 1990; **58**: 2438–2445.

991 89 Kurien BT, Hal Scofield R. Mouse urine collection using clear plastic wrap. *Lab*
992 *Anim* 1999; **33**: 83–86.

993 90 Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Lumley T *et al.* R
994 Package ' gplots '. 2016.[https://cran.r-](https://cran.r-project.org/web/packages/gplots/gplots.pdf)
995 [project.org/web/packages/gplots/gplots.pdf](https://cran.r-project.org/web/packages/gplots/gplots.pdf).

996 91 R Core Team. *R: A Language and Environment for Statistical Computing*. R
997 Foundation for Statistical Computing. Vienna, Austria., 2014 [http://www.r-](http://www.r-project.org)
998 [project.org](http://www.r-project.org).

999 92 Guruprasad K, Reddy BVB, Pandit MW. Correlation between stability of a
1000 protein and its dipeptide composition: A novel approach for predicting *in vivo*
1001 stability of a protein from its primary sequence. *Protein Eng Des Sel* 1990; **4**:
1002 155–161.

1003
1004
1005
1006
1007
1008
1009
1010

Figure Legends

Fig. 1. Experimental design for the development and evaluation of the chimeric-based STEC vaccine. **a)** We selected the OmpT, Cah and Hes proteins as suitable targets against STEC based on their antigenic properties, frequency of detection among clinical STEC strains and absence among commensal *E. coli* strains. **b)** The proteins were analyzed by immunoinformatics tools and a peptide microarray assay for B-cell epitope prediction and mapping, respectively. MHC-II epitopes were also predicted *in silico*. Several B-cell epitopes were validated by ELISA using a collection of HUS sera. **c)** Selected epitopes were used to design two chimeric proteins that were expressed and purified. The reactivity of the chimeric proteins to IgG and IgA present in HUS sera was also confirmed. **d)** BALB/c mice were immunized with different vaccine formulations by intramuscular or intranasal route using Imject Alum (IA) or Sigma adjuvant (SA), respectively. Systemic and local humoral responses were subsequently determined. The protection conferred by immunizations was evaluated in the streptomycin-treated mouse model by challenge with STEC O157:H7 and O91:H21 strains. Bacterial shedding and intestinal colonization were determined for the STEC O157:H7 infected mice. Renal damage was examined in the STEC O91:H21 infected mice.

Fig. 2. Identification and validation of linear B-cell epitopes of the OmpT, Cah and Hes proteins. **a, b and c)** Peptide microarray assays. Peptide slides containing 15, 12 and 10 AA peptides derived from de OmpT (a), Cah (b) and Hes (c) proteins were incubated with a mix of three HUS sera at a dilution of 1:100. After washing, staining was performed with secondary anti-human IgA DyLight800 antibody at a dilution of 1:1000. Control peptides (red spots) framing the peptide slides were staining with specific monoclonal DyLight680 antibody at a dilution of 1:2000. Control peptide slides incubated with anti-human IgA DyLight800 antibody and the specific monoclonal DyLight680 antibody are shown in the left panels. Epitope-like spot patterns are indicated by white arrows. **d-i)** Tukey box plots showing concentrations of IgG and IgA present in individual HUS sera (n=20) that are reactive to short peptides containing B-cell epitopes of the OmpT, Cah and Hes proteins. A lower number of HUS sera and Hes epitopes were assessed due to sera availability. Tukey box plots show the 25th to 75th percentiles, with the median indicated by the horizontal line inside the box. Data analysis was by Kruskal-Wallis test, followed by Dunn's multiple comparison test. *P < 0.05 was considered significant. **j and k)** Heatmaps show the logarithm of the IgG and IgA concentration of each serum (HUS or control sera) that recognizes a specific short peptide, respectively. Data were clustered hierarchically using Euclidean distance and complete linkage analyses. Each row represents a different serum and each column a specific epitope. The average (dotted line) and histogram (solid line) of the values obtained by each peptide are indicated in the columns. The color key indicates the value of the logarithm of the antibody concentration. The figure was made using the "gplots" ⁹⁰ package in R ⁹¹.

Figure 3. Design and production of the chimeric antigens. **a)** Predicted 3D structure of the Chimera 1 (Chi1) antigen. OmpT- and Hes-derived peptides are shown as indicated in the legend at the left. **b and c)** Quality validation of modeled Chi1 structure. Ramachandran plot (b) shows that 94.6% residues were in favored regions. Z-score plot (c) for Chi1 model obtained by ProSA-web.⁸¹ Dark blue and light blue regions represent z-scores of native protein structures determined by NMR and X-ray, respectively. Black spot shows z-score for the Chi1 model. **d)** Predicted 3D structure of the Chimera 2 (Chi2) antigen. OmpT- and Hes-derived peptides are shown as

indicated in the legend at the left. **e and f**) Quality validation of modeled Chi2 structure. Ramachandran plot (e) shows that 86.6% residues were in favored regions. Z-score plot (f) for Chi2 model. **g**) SDS-PAGE of purified Chi1 and Chi2 proteins. M, Molecular weight standard. **h-i**) Tukey box plots showing concentrations of IgG (h) and IgA (i) present in individual HUS sera (n=20) that are reactive to Chi1, Chi2 and α Cah proteins. Tukey box plots show the 25th to 75th percentiles, with the median indicated by the horizontal line inside the box. Data analysis was by Kruskal-Wallis test, followed by Dunn's multiple comparison test. *P < 0.05 was considered significant.

Figure 4. Humoral immune responses triggered by immunization with the chimeric antigens. Sera obtained at days -2, 14, 28 and 42 post-immunization were diluted 1:100 and used for the determination of specific IgG (a), IgA (b) and IgM (c) by ELISA. Fecal sIgA (d) was also determined from feces collected on days -2, 14, 28 and 42. The ELISA plates were coated with 100 μ l of each antigen (Chi1, Chi2 and an equimolar mix Chi1+Chi2) at a final concentration of 1 μ g/ml in carbonate-bicarbonate buffer (pH 9.6) and incubated overnight at 4 °C. Then, the plates were blocked, washed and incubated with different dilutions of each serum. The results are expressed as means $\pm$ SD of absorbance values at 450 nm (A_{450}), which were obtained from individual sera or fecal suspensions of five mice per group. Experimental groups are shown as indicated by legend at the bottom. Data analysis was by a two-way ANOVA, followed by Tukey's multiple comparison test. P < 0.05 was considered significant. Asterisks (*) indicate significant differences between the immunized mice and the PBS control group. Number signs (#) indicate significant differences between administration routes.

Figure 5. Protection conferred by immunizations with the chimeric antigens. a and b) Determination of fecal shedding of STEC O157:H7. Eight mice per group were orally inoculated with 10⁹ CFU of the challenge strain. Fecal pellets were collected daily, weighed, homogenized, and plated on MacConkey agar containing streptomycin. Data are showed as the number of CFU of the challenge strain per 100 mg feces. Error bars represent the standard deviations (s.d.). Differences between immunized mice and the PBS control group were analyzed by a two-way ANOVA with Tukey's multiple comparison test. Experimental groups are shown as indicated by legend at the bottom. **c and d)** Tukey box plots showing creatinine concentrations (mg/dl) in urine determined from five mice per group on days 7 (c) and 12 (d) post-infection with STEC O91:H21. Differences between experimental groups and uninfected mice were analyzed by Mann-Whitney U test. Dotted green line indicates normal creatinine concentration of 1.5 mg/dl. **e)** Histopathology analysis from kidney tissue obtained from five mice per group on day 12 post-infection with STEC O91:H21. Cellular injuries were classified as not evident, mild, moderate or severe as described in the Methods. Cellular injuries are color coded as indicated in the legend at the top. Differences between experimental groups and uninfected mice were analyzed by a two-way ANOVA followed by Tukey's multiple comparison test. For all statistical analyses P < 0.05 was considered significant.

(a) Antigens

- Reactive to IgG and IgA from HUS sera
- Expressed during human infection
- Widespread among clinical STEC strains
- Mainly absent in commensal *E. coli* strains

(b) Identification and validation of epitopes

- Epitope mapping

Table 1. Linear B-cell epitopes identified in the OmpT, Cah and Hes proteins

Protein	Epitope name	Epitope Mapping	<i>In silico</i> B-cell prediction tools			Reactivity to HUS sera by ELISA. No (%)		Reactivity to control sera by ELISA. No (%)	
			BepiPred 2.0	Kolaskar and Tongaonker	Ellipro	IgG	IgA	IgG	IgA
OmpT	OmpT_1	Yes	Yes	No	Yes	17/20 (85)	5/7 (71)	0/3 (0)	0/3 (0)
	OmpT_2	Yes	Yes	No	Yes	18/20 (90)	5/7 (71)	0/3 (0)	0/3 (0)
	OmpT_3	Yes	Yes	Yes	Yes	16/20 (80)	6/7 (86)	0/3 (0)	0/3 (0)
	OmpT_4	Yes	Yes	Yes	Yes	18/20 (90)	3/7 (43)	0/3 (0)	0/3 (0)
	OmpT_5	Yes	Yes	No	Yes	16/20 (80)	4/7 (57)	0/3 (0)	0/3 (0)
	OmpT_6	Yes	Yes	No	Yes	18/20 (90)	5/7 (71)	0/3 (0)	0/3 (0)
Cah	Cah_1	Yes	Yes	Yes	Yes	18/20 (90)	3/7 (43)	0/3 (0)	0/3 (0)
	Cah_2	Yes	No	No	No	16/20 (80)	2/7 (29)	0/3 (0)	0/3 (0)
	Cah_3	Yes	Yes	No	Yes	19/20 (95)	5/7 (71)	0/3 (0)	0/3 (0)
	Cah_4	Yes	No	No	No	16/20 (80)	5/7 (71)	0/3 (0)	0/3 (0)
	Cah_5	Yes	No	No	No	20/20 (100)	4/7 (57)	0/3 (0)	0/3 (0)
	Cah_6	Yes	No	No	No	20/20 (100)	6/7 (86)	0/3 (0)	0/3 (0)
Hes	Hes_1	Yes	Yes	No	Yes	14/20 (70)	5/7 (71)	0/3 (0)	0/3 (0)
	Hes_2	Yes	Yes	Yes	Yes	17/20 (85)	5/7 (71)	0/3 (0)	0/3 (0)
	Hes_3	Yes	Yes	No	Yes	14/20 (70)	N.E.	0/3 (0)	N.E.
	Hes_4	Yes	No	No	Yes	12/20 (60)	N.E.	0/3 (0)	N.E.
	Hes_5	Yes	Yes	No	Yes	20/20 (100)	7/7 (100)	0/3 (0)	0/3 (0)

N.E. Not evaluated

Table 2. Epitopes and chemical and physical properties of the chimeric proteins

Protein	B-cell epitopes	Predicted T-cell epitopes	MW (kDa)	Theoretical pI	Solubility ¹	Estimated half-life ²	Instability index ³
Chimera 1	OmpT_1, OmpT_2, OmpT_3, OmpT_4, Hes_1, Hes_2, Hes_3 and Hes_5	Two of OmpT	29.6	4.51	0.693	>10 hours	21.30
Chimera 2	OmpT_1, OmpT_6, Cah_1, Cah_2, Cah_3, Cah_4, Cah_5, Cah_6, Hes_1, Hes_2 and Hes_5	Two of Cah	56.7	4.52	0.666	>10 hours	8.02

¹ Predicted by Protein-Sol tool ⁶⁹. Scaled solubility value (0-1). A value greater than 0.45 predicts that the protein is soluble.

² Prediction of the time it takes for half of the amount of protein in *E. coli* to disappear after its synthesis. ³ The instability index provides an estimate of the stability of a protein in a test tube. A protein whose instability index is smaller than 40 is predicted as stable ⁷⁷.

Table 3. Intestinal colonization of STEC O157:H7 and protection conferred by immunization with chimeric proteins

Chimeric protein + Adjuvant	Log10 CFU of STEC O157:H7 in Cecum (mean $\pm$ SD)	Log10 units of protection
Chi1 + Imject Alum	5.45 $\pm$ 0.24	0
Chi1 + Sigma Adjuvant	4.70 $\pm$ 0.56	0.68
Chi2 + Imject Alum	5.08 $\pm$ 0.54	0.30
Chi2 + Sigma Adjuvant	4.70 $\pm$ 0.86	0.68
Chi1 + Chi2 + Imject Alum	3.28 $\pm$ 0.10	2.10 *
Chi1 + Chi2 + Imject Alum	4.94 $\pm$ 0.81	0.44
PBS + Imject Alum + Sigma Adjuvant	5.38 $\pm$ 0.24	0

* P < 0.001 as compared to the PBS control group.

[illegible]