Emergence of *optrA*-mediated linezolid resistance in multiple lineages and plasmids of *Enterococcus faecalis* revealed by long read sequencing

Martin P McHugh^{1,2*}, Benjamin J Parcell^{1,3,#a}, Kerry A Pettigrew^{1,#b}, Geoff Toner², Elham Khatamzas^{2,#c}, Anne Marie Karcher^{3,#a}, Joanna Walker³, Robert Weir⁴, Danièle Meunier⁵, Katie L Hopkins⁵, Neil Woodford⁵, Kate E Templeton², Stephen H Gillespie¹, Matthew TG Holden^{1*}

¹School of Medicine, University of St Andrews, St Andrews, UK; ²NHS Lothian Infection Service, Royal Infirmary of Edinburgh, Edinburgh, UK; ³Medical Microbiology, Aberdeen Royal Infirmary, Aberdeen, UK; ⁴Medical Microbiology, Forth Valley Royal Hospital, Larbert, UK; ⁵Antimicrobial Resistance and Healthcare Associated Infections (AMRHA) Reference Unit, National Infection Service, Public Health England, London, UK

^{#a} Present address: Medical Microbiology, Ninewells Hospital, Dundee, UK

^{#b} Present address: Bristol Medical School, University of Bristol, Bristol, UK

^{#c} Present address: Department of Medicine III, University Hospital, LMU Munich, Germany

***Corresponding authors**

Matthew Holden mtgh@st-andrews.ac.uk

Martin McHugh mpm20@st-andrews.ac.uk

ABSTRACT

Objectives

To characterise the genetic environment of *optrA* in linezolid-resistant *Enterococcus faecalis* isolates from Scotland.

Methods

Linezolid-resistant *E. faecalis* were identified in three Scottish Health Boards and confirmed to carry the *optrA* gene at the national reference laboratory. WGS was performed with short read (Illumina MiSeq) and long read (Oxford Nanopore MinION) technologies to generate complete genome assemblies. Illumina reads for 94 *E. faecalis* bloodstream isolates were used to place the *optrA*-positive isolates in a larger UK phylogeny.

Results

Six *optrA*-positive linezolid-resistant *E. faecalis* were isolated from urogenital samples in three Scottish Health Boards (2014-2017). No epidemiological links were identified between the patients, four were community-based, and only one had recent linezolid exposure. Reference-based mapping confirmed the isolates were genetically distinct (>13,900 core SNPs). *optrA* was located on a plasmid in each isolate and these plasmids showed limited nucleotide similarity. There was variable presence of transposable elements surrounding *optrA*, (including IS1216, IS3, and Tn3) and not always as a recognisable gene cassette. *OptrA* amino acid sequences were also divergent, resulting in four protein variants differing in 1-20 residues. One isolate belonged to ST16 and clustered with three other isolates in the UK collection (76-182 SNPs), otherwise the *optrA*-positive isolates were genetically distinct from the bloodstream isolates (>6,000 SNPs).

Conclusions

50 We report multiple variants of the linezolid resistance gene *optrA* in diverse *E.*
51 *faecalis* strain and plasmid backgrounds, suggesting multiple introductions of the
52 gene into the *E. faecalis* population and selection driving recent emergence.

53

INTRODUCTION

Enterococcus faecalis and *Enterococcus faecium* are carried in the intestinal tract and are important opportunistic pathogens in humans.¹ Treatment of enterococcal infections is challenging due to intrinsic or acquired resistance to multiple antimicrobials including aminoglycosides, benzylpenicillin, cephalosporins, fluoroquinolones, macrolides, tetracyclines, and trimethoprim. Among the remaining treatment options, clinical *E. faecium* isolates are usually resistant to amoxicillin and resistance to vancomycin is increasingly common.² In contrast, *E. faecalis* typically remains susceptible to amoxicillin and vancomycin but can acquire significant resistance and has been implicated in the transfer of antimicrobial resistance genes to other Gram-positive pathogens, for example transmitting *vanA*-mediated vancomycin resistance to methicillin-resistant *Staphylococcus aureus*.³

The main treatment options for multi-drug resistant Gram-positive bacteria are the oxazolidinones linezolid or tedizolid, or the lipopeptide daptomycin. Daptomycin therapy is challenging due to significant side effects, limited efficacy in pulmonary infections, uncertain dosing regimens, and challenges with *in vitro* susceptibility determination.⁴ Linezolid blocks protein synthesis by binding to the 50S ribosomal subunit and inhibiting formation of the initiation complex.⁵ Linezolid resistance is uncommon, reported in $\leq 1\%$ of bloodstream enterococcal isolates in the UK.^{6,7} The G2576T mutation in the 23S rRNA genes can arise *de novo* during extended linezolid therapy,⁸ although strict infection control and antimicrobial stewardship have been successful in limiting incidence.⁹ The methyltransferases Cfr and Cfr(B), and ABC-F ribosomal protection proteins OptrA and PoxA also confer resistance to linezolid but are carried on mobile genetic elements, raising the prospect of rapid spread of linezolid resistance across genetically distinct lineages.^{10–12} In 2015, *optrA*

was first reported as conferring resistance to oxazolidinones and phenicols.¹³ Recent international surveillance shows that although linezolid resistance remains rare, *optrA* has spread to every continent and is the dominant mechanism of linezolid resistance in *E. faecalis*.¹⁴ Surveillance has also detected *optrA* in the UK.¹⁵ Studies into the genetic context of *optrA* have identified the gene on both the chromosome and plasmids, often associated with insertion sequence IS1216, a possible explanation for the rapid spread of *optrA*.^{16,17} However, few studies have generated complete genome assemblies of *optrA*-carrying *E. faecalis*, which would provide high precision information on the genetic context of *optrA*.

Here, we investigate the epidemiological and clinical background of *optrA*-carrying *E. faecalis* isolates from human clinical samples collected in Scotland. We used whole genome sequencing to determine whether these isolates represent transmission of a single clonal lineage. We hypothesised the spread of *optrA* is driven by a single mobile genetic element, and to investigate this we made hybrid assemblies of short and long read sequencing data to generate complete genomes and to reconstruct the genetic environment of *optrA*. This study describes the first use of nanopore-based long read sequencing to investigate *optrA*-containing mobile genetic elements.

MATERIALS AND METHODS

Bacterial strains

Isolates were selected for this study based on the presence of the *optrA* gene as determined by Public Health England's Antimicrobial Resistance and Healthcare Associated Infections (AMRHAI) Reference Unit, either as part of non-structured retrospective screening of stored isolates (prior to 2016) or as part of the reference

laboratory service (2016 onwards). Isolates were originally collected in three Scottish Health Boards, and as such represent a subset of Scottish *optrA*-positive isolates identified by AMRHAI. Linezolid- and chloramphenicol-resistant *E. faecalis* were isolated from six clinical samples (Table 1) using standard methods and identified with matrix-assisted laser-desorption ionisation time-of-flight mass spectrometry or the Vitek-2 GP-ID card (bioMérieux, Marcy L'Etoile, France). Antimicrobial susceptibility testing was performed with the Vitek-2 AST-607 card and interpreted with EUCAST breakpoints.¹⁸ Isolates were referred to AMRHAI for characterisation of linezolid resistance mechanisms. Detection of the G2576T mutation (*Escherichia coli* numbering) in the 23S rRNA genes was investigated by PCR-RFLP and, from 2016, by a real-time PCR-based allelic discrimination assay.^{19,20} The *cfr* and *optrA* genes were sought by a multiplex PCR using primers for the detection of *cfr* (*cfr-fw*: 5'-TGA AGT ATA AAG CAG GTT GGG AGT CA-3' and *cfr-rev*: 5'-ACC ATA TAA TTG ACC ACA AGC AGC-3')²¹ and for the detection of *optrA* (*optrA-F*: 5'-GAC CGG TGT CCT CTT TGT CA-3' and *optrA-R*: 5'-TCA ATG GAG TTA CGA TCG CCT-3') (AMRHAI, unpublished data).

An *E. coli* transformant harbouring a plasmid bearing *cfr* (kindly provided by Pr S. Schwarz) was used as a control strain for the detection of *cfr*. This was replaced from 2016 by *Staphylococcus epidermidis* NCTC 13924 harbouring both *cfr* and the G2576T mutation. *E. faecium* NCTC 13923 was used as a control strain for the detection of *optrA*.

Access to isolates and clinical data was approved by the NHS Scotland Biorepository Network (Ref TR000126).

Whole genome sequencing and genomic analysis

Single colonies were inoculated into brain heart infusion broth (Oxoid, Basingstoke, UK) and incubated overnight at 37°C. Genomic DNA was extracted from cell pellets using the Wizard Genomic DNA Purification Kit (Promega, Wisconsin, USA), or QiaSymphony DSP DNA Mini Kit (Qiagen, Hilden, Germany). Short read barcoded libraries were prepared using the Nextera XT kit (Illumina, San Diego, USA) and sequenced with a MiSeq instrument (Illumina) using 250 bp paired-end reads on a 500-cycle v2 kit. Short reads were quality trimmed with Trimmomatic v0.36 and settings [LEADING:5 TRAILING:5 SLIDINGWINDOW:4:15 MINLEN:100].²² Barcoded long read libraries were generated with the 1D Ligation Sequencing Kit (Oxford Nanopore Technologies, Oxford, UK) and sequenced with an R9.4 flow cell on a MinION sequencer (Oxford Nanopore Technologies). Base-calling and barcode de-multiplexing was performed with Albacore v2.1.3 (Oxford Nanopore Technologies) and the resulting fast5 files converted to fastq with Poretools v0.6.0,²³ or basecalled and de-multiplexed with Albacore v2.3.3 with direct fastq output. Porechop v0.2.3 (<https://github.com/rrwick/Porechop>) was used to remove chimeric reads and trim adapter sequences. The data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB36950 (<https://www.ebi.ac.uk/ena/data/view/PRJEB36950>).

To generate a UK-wide *E. faecalis* phylogenetic context for the *optrA*-positive isolates, raw sequence data was downloaded from the ENA (www.ebi.ac.uk/ena) under study accession numbers PRJEB4344, PRJEB4345, and PRJEB4346.²⁴ Short reads were mapped to the *E. faecalis* reference genome V583 (accession number AE016830) using SMALT v0.7.4.²⁵ Mapped assemblies were aligned and regions annotated as mobile genetic elements in the V583 genome (transposons, integrases, plasmids, phages, insertion sequences, resolvases, and recombinases; tab file of

regions available in Table S1) were removed from the assembly
(https://github.com/sanger-pathogens/remove_blocks_from_aln). All sites in the
alignment with single nucleotide polymorphisms (SNPs) were extracted using SNP-
sites v2.4.0²⁶ and a phylogeny was created from the core-genome SNP alignment
using RAxML v8.2.8²⁷ with 100 bootstrap replicates and visualised with iTOL.²⁸
Recombination blocks were removed from ST16 isolates using Gubbins v1.4.10.²⁹
Hybrid assembly was performed with Illumina short reads and Nanopore long
reads using Unicycler v0.4.7³⁰ in standard mode. The resulting assemblies were
annotated with Prokka v1.5.1 using a genus specific RefSeq database.³¹ Hybrid
assemblies were checked for indel errors using Ideel
(<https://github.com/mw55309/ideel>) and UniProtKB TrEMBL database v2019_1.
Plasmid comparisons were generated and visualised with EasyFig v2.2.2³² and
BRIG v0.95.³³
MLST typing was performed using SRST2 v0.2.0³⁴ and the *E. faecalis* MLST
database (<https://pubmlst.org/efaecalis/>) sited at the University of Oxford.^{35,36}
Antimicrobial resistance mechanisms were detected using ARIBA v2.12.1³⁷ and the
ResFinder database v3.0³⁸ with the addition of linezolid resistance mutations in the
23S rRNA (G2505A and G2576T based on *E. coli* numbering).

RESULTS AND DISCUSSION

Detection of optrA-positive E. faecalis

Six *E. faecalis* isolated from urogenital samples were initially identified as linezolid-
and chloramphenicol-resistant in routine diagnostic laboratories and confirmed to
carry *optrA* at the AMRHA Reference Unit (Table 1). The earliest isolates in this

collection were from the Grampian region of Scotland in 2014, 2015, and 2016. Three more isolates were identified in 2017 from other regions of Scotland (Forth Valley and Lothian, Table 1), with no clear epidemiological links between the patients. Prior to isolation of *optrA*-positive *E. faecalis*, patients 1-3 were treated with trimethoprim for recurrent urinary tract infections, with patient 3 also receiving cefalexin. Patients 5 and 6 were managed in general practice and it was not possible to determine their antimicrobial exposure. Patient 4 was the only patient with known exposure to linezolid, a two-week course prior to the isolation of *optrA*-positive *E. faecalis*. Patient 4 was a surgical inpatient and another patient on the same ward had *optrA*-positive *E. faecalis* isolated from an abdominal wound, indicating possible transmission between this patient and patient 4. The *optrA*-positive *E. faecalis* from the contact of patient 4 was not available for study. Further screening of the ward environment and patients found no further linezolid-resistant enterococci or staphylococci over a two-month period, although the contact of patient 4 continued to have *optrA*-positive *E. faecalis* isolated from their abdominal wound for a month until discharge.

***optrA* is carried by distinct strains**

Whole genome sequencing was performed to investigate the genetic relationship between the isolates. *In silico* MLST showed the six isolates belonged to different sequence types (STs), suggesting they were genetically distinct (Table 1). To further confirm this, we analysed SNPs in the core genomes of the *optrA*-positive isolates and found the isolates differed by a median 18,806 SNPs (range 13,909 – 22,272). Previous estimates suggest a genetic diversification rate of 2.5-3.4 SNPs/year for *E. faecalis*, highlighting the *optrA*-positive strains share a very distant common ancestor.²⁴

***optrA* is carried on diverse plasmids**

We then examined the genetic context of *optrA* in each isolate. Initial *de novo* assembly of short-read data generated fragmented assemblies (72-135 contigs, mean N50 198 kb), but with *optrA* present on moderate sized contigs (11-44 kb). Three *optrA*-positive contigs carried plasmid-associated replication or transfer genes, but none represented a complete plasmid, or had increased read depth coverage compared to core genes indicative of being multicopy. Therefore, it was unclear if *optrA* was carried on plasmids (often present in multiple copies within a cell) or the chromosome, and how similar these regions were between the six isolates. To resolve repetitive regions and try to complete the genome assemblies we utilised Nanopore sequencing to generate long reads, and then combined these with Illumina short reads to produce high quality hybrid assemblies. In four of the isolates completed genomes were obtained, with the other two generating near-complete genomes (Table S2). Analysis of the six hybrid assemblies showed <3 % putative coding sequences were shorter than the closest reference match (Table S2) indicating the hybrid assembly process removed most indel errors and the short coding sequences were likely to be true pseudogenes.³⁹ The hybrid assemblies contained between one and three plasmids ranging in size from 11-80 kb, with *optrA* present on a single complete plasmid in each isolate (Table 2).

In general, the *optrA*-positive plasmids had limited sequence identity, although pBX5936-1 (69 kb) and pTM6294-2 (53 kb) had 97% average nucleotide identity over 40 kb aligned sequence. These two plasmids also had several unique regions indicating the plasmids shared a backbone but had distinct additional content (Figure S1). *optrA* and the phenicol resistance gene, *fexA*, were located in the same orientation and within 550-750 nucleotides of each other, with a short (~200

nucleotides) hypothetical coding sequence in the intervening region (Figure 1). Additionally, limited similarity was seen between the Scottish *optrA*-positive plasmids and the first identified *optrA*-positive plasmid from China (pE394, accession KP399637; Figure S2).

A number of insertion sequence transposases were identified in the *optrA*-positive plasmids, although we were unable to identify many beyond the family level due to limited matches in public databases (Table 2). We found evidence of IS1216 in all the *optrA*-positive plasmids, although only pBX5936-1 and pBX8117-2 had IS1216 flanking the *optrA* and *fexA* region as a cassette (Figure 1). BLASTn comparison of pWE0254-1 with the other *optrA*-positive plasmids highlighted a partial IS1216 transposase that was not identified by automated annotation. Immediately upstream of the partial IS1216 was an IS3-family transposase, the insertion of which likely disrupted the IS1216 (Figure S1). pWE0438 had Tn3-family transposases surrounding *optrA* and *fexA*, as well genes encoding resolvases, which may represent a transposable unit (Figure 1). pWE0851-1 carried one IS1216 transposase upstream and one Tn3-transposase downstream of *optrA/fexA*, but also had multiple copies of IS3-family transposases throughout the plasmid so multiple possible mechanisms of transposition exist (Table 2, Figure S1). pTM6294-2 had one IS1216 transposase and one ISL3-family transposase surrounding *optrA/fexA*. The variable presence of IS1216 in these isolates suggest other means of transposition may also be important in the spread of *optrA*, including IS3-family and Tn3-family transposases.

***optrA* sequences vary between isolates**

Comparison of the OptrA amino acid sequence from each isolate revealed different variants of the resistance protein: two isolates had the same sequence as the first identified OptrA from pE394, BX5936 had a single substitution, WE0851 had two substitutions, WE0348 had three substitutions, and BX8117 had 20 substitutions (Table 2). BX5936 and BX8117 had novel OptrA sequences not yet described in the literature. The OptrA sequence from BX8117 was similar to E35048 detected in an *E. faecium* isolated in Italy in 2015 with the two OptrA sequences differing at three amino acid positions.⁴⁰ Over 40 OptrA sequence variants have been described, although the role of this sequence variation is unclear as they do not significantly differ in their linezolid minimum inhibitory concentration *in vitro*.^{41,42}

The degree of sequence variation between the six FexA proteins was less than that seen in OptrA. Comparison to the first reported FexA (AJ549214) showed four common variants in all strains (A34S, L39S, I131V, and V305I), with all but BX8117 having an additional D50A variant. This suggests there is a more diverse background of *optrA* sequences compared to *fexA*, and/or there is ongoing diversifying selective pressure applied only to *optrA* despite the close genetic linkage of the two genes.

Of note, all six isolates had an inferred OptrA sequence 18 amino acids shorter than most public sequences. Inclusion of the 18 upstream amino acids showed that all six isolates had an M1L variant compared to OptrA_{pE394}. We believe this is an artefact introduced during coding sequence prediction. The first *optrA* genes were identified *in silico* using ORFfinder (<https://www.ncbi.nlm.nih.gov/orffinder>), which detects putative coding sequences based on the presence of in-frame start and stop codons only. We used Prokka for genome annotation which implements Prodigal to score potential coding sequences

based on start/stop codon position, coding sequence length, and upstream promoter regions and outputs the highest confidence coding sequences. Indeed, most published *optrA* sequences start with nucleotide codons TTG (usually encoding leucine), but the corresponding amino acid sequences start with methionine indicating this codon has been designated as a start codon. The only other report of the M1L variant is from a study that also used Prokka for annotation.⁴³ Given the possible effect of methodology on identification of the first amino acid we have not reported the M1L variant in our results but mention it here for completeness. The true *optrA* start codon should be confirmed to aid ongoing surveillance efforts.

***optrA*-positive strains are distantly related to bloodstream isolates**

To investigate whether the *optrA*-positive isolates represented common *E. faecalis* strains in the UK, publicly available sequence data of 94 *E. faecalis* isolates from the British Society for Antimicrobial Chemotherapy (BSAC) bacteraemia surveillance programme (isolated between 2001 and 2011) were analysed together with the six known *optrA*-positive isolates.²⁴ We first looked for determinants of linezolid resistance in the 94 sequences, and found no evidence of *cfr*, *cfr*(B), *cfr*(D), *optrA*, *poxA*, or the G2505A 23S rRNA gene mutation. Only one of the BSAC isolates (accession ERS324700) carried the G2576T 23S rRNA gene mutation conferring linezolid resistance. Core genome phylogeny showed BX8117 was related to three other ST16 isolates from the UK, after removal of putative recombination blocks there were 76, 81, and 182 SNPs between these isolates suggesting they diverged from a common background but are not linked to recent transmission (Figure 2). ST16 has been associated with multidrug-resistant infections in humans and animals, highlighting the potential for the emergence of linezolid resistance in invasive enterococcal infections.⁴⁴ The other five *optrA*-positive isolates have no

close genetic links in this phylogeny (minimum pairwise SNPs 12,314 – 17,891). Our study is not designed to infer patterns across Scotland and the rest of the UK, but our findings suggest the *optrA*-positive isolates are generally distinct from those recently causing bloodstream infections in the UK.

***optrA*-positive *E. faecalis* harbour multiple resistance mechanisms**

Looking at all assembled plasmids, the isolates carried genes conferring resistance to aminoglycosides (*ant*(6)-Ia, *aph*(3')-IIIA, *aac*(6')-Ie-*aph*(2'')-Ia, and others listed in Table 2), chloramphenicol (*catA8*), bacitracin (*bcrA*), macrolides (*ermA*-like, *ermB*), tetracyclines (*tet*(L), *tet*(M)), trimethoprim (*dfrG*), and the heavy metals cadmium (*cadA*) and copper (*copZ*). However, the pattern of carried genes differed between isolates with only *optrA* and *fexA* found in all isolates (Table 2).

pBX8117-2 carried a gene with 100% nucleotide identity and coverage to *cfr*(D) from *Enterococcus faecium* isolated in France in 2015, and in Australia in 2019.^{45,46} In both isolates, *optrA* and *cfr*(D) genes were present on different contigs based on short-read sequencing. Our study is the first to detect *cfr*(D) in *E. faecalis* and using hybrid assembly we identified co-carriage of *optrA* and *cfr*(D) on the same plasmid. The French and Australian *cfr*(D)-positive isolates also carried *vanA*-type vancomycin resistance genes, although the Australian isolate was phenotypically vancomycin sensitive due to the loss of the regulatory genes *vanR* and *vanS*. At present, no *in vitro* work has described the impact of *cfr*(D) on antimicrobial resistance in enterococci so it is unclear whether or not *cfr*(D) confers the PhLOPS_A multiresistance phenotype originally described with Cfr.⁴⁷

There is evidence of *optrA* being more common in particular *E. faecalis* lineages, with ST16, ST330, ST480, and ST585 in particular being described here

and in other studies.^{14,48–50} These *optrA*-positive lineages are not specific to one host species and have been isolated from humans, animals, and the environment.^{43,51} Florfenicol use in food animals is associated with the presence of *optrA* in animal waste and the environment surrounding livestock farms.^{52,53} Additionally, *optrA*-positive enterococci and staphylococci have been isolated from raw food purchased from retail stores in China, Columbia, Denmark, and Tunisia.^{51,54–56} Wu *et al.* (2019) found evidence of transmission of *optrA*-positive *E. faecalis* from raw meat to a dog in China.⁵⁶ However, the incidence of *optrA*-positive isolates in raw foods was low in the available studies, and there is currently no direct evidence to suggest *optrA*-positive strains are transmitted to humans via the food chain.^{57,58} Increasing use of linezolid in human medicine may also select for *optrA*-positive strains, and once carried in the gut may be co-selected by other antimicrobials given the multidrug resistance phenotype of these isolates. The role of antimicrobial use, animal contact, food hygiene, and the environment in transmission of *optrA*-positive strains should be investigated further.

Our finding that *optrA* is present as different gene variants, carried on different mobile genetic elements, in unrelated strains of *E. faecalis* suggest a diverse *optrA* reservoir that is only partly investigated in this study. As well as *optrA*, the *cfr* and *poxA* genes are emerging transferable linezolid resistance mechanisms. Further studies from a One Health perspective are warranted to understand the selection pressures driving transferable linezolid resistance, and the transmission dynamics of these strains to avoid further spread of linezolid resistance within *E. faecalis* and other Gram-positive bacteria.

ACKNOWLEDGEMENTS

The authors would like to thank the Bioinformatics Unit at the University of St Andrews and Pathogen Informatics at the Wellcome Sanger Institute for access to high performance computing clusters.

FUNDING

This work was supported by the Chief Scientist Office (Scotland) through the Scottish Healthcare Associated Infection Prevention Institute (Reference SIRON/10).

TRANSPARENCY DECLARATION

The authors report no conflicts of interest related to this work.

REFERENCES

1. García-Solache M, Rice LB. The Enterococcus: A model of adaptability to its environment. *Clin Microbiol Rev* 2019; **32**: e00058-18.
2. ECDC. Surveillance of antimicrobial resistance in Europe – Annual report of the European Antimicrobial Resistance Surveillance Network (EARS-Net) 2017. 2018. Available at: <http://www.ecdc.europa.eu/sites/portal/files/documents/EARS-Net-report-2017-update-jan-2019.pdf>. Accessed September 19, 2019.
3. Weigel LM, Clewell DB, Gill SR, *et al*. Genetic analysis of a high-level vancomycin-resistant isolate of *Staphylococcus aureus*. *Science* 2003; **302**: 1569–71.
4. Humphries RM. The new, new daptomycin breakpoint for *Enterococcus* spp. *J Clin Microbiol* 2019; **57**: e00600-19.
5. Zahedi Bialvaei A, Rahbar M, Yousefi M, *et al*. Linezolid: a promising option in the treatment of Gram-positives. *J Antimicrob Chemother* 2017; **72**: 354–64.
6. Public Health England. *English surveillance programme for antimicrobial utilisation and resistance (ESPAUR) Report 2018-2019*. London, UK: PHE; 2019. Available at:

- 376 <https://www.gov.uk/government/publications/english-surveillance-programme-antimicrobial->
377 [utilisation-and-resistance-espaur-report.](https://www.gov.uk/government/publications/english-surveillance-programme-antimicrobial-)
- 378 7. Health Protection Scotland. *Scottish One Health Antimicrobial Use and Resistance in*
379 *2018 Annual Report*. Glasgow, UK: HPS; 2019. Available at:
380 <https://www.hps.scot.nhs.uk/web-resources-container/scottish-one-health-antimicrobial-use->
381 [and-antimicrobial-resistance-in-2018/](https://www.hps.scot.nhs.uk/web-resources-container/scottish-one-health-antimicrobial-use-).
- 382 8. Mendes RE, Deshpande LM, Jones RN. Linezolid update: Stable *in vitro* activity following
383 more than a decade of clinical use and summary of associated resistance mechanisms.
384 *Drug Resist Updat* 2014; **17**: 1–12.
- 385 9. Abbo L, Shukla BS, Giles A, *et al*. Linezolid and vancomycin-resistant *Enterococcus*
386 *faecium* in solid organ transplant recipients: Infection control and antimicrobial stewardship
387 using whole genome sequencing. *Clin Infect Dis* 2019; **69**: 259–65.
- 388 10. Antonelli A, D’Andrea MM, Brenciani A, *et al*. Characterization of *poxtA*, a novel
389 phenicol–oxazolidinone–tetracycline resistance gene from an MRSA of clinical origin. *J*
390 *Antimicrob Chemother* 2018; **73**: 1763–9.
- 391 11. Deshpande LM, Ashcraft DS, Kahn HP, *et al*. Detection of a new *cfr*-like gene, *cfr*(B), in
392 *Enterococcus faecium* isolates recovered from human specimens in the United States as
393 part of the SENTRY Antimicrobial Surveillance Program. *Antimicrob Agents Chemother*
394 2015; **59**: 6256–61.
- 395 12. Diaz L, Kiratisin P, Mendes RE, *et al*. Transferable Plasmid-Mediated Resistance to
396 Linezolid Due to *cfr* in a Human Clinical Isolate of *Enterococcus faecalis*. *Antimicrob Agents*
397 *Chemother* 2012; **56**: 3917–22.
- 398 13. Wang Y, Lv Y, Cai J, *et al*. A novel gene, *optrA*, that confers transferable resistance to
399 oxazolidinones and phenicols and its presence in *Enterococcus faecalis* and *Enterococcus*
400 *faecium* of human and animal origin. *J Antimicrob Chemother* 2015; **70**: 2182–90.
- 401 14. Deshpande LM, Castanheira M, Flamm RK, *et al*. Evolving oxazolidinone resistance
402 mechanisms in a worldwide collection of enterococcal clinical isolates: Results from the
403 SENTRY Antimicrobial Surveillance Program. *J Antimicrob Chemother* 2018; **73**: 2314–22.
- 404 15. Health Protection Scotland. Oxazolidinone-resistance due to *optrA* in *Enterococcus*
405 *faecalis*. *HPS Wkly Rep* 2016; **50**: 230–1.
- 406 16. Cai J, Wang Y, Schwarz S, *et al*. High detection rate of the oxazolidinone resistance
407 gene *optrA* in *Enterococcus faecalis* isolated from a Chinese anorectal surgery ward. *Int J*
408 *Antimicrob Agents* 2016; **48**: 757–9.
- 409 17. He T, Shen Y, Schwarz S, *et al*. Genetic environment of the transferable
410 oxazolidinone/phenicol resistance gene *optrA* in *Enterococcus faecalis* isolates of human
411 and animal origin. *J Antimicrob Chemother* 2016; **71**: 1466–73.
- 412 18. EUCAST. *Breakpoint tables for interpretation of MICs and zone diameters*. 2018.
- 413 19. Woodford N, Tysall L, Auckland C, *et al*. Detection of oxazolidinone-resistant
414 *Enterococcus faecalis* and *Enterococcus faecium* strains by real-time PCR and PCR-
415 restriction fragment length polymorphism analysis. *J Clin Microbiol* 2002; **40**: 4298–300.

416 20. Werner G, Strommenger B, Klare I, *et al.* Molecular detection of linezolid resistance in
417 *Enterococcus faecium* and *Enterococcus faecalis* by use of 5' nuclease real-time PCR
418 compared to a modified classical approach. *J Clin Microbiol* 2004; **42**: 5327–31.

419 21. Kehrenberg C, Schwarz S. Distribution of florfenicol resistance genes *fexA* and *cfr*
420 among chloramphenicol-resistant *Staphylococcus* isolates. *Antimicrob Agents Chemother*
421 2006; **50**: 1156–63.

422 22. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence
423 data. *Bioinforma Oxf Engl* 2014; **30**: 2114–20.

424 23. Loman NJ, Quinlan AR. Poretools: a toolkit for analyzing nanopore sequence data.
425 *Bioinformatics* 2014; **30**: 3399–401.

426 24. Raven KE, Reuter S, Gouliouris T, *et al.* Genome--based characterization of hospital-
427 adapted *Enterococcus faecalis* lineages. *Nat Microbiol* 2016; **1**.

428 25. Ponstingl H, Ning Z. SMALT. *Wellcome Trust Sanger Inst* 2014. Available at:
429 <http://www.sanger.ac.uk/science/tools/smalt-0>. Accessed June 21, 2017.

430 26. Page AJ, Taylor B, Delaney AJ, *et al.* SNP-sites: Rapid efficient extraction of SNPs from
431 multi-FASTA alignments. *Microb Genomics* 2016; **2**: e000056.

432 27. Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of
433 large phylogenies. *Bioinformatics* 2014; **30**: 1312–3.

434 28. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and
435 annotation of phylogenetic and other trees. *Nucleic Acids Res* 2016; **44**: W242–5.

436 29. Croucher NJ, Page AJ, Connor TR, *et al.* Rapid phylogenetic analysis of large samples
437 of recombinant bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* 2015;
438 **43**: e15.

439 30. Wick RR, Judd LM, Gorrie CL, *et al.* Unicycler: Resolving bacterial genome assemblies
440 from short and long sequencing reads. *PLOS Comput Biol* 2017; **13**: e1005595.

441 31. Pruitt KD, Tatusova T, Brown GR, *et al.* NCBI Reference Sequences (RefSeq): current
442 status, new features and genome annotation policy. *Nucleic Acids Res* 2012; **40**: D130–5.

443 32. Sullivan MJ, Petty NK, Beatson SA. Easyfig: A genome comparison visualizer.
444 *Bioinformatics* 2011; **27**: 1009–10.

445 33. Alikhan N-F, Petty NK, Ben Zakour NL, *et al.* BLAST Ring Image Generator (BRIG):
446 simple prokaryote genome comparisons. *BMC Genomics* 2011; **12**: 402.

447 34. Inouye M, Dashnow H, Raven L-A, *et al.* SRST2: Rapid genomic surveillance for public
448 health and hospital microbiology labs. *Genome Med* 2014; **6**: 1–16.

449 35. Jolley KA, Maiden MC. BIGSdb: Scalable analysis of bacterial genome variation at the
450 population level. *BMC Bioinformatics* 2010; **11**: 595.

451 36. Ruiz-Garbajosa P, Bonten MJM, Robinson DA, *et al.* Multilocus sequence typing scheme
452 for *Enterococcus faecalis* reveals hospital-adapted genetic complexes in a background of
453 high rates of recombination. *J Clin Microbiol* 2006; **44**: 2220–8.

37. Hunt M, Mather AE, Sánchez-Busó L, *et al.* ARIBA: Rapid antimicrobial resistance genotyping directly from sequencing reads. *Microb Genomics* 2017; **3**: e000131.
38. Zankari E, Hasman H, Kaas RS, *et al.* Genotyping using whole-genome sequencing is a realistic alternative to surveillance based on phenotypic antimicrobial susceptibility testing. *J Antimicrob Chemother* 2013; **68**: 771–7.
39. Goodhead I, Darby AC. Taking the pseudo out of pseudogenes. *Curr Opin Microbiol* 2015; **23**: 102–9.
40. Brenciani A, Morroni G, Vincenzi C, *et al.* Detection in Italy of two clinical *Enterococcus faecium* isolates carrying both the oxazolidinone and phenicol resistance gene *optrA* and a silent multiresistance gene *cfr*. *J Antimicrob Chemother* 2016; **71**: 1118–9.
41. Cai J, Schwarz S, Chi D, *et al.* Faecal carriage of *optrA*-positive enterococci in asymptomatic healthy humans in Hangzhou, China. *Clin Microbiol Infect* 2019; **25**: 630.e1-630.e6.
42. Morroni G, Brenciani A, Simoni S, *et al.* Nationwide surveillance of novel oxazolidinone resistance gene *optrA* in *Enterococcus* Isolates in China from 2004 to 2014. *Front Microbiol* 2017; **8**: 1631.
43. Freitas AR, Elghaieb H, León-Sampedro R, *et al.* Detection of *optrA* in the African continent (Tunisia) within a mosaic *Enterococcus faecalis* plasmid from urban wastewaters. *J Antimicrob Chemother* 2017; **72**: 3245–51.
44. Larsen J, Schønheyder HC, Lester CH, *et al.* Porcine-origin gentamicin-resistant *Enterococcus faecalis* in humans, Denmark. *Emerg Infect Dis* 2010; **16**: 682–4.
45. Pang S, Boan P, Lee T, *et al.* Linezolid-resistant ST872 *Enterococcus faecium* harbouring *optrA* and *cfr* (D) oxazolidinone resistance genes. *Int J Antimicrob Agents* 2020; **55**: 105831.
46. Sassi M, Guerin F, Zouari A, *et al.* Emergence of *optrA*-mediated linezolid resistance in enterococci from France, 2006–2016. *J Antimicrob Chemother* 2019; **74**: 1469–72.
47. Long KS, Poehlsgaard J, Kehrenberg C, *et al.* The Cfr rRNA methyltransferase confers resistance to phenicols, lincosamides, oxazolidinones, pleuromutilins, and streptogramin A antibiotics. *Antimicrob Agents Chemother* 2006; **50**: 2500–5.
48. Cai J, Wang Y, Schwarz S, *et al.* Enterococcal isolates carrying the novel oxazolidinone resistance gene *optrA* from hospitals in Zhejiang, Guangdong, and Henan, China, 2010–2014. *Clin Microbiol Infect* 2015; **21**: 1095.e1-1095.e4.
49. Bender JK, Fleige C, Lange D, *et al.* Rapid emergence of highly variable and transferable oxazolidinone and phenicol resistance gene *optrA* in German *Enterococcus* spp. clinical isolates. *Int J Antimicrob Agents* 2018; **52**: 819–27.
50. Hua R, Xia Y, Wu W, *et al.* Molecular epidemiology and mechanisms of 43 low-level linezolid-resistant *Enterococcus faecalis* strains in Chongqing, China. *Ann Lab Med* 2019; **39**: 36–42.
51. Elghaieb H, Freitas AR, Abbassi MS, *et al.* Dispersal of linezolid-resistant enterococci carrying *poxTA* or *optrA* in retail meat and food-producing animals from Tunisia. *J Antimicrob Chemother* 2019; **75**: 2865–9.

52. Munk P, Knudsen BE, Lukjancenko O, *et al.* Abundance and diversity of the faecal resistome in slaughter pigs and broilers in nine European countries. *Nat Microbiol* 2018; **3**: 898–908.
53. Zhao Q, Wang Y, Wang S, *et al.* Prevalence and abundance of florfenicol and linezolid resistance genes in soils adjacent to swine feedlots. *Sci Rep* 2016; **6**: 1–7.
54. Cavaco LM, Bernal JF, Zankari E, *et al.* Detection of linezolid resistance due to the *optrA* gene in *Enterococcus faecalis* from poultry meat from the American continent (Colombia). *J Antimicrob Chemother* 2017; **72**: 678–83.
55. Cavaco LM, Korsgaard H, Kaas RS, *et al.* First detection of linezolid resistance due to the *optrA* gene in enterococci isolated from food products in Denmark. *J Glob Antimicrob Resist* 2017; **9**: 128–9.
56. Wu Y, Fan R, Wang Y, *et al.* Analysis of combined resistance to oxazolidinones and phenicols among bacteria from dogs fed with raw meat/vegetables and the respective food items. *Sci Rep* 2019; **9**: 15500.
57. Chang Q, Wang W, Regev-Yochay G, *et al.* Antibiotics in agriculture and the risk to human health: How worried should we be? *Evol Appl* 2015; **8**: 240–7.
58. Bortolaia V, Espinosa-Gongora C, Guardabassi L. Human health risks associated with antimicrobial-resistant enterococci and *Staphylococcus aureus* on poultry meat. *Clin Microbiol Infect* 2016; **22**: 130–40.

516
517

Table 1. Details of the *optrA*-positive *E. faecalis* characterized in this study

Patient ID	Isolate	Year	Region	Patient Sex	Patient Age	Sample	Source	MLST
1	WE0851	2014	Grampian	Female	21	Urine	Outpatient	480
2	WE0254	2015	Grampian	Male	71	Urine	Outpatient	19
3	WE0438	2016	Grampian	Female	58	Urine	Inpatient	330
4	TM6294	2017	Forth Valley	Female	74	Urine	Inpatient	585
5	BX5936	2017	Lothian	Male	60	Semen	Outpatient	894
6	BX8117	2017	Lothian	Female	19	Urine	Outpatient	16

518
519

Table 2. Plasmids from Hybrid Assemblies

Isolate	Element	Copy Number ^a	Size (bp)	Plasmid rep type	Best NCBI match	Resistance genes	Variation compared to <i>optrA</i> _{pE394^b}	Transposases (n)
BX5936	pBX5936-1	1	68656	rep9	Efs pE035, coverage 65%, 98% ID (MK140641)	<i>fexA</i> , <i>optrA</i>	S2F	ISEf1 (2) IS1216 (2)
	pBX5936-2	1	51669	rep9	Efs FC unnamed plasmid1, coverage 85%, 100% ID (CP028836)	<i>catA8</i> , <i>tet(L)</i> , <i>tet(M)</i> , <i>ant(6)-la</i> , <i>cadA</i> , <i>copZ</i> , <i>ermB</i>	-	NA
BX8117	pBX8117-1	1	68773	rep9	Efs FDAARGOS_324 unnamed plasmid2, coverage 100%, 100% ID (CP028284)	None	-	NA
	pBX8117-2	1	41839	rep9	Efs pEF123, coverage 64%, 98% ID (KX579977)	<i>catA8</i> , <i>cfr(D)</i> , <i>optrA</i> , <i>fexA</i>	K3E, N12Y, E37K, N122K, Y135C, Y176D, A350V, V395A, A396S, Q509K, Q541E, M552L, N560Y, K562N, Q565K, E614Q, I627L, D633E, N640I, R650G	IS1216 (5)
TM6294	pTM6294-1	1	75362	rep9	Efs FC unnamed plasmid1, coverage 74%, 99% ID (CP028836)	<i>catA8</i> , <i>tet(L)</i> , <i>tet(M)</i> , <i>ant(6)-la</i> , <i>cadA</i> , <i>copZ</i> , <i>ermB</i> , <i>aph(3')-IIIa</i> , <i>sat4</i> , <i>ant(6)-la</i> , <i>lnuB</i> , <i>lsaE</i> , <i>ant(9)</i> , <i>ant(6)-la</i> ,	-	NA

						<i>aac(6')-le-aph(2'')-la, aadK, ermB, dfrG</i>		
	pTM6294-2	1	52776	rep9	Efs pE035, coverage 87%, 99% ID (MK140641)	<i>fexA, optrA</i>	None	ISL3-family (1) IS1216 (1)
WE0254	pWE0254-1	1	80496	repUS11	Efs FDAARGOS_324 unnamed plasmid3, coverage 49%, 99% ID (CP028283)	<i>ant(9)-la, ermA-like, fexA, optrA</i>	None	IS3-family (8) IS1216-partial (1)
	pWE0254-2 ^c	1	79293	NA	Efs NCTC8732 chromosome, coverage 99%, 100% ID (LR594051)	None	-	NA
WE0438	pWE0438	1	61284	rep9	Efs pEF123, coverage 76%, 99% ID (KX579977)	<i>tet(L), tet(M), bcrA, cadA, copZ, ant(6)-la, optrA, fexA, ermB</i>	K3E, Y176D, I622M	IS1216 (1) ISEnfa1 (2) IS3-family (4) IS6-family (1) Tn3-family (2)
WE0851	pWE0851-1	1	59708	repUS11	Efs pEF123, coverage 22%, 100% ID (KX579977)	<i>fexA, optrA, ermA-like</i>	T112K, Y176D	IS1216 (1) IS3-family (6) Tn3-family (1)
	pWE0851-2	1	26996	repUS11	Efs pKUB3007-3, coverage 63%, 100% ID (AP018546)	<i>aac(6')-le-aph(2'')-la</i>	-	NA
	pWE0851-3	3	10826	NA	Efs pE035, coverage 63%, 99% ID (MK140641)	<i>aac(6')-le-aph(2'')-la, aac(6')-le-aph(2'')-la, aadK, ermB, ant(6)-la, aph(3')-IIIa, sat4</i>	-	NA

520 bp, base pairs; Efs, *E. faecalis*; ID, identity; NA, not analysed

521 a Inferred from depth of coverage relative to chromosomal fragment in hybrid assembly

522 b Amino acid sequence variants compared to the first described *optrA* sequence from pE394 (KP399637)

523 c Incompletely assembled, in four contigs

524

525

Figure 1. Comparison of *optrA* genetic environments. Alignment of *optrA*-carrying regions shows limited shared sequence identity, apart from around the *optrA* and *fexA* genes. There is evidence of IS1216 near to the *optrA* gene in all isolates (partial sequence in pWE0254-1 indicated by asterisk), but other insertion sequences are also present suggesting multiple means of *optrA* transmission and ongoing diversification of the element. Arrows indicate coding sequences, blocks between each sequence indicate regions with BLASTn sequence identity >90% and length >100bp.

Figure 2. *optrA*-positive *E. faecalis* isolates in a national perspective. Phylogenetic analysis of the six *optrA*-positive isolates and 94 isolates from bloodstream infections in the UK shows the *optrA*-positive isolates are generally unrelated to others in the collection. Illumina reads were mapped to *E. faecalis* V583 reference genome, mobile genetic elements removed, and a maximum likelihood phylogeny performed on SNP alignment. Scale bar shows ~9500 SNPs, linezolid-resistant isolates are indicated by circles, the outer ring indicates isolate source.

Figure S1. Alignment of full *optrA*-positive plasmid sequences. While some sequence similarity is seen between pTM6294-2 and pBX5936-1, in general identity is low between the *optrA*-positive plasmids, indicating *optrA* has mobilised to multiple plasmid backbones. Arrows indicate coding sequences, blocks between each sequence indicate regions with BLASTn sequence identity $\geq 80\%$ and length $> 100\text{bp}$.

Figure S2. Alignment of full *optrA*-positive plasmid sequences against pE394. Sequence similarity confined to the *optrA/fexA* region. Inner ring indicates GC content of pE394, then alignment of pWE0438, pBX8117-2, pWE0851-1, pWE0254-1, pBX5936-1, pTM6294-2, and outer ring indicating coding sequences in pE394 (accession KP399637). Figure made with BRIG v0.95