1 Birth weight associations with psychiatric and physical health, cognitive function, and DNA methylation
2 differences in an adult population

3

4 Rebecca A. Madden¹, Daniel L. McCartney², Rosie M. Walker^{2,4}, Robert F. Hillary², Mairead L. Bermingham²,
5 Konrad Rawlik³, Stewart W. Morris², Archie Campbell², David J. Porteous^{2,4}, Ian J. Deary^{4,5}, Kathryn L.
6 Evans^{2,4}, Jonathan Hafferty^{1,†}, Andrew M. McIntosh^{1,2,4,†}, Riccardo E. Marioni^{2,4,†,*}

7

8 1. Division of Psychiatry, University of Edinburgh, Royal Edinburgh Hospital, Edinburgh, EH10 5HF, UK

9 2. Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine, University of
10 Edinburgh, Edinburgh, EH4 2XU, UK

11 3. The Roslin Institute, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Midlothian, EH25
12 9RG, UK

13 4. Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, EH8 9JZ,
14 UK

15 5. Department of Psychology, University of Edinburgh, Edinburgh, EH8 9JZ, UK

16

17 * Corresponding Author: Riccardo E Marioni, Centre for Genomic and Experimental Medicine, Institute of
18 Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, EH4 2XU, UK. Email:
19 riccardo.marioni@ed.ac.uk Telephone: +44 (0)131 651 8528

20 [†] These authors contributed equally

21 **Abstract**

22 **Background:**

23 The Developmental Origins of Adult Disease (DOAD) theory predicts that prenatal and early life events shape
24 adult health outcomes. Birth weight is a useful indicator of the foetal experience, and has been associated with
25 multiple adult health outcomes. DNA methylation (DNAm) is one plausible mechanism behind the relationship
26 of birth weight to adult health.

27 **Methods:**

28 The Generation Scotland study allows data linkage to historic Scottish birth cohorts, and birth records held
29 through the NHS Information and Statistics Division. Data linkage with these sources yielded a sample of 4, 710
30 individuals. Health measures were related to birth weight in regression models. An epigenome-wide association
31 study (EWAS) was performed in a subgroup (n=1, 395), relating adult DNAm from whole blood to birth weight,
32 with replication in an independent sample (n=362). Associations between birth weight and epigenetic clocks
33 were also assessed.

34 **Findings:**

35 Higher birth weight was significantly associated with reduced incidence of depression and osteoarthritis, higher
36 body mass index, and higher general intelligence (absolute standardised effect size range 0.04 to 0.30,
37 $p_{(FDR)} < 0.05$). Meta-analysis of discovery and replication EWAS studies yielded one genome-wide significant
38 CpG site ($p = 5.97 \times 10^{-9}$), cg00966482. Significant associations between birth weight and Grim Age ($p = 0.0014$)
39 and DNAm-derived telomere length ($p = 3.3 \times 10^{-4}$) are also described.

40 **Interpretation:**

41 Our results demonstrate associations between birth weight and adult health outcomes, with particularly striking
42 effects for depression risk. It also provides support for an association between birth weight and DNAm,
43 describing the first significant EWAS site associated with birth weight in an adult sample.

44 **Funding:**

45 Wellcome Trust Strategic Award 104036/Z/14/Z

47 **Keywords:** DNA methylation, Generation Scotland, birth weight, depression, EWAS

49 **Research in Context:**

50 *Evidence before this study*

51 The associations between birth weight and various adult health outcomes have been well established. DNA
 52 methylation is a plausible mechanism through which early life experiences may continue to affect health
 53 throughout the lifecourse; however, evidence for birth weight associations with DNA methylation in adulthood
 54 has not yet been robustly established. This is likely due to small sample sizes of previous samples, as well as the
 55 use of poor-quality birth weight data, such as binary ‘low/normal’ variables or retrospective self-report.
 56 Alternatively, work has attempted to describe the persistence into adulthood of DNA methylation at sites
 57 identified at birth.

58 *Added value of this study*

59 We investigated genome-wide differential DNA methylation patterns from whole blood using data linkage-
 60 derived, continuous birth weight data, in the largest reported adult sample (n=1, 395) with replication (n=362)
 61 and meta-analysis. Meta-analysis revealed one epigenome-wide significant CpG site, to our knowledge the first
 62 significant EWAS result reported for birth weight in an adult sample. In addition, we found associations
 63 between birth weight and GrimAge and a DNA methylation-derived measure of telomere length, demonstrating
 64 accelerated biological ageing in lower birth weight individuals. Together, these results suggest differential
 65 methylation exists in adulthood related to birth weight, and this may be relevant to health and mortality.

66 *Implications of all the available evidence*

67 Although CpG sites differentially methylated with birth weight at parturition may not remain so throughout life,
 68 the adult epigenome may still provide information on the impact of birth weight on health outcomes. The adult
 69 epigenome, therefore, may represent a useful archive of the foetal experience which results in birth weight
 70 variability, and this information may provide clinically useful information in mid-life.

Introduction

The Developmental Origins of Adult Disease theory (DOAD) states that through developmental plasticity, the foetal experience can permanently influence adult health (1). The theory's main proponent, David Barker, originally relied on birth weight as an index of foetal nutrition – an assumption that has been contested by the awareness that multiple factors can influence birth weight (2). Maternal stress, illness, and socioeconomic status (3-5) are among modifiable influences over offspring birth weight; in addition, maternal-specific genetic variation has been found to influence birth weight, acting through the intrauterine environment (6). Thus, birth weight can be seen as an index of the general foetal experience.

There is evidence of a U-shaped association between birth weight and adult health. Birth weight is clinically conceptualised as 'low' below 2.5kg, and 'high' above 4.5kg (7), however it can also be analysed on a continuous scale. Lower birth weight has been associated with an increased risk of heart disease, type II diabetes, stroke, and hypertension (8-11). Low birth weight is also associated with poorer cognitive ability, and a raised risk for mood disorders (12, 13). At the higher end of the spectrum, birth weight is associated with higher body mass index, and higher risk of breast cancer (14, 15). These associations are found after accounting for adult lifestyle factors, such as smoking and BMI, indicating a residual effect of birth weight on health outcomes. Since birth weight information is recorded as standard practice, it could be considered a useful and readily available predictive tool for adult health risks.

The DOAD theory is a plausible explanation for these observations. Prenatal factors affect the foetus in its highly plastic state, giving rise to birth weight variability, and also to developmental changes which permanently affect the function and health of organs and systems (1). Some of these changes may be structural, affecting for example, vascular development and function (16). DNA methylation (DNAm) is an epigenetic modification that can be influenced by genetics or by environmental factors throughout life. CpG methylation is characterised by the addition of a methyl group to cytosine nucleotides in the context of cytosine-guanine dinucleotides. DNAm changes are linked to the regulation of gene expression, providing a possible mechanism through which environmental influences may have lasting biological effects (17). Therefore, DNAm is one putative mechanism through which developmental experience may influence adult health.

Birth weight effects on the epigenome have been previously described in cord blood (18, 19) and during childhood (20), seeming to diminish into adolescence and beyond (18, 21). It is possible, however, that examining only the persistence of DNAm patterns associated with birthweight at the point of birth ignores the full adult epigenome which, despite having changed since birth, nevertheless may provide information on the impact of birthweight on downstream health. Here, we perform the first large EWAS study of birth weight on adult DNAm. In addition, recent work has demonstrated the utility of so-called “epigenetic clocks” in demonstrating elements of vulnerability to disease and mortality (22), conceptualised as accelerated biological age in comparison to chronological age. These analyses together have the potential to describe the known associations between birthweight and many aspects of whole-body health in adulthood.

Here, we describe associations between birth weight and a range of adult health conditions and risk factors, using high-quality data that improves on previous work in the field (e.g. (13)); it also reports an Epigenome-Wide Association Study (EWAS) of birthweight using whole-blood DNA derived in an adult sample, and associations between birth weight and five epigenetic clocks. While previous work has looked at DNAm relationships to birth weight in adults, none have performed EWAS studies in this age group, looking instead at persistence of DNAm effects from birth (18, 21). We, therefore, hypothesise that while birth weight associated DNAm patterns may change over time, differences will still exist in adulthood.

Materials and Methods

Generation Scotland and other Cohorts

Generation Scotland (GS) is a Scottish family-based cohort n=23, 690 (23). Data were collected from participants between 2006 and 2011. GS is a deeply-phenotyped cohort, allowing examination of many aspects of adult health, alongside genomic and biometric measures. In addition, 98% of GS participants gave informed consent for data linkage to routinely collected health data and to information from other Scottish population cohort studies, both current and historical. These include several with neonatal and maternity information: the Aberdeen Children of the 1950s ((24), ACONF); the Aberdeen Maternity and Neonatal Databank ((25), AMND); the Walker Birth Cohort (26); and the Scottish Morbidity Records ((27), SMR02 – the Maternity Inpatient and Day Case record, and SMR11 – the Neonatal Inpatient dataset). Birth weight in grams, alongside gestational age at birth and twin information, was collated from these sources and linked to adult GS records for

4, 713 participants (**Supplementary File 1, Figure 1**). Birth weight data derived from health records taken at birth improves on the self-reported nature of birth weight variables found in other population cohorts.

Statistical Analyses

All analyses were conducted in R version 3.5.1 (28).

To control for the known effects of gestational age and sex on birthweight (29), we considered the residuals from a regression model in place of raw birth weight throughout:

$$\text{Birth weight (g)} \sim \text{sex} + \text{gestational age}$$

Self-reported diseases (yes/no) that were commonly reported in the cohort (prevalence >1%) included depression (prevalence 6.6%), osteoarthritis (1.6%), high blood pressure (3.4%), diabetes (1.3%), and asthma (16.6%). These were modelled against the birth weight residuals using logistic regression. Average diastolic and systolic blood pressure measurements, body mass index, high-density lipoprotein (HDL) cholesterol, and general intelligence score (derived from multiple cognitive tests - see **Supplementary File 1**) were examined as continuous variables in linear regression models. Depression diagnosed using the Structured Clinical Interview for DSM-IV (SCID;(30) was also examined, as a binary yes/no for a diagnosis of major depressive disorder (**Supplementary File 1**). Three individuals who had responded 'yes' to all self-reported illnesses were excluded from analyses, leaving a final population of n=4, 710. Some traits had missing values, which resulted in case-wise exclusion from the relevant regression model (**Table 1**).

Adult health and functional domains were regressed against birth weight residuals and relevant covariates in a simple model:

$$\text{Trait} \sim \text{birth weight residuals} + \text{sex} + \text{age}$$

And in a fully-adjusted model:

$$\text{Trait} \sim \text{birth weight residuals} + \text{sex} + \text{age} + \text{Scottish Index of Multiple Deprivation (SIMD)} \\ \text{quintile} + \text{lifetime smoking (ever/never)} + \text{years of education}$$

For the general intelligence trait, years of education was removed from this fully-adjusted model as the two are highly collinear (Pearson $r=0.36$).

Correction for multiple testing was carried out using a false discovery rate $P<0.05$.

Epigenome-wide association study

Genome-wide DNA methylation was profiled in 5,190 individuals in GS, taken from peripheral whole blood, using the Illumina HumanMethylationEPIC BeadChip (Illumina Inc., San Diego, CA). Quality control and normalisation were carried out as described elsewhere ((31, 32); **Supplementary File 2**). For the subgroup of GS for whom birth weight and gestational age information were available, DNAm data was available for $n=1,395$ (**Figure 1**).

In a second release of data generated during this analysis, using a near identical protocol (**Supplementary File 2**), an independent set of methylation data became available for an additional 4,450 GS participants. In this replication sample, a further 362 participants with both birth weight and gestational age information were used.

The birth weight residuals described above were used in the EWAS model, which was run using the ‘limma’ package in R (empirical Bayes moderated t-statistics). The discovery EWAS model used CpGs corrected for relatedness (**Supplementary File 2**), as the first batch of DNAm data was collected on related individuals:

$$\text{CpG} \sim \text{birth weight residuals} + \text{age} + \text{sex} + \text{smoking [ever smoke and pack years]} + 20 \text{ methylation PCs}$$

Additional covariates (estimated white blood cell proportions – CD4T, CD8T, Granulocytes, BCells, Natural Killer cells – and methylation batch), which were regressed out during the relatedness pre-correction for the discovery dataset, were also included in the replication dataset of unrelated individuals.

Meta-Analysis of discovery and replication samples

A standard error-weighted meta-analysis of the discovery and replication EWASs was performed using the METAL software package. Summary statistics from the discovery, replication and meta-analysis EWASs are available at [**weblink to be inserted upon acceptance**].

Sensitivity analysis

As a sensitivity analysis, the 20 CpGs most strongly associated with birth weight in the discovery EWAS were re-examined with a maximally-corrected model to account for additional lifestyle factors which may affect both DNAm and health. These were body mass index, socioeconomic status (Scottish Index of Multiple Deprivation - SIMD), and number of years of education, on top of the correction for smoking already included in the discovery EWAS model. DNAm data were pre-corrected for batch, cell counts, and relatedness. The maximally corrected model was:

$$\text{CpG} \sim \text{birth weight residuals} + \text{age} + \text{sex} + \text{BMI} + \text{SIMD} + \text{years of education} + \text{smoking [ever smoke and pack years]} + 20 \text{ methylation PCs}$$

Epigenetic clock analyses

Detail on the estimation of the five epigenetic clocks used has previously been reported elsewhere (22). Briefly, penalised regression models have used to identify subsets of CpG sites that predict chronological age (33, 34), biological age (35), telomere length (36), and survival (37). The clocks either trained predictors directly on the outcome (e.g., chronological age or telomere length), or via surrogate markers (e.g., protein levels and biomarkers that are known to associated with biological/health processes). The residuals from regressions of these predictors on chronological age gives an index of age acceleration. Here, we considered the association (linear regression) between birth weight (outcome) and the five sets of age acceleration residuals, with covariate adjustment for age, sex, methylation set, and estimated white cell proportions.

Ethics approval and consent to participate

All components of GS received ethical approval from the NHS Tayside Committee on Medical Research Ethics (REC Reference Number: 05/S1401/89). GS has also been granted Research Tissue Bank status by the Tayside Committee on Medical Research Ethics (REC Reference Number: 10/S1402/20), providing generic ethical approval for a wide range of uses within medical research.

Results

Phenotype population characteristics

There were 4,710 Generation Scotland (GS) participants with birth weight and gestational age information available (**Table 1**). The population was 57% female, with a mean birth weight of 3.40kg (SD=0.52), and gestational age of 39.8 weeks (SD=1.7). The minimum birth weight in the sample was 0.7kg, the maximum was 5.1kg. Using the clinical cut-off of 2.5kg, 3.99% (n=188) of the sample were of clinically 'low' birth weight. This means the data describe phenotypic relationships to birth weight across pathological and non-pathological cases.

Relationship of adult outcomes to birth weight

Regression models identified significant associations between higher birth weight (effect sizes are reported per SD) and lower risk of SCID-diagnosed depression (OR=0.86; 95% CI 0.78-0.95, p=0.018); lower risk of self-reported osteoarthritis (OR=0.74; 95% CI 0.58-0.94; p=0.034); higher BMI (β =0.072; SE=0.016, p=3.7x10⁻⁵) and higher general intelligence (β =0.042; SE=0.016, p=0.027; **Table 2**).

In minimally-adjusted regression models (covarying for age and sex alone), higher birth weight showed a relationship with lower self-reported depression risk (OR=0.85; 95% CI 0.75-0.96; p=0.018). Associations were also found for the four traits described above, but no other adult health or functional outcomes (**Supplementary Table 1**).

Epigenome-wide association study of birthweight

The epigenome-wide association study of birth weight revealed no CpGs significant at the genome-wide level (p<3.6x10⁻⁸; (38)), although 19 CpGs had P<1x10⁻⁵ (minimum FDR corrected P-value of 6.05x10⁻⁸ for cg00966482) (**Figure 2a; Supplementary Table 2**). The 19 CpGs were largely uncorrelated, with the exception of three CpGs (located within *CASZ1* – two within 200 base pairs of each other, with the third site around 11kb away – **Supplementary Table 2**) that had absolute $r \geq 0.6$ (**Figure 3**).

Replication in an independent DNAm sample

There were no genome-wide significant associations in the replication sample (n=362) (**Figure 2b; Supplementary Table 3**).

Of the 19 CpGs that exceeded the suggestive significance threshold in the discovery EWAS, two reached nominal significance ($p < 0.05$) in the replication analysis: cg00590817 ($p = 0.0069$), and cg00966482 ($p = 0.031$). The latter CpG was the most significantly associated site from the discovery EWAS ($p = 6.05 \times 10^{-8}$) and is found within the *HERV-FRD* (also referred to as *ERVFRD-1*) gene. There was good concordance between the effect sizes of DNAm associations with birthweight in the discovery and replication studies ($r = 0.59$) (**Figure 4**).

Meta-analysis of discovery and replication samples

Meta-analysis of the original discovery EWAS sample with the replication sample found a genome-wide significant effect of birthweight on DNAm at the CpG site cg00966482 located in the *ERVFRD-1* gene ($p = 5.97 \times 10^{-9}$; **Figure 2c; Supplementary Table 4**).

Adjusting the EWAS model for lifestyle factors

As a sensitivity analysis, we re-ran the EWAS pipeline for the top CpG sites, with additional corrections for BMI, education, and SIMD. This resulted in some minor changes to the effect sizes (mean attenuation of 1.35%, range 11.12% attenuation to 12.67% increase in effect size; **Supplementary Table 5**).

Relationship of birth weight to epigenetic signatures of age and telomere length

The mean values for the five epigenetic clocks were: Horvath age 44.7 yrs (SD=10.6); Hannum age 36.1 yrs (SD=10.2); PhenoAge 32.8 yrs (SD=12); GrimAge 45.7 yrs (SD=14.1); DNAm telomere length 7.6 kilobase pairs (SD=0.29). Linear regression models of birth weight residuals (predictor) against five signatures of epigenetic age acceleration revealed significant associations between higher birth weight and both lower GrimAge ($\beta = -0.083$; SE=0.026, $p = 0.0014$) and shorter DNAm telomere length ($\beta = 0.098$; SE=0.027, $p = 3.3 \times 10^{-4}$; **Table 3**). Associations between birth weight and the other three measure of epigenetic age acceleration were non-significant.

Discussion

We observed a strong association between birth weight and adult depression, where a 1 standard deviation (SD; 0.52kg) increase in birth weight was linked to 15% lower odds of future depression. A 1SD increase in birth weight was also linked to a 0.072 SD ($\sim 0.37\text{kg/m}^2$) higher adult BMI, a 0.042 SD higher general intelligence

score, and 30.7% lower odds for self-reported osteoarthritis. We also identified one genome-wide significant association between birth weight and blood-based DNA methylation in adulthood.

Previous studies have found both positive and negative associations between birth weight and depression. A meta-analysis of 18 studies ($n \sim 50,000$) found support for a modest effect of low birthweight ($<2.5\text{kg}$) on depression risk ($\text{OR}=1.15$, $95\% \text{ CI}=1.00-1.32$), but suggested this was a result of publication bias (13). We improved upon the design of many of the papers within that meta-analysis by utilising clinician-diagnosed depression and birth weight as a continuous variable (compared to self-reported measures of depression or psychological distress and use of a binary ($<2.5\text{kg}$ vs normal) birth weight variable). Compared to SCID-diagnosed depression, we found self-reported depression had a weaker association with birth weight. Furthermore, the association with the self-reported measure was statistically significant in the minimally-adjusted but not fully-adjusted regression models. This emphasises the importance of using high quality, clinically meaningful tools for measuring psychiatric disorders. The aetiology of depression is complex, with multiple biological and psychosocial factors contributing increments of risk, thought to sum towards illness. This study indicates that birth weight may be considered as one of these contributing risk factors, as its association with adult diagnosis remained after accounting for many adult lifestyle factors.

Consistent with the published literature (11, 14), we identified an association between higher birth weight and higher adult BMI. Previous work has used Mendelian randomisation techniques to show that birthweight is a causal factor in determining adult BMI (9). We also found a significant relationship between higher birth weight and lower prevalence of osteoarthritis in this sample. Developmental origins of osteoarthritis have been described before, with higher disease prevalence in low birth weight individuals (39), and those with low weight at age one (40).

A positive association between birth weight and general intelligence is also reported here (1SD increase in birthweight results in a 0.042 SD higher general intelligence score). Impaired cognitive ability after low birth weight has been shown in several samples in childhood (12) and into adulthood (41).

Some previously-established effects of birth weight on adult health are not supported here, for instance, links to type 2 diabetes and cardiovascular disease (9). This may be partly due to the self-report nature of the health

measures used. Our sample size was also smaller than previous investigations into birth weight and cardiovascular outcomes (9, 11).

In the EWAS meta-analysis we observed a genome-wide significant association between higher birthweight and higher methylation levels at cg00966482 (*ERVFRD-1*). *ERVFRD-1* encodes syncitin-2, a protein involved in placental embedding (42). This site was not identified in a previous meta-analysis EWAS of birthweight (21). Moreover, this is the first report of any genome-wide significant EWAS association for birth weight in an adult sample. DNAm data was derived from whole blood taken during adulthood, so our findings might not generalise to other tissue types. However, there are many advantages to interrogating blood-based methylation levels: first, it is the only tissue type that is readily available in large epidemiological studies – it would be expensive and invasive to obtain adipose tissue samples on over 1,000 individuals; second, it is systemic and tracks multiple biological processes including biomarkers of inflammation, cardiovascular disease, cardiometabolic disease, all of which are relevant processes related to birthweight; third, the DOAD theory posits that birth weight is related to health outcomes across the whole body.

Of the 19 CpG sites with $P < 1 \times 10^{-5}$ in the discovery EWAS, 10 were located within known genes. Some of these genes contain SNPs that have genome-wide associations (with $P < 5 \times 10^{-8}$) with cardiovascular, psychiatric, and developmental pathways (**Supplementary Table 6**). Three of the 19 nominally significant CpG sites identified in the discovery EWAS were highly correlated (min $r = 0.69$). Higher birthweight was associated with higher methylation levels at these sites, which were located within *CASZI*, a gene encoding the zinc finger protein castor homolog 1, a transcriptional activator involved in vascular morphogenesis (43). A differentially methylated region (DMR) in the *CASZI* gene was recently identified in placental tissue between infants born small vs. large for gestational age (19), thus this study suggests the persistence of differential methylation of this gene into adulthood. Genetic variants in *CASZI* have previously been implicated in GWAS studies on various aspects of cardiovascular health (**Supplementary Table 6**). These have included studies in multi-ethnic populations on blood pressure (44, 45), and on other cardiovascular health issues such as atrial fibrillation (46) and stroke (47).

A recent longitudinal meta-analysis (21) of DNAm relationships to birth weight identified 914 CpG sites associated with birth weight in 24 EWAS studies from neonatal blood (total $n = 8,825$). The persistence of

methylation differences at these sites was then examined in other cohort data from childhood (total n=2, 756 from 10 studies; 2-12y), adolescence (total n=2, 906 from 6 studies; 16-18y), and adulthood (1, 616 from 3 studies; 30-45y). Nominally significant methylation differences were found to persist into childhood, adolescence, and adulthood at 87, 49, and 42 sites respectively. This supports evidence that birth weight-related methylation differences may attenuate over time (18, 20). It has, however, been demonstrated that some methylation patterns persist into late adulthood after prenatal famine exposure (48). It is therefore plausible that other prenatal factors may continue to affect DNAm into adulthood. This study adds to the available literature, demonstrating that DNAm differences may exist in adulthood which are not persistent from birth, but which nonetheless have the potential to describe part of the ongoing association between birth weight and adult health. Whereas there was good concordance in effect sizes for the top discovery EWAS findings in our replication dataset, none were significant after correction for multiple testing. Further EWAS studies of birth weight and DNAm in mid-life are therefore required to replicate our genome-wide significant finding from the combined discovery and replication meta-analysis.

Significant associations are demonstrated between birth weight and epigenetic predictors of ageing, mortality, and cellular senescence. Higher birth weight is associated with lower GrimAge acceleration – a measure of epigenetic age with validated relationships to health and all-cause mortality (22) – and longer telomeres as estimated by DNAm data, known to associate with a variety of health outcomes (36). These findings further validate the rationale behind the current study, by indicating longitudinal associations between birth weight and DNAm predictors of health in mid-life.

This study exploited rarely available data linkage capacity to acquire neonatal information from birth medical records. The phenotyping and DNAm data available within the cohort have allowed the association of both health traits and DNAm data with birth weight. There are, however, some limitations inherent to the cross-sectional design of this study. Longitudinal data would allow analysis of the persistence of DNAm signatures across time. In addition, the number of individuals for whom accurate birth weight and gestational information could be identified in historical health cohorts limited the sample size for the EWAS analyses.

This study presents the first epigenome wide association study of birth weight on DNA methylation in adulthood, alongside associations between birthweight and both epigenetic age acceleration and telomere length.

We also present a comprehensive study of birth weight in relation to cognitive, psychiatric, and disease traits in mid-life. The Developmental Origins of Adult Disease theory predicts that birth weight can affect many domains of adult physical and psychological health. We found evidence to support this at both the molecular and broad disease/health phenotype levels.

Abbreviations:

DOAD - Developmental Origins of Adult Disease

DNAm - DNA methylation

EWAS - epigenome-wide association study

BMI - body mass index

GS - Generation Scotland

OR - Odds Ratio

SD - Standard Deviation

SE - Standard Error

CI - confidence interval

SCID - Structured Clinical Interview for DSM-IV

ACONF - Aberdeen Children of the 1950s

AMND - Aberdeen Maternity and Neonatal Databank

SMR - Scottish Morbidity Records

HDL - high-density lipoprotein

Acknowledgements

The authors thank all individuals and project team members who have contributed to both GS:SFHS and to the ‘STRADL: Stratifying Resilience and Depression Longitudinally’ follow-up study.

Funding Sources

GS received core support from the Chief Scientist Office of the Scottish Government Health Directorates (CZD/16/6) and the Scottish Funding Council (HR03006). Genotyping and DNA methylation profiling of the GS samples was carried out by the Genetics Core Laboratory at the Wellcome Trust Clinical Research Facility, Edinburgh, Scotland and was funded by the Medical Research Council UK and the Wellcome Trust (Wellcome

Trust Strategic Award “STratifying Resilience And Depression Longitudinally” ((STRADL) Reference 104036/Z/14/Z). Data linkage was supported by MRC Mental Health Pathfinder Award (Reference MC_PC_17209). This work was conducted in the Centre for Cognitive Ageing and Cognitive Epidemiology, which is supported by the Medical Research Council and Biotechnology and Biological Sciences Research Council (MR/K026992/1). DLM and REM are supported by Alzheimer’s Research UK major project grant ARUK-PG2017B-10. RAM and RFH are supported by funding from the Wellcome Trust 4-year PhD in Translational Neuroscience – training the next generation of basic neuroscientists to embrace clinical research [108890/Z/15/Z].

Declaration of Interests

AMM has received research support from Eli Lilly, Janssen and the Sackler Trust. AMM has also received speaker fees from Janssen and Illumina. The other authors declare that they have no competing interests.

Author contributions

Conception and design: RAM, JH, AMM, and REM. Data analysis: RAM and DLM. Interpretation: RAM, DLM, JH, AMM, REM. Drafting the article: RAM. Revision of the article: all authors.

Availability of data and materials

According to the terms of consent for Generation Scotland participants, access to data must be reviewed by the Generation Scotland Access Committee. Applications should be made to access@generationscotland.org. Summary statistics for the EWAS models will be made available on the University of Edinburgh DataShare facility upon acceptance [doi to be added]

References

1. Barker DJ. The developmental origins of adult disease. *Journal of the American College of Nutrition*. 2004;23(6 Suppl):588s-95s.
2. Paneth N, Susser M. Early origin of coronary heart disease (the "Barker hypothesis") *British Medical Journal*. 1995;310(6977):411-2.
3. Rondo PHC, Ferreira RF, Nogueira F, Ribeiro MCN, Lobert H, Artes R. Maternal psychological stress and distress as predictors of low birth weight, prematurity and intrauterine growth retardation. *European Journal of Clinical Nutrition*. 2003;57(2):266-72.
4. Grote NK, Bridge JA, Gavin AR, Melville JL, Iyengar S, Katon WJ. A Meta-analysis of Depression During Pregnancy and the Risk of Preterm Birth, Low Birth Weight, and Intrauterine Growth Restriction. *Archives of General Psychiatry*. 2010;67(10):1012-24.
5. Rosen D, Seng JS, Tolman RM, Mallinger G. Intimate partner violence, depression, and posttraumatic stress disorder as additional predictors of low birth weight infants among low-income mothers. *Journal of Interpersonal Violence*. 2007;22(10):1305-14.
6. Warrington NM, Beaumont RN, Horikoshi M, Day FR, Helgeland Ø, Laurin C, et al. Maternal and fetal genetic effects on birth weight and their relevance to cardio-metabolic risk factors. *Nature Genetics*. 2019;51(5):804-14.
7. Kiserud T, Piaggio G, Carroli G, Widmer M, Carvalho J, Neerup Jensen L, et al. The World Health Organization Fetal Growth Charts: A Multinational Longitudinal Study of Ultrasound Biometric Measurements and Estimated Fetal Weight. *PLoS Medicine*. 2017;14(1):1-36.
8. Huxley R, Owen CG, Whincup PH, Cook DG, Rich-Edwards J, Smith GD, et al. Is birth weight a risk factor for ischemic heart disease in later life? *American Journal of Clinical Nutrition*. 2007;85(5):1244-50.
9. Zanetti D, Tikkanen E, Gustafsson S, Priest JR, Burgess S, Ingelsson E. Birthweight, Type 2 Diabetes Mellitus, and Cardiovascular Disease Addressing the Barker Hypothesis With Mendelian Randomization. *Circulation-Genomic and Precision Medicine*. 2018;11(6).
10. Mu M, Wang SF, Sheng J, Zhao Y, Li HZ, Hu CL, et al. Birth weight and subsequent blood pressure: A meta-analysis. *Archives of Cardiovascular Diseases*. 2012;105(2):99-113.
11. Rich-Edwards JW, Kleinman K, Michels KB, Stampfer MJ, Manson JE, Rexrode KM, et al. Longitudinal study of birth weight and adult body mass index in predicting risk of coronary heart disease and stroke in women. *British Medical Journal*. 2005;330(7500):1115-8A.
12. Shenkin SD, Starr JM, Deary IJ. Birth weight and cognitive ability in childhood: A systematic review. *Psychological Bulletin*. 2004;130(6):989-1013.
13. Wojcik W, Lee W, Colman I, Hardy R, Hotopf M. Foetal origins of depression? A systematic review and meta-analysis of low birth weight and later depression. *Psychological Medicine*. 2013;43(1):1-12.
14. Curhan GC, Chertow GM, Willett WC, Spiegelman D, Colditz GA, Manson JE, et al. Birth weight and adult hypertension and obesity in women. *Circulation*. 1996;94(6):1310-5.
15. Xu XH, Dailey AB, Peoples-Sheps M, Talbott EO, Li N, Roth J. Birth Weight as a Risk Factor for Breast Cancer: A Meta-Analysis of 18 Epidemiological Studies. *Journal of Womens Health*. 2009;18(8):1169-78.
16. Leeson CPM, Whincup PH, Cook DG, Donald AE, Papacosta O, Lucas A, et al. Flow-mediated dilation in 9- to 11-year-old children - The influence of intrauterine and childhood factors. *Circulation*. 1997;96(7):2233-8.
17. Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nature Genetics*. 2003;33:245-54.
18. Simpkin AJ, Suderman M, Gaunt TR, Lyttleton O, McArdle WL, Ring SM, et al. Longitudinal analysis of DNA methylation associated with birth weight and gestational age. *Human Molecular Genetics*. 2015;24(13):3752-63.
19. Chen P-Y, Chu A, Liao W-W, Rubbi L, Janzen C, Hsu F-M, et al. Prenatal Growth Patterns and Birthweight Are Associated With Differential DNA Methylation and Gene Expression of Cardiometabolic Risk Genes in Human Placentas: A Discovery-Based Approach. *Reproductive Sciences*. 2018;25(4):523-39.
20. Agha G, Hajj H, Rifas-Shiman SL, Just AC, Hivert MF, Burris HH, et al. Birth weight-for-gestational age is associated with DNA methylation at birth and in childhood. *Clinical Epigenetics*. 2016;8.
21. Küpers LK, Monnereau C, Sharp GC, Yousefi P, Salas LA, Ghantous A, et al. Meta-analysis of epigenome-wide association studies in neonates reveals widespread differential DNA methylation associated with birthweight. *Nature Communications*. 2019;10(1):1893.
22. Hillary RF, Stevenson AJ, McCartney DL, Campbell A, Walker RM, Howard DM, et al. Epigenetic clocks predict prevalence and incidence of leading causes of death and disease burden. *bioRxiv*. 2020:2020.01.31.928648.

23. Smith BH, Campbell A, Linksted P, Fitzpatrick B, Jackson C, Kerr SM, et al. Cohort Profile: Generation Scotland: Scottish Family Health Study (GS:SFHS). The study, its participants and their potential for genetic research on health and illness. *International Journal of Epidemiology*. 2013;42(3):689-700.
24. Leon DA, Lawlor DA, Clark H, Macintyre S. Cohort profile: The Aberdeen children of the 1950s study. *International Journal of Epidemiology*. 2006;35(3):549-52.
25. Ayorinde AA, Wilde K, Lemon J, Campbell D, Bhattacharya S. Data Resource Profile: The Aberdeen Maternity and Neonatal Databank (AMND). *International Journal of Epidemiology*. 2016;45(2):389-94.
26. Libby G, Smith A, F McEwan N, Chien P, Greene S, Forsyth S, et al. The Walker Project: A longitudinal study of 48 000 children born 1952-1966 (aged 36-50 years in 2002) and their families 2004. 302-12 p.
27. Ndc.scot.nhs.uk. ISD Scotland | Information Services Division. 2018 [[Accessed 22 Nov. 2018]]. Available from: <http://www.ndc.scot.nhs.uk/Dictionary-A-Z/index.asp>.
28. Team RC. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria 2018. p. <https://www.R-project.org/>.
29. Norris T, Seaton SE, Manktelow BN, Baker PN, Kurinczuk JJ, Field D, et al. Updated birth weight centiles for England and Wales. *Archives of Disease in Childhood-Fetal and Neonatal Edition*. 2018;103(6):F577-F82.
30. Kübler U. Structured Clinical Interview for DSM-IV (SCID). In: Gellman MD, Turner JR, editors. *Encyclopedia of Behavioral Medicine*. New York, NY: Springer New York; 2013. p. 1919-20.
31. McCartney DL, Stevenson AJ, Walker RM, Gibson J, Morris SW, Campbell A, et al. Investigating the relationship between DNA methylation age acceleration and risk factors for Alzheimer's disease. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*. 2018;10:429-37.
32. Bermingham ML, Walker RM, Marioni RE, Morris SM, Rawlik K, Zeng Y, et al. Identification of novel differentially methylated sites with potential as clinical predictors of impaired respiratory function and COPD. *bioRxiv*. 2018:473629.
33. Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sada S, et al. Genome-wide Methylation Profiles Reveal Quantitative Views of Human Aging Rates. *Molecular Cell*. 2013;49(2):359-67.
34. Horvath S. DNA methylation age of human tissues and cell types. *Genome Biology*. 2013;14(10):3156.
35. Levine ME, Lu AT, Quach A, Chen BH, Assimes TL, Bandinelli S, et al. An epigenetic biomarker of aging for lifespan and healthspan. *Aging (Albany NY)*. 2018;10(4):573-91.
36. Lu AT, Seebach A, Tsai P-C, Sun D, Quach A, Reiner AP, et al. DNA methylation-based estimator of telomere length. *Aging (Albany NY)*. 2019;11(16):5895-923.
37. Lu AT, Quach A, Wilson JG, Reiner AP, Aviv A, Raj K, et al. DNA methylation GrimAge strongly predicts lifespan and healthspan. *Aging (Albany NY)*. 2019;11(2):303-27.
38. Saffari A, Silver MJ, Zavattari P, Moi L, Columbano A, Meaburn EL, et al. Estimation of a significance threshold for epigenome-wide association studies. *Genetic Epidemiology*. 2018;42(1):20-33.
39. Hussain SM, Wang YY, Wluka A, Shaw JE, Magliano DJ, Graves S, et al. Association of Low Birth Weight and Preterm Birth With the Incidence of Knee and Hip Arthroplasty for Osteoarthritis. *Arthritis Care & Research*. 2015;67(4):502-8.
40. Clynes MA, Parsons C, Edwards MH, Jameson KA, Harvey NC, Sayer AA, et al. Further evidence of the developmental origins of osteoarthritis: results from the Hertfordshire Cohort Study. *Journal of Developmental Origins of Health and Disease*. 2014;5(6):453-8.
41. Richards M, Hardy R, Kuh D, Wadsworth MEJ. Birth weight and cognitive function in the British 1946 birth cohort: longitudinal population based study. *British Medical Journal*. 2001;322(7280):199-203.
42. Chen C-P, Chen L-F, Yang S-R, Chen C-Y, Ko C-C, Chang G-D, et al. Functional Characterization of the Human Placental Fusogenic Membrane Protein Syncytin 21. *Biology of Reproduction*. 2008;79(5):815-23.
43. Charpentier Marta S, Christine Kathleen S, Amin Nirav M, Dorr Kerry M, Kushner Erich J, Bautch Victoria L, et al. CASZ1 Promotes Vascular Assembly and Morphogenesis through the Direct Regulation of an EGFL7/RhoA-Mediated Pathway. *Developmental Cell*. 2013;25(2):132-43.
44. Kato N, Takeuchi F, Tabara Y, Kelly TN, Go MJ, Sim XL, et al. Meta-analysis of genome-wide association studies identifies common variants associated with blood pressure variation in east Asians. *Nature Genetics*. 2011;43(6):530-U57.
45. Ehret GB, Ferreira T, Chasman DI, Jackson AU, Schmidt EM, Johnson T, et al. The genetics of blood pressure regulation and its target organs from association studies in 342,415 individuals. *Nature Genetics*. 2016;48(10):1171-84.
46. Roselli C, Chaffin MD, Weng LC, Aeschbacher S, Ahlberg G, Albert CM, et al. Multi-ethnic genome-wide association study for atrial fibrillation. *Nature Genetics*. 2018;50(9):1225-+.
47. Malik R, Chauhan G, Traylor M, Sargurupremraj M, Okada Y, Mishra A, et al. Multiancestry genome-wide association study of 520,000 subjects identifies 32 loci associated with stroke and stroke subtypes. *Nature Genetics*. 2018;50(4):524-+.

535 48. Heijmans BT, Tobi EW, Stein AD, Putter H, Blauw GJ, Susser ES, et al. Persistent epigenetic
536 differences associated with prenatal exposure to famine in humans. Proceedings of the National Academy of
537 Sciences. 2008;105(44):17046-9.

538

539

Tables

Table 1: Population characteristics of the phenotyped and EWAS samples.

	Phenotype Sample			EWAS Sample		
	N	Mean	SD	N	Mean	SD
Age (years)	4, 710	29.5	10.9	1, 395	37.1	14.7
Birthweight (g)	4, 710	3, 399	516	1, 395	3, 377	518
Gestation (weeks)	4, 710	39.8	1.7	1, 395	40	1.8
BMI (kg/m ²)	4, 385	25.2	5.2	1, 387	26.1	5.4
Education*	4, 411	5	4-6	1, 317	5	3-6
	N	%		N	%	
Sex - Male	2, 025	43.0		569	40.8	
Female	2, 688	57.0		826	59.2	
Socioeconomic Status**	(4, 389)			(1, 310)		
Quintile 1 (most deprived)	621	14.2		207	15.8	
Quintile 2	717	16.3		210	16.0	
Quintile 3	715	16.3		183	14.0	
Quintile 4	1, 064	24.2		301	23.0	
Quintile 5 (least deprived)	1, 272	29.0		409	31.2	
Smoking	(4, 525)			(1, 344)		
Current Smoker	918	20.3		257	19.1	
Ex-Smoker (<12 months)	237	5.2		56	4.2	
Ex-Smoker (>12 months)	662	14.6		273	20.3	
Never Smoker	2, 708	59.8		758	56.4	

* Median and Interquartile range reported. Education was coded as an ordinal variable: 0 = 0yrs, 1 = 1-4yrs, 2 = 5-9yrs, 3 = 10-11yrs, 4 = 12-13yrs, 5 = 14-15yrs, 6 = 16-17yrs, 7 = 18-19yrs, 8 = 20-21yrs, 9 = 22-23yrs, 10 = ≥24yrs.

**SIMD Quintile. SIMD is the Scottish Index of Multiple Deprivation, a postcode-derived index of socioeconomic status. The quintiles derived on the full Generation Scotland cohort ranged from 1 (most deprived) to 5 (least deprived).

Table 2: Outputs of logistic and linear regression models of health traits ~ birth weight residuals + age + sex + socioeconomic status (SIMD) + ever smoke + education years, with FDR-correction for multiple testing.

Trait	Odds Ratio	95% CI	P value	P (FDR)
SCID depression	0.86	0.78 – 0.95	0.0033	0.018
SR depression	0.92	0.81 – 1.05	0.21	0.38
SR hypertension	0.93	0.78 – 1.09	0.36	0.44
SR diabetes	0.96	0.74 – 1.25	0.76	0.76
SR osteoarthritis	0.74	0.58 – 0.94	0.012	0.034
SR asthma	0.96	0.88 – 1.04	0.3	0.43
Trait	Beta	SE	P value	P (FDR)
Body Mass Index (kg/m ²)	0.072	0.016	3.4x10 ⁻⁶	3.7x10 ⁻⁵
HDL cholesterol	0.025	0.016	0.12	0.27
Average systolic BP	-0.014	0.014	0.32	0.43
Average diastolic BP	-0.011	0.015	0.47	0.51
General intelligence*	0.042	0.016	0.0074	0.027

*For the ‘g’ general intelligence factor, education years was removed from the model as it resulted in over-correction of the effect.

Table 3: Outputs of linear regression models between birth weight residuals and five epigenetic signatures of accelerated biological ageing – Horvath, Hannum, PhenoAge, GrimAge, and DNAmTL.

Clock	Beta	SE	P value
Horvath	0.0032	0.025	0.9
Hannum	-0.0062	0.027	0.82
PhenoAge	-0.042	0.027	0.12
GrimAge	-0.083	0.026	0.0014
DNAmTL	0.098	0.027	3.3×10^{-4}

Figures

Figure 1: Inclusion flow diagram detailing the selection of samples and subsamples for the current study.

Figure 2: Manhattan plots for the epigenome-wide association study of birth weight (a); the replication sample EWAS (b); and the meta-analysis of the main EWAS and the replication (c). The black and red lines represent the suggestive, and Bonferroni corrected P-value thresholds of $P=1 \times 10^{-5}$ and 3.6×10^{-8} , respectively.

Figure 3: Correlation plot between methylation at the top 19 CpG sites from the epigenome-wide association study. The shade and scale of the dots represent the magnitude and direction of the correlation between pairs of CpGs.

Figure 4: Effect sizes for 18 of the top 19 CpG sites in the discovery sample plotted against the effect sizes in the replication sample (cg04988918 was not included in the replication array). The point size is determined by the $-\log_{10}$ of the p-values for these hits in the replication analysis. The two points labelled in black are the two CpG sites which achieved nominal significance in the replication study, and the three highlighted in red are the three co-methylated CpG sites within the *CASZ1* gene.

Figure 1: Inclusion flow diagram detailing the selection of samples and subsamples for the current study.

Figure 2: Manhattan plots for the epigenome-wide association study of birth weight (a); the replication sample EWAS (b); and the meta-analysis of the main EWAS and the replication (c). The black and red lines represent the suggestive, and Bonferroni corrected P-value thresholds of $P=1 \times 10^{-5}$ and 3.6×10^{-8} , respectively.

Figure 4: Effect sizes for 18 of the top 19 CpG sites in the discovery sample plotted against the effect sizes in the replication sample (cg04988918 was not included in the replication array). The point size is determined by the $-\log_{10}$ of the p-values for these hits in the replication analysis. The two points labelled in black are the two CpG sites which achieved nominal significance in the replication study, and the three highlighted in red are the three co-methylated CpG sites within the *CASZ1* gene.

Figure 3: Correlation plot between methylation at the top 19 CpG sites from the epigenome-wide association study. The shade and scale of the dots represent the magnitude and direction of the correlation between pairs of CpGs.