1 **Characterization of interstitial heterogeneity in the developing kidney**

2
3 Alicia R. England^{1,2,3,*}, Christopher P. Chaney^{2,*}, Amrita Das^{4,*}, Mohita Patel⁵, Alicia
4 Malewsak⁶, Daniel Armendariz^{7,3}, Gary Hon^{7,3}, Douglas Strand⁷, Keri Drake⁵, Thomas J.
5 Carroll^{1,2,3 ^}
6

7 ¹Department of Molecular Biology, Center for Regenerative Science and Medicine,
8 University of Texas Southwestern Medical Center, Dallas, TX, 75390, USA

9 ²Department of Internal Medicine, Division of Nephrology, University of Texas
10 Southwestern Medical Center, Dallas, TX, 75390, USA

11 ³Hamon Center for Regenerative Science and Medicine, Dallas, TX, 75390, USA.

12 ⁴Amgen, Inc., San Francisco, CA

13 ⁵Division of Pediatric Nephrology, University of Texas Southwestern Medical Center,
14 Dallas, TX, 75390, USA

15 ⁶Department of Urology, UT Southwestern Medical Center, Dallas, TX 75390, USA.

16 ⁷Laboratory of Regulatory Genomics, Cecil H. and Ida Green Center for Reproductive
17 Biology Sciences, Division of Basic Reproductive Biology Research, Department of
18 Obstetrics and Gynecology, University of Texas Southwestern Medical Center, Dallas,
19 TX, 75390, USA.

20 [^]Corresponding author, Thomas.Carroll@UTSoutwestern.edu

21 ^{*}Denotes equal contribution
22

ABSTRACT

Kidney formation requires the coordinated growth of multiple cell types including the collecting ducts, nephrons, vasculature and interstitium. There has been a long-held belief that interactions between the progenitors of the collecting ducts and nephrons are primarily responsible for development of this organ. However, over the last several years, it has become increasingly clear that multiple aspects of kidney development require signaling from the interstitium. How the interstitium orchestrates these various roles is still poorly understood. We show that during development, the interstitium is a highly heterogeneous, patterned population of cells that occupies distinct positions correlated to the adjacent parenchyma. Our analysis indicates that the heterogeneity is not a mere reflection of different stages in a linear developmental trajectory but instead represents several novel differentiated cell states. Further, we find that beta-catenin has a cell autonomous role in the development of a medullary subset of the interstitium and that this non-autonomously affects the development of the adjacent epithelia. These findings suggest the intriguing possibility that the different interstitial subtypes may create microenvironments that play unique roles in development of the adjacent epithelia and endothelia.

INTRODUCTION

Development of the kidney relies on interactions between the metanephric mesenchyme (MM) and an epithelial structure known as the ureteric bud (UB) [1, 2]. The MM is a heterogeneous population of cells containing at least two cell type specific progenitor populations. A Six2/Cited1+ nephron progenitor cell (NPC) population is located within the MM directly adjacent to the tips of the UB [3, 4]. NPCs undergo mesenchymal-to-epithelial transition (MET) to ultimately form the nephron, the functional unit of the kidney, which is patterned into functionally distinct segments: the glomerulus, proximal tubule, loop of Henle, distal tubule and connecting segment [3, 4]. There is a second molecularly distinct progenitor population within the MM that surrounds the NPCs. These Forkhead box D1- (*Foxd1*, formerly known as *BF2*) expressing cells have been shown to give rise to a significant percentage of the interstitium and the renal capsule [5-7].

During development, the UB coordinates the proliferation and differentiation of nephron progenitors into the precursor of the nephron, the renal vesicle [8, 9]. Reciprocally, the MM (both the NPC and interstitial populations) induces the outgrowth and branching of the UB until the UB has formed an arborized epithelial network of tubules referred to as the collecting ducts [1, 10]. In addition to its role in regulating branching, the interstitium plays further roles in differentiation of the nephron progenitor population and patterning of the vasculature [5, 11-13]. Although frequently referred to in broad terms, the adult interstitial cell population includes renal fibroblasts and various smooth muscle cell types including vascular smooth muscle, pericytes, mesangial cells and the smooth muscle of the ureter and renal pelvis [6, 7]. Moreover, much of the

endocrine function of the kidney, including the production of renin and erythropoietin, is performed by interstitial cells [14, 15].

Given that the interstitium has diverse roles in renal development, structure and function, it seems likely that this cell population is molecularly heterogeneous. However, extensive molecular characterization has been lacking. Here, we perform single cell RNA sequencing of Foxd1-derived interstitial cells from E18.5 mouse kidneys in order to define the heterogeneity and thus facilitate further inquiry into the development and function of these cells.

Our analysis revealed striking transcriptional heterogeneity in the renal interstitium, identifying 17 unique cellular clusters. Antisense mRNA *in situ* hybridization analysis demonstrated unexpected regionalization, uncovering at least 12 histologically similar but anatomically distinct domains along the cortical medullary axis. Importantly, comparison of mouse and human single cell data reveals that the interstitial heterogeneity is conserved. Analysis of transcription factor activity (regulons) showed cluster specificity, with the transcriptional regulator beta-catenin active within a medullary sub-population of stroma. Genetic ablation studies showed beta-catenin played a cell autonomous role in formation of this medullary interstitial region and a non-autonomous role in the development of the adjacent medullary epithelia. These findings stimulate multiple questions regarding the nature of interstitial-parenchymal crosstalk in development, physiology, homeostasis and disease of the embryonic and adult kidney.

RESULTS

Transcriptional analysis of the renal interstitium reveals molecular heterogeneity

86 Interstitial/stromal cells are present throughout the kidney extending from the
 87 cortex to the most medullary regions of the renal papillae and surrounding the ureter.
 88 Within the interstitium, three molecularly and anatomically distinct regions of interstitial
 89 cells have previously been defined. These populations were annotated as nephrogenic,
 90 cortical and medullary interstitium based on their unique anatomical positions [16].
 91 However, more detailed examination of kidneys stained with various antibodies to
 92 proteins expressed in interstitial cells suggests that the degree of heterogeneity may be
 93 under-estimated [7]. For example, Foxd1 is expressed in a small subset of the interstitial
 94 cells cortical and lateral to the Six2-positive cap mesenchyme. Expression ceases
 95 adjacent to the renal vesicles (Figure 1a-b). In comparison, Tenascin-C is expressed in a
 96 subset of Foxd1 expressing cells lateral to the Six2-positive cap mesenchyme but is not
 97 expressed in the cells cortical to the cap. Tenascin-C expression extends into the cells
 98 adjacent to the proximal tubule (Figure 1c-d). Slug expression appears to be largely, if
 99 not completely, non-overlapping with Foxd1, marking a subset of interstitial cells just
 100 medial to the cap mesenchyme and extending medially to the level of the proximal
 101 tubules (Figure 1 e-f). Acta2 is detectable in the interstitial cells adjacent to the proximal
 102 tubules but is not detectable in the medullary interstitium of the renal papillae (Figure
 103 1g-h). CDKN1c is expressed in the majority, if not all, interstitial cells of the renal papilla.
 104 Lef1 is expressed in interstitial cells extending from just medullary to the renal vesicles
 105 through the entire papillae (Figure 1i-j). However, in contrast to CDKN1c, Lef1 appears
 106 to only be expressed in a single layer of interstitial cells that lie directly adjacent to the
 107 collecting ducts (Figure 1k-l). Tbx18 is only expressed in interstitial cells surrounding the
 108 renal pelvis and ureter (Figure 1m). Interestingly, of all the proteins discussed, only

Acta2 is expressed in the smooth muscle surrounding the vasculature and mesangial cells, both of which are Foxd1-derived interstitial cell types. Based on the spatial differences in expression of the proteins described above and recent studies, one would predict that there are may be as many as 10 distinct interstitial cell types within the kidney.

To gain a more complete understanding of interstitial heterogeneity within the developing kidney, we performed single cell RNA sequencing (scRNA-seq) on dissociated E18.5 wild type mouse kidneys. Although previous single cell analyses have been performed on both adult and embryonic kidneys [17-20], the interstitium represents a relatively small percentage of the total number of cells and thus has been under-represented. To enrich for interstitium, we purified cells via fluorescence activated cell sorting (FACS) from Foxd1Cre;Rosa-Tomato kidneys as Foxd1-positive cells have been shown to give rise to the majority of the renal interstitium [7, 21]. Further, because recent studies suggest that a sub-population of the interstitium (in particular the ureteric fibroblasts/smooth muscle) is derived from a distinct, Tbx18-positive progenitor, and thus might not be present in our isolated cells, we bioinformatically isolated interstitial cells from published datasets derived from whole kidneys (1,482 total cells) and included them in our analysis [17]. Unsupervised clustering was performed on all sequenced cells that met quality control standards. After pre-processing, quality control, normalization for cell cycle phase and lineage filtering, 8,683 interstitial cells were selected for further analysis.

Shared nearest neighbor clustering based on similarity between gene expression profiles revealed 17 distinct clusters of cells that were identified as “interstitial” (Figure 2,

Supplemental Figure 1a, Supplemental Video 1 and Supplemental Video 2). Clusters ranged in size from 16 cells (cluster 13) to 1564 cells (cluster 10).

Importantly, the only cluster that was not derived from the Foxd1-Cre isolated cells was cluster 13, which represents the Tbx18-derived ureteric interstitium. These data suggest that our analysis includes most, if not all, renal interstitial cells, and that this population shows a high level of heterogeneity. We next sought to validate this heterogeneity in situ in order to gain insight into the significance and nature of the cellular diversity.

Renal interstitium shows spatial heterogeneity

To validate our results, we performed section in situ hybridization with over 50 differentially expressed genes (DEGs) from each cluster (Figure 3-5, Supplemental Figure 1c-k and Supplemental Table 1, and Supplemental Table 2). Candidates were chosen based on relative abundance and any identified regionalized expression observed in publicly available databases [22-24]. Although all DEGs examined were expressed in the interstitium, some that were indicated as being differentially expressed between clusters did not show cell type specific expression by in situ hybridization (Supplemental Table 1 and Supplemental Table 2). Instead, these genes appeared to be ubiquitously expressed suggesting that their being called as a DEG was due to differences in mRNA levels between different cell types rather than cell type specificity. A subset of the queried genes were expressed broadly in epithelial and/or endothelial populations as well as the interstitium. Both of these classes of genes were largely excluded from further analysis. Several DEGs appeared to be enriched in the interstitium over other cell types and showed regionalized expression (see below). It is

important to note that very few DEGs were detected in only a single cluster. Although individual examples for each cluster are presented in the main figures, anatomical assignment of clusters is based on in situ data from multiple DEGs (see supplemental Table 2). All in situ data will be available at the Re-Building a Kidney website (<https://www.rebuildingakidney.org/>).

Clusters 1-3 are highlighted by expression of *Foxd1*, a gene known to be expressed in the cortical interstitium (Figure 4a-b, Supplemental Figure 1d, Supplemental Table 1) and podocytes, a non-interstitial cell type. *Foxd1* expressing cells that were disjoint from interstitial cells and expressed typical podocyte markers, e.g. *Podxl* and *Nphs2*, were excluded from further analysis.

To gain insight into the nature of clusters 1-3, we assayed the expression of several mRNAs that were differentially expressed between the clusters including *Netrin1* (clusters 1 and 3), *Fibin* (clusters 2 and 3), *Smoc2* (cluster 1) and *Dlk1* (cluster 1) (Figure 3, Figure 4a-b', Supplemental Figure 1c-g, Supplemental Table 1). *Netrin1* (clusters 1 and 3) was uniformly expressed in the interstitial cells cortical but not lateral to the cap mesenchyme (Figure 3c-d). *Fibin* (clusters 2 and 3), shows expression in a subset of the cortical interstitium as well as expression in cells that lie lateral to the cap mesenchyme (Figure 3e-f). *Smoc2* (clusters 1 and 9) and *Dlk1* (cluster 1, 13 and 16) both showed mosaic expression in the cortical most interstitium (Figure 3g-j) as well as other, non-cortical cell types (described below). Due to the limited resolution of in situ hybridization, we could not determine whether *Fibin* was expressed in distinct cell types from *Smoc2* and *Dlk1*. However, this analysis suggests that the cortical *Foxd1* expressing cells are not molecularly homogeneous and these observations suggest that

clusters 1 and 3 are unique subsets within the cortical-most subset of Foxd1-expressing cells and cluster 2 represents a lateral sub-population of Foxd1-expressing cells. Thus, the region of stroma previously annotated as “cortical” appears to have at least 3 molecularly distinct cell types.

Genes present in clusters 4-5 were expressed on the medullary side of the ureteric bud tips surrounding the newly forming renal vesicles and correlated with a region of interstitium previously referred to as the nephrogenic interstitium (Supplemental Table 1 and data not shown). These clusters were highlighted by the expression of multiple genes involved in cell division (even after controlling for cell cycle phase).

Lysyl oxidase (LOX, clusters 6-8) was enriched in the interstitium adjacent to the most-medullary proximal tubules (Figure 4c-d', Supplemental Figure 1h Supplemental Table 1). Clca3a1 (cluster 7) was also enriched in cells adjacent to the proximal tubules although it appeared not to be expressed adjacent to the more medullary proximal tubules and its expression appeared more mosaic than Lox (Figure 4e-f').

Smoc2 (Cluster 9) is enriched in a population of interstitium that lies just medullary to the proximal tubule in the outer medulla (Figure 4g-h', Supplemental Figure 1g, Supplemental Table 1). Smoc2 expressing cells do not appear to expand cortically into the region adjacent to the proximal tubules.

The mRNA for Proenkephalin (*Penk*, clusters 9-12) is most intense in a population of interstitium just medullary to Smoc2 in the outer stripe of the inner medulla (Figure 4i-j', Supplemental Figure 1i, Supplemental Table 1). Although its expression

does appear to expand into more cortical populations of cells, it does not appear to overlap with *Smoc2*, *Lox* or *Clca3a1*.

Wnt4 is enriched in clusters 11-12. In situ analysis shows that along with expression in the pre-tubular aggregates, it is highly expressed in the interstitium of the papillary region of the kidney, a region previously referred to as the medullary interstitium. (Figure 4k-l', Supplemental Figure 1j, Supplemental Table 1 and data not shown).

Cluster 13 is the only population that was not derived from the *Foxd1* lineage sorted cells. All cluster 13 cells were derived from the whole kidney single cell data generated by Combes and colleagues [17]. Of note, cluster 13 was composed of only 16 cells, most likely a reflection of the relatively small numbers of interstitial cells included in the whole organ studies. As expected, genes present in cluster 13 (e.g. *Dlk1* and *Tbx18*) were predominantly expressed in the interstitium adjacent to the ureter (Figure 1m, Figure 5a-b, Supplemental Figure 1c, Supplemental Table 1).

Clusters 14-16 showed a high degree of overlap in gene expression. All three clusters expressed several genes that are accepted markers of perivascular cells. To determine if the clusters represented unique perivascular cell types, we performed in situ hybridization with cluster specific DEGs. *Akr1b7* and *Ren1* are both enriched in cluster 14 over other clusters and show expression in vascular smooth muscle as well as cells within the juxtaglomerular apparatus (Figure 5c-d, Supplemental Figure 1b, Supplemental Table 1, and data not shown). *Dlk1* (clusters 15 and 16 along with clusters 1 and 13) was detectable in the glomerular mesangial cells (Figure 5a-b, Supplemental Figure 1c, Supplemental Table 1). *Igf1* was enriched in cluster 15 and

showed expression in a subset of interstitial cells in the papillary region of the kidney, similar to the anatomical location assigned to cluster 10. Based on these observations along with gene set enrichment analysis [28] based on automated text-mining of protein-cell type associations from the biomedical literature [29] (Supplemental Figure 2), we suggest that cluster 14 represent vascular smooth muscle, cluster 15 represents a papillary pericyte population and cluster 16 represents mesangial cells. However, given the molecular similarity between these three cell types, it is possible that greater sequencing depth/clustering will alter these annotations.

Cluster 17 contained many genes that were also DEGs in other clusters. Although we did identify a specific DEG list for cluster 17, the most differentially expressed genes encoded mitochondrial mRNAs and we have not been able to produce strong signal with any of these probes (Supplemental Table 1).

In all, analysis of the transcriptome of single interstitial cells generated 17 clusters of cells that have been assigned to at least 12 anatomically distinct cell types.

Pseudo-time analysis suggests multiple distinct developmental trajectories within the interstitium

Although interstitial heterogeneity is not completely unexpected, the heterogeneity in the histologically indistinguishable cells spanning the cortical-medullary axis was surprising. A simple explanation of the identity of these cells is that they merely represent different stages in a linear developmental trajectory from a cortical-most progenitor to a more medullary, differentiated cell type. To gain insight into the lineage relationship of the different clusters, we employed the complementary approaches of

RNA velocity [25] and pseudotemporal ordering [26]. Briefly, RNA velocity is determined by modeling the relationship between the unspliced and spliced forms of a transcript, with the reasonable assumption that a cell actively transcribing a gene will have a higher ratio of unspliced to spliced transcripts and so a higher “RNA velocity”. Conversely, cells that are not actively transcribing a gene (but still maintain its expression) will have a lower ratio of unspliced to spliced transcripts as the majority of mRNA will represent processed transcripts (lower RNA velocity). Using single genes as representative examples, this analysis indicates that Lef1 is actively transcribed in cluster 10 and stabilizes in clusters 11 and 12 suggesting that clusters 11 and 12 represent cells derived from cluster 10 (Supplemental Figure 3a-b). In comparison, Smoc2 is actively transcribed and processed in cells present within cluster 9 but not in any other clusters (other than the Foxd1+ progenitor cells), suggesting cluster 9 represents a terminal differentiation state (Supplemental Figure 3c-d). Extending this model to all genes and all cells within the purified interstitium, we created an RNA velocity field to predict transitions between clusters (Supplemental Figure 3e). It is important to note that the kidney capsule, which is derived from Foxd1-expressing cells [7], was removed from our kidneys prior to cell sorting and thus is not included in our analysis. Therefore, we cannot comment on the derivation of this cell type at this time. We also ultimately excluded the ureteral smooth muscle cells (cluster 13) from this analysis as they did not show a relationship with any of the other clusters, as expected from their independent origins [21].

Simulation of a reverse Markov process with RNA velocity-based transition probabilities (please refer to Materials and Methods for a more complete description)

allowed us to identify a group of cells that likely represented the ontological parent of the analyzed interstitial cells (Figure 6a). After selecting one such parent or “root” cell, we were able to pseudotemporally order the cells. Annotating the diffusion map with pseudotime demonstrated that the parent/root population ramifies into multiple distinct branches that develop simultaneously rather than following a single linear differentiation trajectory (Supplemental figure 3f).

The integration of the RNA-velocity and pseudotime analyses generates a model wherein at least two of the distinct clusters of Foxd1-positive “progenitor” cells undergo independent branching events to give rise to the cycling/proliferating cells and at least 6 different trajectories that we refer to as the inner medullary fibroblast, outer medullary fibroblast, vascular smooth muscle, proximal tubule interstitium, pericyte and mesangial subtypes (Figure 6b). The vascular smooth muscle cluster appears more distantly related than the other cell types consistent with its distinct differentiation profile (Figure 6b and Supplemental video 3). Interestingly, clusters 6 appears to be located at a bifucation or trifucation point that gives rise to clusters 14, 15 and 16. This data suggests that the distinct interstitial subtypes do not represent transient stages in a linear developmental trajectory. Instead, these cells appear to represent previously undescribed, anatomically distinct interstitial cell types of distinct lineages.

Although frequently referred to as a homogeneous population of cells that functions primarily as a scaffold, several recent studies have shown that in various systems, the interstitium sets up unique microenvironmental niches that direct tissue development and/or maintenance and can contribute to pathological conditions [27-34]. Indeed, within the kidney, studies have shown that the interstitium is important in

numerous developmental processes affecting both the epithelia and endothelia [5, 11-13]. We next sought to re-visit previous work in light of our current findings.

Interstitial pattern affects development of the renal parenchyma

Several transcriptional regulators have been shown to be expressed in and necessary for the development of the stroma. To determine if any of these factors were active in specific stromal subtypes, we employed the SCENIC package to reconstruct gene regulatory networks and measure their activity within cells and so within the cells' parent clusters (Figure 7) [35]. Strikingly, certain transcriptional signatures (aka regulons) were active in distinct clusters or groups of clusters while inactive in others. Interestingly, we found that the Lef1 regulon was predominantly active in clusters 8-12 (Figure 8a), which represents much of the medullary interstitium (Figure 5). Previous studies have shown that inactivation of beta-catenin in the interstitial progenitor population (using Foxd1Cre) leads to defects in the expression of several genes within the stroma, especially beta-catenin targets [36]. To test whether beta-catenin was necessary for formation of specific stromal subtypes (versus a general stromal defect), we assessed the expression of regionalized genes in Foxd1Cre;catnb^{-flox} kidneys. Expression of genes normally expressed in clusters 1-7 (clusters where beta-catenin is not active) appeared unaffected or even expanded in Foxd1Cre;catnb^{-flox} mutants (Figure 8b-d, h-j). In contrast, genes expressed in clusters 8-12 were markedly reduced or undetectable (Figure 8e-f, k-l) in mutants. The absence of medullary interstitium (clusters 8-12) correlated with a severe deficit in the formation of the epithelia (including the loop of Henle) that lay adjacent to these zones while the epithelia that lay adjacent

to the unaffected cortical interstitium (including the proximal tubules) appeared to form normally, as previously reported by Yu et al. [36] (Figure 8g, m).

Human fetal kidney interstitium shows a similar degree of heterogeneity to the mouse

Recently, several groups have employed single cell RNA sequencing on human fetal kidney at different stages [37]. While analyzing the nephrogenic region of fetal kidneys, which contains cells from multiple lineages, only 5 unique interstitial sub-clusters were identified [37]. Thus, we wondered whether extensive interstitial heterogeneity is unique to the mouse.

To understand whether interstitial heterogeneity is a generalizable phenomenon between these two species, we reanalyzed previously published [38, 39] week 17 human fetal kidney scRNA seq data. After identifying the major cell populations within the data (epithelia, endothelia, leukocytes, etc.), we bioinformatically isolated the cells defined as interstitium (see methods). We then deployed the same clustering technique used to cluster mouse interstitium on the interstitium of the human fetal kidney. We find that the interstitium of the cortical region of human fetal kidney segments into 13 molecularly distinct clusters (Figure 9a). Although the human data set was limited to more cortical cell populations, we were able to identify almost as many unique clusters as found in whole embryonic kidney. When the cluster assignments from E18.5 mouse interstitial cells were mapped onto the 17-week human interstitial cells, the majority were found to be represented in the human data although they did not resolve as clearly (Figure 9b). There are likely several factors underlying this imperfect alignment

including incomplete sampling in the human data, divergence between developmental time scales and time points between mouse and human and fewer cells available for analysis in the human dataset resulting in less resolving power. Evidence of the sampling bias is evident in that we did not identify any human cells that were analogous to cluster 13, the most medullary interstitium. These data indicate that renal interstitial heterogeneity is a generalizable characteristic between mouse and human.

DISCUSSION

Although known to play a role in providing physical support, growing evidence from multiple systems indicates that interstitial cells play active roles in tissue development, maintenance and disease. Within the developing kidney, non-autonomous roles for the interstitium have been identified in ureteric bud branching, nephron differentiation and blood vessel formation [5, 11-13]. Mechanistic insight into how the interstitium has so many distinct functions has been hampered by a poorly defined transcriptome. Here, using single cell RNA-sequencing combined with *in situ* hybridization, we have generated a map of interstitial gene expression in the E18.5 kidney. Our analysis shows a previously unappreciated level of heterogeneity in the interstitium of the developing mouse kidney. Analysis of human embryonic interstitial cells show correspondence in heterogeneity found in the embryonic mouse.

While previous work characterizing the heterogeneity of the developing mouse kidney uncovered 4 interstitial clusters (1,482 cells) [17], our informed analysis identified 17 distinct clusters that we were able to spatially resolve into at least 12 anatomically distinct subtypes. Additional analysis will be required to determine whether the 5

remaining clusters represent additional unique cell types or the data is currently overclustered. For example, we were not able to obtain signal from in situ hybridization with any DEGs specific to cluster 17, a population enriched for mitochondrial genes. Thus, it is possible that this cluster represents an artifact of the dissociation protocol. However, it is also worth noting that although immunostaining identified an interstitial cell type associated with the collecting ducts, our clustering did not conclusively identify these cells. It is possible that they lie within one of the medullary populations (12 or 13) or alternatively, that rather than being over-clustered, our data is under-clustered. We think it is likely that higher resolution techniques (e.g. single cell resolution in situ hybridization, antibody staining with or reporter gene generation from cluster specific DEGs) will resolve additional unique cell types. For example, there are three clusters containing genes expressed in the interstitium surrounding the proximal tubules. Although there do appear to be spatial differences in the expression of some proximal tubule interstitial genes, without cellular resolution, we cannot at this point determine whether these cells represent unique cell types. The fact that one of the, cluster 6, shows similarity to and is predicted to be the parent of clusters 14, 15 and 16, suggests that this cell type may represent a spatially distinct mural cell.

Unexpectedly, we found at least 3 molecularly distinct clusters within the *Foxd1* expression domain. This observation raises the question of the identity of the true interstitial progenitor cell. One possibility that is supported by our RNA velocity analysis, is that rather than containing a single multipotent progenitor cell, the *Foxd1* domain is comprised of several lineage restricted progenitors. Testing this will require

more detailed molecular characterization as well as lineage tracing with cell type specific Cre drivers.

Interestingly, our analysis identified several zones of histologically indistinguishable but molecularly distinct fibroblast-like cell types that occupy unique spatial locations along the cortical medullary axis, where they correlate with distinct anatomical regions in the adjacent parenchyma. By analyzing our transcriptomic data, we were able to identify signaling pathways unique to the distinct interstitial clusters. Reanalysis of beta-catenin mutant interstitium reveals a unique role for this factor in the development of the papillary stroma, which secondarily affected the development of the adjacent epithelia. Of note, previous work has shown that mesenchymal cells play instructive roles in patterning the adjacent epithelia during development of various organ systems including patterning of the vertebrate gut tube [40-43]. The close correlation of the distinct kidney interstitial cell types with the functional subdomains within the nephron raises the intriguing possibility of stromal-epithelial cross talk that is involved in the patterning and/or differentiation of the kidney parenchyma. Although several studies have revealed cell autonomous mechanisms underlying nephron patterning, it is possible that distinct interstitial subpopulations produce factors that interact with intrinsic pathways to assure proper position and relative size and spacing of the nephron segments with the adjacent collecting ducts and associated renal vasculature. RNA-seq data reveals multiple growth factors, small molecules, extracellular matrix components and metabolites that appear to be regionally produced. Further genetic analysis will be required to test specific roles.

404 Finally, as the adult kidney shows exquisite patterning along the
405 cortical/medullary axis, it will be of great interest to determine whether a similar degree
406 of heterogeneity and pattern exists in the adult interstitium and how this pattern
407 correlates to normal anatomy, physiology, injury, regeneration and disease state. Given
408 the growing evidence of the essential nature of the interstitium in multiple processes
409 [27-34], a similar analysis of interstitial heterogeneity in different organ systems at
410 different stages may reveal that the interstitium's role in patterning and morphogenesis
411 is a generalizable principle.

MATERIALS and METHODS

Mice

All animals were housed, maintained and used according to protocols approved by the Institutional Animal Care and Use Committees at the University of Texas Southwestern Medical Center and following the guidelines from the NCI-Frederick Animal Care and Use Committee. For each experiment, female mice of 7–8 weeks of age were crossed with a male of 9–10 weeks of age. Plugs were checked and the embryos were collected at the desired time points for further analysis. Noon of the day on which the mating plug was observed was designated embryonic day (E) 0.5. The following mice were used in the studies described: *Foxd1*Cre (JAX Stock #012463), *Rosa26*Tomato (JAX Stock #007909), *Rosa26*DTA (JAX Stock #006331), *Rosa26*YFP (JAX Stock #006148), *catnb* null and *catnb* flox [44].

In situ hybridization

For section *in situ* hybridization, kidneys isolated at specific stages were fixed overnight in 4% PFA (in PBS) at 4 °C and cryopreserved in 30% sucrose. Tissues were frozen in OCT (Tissue Tek) and sectioned at 10 µm. Sections were subjected to *in situ* hybridization as previously described [9]. The following antisense RNA probes against *Foxd1*, *Lox*, *Smoc2*, *Penk*, *Wnt4*, *Lef1*, *Tgfb1i1*, were linearized and transcribed as previously described. Plasmids were unavailable for *Dlk1* and *Akr1b7*; thus, single stranded DNA for each gene was purchased with the T7 RNA polymerase binding site in the reverse orientation added to 3' end of the gene sequence. These probes were made through RNA transcription of these single stranded DNA gblocks using T7 polymerase.

435 *Histology, immunohistochemistry and immunocytochemistry*

436 Kidneys isolated at birth were formaldehyde fixed and paraffin embedded. Sections
 437 (5 µm) from paraffin-embedded kidneys were subjected to haematoxylin and eosin
 438 staining. For immunohistochemistry, fixed kidneys were embedded in OCT and
 439 sectioned on a cryostat (10 µm). Frozen sections were washed with PBS and blocked
 440 with 5% serum for an hour at room temperature and incubated with primary antibodies
 441 at 4 °C overnight. After primary incubation, sections were washed and incubated with
 442 HRP-tagged secondary antibodies for 1 h at room temperature. Further, signal was
 443 detected with tyramide amplification. Slides were washed and re-stained with additional
 444 markers according to the above-mentioned immunohistochemistry protocol. Slides were
 445 then mounted with Vectashield and images were captured with a Zeis LSM500,
 446 ZeisLSM700 or a Nikon A1R confocal microscope. The following antibodies were used:

Protein Target	Antibody Product ID	Dilution
Acta2 (aSMA)	Sigma-Aldrich Cat# A2547, RRID:AB_476701	1:250
Cdkn1c	ThermoFisher Cat# PA5-34432, RRID:AB_2551784	1:250
DBA	Vector Laboratories Cat# B-1035, RRID:AB_2314288	1:500
E-Cadherin	Invitrogen Cat#13-1900, RRID:AB_2533005	1:250
E-Cadherin	BD Biosciences Cat# 610182, RRID:AB_397581	1:250
Foxd1	Santa Cruz Cat# SC-47585; RRID:AB_2105295	1:750
GFP	Aves Labs Cat# GFP-1020, RRID:AB_10000240	1:500
Laminin	Sigma-Aldrich Cat# L9393, RRID:AB_477163	1:500
Lef1	Cell Signaling Technology Cat# 2230, RRID:AB_823558	1:250
LTL	Vector Laboratories Cat# B-1325, RRID:AB_2336558	1:500
Meis1/2	Santa Cruz Cat# SC-10599, RRID:AB_2143020	1:750
Meis1/2/3	Active Motif Cat# 39795, RRID:AB_2750570	1:100
pan Cytokeratin (CK)	Sigma-Aldrich Cat# C2562, RRID:AB_476839	1:500
PDGFrβ	Cell Signaling Cat# 3169S, RRID:AB_2162497	1:250
Six2	Proteintech Group Cat# 11562-1-AP, RRID:AB_2189084	1:500

Slug	Cell Signaling Technology Cat# 9585, RRID:AB_2239535	1:250
Tbx18	Santa Cruz Cat# SC-17869, RRID:AB_2200374	1:100
THP	Alfa Aesar Cat# J65429	1:500

Single cell sample preparation and sequencing

E18.5 Foxd1Cre;Rosa26Tomato mouse kidneys were dissected in cold PBS without calcium or magnesium, Kidneys were cleaned and adrenal gland, capsule and ureters were removed.. Kidneys were washed in HBSS for 2 minutes at 37°C then minced using two razorblades on ice. The minced kidneys were digested for 8 minutes at 37C in 2mL of 0.25% w/v Collagenase A/1% w/v Pancreatin (Sigma-Aldrich; Cat. 101378001, P1750), with manual dissociation via pipetting through a P1000 tip every 2 minutes. No more than 4 pairs of kidneys were dissociated in 2mL enzyme digest. Digestion was inactivated by adding 125ul serum. After pelleting the cells at 400g for 5 minutes, cells were resuspended in 1mL of AutoMACS Running Buffer (Miltenyi Biotec, Cat. 130-091-221) and passed through a 30 um pre-separation filter (Milteni Biotec, Cat. 130-041-407). Filters were immediately washed with 500ul AutoMACS Running Buffer. Cells were resuspended in 500uL AutoMACS Running Buffer and filtered at least 2 more times through a cell-strainer cap (Falcon, Cat. 352235) attached to a 5mL Polypropylene round bottom tube (Globe Scientific, Cat. 110428). Ten thousand Foxd1Cre;Rosa26Tomato+ cells, isolated via FACS, were run on a chromium 10x Single Cell Chip (10x Genomics). Libraries were prepared using Chromium Single Cell Library kit V2, and sequenced on an Illumina NextSeq using 75pb paired-end sequencing. Sequencing resulted in an average of 12,000 reads/cell and 3,000 genes/cell. Upon acceptance, the single cell data presented in this manuscript will be deposited onto Gene Expression Omnibus and Rebuilding a Kidney databases.

Single cell data analysis

When analyzing the single cell data collected from the experiments outlined above, we included, where possible, the dataset generated by Combes et al. obtained via NCBI's Gene Expression Omnibus under accession GSE108291 [17]. Each batch was processed independently using the *scrn* Bioconductor package [45]. Unfiltered feature-barcode matrices were generating by running the CellRanger count pipeline. Cells were called from empty droplets by testing for deviation of the expression profile for each cell from the ambient RNA pool [46]. Cells with large mitochondrial proportions, i.e. more than 3 mean-absolute deviations away from the median, were removed. Cells were pre-clustered, a deconvolution method was applied to compute size factors for all cells [47] and normalized log-expression values were calculated. Variance was partitioned into technical and biological components by assuming technical noise was Poisson-distributed and attributing any estimated variance in excess of that accounted for by a fitted Poisson trend to biological variation. The dimensionality of the data set was reduced by performing principal component analysis and discarding the later principal components for which the variance explained was less than variance attributable to technical noise.

Masking of biological effects by expression changes due to cell cycle phase were mitigated by blocking on this covariate. The cell cycle phase was inferred using the pair-based classifier implemented in the *cyclone* function of *scrn*. Corrected log-normalized expression counts were obtain by calling the `removeBatchEffect` from the *limma* [48] Bioconductor package with a design formula including G1 and G2M cell cycle phase scores as covariates.

A single set of features for batch correction were obtained by computing the average biological component of variation across batches and retaining those genes with a positive biological component. The batches were rescaled and log-normalized expression values recomputed after the size factors were adjusted for systemic differences in sequencing depth between batches. Batch effects were corrected by matching mutual nearest neighbors in the high-dimensional expression space [49]. The resulting reduced-dimensional representation of the data was used for all subsequent embeddings including t-SNE and UMAP.

Cells were clustered by building a shared nearest neighbor graph[50] and executing the Walktrap algorithm [51]. Differential gene expression analysis was performed using the two-part generalized linear model that concurrently models expression rate above background and expression mean implemented in MAST[52]. A one-versus-all strategy was employed comparing each cluster to all other identified interstitial clusters.

Gene sets for enrichment analysis were obtained from the TISSUES Text-mining Tissue Protein Expression Evidence Scores dataset [53] located at <http://amp.pharm.mssm.edu/Harmonizome/>. The gene sets were filtered to include only those genes with a standardized value greater than 1. Enrichment analysis was performed using the fgsea [54] Bioconductor package.

Week 17 human fetal kidney scRNA seq data was obtained from the Gene Expression Omnibus under series accession numbers GSE112570 and GSE124472 [38, 39]. Cluster assignments were transferred from the E18.5 mouse kidney dataset to the 17-week human fetal kidney dataset using a neural network classifier constructed

using the Tensorflow system [55]. Graph regularization [56] considering eight shared nearest neighbors of each cell was used during training of a sequential network with two hidden layers, each containing 1024 hidden nodes, to classify the expression profiles of cells from the mouse dataset by cluster. Orthologue-mapped expression profiles of human cells as input to the classifier to assign each cell in the human dataset to mouse cluster with greatest similarity. Expression profiles were cosine-normalized prior to training or prediction. Only genes with a biological-to-technical variance ratio greater than zero were utilized for classification.

RNA velocity was calculated to analyze the dynamic relationships between identified cell states [25]. The analysis was limited to the replicate single cell datasets produced for this paper as BAM files were not available for the Combes, *et al.* dataset [17]. Read counts were partitioned between spliced, unspliced and ambiguous sets by calling `velocyto run10x` with an mm10 repeat mask obtained from the UCSC genome browser and the genome annotation file that came prepackaged with `cellranger`. The results of previously conducted cell filtering and feature selection were applied to these expression matrices. Normalization, principal component analysis, k-nearest neighbor smoothing, gamma fit, extrapolation, Markov process modelling and projection onto pre-defined low-dimensional embeddings were executed by calling the relevant functions from the `velocyto.analysis` module.

To reconcile clustering with trajectory inference, we performed partition-based graph abstraction (PAGA) to model the connectivity between clusters [57]. At cluster resolution, the edge-score threshold was varied until the graph was decomposed into connected components. Evaluation of marker gene expression within the components

permitted assignment of components to the categories epithelium, leukocyte, erythrocyte, endothelium and interstitium. In many cases, RNA velocity allowed for assigning a direction to the edges between clusters indicating a tendency of transition between the clusters.

Gene regulatory network reconstruction and measurement of regulon activity within each was conducted using SCENIC [35]. All cells of acceptable quality were used for network inference. The regulon activity was binarized using the following strategy. An attempt was made to model the regulon's activity as a mixture of two normal distributions using the mixtools R package. If such a model could be fit using expectation maximization, then those cells for which the regulon activity were assigned higher probability by the distribution with the greater mean were assigned a binarized regulon activity of one and zero otherwise. If a two-component mixture of normal distributions could not be fit, then a beta distribution was fit to the regulon's activity and cells for which the regulon's activity was greater than one mean absolute deviation above the mean were assigned a binarized regulon activity of one and zero otherwise.

Acknowledgements: The authors would like to thank Ondine Cleaver, Denise Marciano, and Phoebe Carter for their insight in preparing this manuscript.

Declaration of Interests: The authors have no competing interests

Funding: This work was supported by a fellowship from the UT Southwestern Hamon Center for Regenerative Science and Medicine to ARF, NIH grants DK095057, DK106743, DK090127 to TJC and F31DK122670 to ARF and the UT Southwestern George O'Brien Kidney Research Core DK079328.

Data Availability: Upon acceptance, the single cell data presented in this manuscript will be deposited onto Gene Expression Omnibus and Rebuilding a Kidney databases with DOIs included.

Author Contributions: ARF designed experiments, performed experiments, analyzed data and wrote the manuscript. CPC designed experiments, analyzed data and wrote the manuscript. AD designed experiments, performed experiments and analyzed data, KD and MP designed experiments. CSK designed experiments, performed experiments, analyzed data and wrote the manuscript. AM, DA, DS, and GH prepared single cell libraries. TJC designed experiments, analyzed data and wrote the manuscript.

REFERENCES

1. Grobstein, C., *Inductive interaction in the development of the mouse metanephros*.
Journal of Experimental Zoology, 1955. **130**(2): p. 319-339.
2. Grobstein, C., *Morphogenetic interaction between embryonic mouse tissues separated
by a membrane filter*. Nature, 1953. **172**(4384): p. 869-70.
3. Boyle, S., et al., *Fate mapping using Cited1-CreERT2 mice demonstrates that the cap
mesenchyme contains self-renewing progenitor cells and gives rise exclusively to
nephronic epithelia*. Dev Biol, 2008. **313**(1): p. 234-45.
4. Kobayashi, A., et al., *Six2 defines and regulates a multipotent self-renewing nephron
progenitor population throughout mammalian kidney development*. Cell Stem Cell, 2008.
3(2): p. 169-81.
5. Hatini, V., et al., *Essential role of stromal mesenchyme in kidney morphogenesis
revealed by targeted disruption of Winged Helix transcription factor BF-2*. Genes Dev,
1996. **10**(12): p. 1467-78.
6. Humphreys, B.D., et al., *Fate tracing reveals the pericyte and not epithelial origin of
myofibroblasts in kidney fibrosis*. Am J Pathol, 2010. **176**(1): p. 85-97.
7. Kobayashi, A., et al., *Identification of a multipotent self-renewing stromal progenitor
population during mammalian kidney organogenesis*. Stem Cell Reports, 2014. **3**(4): p.
650-62.
8. Carroll, T.J., et al., *Wnt9b plays a central role in the regulation of mesenchymal to
epithelial transitions underlying organogenesis of the mammalian urogenital system*. Dev
Cell, 2005. **9**(2): p. 283-92.
9. Karner, C.M., et al., *Canonical Wnt9b signaling balances progenitor cell expansion and
differentiation during kidney development*. Development, 2011. **138**(7): p. 1247-57.
10. Grobstein, C., *Inductive epitheliomesenchymal interaction in cultured organ rudiments of
the mouse*. Science, 1953. **118**(3054): p. 52-5.

- 599 11. Das, A., et al., *Stromal-epithelial crosstalk regulates kidney progenitor cell differentiation*.
600 Nat Cell Biol, 2013. **15**(9): p. 1035-44.
- 601 12. Hum, S., et al., *Ablation of the renal stroma defines its critical role in nephron progenitor*
602 *and vasculature patterning*. PLoS One, 2014. **9**(2): p. e88400.
- 603 13. Hurtado, R., et al., *Pbx1-dependent control of VMC differentiation kinetics underlies*
604 *gross renal vascular patterning*. Development, 2015. **142**(15): p. 2653-64.
- 605 14. Maxwell, P.H., et al., *Sites of erythropoietin production*. Kidney Int, 1997. **51**(2): p. 393-
606 401.
- 607 15. Berg, A.C., et al., *Pericytes synthesize renin*. World J Nephrol, 2013. **2**(1): p. 11-6.
- 608 16. Little, M.H., et al., *A high-resolution anatomical ontology of the developing murine*
609 *genitourinary tract*. Gene Expr Patterns, 2007. **7**(6): p. 680-99.
- 610 17. Combes, A.N., et al., *Single cell analysis of the developing mouse kidney provides*
611 *deeper insight into marker gene expression and ligand-receptor crosstalk*. Development,
612 2019. **146**(12).
- 613 18. Wu, H., et al., *Advantages of Single-Nucleus over Single-Cell RNA Sequencing of Adult*
614 *Kidney: Rare Cell Types and Novel Cell States Revealed in Fibrosis*. J Am Soc Nephrol,
615 2019. **30**(1): p. 23-32.
- 616 19. Lindstrom, N.O., et al., *Single-Cell RNA Sequencing of the Adult Mouse Kidney: From*
617 *Molecular Cataloging of Cell Types to Disease-Associated Predictions*. Am J Kidney Dis,
618 2019. **73**(1): p. 140-142.
- 619 20. Park, J., et al., *Single-cell transcriptomics of the mouse kidney reveals potential cellular*
620 *targets of kidney disease*. Science, 2018. **360**(6390): p. 758-763.
- 621 21. Bohnenpoll, T., et al., *Tbx18 expression demarcates multipotent precursor populations in*
622 *the developing urogenital system but is exclusively required within the ureteric*
623 *mesenchymal lineage to suppress a renal stromal fate*. Dev Biol, 2013. **380**(1): p. 25-36.

- 624 22. McMahon, A.P., et al., *GUDMAP: the genitourinary developmental molecular anatomy*
625 *project*. J Am Soc Nephrol, 2008. **19**(4): p. 667-71.
- 626 23. Harding, S.D., et al., *The GUDMAP database--an online resource for genitourinary*
627 *research*. Development, 2011. **138**(13): p. 2845-53.
- 628 24. Visel, A., C. Thaller, and G. Eichele, *GenePaint.org: an atlas of gene expression*
629 *patterns in the mouse embryo*. Nucleic Acids Res, 2004. **32**(Database issue): p. D552-6.
- 630 25. La Manno, G., et al., *RNA velocity of single cells*. Nature, 2018. **560**(7719): p. 494-498.
- 631 26. Haghverdi, L., et al., *Diffusion pseudotime robustly reconstructs lineage branching*.
632 *Nature Methods*, 2016. **13**: p. 845.
- 633 27. Benias, P.C., et al., *Structure and Distribution of an Unrecognized Interstitium in Human*
634 *Tissues*. Sci Rep, 2018. **8**(1): p. 4947.
- 635 28. Liotta, L.A. and E.C. Kohn, *The microenvironment of the tumour-host interface*. Nature,
636 2001. **411**(6835): p. 375-9.
- 637 29. Yu, V.W. and D.T. Scadden, *Hematopoietic Stem Cell and Its Bone Marrow Niche*. Curr
638 *Top Dev Biol*, 2016. **118**: p. 21-44.
- 639 30. Greicius, G. and D.M. Virshup, *Stromal control of intestinal development and the stem*
640 *cell niche*. Differentiation, 2019.
- 641 31. Driskell, R.R. and F.M. Watt, *Understanding fibroblast heterogeneity in the skin*. Trends
642 *Cell Biol*, 2015. **25**(2): p. 92-9.
- 643 32. Driskell, R.R., et al., *Distinct fibroblast lineages determine dermal architecture in skin*
644 *development and repair*. Nature, 2013. **504**(7479): p. 277-281.
- 645 33. De Palma, M., D. Biziato, and T.V. Petrova, *Microenvironmental regulation of tumour*
646 *angiogenesis*. Nat Rev Cancer, 2017. **17**(8): p. 457-474.
- 647 34. Hepler, C., et al., *Identification of functionally distinct fibro-inflammatory and adipogenic*
648 *stromal subpopulations in visceral adipose tissue of adult mice*. Elife, 2018. **7**.

35. Aibar, S., et al., *SCENIC: single-cell regulatory network inference and clustering*. Nat Methods, 2017. **14**(11): p. 1083-1086.
36. Yu, J., et al., *A Wnt7b-dependent pathway regulates the orientation of epithelial cell division and establishes the cortico-medullary axis of the mammalian kidney*. Development, 2009. **136**(1): p. 161-71.
37. Lindström, N.O., et al., *Conserved and Divergent Features of Mesenchymal Progenitor Cell Types within the Cortical Nephrogenic Niche of the Human and Mouse Kidney*. Journal of the American Society of Nephrology : JASN, 2018. **29**(3): p. 806-824.
38. Lindstrom, N.O., et al., *Progressive Recruitment of Mesenchymal Progenitors Reveals a Time-Dependent Process of Cell Fate Acquisition in Mouse and Human Nephrogenesis*. Dev Cell, 2018. **45**(5): p. 651-660.e4.
39. Tran, T., et al., *In Vivo Developmental Trajectories of Human Podocyte Inform In Vitro Differentiation of Pluripotent Stem Cell-Derived Podocytes*. (1878-1551 (Electronic)).
40. Golosow, N. and C. Grobstein, *Epitheliomesenchymal interaction in pancreatic morphogenesis*. Dev Biol, 1962. **4**: p. 242-55.
41. Kim, T.H., et al., *Endodermal Hedgehog signals modulate Notch pathway activity in the developing digestive tract mesenchyme*. Development, 2011. **138**(15): p. 3225-33.
42. Le Guen, L., et al., *Mesenchymal-epithelial interactions during digestive tract development and epithelial stem cell regeneration*. Cell Mol Life Sci, 2015. **72**(20): p. 3883-96.
43. McCulley, D., M. Wienhold, and X. Sun, *The pulmonary mesenchyme directs lung development*. Curr Opin Genet Dev, 2015. **32**: p. 98-105.
44. Brault, V., et al., *Inactivation of the beta-catenin gene by Wnt1-Cre-mediated deletion results in dramatic brain malformation and failure of craniofacial development*. Development, 2001. **128**(8): p. 1253-64.

45. Lun, A.T., D.J. McCarthy, and J.C. Marioni, *A step-by-step workflow for low-level analysis of single-cell RNA-seq data with Bioconductor*. F1000Res, 2016. **5**: p. 2122.
46. Lun, A.T.L., H. Pages, and M.L. Smith, *beachmat: A Bioconductor C++ API for accessing high-throughput biological data from a variety of R matrix types*. PLoS Comput Biol, 2018. **14**(5): p. e1006135.
47. L. Lun, A.T., K. Bach, and J.C. Marioni, *Pooling across cells to normalize single-cell RNA sequencing data with many zero counts*. Genome Biology, 2016. **17**(1): p. 75.
48. Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing and microarray studies*. Nucleic Acids Res, 2015. **43**(7): p. e47.
49. Haghverdi, L., et al., *Batch effects in single-cell RNA-sequencing data are corrected by matching mutual nearest neighbors*. Nat Biotechnol, 2018. **36**(5): p. 421-427.
50. Xu, C. and Z. Su, *Identification of cell types from single-cell transcriptomes using a novel clustering method*. Bioinformatics, 2015. **31**(12): p. 1974-1980.
51. Pons, P.L., Matthieu, *Computing Communities in Large Networks Using Random Walks*. Journal of Graph Algorithms and Applications, 2006. **10**(2): p. 191-218.
52. Finak, G., et al., *MAST: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data*. Genome Biology, 2015. **16**(1): p. 278.
53. Santos, A., et al., *Comprehensive comparison of large-scale tissue expression datasets*. PeerJ, 2015. **3**: p. e1054.
54. Sergushichev, A.A., *An algorithm for fast preranked gene set enrichment analysis using cumulative statistic calculation*. bioRxiv, 2016: p. 060012.
55. Nagalakshmi, V.K. and J. Yu, *The ureteric bud epithelium: morphogenesis and roles in metanephric kidney patterning*. Mol Reprod Dev, 2015. **82**(3): p. 151-66.
56. Bui, T.D., S. Ravi, and V. Ramavajjala, *Neural Graph Machines: Learning Neural Networks Using Graphs*. 2017.

- 700 57. Wolf, F.A., et al., *PAGA: graph abstraction reconciles clustering with trajectory inference*
701 *through a topology preserving map of single cells*. Genome Biol, 2019. **20**(1): p. 59.
702

FIGURE LEGENDS

Graphical Abstract: The developing interstitium is a highly heterogeneous, patterned population of cells that occupies distinct positions correlated to the adjacent parenchyma.

Figure 1: Unexpected heterogeneity within the renal interstitium revealed through antibody staining and single cell RNA sequencing. Wildtype E18.5 kidneys stained for Six2 and CK (red, blue respectively in a-m), and Foxd1 (green in a-b), Tenascin C (green in c-d), Slug (green in e-f), Acta2 (green in g-h), CDKN1c (green in i-j), Lef1 (green in k-l), Tbx18 (green in m). Scale bar 100um.

Figure 2: Interstitial heterogeneity revealed through single cell RNA sequencing: UMAP of E18.5 mouse interstitium. For a 3-dimensional representation of the UMAP see supplemental video 1 (a).

Figure 3: Single cell RNA sequencing reveals unexpected cortical heterogeneity within the renal interstitium that is validated through mRNA in situ hybridization. Specific gene expression displayed as a UMAP (a, c, e, g, i) for Foxd1(a), Netrin (c), Fibin (e), Smoc2 (g) and Dlk1 (i). E18.5 *in situ* hybridization of Foxd1, expressed in clusters 1-3 (b), Netrin1, expressed in clusters 1, 3 (d), Fibin expressed in clusters 2-3 (f), Smoc2, expressed in clusters 1 (h) and Dlk1, expressed in cluster 1 (j) reveals cortical heterogeneity. Scale bar: 100um

Figure 4: Spatial characterization of scRNA seq-clusters reveals cortico-medullary patterning among interstitial subtypes. Specific gene expression displayed as a UMAP (a, c, e, g, i, k) for Foxd1 (a), Lox (c), Clca3a1 (e), Smoc2 (g), Penk (i), and Wnt4 (k). E18.5 *in situ* hybridization of Foxd1: clusters 1-3 interstitial expression indicated with arrow, epithelial lineage expression in podocytes indicated with asterisk) (b-b'), Lox: cluster 7 (d-d'), Clca3a1: clusters 6-8 (f-f'), Smoc2: cluster 9 (h-h'), Penk: cluster 9-12 (j-j'), Wnt4: cluster 11-12 (interstitial expression indicated with arrow, epithelial lineage expression in PTA indicated with asterisk) (l-l'), reveals a molecular pattern which spans the cortico-medullary axis. b, d, f, h, j, l Scale bar: 500um. b', d', f', h', j', l' Scale bar: 100um.

Figure 5: Mesangium and pericyte populations detected using single cell RNA sequencing. mRNA *in situ* hybridization of Dlk1 expression in clusters 15-16 (a) and Akr1b7 expression in cluster 14 (c). Dotted line outlines the glomerulus. Scale bar:100um Specific gene expression displayed as a tSNE (b, d) for Dlk1 (b) and Akr1b7 (d).

Figure 6: Foxd1 expressing cells may contain multiple, lineage-restricted progenitor cells. Differentiation roots identified by simulation of a Markov process with velocity-based transition probabilities in the reverse direction (a). Diffusion map of interstitial cells demonstrating dispersion of identified cell types along multiple distinct trajectories (b). For 3-dimensional representation of the diffusion map see supplemental video 3. The root cell is indicated by a diamond.

749

750 **Figure 7: Distinct clusters show unique transcriptional activity.** Heat map showing
751 each cell organized by interstitial cluster (abscissa) and binarized regulon activity
752 (ordinate) where black indicates an active regulon, while white indicates an inactive
753 regulon.

754

755 **Figure 8: Perturbed patterning of the interstitial zones disrupts nephron**
756 **differentiation.** The Lef1 regulon is most active in clusters 9-12 (a). mRNA *in situ*
757 hybridization (b, e, f, h, k, l) or immunofluorescence staining (c, d, g, i, j, m) of E18.5
758 wildtype (b-g) and Foxd1Cre;Beta-Catenin^{o/-} mutant (h-m) kidneys with probes to
759 regional interstitial markers Foxd1(b, h), LOX (e, k) and TGFB1i1 (f, l) or antibodies to
760 interstitial marker Meis1/2/3 (green in c, i) and Slug (green in d and j), nephron
761 progenitor marker Six2 (red in c, d, i, j), collecting duct marker CK (white in c, d, i, j) or
762 loop of Henle marker THP (green in g, m), and proximal tubule marker LTL (red in g, m).
763

764 **Figure 9: Human fetal renal interstitial heterogeneity revealed through single cell**
765 **RNA sequencing:** UMAP of 17 week fetal human renal interstitium. For a 3-dimensional
766 representation of the UMAP see supplemental video 4 (a). UMAP of cluster
767 assignments from E18.5 mouse interstitial cells mapped onto the 17-week human
768 interstitial cells (b).

769

770 **Supplemental Table 1:** List of top 50 differentially expressed genes (DEGs) in each
771 cluster when compared to all clusters. Our annotations for each cluster as described in

the text are as follows: Cortical Interstitium: Clusters 1-3; Nephrogenic Interstitium: Clusters 4-5; Proximal Tubule Interstitium: Clusters 6-8; Interstitium Medullary to Proximal Tubule (Outer Medulla): Cluster 9; Outer strip of inner medulla Interstitium: Cluster 10; Papillary Interstitium: Clusters 10-12; Ureteric Interstitium: Cluster 13; Vascular Smooth Muscle: Cluster 14; Pericyte: Cluster 15; Mesangium: Cluster 16; Indeterminate Signature: Cluster 17.

Supplemental Table 2: List of stromal gene against which mRNA in situ hybridizations were conducted, the image of the in situ, along with the rank of the gene in the DEG list for each cluster. Yellow indicates the gene was ranked in the top 100 DEGs, while grey indicates the gene was not present in the DEG list for that specific cluster.

Supplemental Figure 1: tSNE of renal interstitium. For a 3-dimensional representation of the tSNE see supplemental video 2 (a). Specific gene expression displayed as a tSNE (b-j) for Akr1b7 (b), Dlk1 (c), Foxd1 (d), Ntn1 (e), Fibin (f), Smoc2 (g), Lox (h), Penk (i) and Wnt4 (j)

Supplemental Figure 2: Normalized enrichment scores (NES) for mesangial, pericyte and vascular smooth muscle cell types specifically for clusters 14, 15 and 16.

Supplemental Figure 3: UMAP with cells colored by abundance of Lef1 (a) or Smoc2 (c) unspliced transcripts. UMAP with cells colored by abundance of Lef1 (b) or Smoc2 (d) spliced transcripts. RNA velocity field visualized using Gaussian smoothing on a regular grid (e). Diffusion map of stoma cells colored by diffusion pseudotime (f).

796 **Supplemental Video 1:** 3-dimensional view of mouse renal interstitial UMAP

797

798 **Supplemental Video 2:** 3-dimensional view of mouse renal interstitial tSNE

799

800 **Supplemental Video 3:** 3-dimensional view of mouse renal interstitial diffusion map

801

802 **Supplemental Video 4:** 3-dimernasional view of fetal human renal interstitial UMAP.

803

Graphical Abstract

Graphical Abstract: The developing interstitium is a highly heterogeneous, patterned population of cells that occupies distinct positions correlated to the adjacent parenchyma.

Figure 1

Figure 1: Unexpected heterogeneity within the renal

interstitium revealed through antibody staining and

single cell RNA sequencing. Wildtype E18.5 kidneys

stained for Six2 and CK (red, blue respectively in a-m),

and Foxd1 (green in a-b), Tenasin C (green in c-d), Slug

(green in e-f), Acta2 (green in g-h), CDKN1c (green in i-

j), Lef1 (green in k-l), Tbx18 (green in m). Scale bar

100um.

Figure 2

Figure 2: Interstitial heterogeneity revealed through single cell RNA sequencing: UMAP of E18.5 mouse intersitium. For a 3-dimensional representation of the UMAP see supplemental video 1 (a).

Figure 3

Figure 3: Single cell RNA sequencing reveals unexpected cortical heterogeneity within the renal interstitium that is validated through mRNA in situ hybridization. Specific gene expression displayed as a UMAP (a, c, e, g, i) for Foxd1(a), Netrin (c), Fibin (e), Smoc2 (g) and Dlk1 (i). E18.5 *in situ* hybridization of Foxd1, expressed in clusters 1-3 (b), Netrin1, expressed in clusters 1, 3 (d), Fibin expressed in clusters 2-3 (f), Smoc2, expressed in clusters 1 (h) and Dlk1, expressed in cluster 1 (j) reveals cortical heterogeneity. Scale bar: 100um

Figure 4

Figure 4: Spatial characterization of scRNA seq-clusters reveals cortico-medullary patterning

among interstitial subtypes. Specific gene expression displayed as a UMAP (a, c, e, g, i, k) for Foxd1 (a), Lox (c), Clca3a1 (e), Smoc2 (g), Penk (i), and Wnt4 (k). E18.5 *in situ* hybridization of Foxd1: clusters 1-3 interstitial expression indicated with arrow, epithelial lineage expression in podocytes indicated with asterisk) (b-b'), Lox: cluster 7 (d-d'), Clca3a1: clusters 6-8 (f-f'), Smoc2: cluster 9 (h-h'), Penk: cluster 9-12 (j-j'), Wnt4: cluster 11-12 (interstitial expression indicated with arrow, epithelial lineage expression in PTA indicated with asterisk) (l-l'), reveals a molecular pattern which spans the cortico-medullary axis. b, d, f, h, j, l Scale bar: 500um. b', d', f', h', j', l' Scale bar: 100um.

Figure 5

Figure 5: Mesangium and pericyte populations detected using single cell RNA sequencing.

mRNA *in situ* hybridization of Dlk1 expression in clusters 15-16 (a) and Akrlb7 expression in cluster 14 (c). Dotted line outlines the glomerulus. Scale bar:100um Specific gene expression displayed as a tSNE (b, d) for Dlk1 (b) and Akrlb7 (d).

Figure 6

Figure 6: Foxd1 expressing cells may contain multiple, lineage-restricted progenitor cells. Differentiation roots identified by simulation of a Markov process with velocity-based transition probabilities in the reverse direction (a). Diffusion map of interstitial cells demonstrating dispersion of identified cell types along multiple distinct trajectories (b). For 3-dimensional representation of the diffusion map see supplemental video 3. The root cell is indicated by a diamond.

Figure 7

Figure 7: Distinct clusters show unique transcriptional activity. Heat map showing each cell organized by interstitial cluster (abscissa) and binarized regulon activity (ordinate) where black indicates an active regulon, while white indicates an inactive regulon.

Figure 8

Figure 8: Perturbed patterning of the interstitial zones disrupts nephron differentiation. The

Lef1 regulon is most active in clusters 9-12 (a). mRNA *in situ* hybridization (b, e, f, h, k, l) or immunofluorescence staining (c, d, g, i, j, m) of E18.5 wildtype (b-g) and Foxd1Cre;Beta-Catenin^{cl-} mutant (h-m) kidneys with probes to regional interstitial markers Foxd1(b, h), LOX (e, k) and TGFB1i1 (f, l) or antibodies to interstitial marker Meis1/2/3 (green in c, i) and Slug (green in d and j), nephron progenitor marker Six2 (red in c, d, i, j), collecting duct marker CK (white in c, d, i, j) or loop of Henle marker THP (green in g, m), and proximal tubule marker LTL (red in g, m).

Figure 9

Figure 9: Human fetal renal interstitial heterogeneity revealed through single cell RNA

sequencing: UMAP of 17 week fetal human renal intersitium. For a 3-dimensional representation of the UMAP see supplemental video 4 (a)._UMAP of cluster assignments from E18.5 mouse interstitial cells mapped onto the 17-week human interstitial cells (b).

Supplemental Figure 1

Supplemental Figure 1: tSNE of renal interstitium. For a 3-dimensional representation of the tSNE see supplemental video 2 (a). Specific gene expression displayed as a tSNE (b-j) for Akr1b7 (b), Dlk1 (c), Foxd1 (d), Ntn1 (e), Fibin (f), Smoc2 (g), Lox (h), Penk (i) and Wnt4 (j)

Supplemental Figure 2

Supplemental Figure 2: Normalized enrichment scores (NES) for mesangial, pericyte and vascular smooth muscle cell types specifically for clusters 14, 15 and 16.

Supplemental Figure 3

Supplemental Figure 3: UMAP with cells colored by abundance of Lef1 (a) or Smoc2 (c) unspliced transcripts. UMAP with cells colored by abundance of Lef1 (b) or Smoc2 (d) spliced transcripts. RNA

880 velocity field visualized using Gaussian smoothing on a regular grid (e). Diffusion map of stoma cells
881 colored by diffusion pseudotime (f).

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5	Cluster 6	Cluster 7	Cluster 8	Cluster 9	Cluster 10	Cluster 11	Cluster 12	Cluster 13	Cluster 14	Cluster 15	Cluster 16	Cluster 17
1	Igfbp6	Mgp	Igfbp5	Igfbp5	Hmgb2	Sept7	G0s2	Dcn	Acta2	Tcf21	Cdkn1c	Penk	Upk3b	Rgs5	Dcn	Dlk1	mt-Atp6
2	Igfbp5	Dkk1	Aldh1a2	Aldh1a2	Tyms	Dcn	Ace2	Hsp90b1	Tagln	Penk	Pfn2	Cdkn1c	Sparc	Ndufa4l2	Dlk1	Meg3	mt-Cytb
3	Dlk1	Aldh1a2	Mest	Mest	Lig1	Sparc	Igfbp3	Tcf21	Hsp90b1	Fbln1	Wnt4	Tmsb4x	Crip1	Mef2c	Meg3	Col1a1	mt-Nd1
4	Meg3	Ptn	Mgp	Ntn1	Dut	Rgs5	Ptn	Hmcn1	Tpm1	Tuba1a	Nts	Neto2	Krt19	Gm13889	Col1a1	Itm2a	mt-Co2
5	Alcam	Shisa3	Fibin	Aprt	Rrm2	Mfap4	Col14a1	Plac8	Penk	Col14a1	Penk	Dcll1	Dlk1	Tm4sf1	Sparc	Col3a1	mt-Co3
6	Rspo3	Mest	Mecom	Cadm1	Smc2	Ndufa4l2	S100a6	Fbln5	Pfn2	Zeb2	Tmsb4x	Arl4a	Rarres2	Crip1	Postn	Dcn	mt-Nd4
7	Ntn1	Igfbp5	Cadm1	Fibin	Ranbp1	Col1a1	Plac8	Col14a1	Myl9	Snai2	Ncam1	Sdc2	Dmkn	Kcnj8	Col1a2	Mfap5	Igfbp5
8	Gas1	Hmgn2	Gpc6	Tgfb1	Dek	Ace2	Clca3a1	Col6a2	Cldn11	H2afz	Nkd1	Hoxa10	Fabp5	Mgp	Col3a1	H19	mt-Nd2
9	Mgp	Nme1	Ntn1	Mecom	Nasp	Cst3	Fbln5	Meg3	Smoc2	S100a10	Acta2	Cd24a	Igfbp6	Sfrp2	Lum	Col1a2	mt-Nd3
10	Gsn	Ran	Aprt	Gdnf	Cks1b	Col3a1	Sept7	Acta2	Dcn	Plac8	Sdc2	Nkd1	Dcn	Ptp4a3	Sfrp2	Tppp3	Ebf1
11	Cdkn1c	Stmn1	Adgrl2	Mgp	Top2a	Crip1	H19	Tshz2	Igfbp2	Tmsb4x	Epha4	Lrriq1	Eln	Cox4i2	Cst3	Cdkn1c	mt-Co1
12	Aprt	Fhl2	Scx	Isyna1	Gmn	Ccl11	Sparc	Cfh	Sfrp1	Tagln2	Gas6	Ncam1	Prss23	Akr1b7	Cthrc1	Igf2	Malat1
13	Ogn	Eif5a	Ptn	Slc40a1	Mcm7	Angpt2	Rasgrp2	Sparc	Cd24a	Ace2	Atp2b1	Nts	Krt14	Col4a1	Gas1	Sparc	Gm42418
14	H19	Slc25a5	Gdnf	Ccnd2	Dctpp1	Sfrp2	Peg3	G0s2	Ncam1	G0s2	Vcan	Vcan	Col1a1	Igfbp7	Mfap4	Cthrc1	Dkk1
15	Serping1	Mecom	Crabp2	Shisa3	Mgp	Col1a2	Tagln2	Igfbp3	Tmsb4x	Bok	Arl4a	Tcf21	Krt7	Heyl	Csrp2	Clec3b	AY036118
16	Aldh1a2	Hmgb1	Alcam	Pkm	Pcna	Bgn	Ndufa4l2	Col3a1	Tpm2	H2afy2	Cd24a	Trps1	Slc9a3r1	Fabp5	Igf1	Igfbp6	Hbb-bs
17	Col1a1	Id2	Rspo3	Scx	Ran	Igfbp3	Meg3	Col1a1	Col1a1	Lbh	Plekha5	Pla2g4a	Msln	Cald1	Cdkn1c	Crip1	Fbn2
18	Cst3	Id1	Igfbp6	Gpc6	Slbp	Ndrp2	Vcam1	Lox	Cnn2	Stmn1	Sema5a	Ccnd2	Sfrp2	Gata3	H19	Fbn1	Son
19	Fgfbp1	Lsm4	Shisa3	Sub1	Mcm5	Pmepa1	Tcf21	Olfrl3	Col1a2	Cdo1	Nfat5	Serpine2	Rspo1	Itga1	Gsn	Postn	Nrp1
20	Smoc2	Fibin	Ptgr1	Mfap4	Hells	Tm4sf1	Serpib6b	Pmp22	Gas6	Ptn	Tagln	Wnt4	Lrrn4	Pdgfrb	Aspn	Plagl1	Emilin1
21	Mest	Ecm1	Col4a4	Tpi1	Tk1	Fhl1	Igfbp7	Col6a1	Col3a1	Basp1	Hoxa10	Cck	Ezr	Csrp2	Agtr2	Ly6a	Meis2
22	Ackr3	Ptma	Foxd1	Adgrl2	Dnajc9	Igfbp7	Nid1	Ptch1	Cxcl14	Nkain4	Myl9	Ntm	Meg3	Sept4	Serpinf1	Col14a1	Mecom
23	Sparc	Gdnf	Ccnd2	Hacd2	Gins2	Eln	Gpc3	Ace2	Meg3	Csrp1	Tpm1	Lef1	Sulf1	Dkk2	Apoe	Prss23	Fbxl7
24	Ptgis	Cacybp	Zbtb20	Col4a4	Tipin	Col14a1	Anxa2	Tpm1	Ptch1	Ppp1r14a	Lef1	Col15a1	S100a6	Rasl11a	Nfib	S100a10	Adgrl2
25	Mfap4	Hnrnpa1	Tgfb1	Acp6	Slc25a5	Heyl	Serping1	Lbh	Jun	Pja1	Grem2	Dach1	Col3a1	Ebf1	Col14a1	Krt19	Ccnd2
26	Crabp2	Foxd1	Polr2m	Polr2m	Rpa2	Col4a1	Sept11	Bok	Atp2b1	Marcks	Dcll1	Pfn2	Hspb1	Gng11	Ogn	Nrk	Ptn
27	Cpe	Snrpf	Reln	Fhl2	Cdk1	Gm13889	Fhl2	Thbs2	Dstn	Rbp1	Palll	Sox4	Gm12840	Sparcl1	S100a6	Fstl1	Lama4
28	Fxyd6	Polr2m	Pbx1	Bex2	H2afz	Tsc22d1	Myl9	Myl9	Timp3	Itga8	Col15a1	Plekha5	Gas1	Col4a2	Col5a1	Col5a1	Ogt

29	Foxd1	Txn1	Meis1	Cenpf	Nme1	Fbln5	Emp2	Peg3	Sostdc1	Hmgn2	Epb41l2	Epha4	Efemp1	Nrp1	Col6a1	Fst	mt-Nd5
30	Ctsl	Tm4sf1	Kcnq1ot1	Ptma	Igfbp5	Ifitm1	Ifitm1	Cldn11	Sparc	Pla2g4a	Mylk	Ctbp2	Bcam	Reln	Timp2	Pi16	Cltc
31	Il33	Prdx1	Isyna1	Rpl12	Fam111a	Ltbp1	Pmp22	Pcolce	Vim	Tuba1b	Ranbp3l	Epb41l2	Osr1	Mustn1	Cxcl12	S100a6	Reln
32	Il17re	Cox5a	Meis2	Zcchc7	Dhfr	Sept7	S100a11	Dpep1	Rarres2	Plcb1	Kif26b	Kif26b	Col1a2	Abcc9	Pcolce	Serpinf1	Nisch
33	Plagl1	Rbm3	H19	Npm1	Mcm3	Foxs1	Mfge8	Tagln	Olfml3	Dlc1	Ltbp1	Sema5a	Cd2ap	Ren1	Osr1	Nbl1	Pbx1
34	Paqr6	Ifitm3	Myh10	Eef1b2	Lsm2	Itga1	Agtr1a	Col1a2	Ckb	Zfp503	Id3	Plcb1	Plp2	Notch3	Akap12	Ahnak	Tcf4
35	Ddah1	Cenpa	Actr3b	Ssbp4	Stmn1	Ctsl	Acta2	Igfbp7	Lgals1	Sept7	Trps1	Hmga2	Lgals3	Ndrg2	Nrk	Timp2	mt-Nd4l
36	C1qtnf2	Arpc1b	Sub1	Hibadh	Dtymk	Colec11	Cnn2	Sfrp1	Mylk	Olfml3	Id4	H2afz	Cpe	Gadd45b	Cpe	Col5a2	Ptprd
37	Nbl1	H2afv	Egr1	Gja1	Dtl	Lmna	S100a10	Zeb2	Hoxa10	Tubb5	Snhg18	Dach2	S100a10	Iqgap1	Fmo1	Fbn2	Mef2c
38	Cldn1	Hnrnpa2b1	Cldn1	Ak2	Aldh1a2	Col4a2	Lox	Cdc42ep3	Id4	Sept11	Ccnd1	Id4	Nbl1	Fos	Matn2	Gsn	Hspa5
39	Rbp1	Fbn2	Pkm	Gapdh	Mcm6	Pdgfrb	Col6a2	Basp1	Bst2	Hmgb2	Pcdh10	Malat1	Nkain4	Blmh	Crip1	Sema3d	Cadm1
40	Tmem176b	Ccnd2	Fbn2	P3h4	Snrpd1	Meg3	Ppp1r14a	Vim	Fn1	Rem1	Igfbp2	Ccnd1	Aebp1	Nrarp	Fst	Akap12	F2r
41	Ybx3	Anp32b	Adamts5	Zbtb20	Anp32b	Nrk	Tshz2	Ccdc80	Cfh	Csrp2	Bdnf	Snhg18	Ptgis	Zfhx3	Tmem158	Htra3	Sptbn1
42	Fosb	Set	Hnrnpa1	Adamts5	Fibin	Cald1	Zeb2	Cxcl12	Tuba1a	Des	Fgfr2	Ogfr1	Alad	Btg1	Tsc22d1	Col6a3	Atp1a1
43	Omd	Hspd1	Rpl12	Ndufa4	Spc24	App	Dpep1	Cpe	Bok	Fblim1	Cpq	Basp1	Emp2	Hmgcs2	Col6a2	Cd34	Hba-a1
44	C1qtnf3	Odc1	Tnnt1	Cfap36	Ung	Nrp1	Tmsb10	Mbnl2	Nexn	6030408B16Rik	Cnn2	Tanc1	Timp2	Mfge8	Aes	Klf2	Aplp2
45	Upk3bl	Pdgfrb	Slc40a1	Tma7	Cdca7	Mmp14	Col6a1	Epha3	Col23a1	Serpine2	Fzd10	Fbln1	Upk1b	Map3k7cl	Ein	Kcnq1ot1	App
46	Col12a1	Serbp1	Ptgis	Ptgr1	Atad2	Aspn	Maf	Pkdcc	Tcf21	Frem1	Csrp1	Lhfp12	Spint2	Tuba1c	Gnas	Csrp2	Trip12
47	Gsta4	Asb4	Nme1	Loxl1	Smc4	S100a10	Tuba1a	Ogn	Col6a2	Abrac1	Dach1	Tmtc4	Fbln2	Angpt2	Lmna	Nid1	Nfib
48	Tmem176a	Loxl1	Ica1	Ybx3	Hat1	Loxl1	Itga1	Ctsl	Zeb2	Mmp2	Corin	Enpep	Cd151	Ier2	Col5a2	Col6a1	Ctnnb1
49	Hspb1	Hmgb2	Hic1	Sertad4	Mrpl18	Cygb	Cald1	Cnn2	Negr1	Hmgb3	Sct	Jun	Cxadr	Ltbp1	Jund	Itih5	Kcnq1ot1
50	Folr1	Rpl12	Eef1b2	Cxcl12	H2afx	Postn	Fabp5	Ccl21a_ENSMUSG00000094686	Wnt4	Lgals1	Cck	Nfat5	Anxa2	Serpini1	Prrx1	Cd248	Gpc6

Supplemental Table 1: List of top 50 differentially expressed genes (DEGs) in each cluster when compared to all clusters. Our

annotations for each cluster as described in the text are as follows: Cortical Interstitium: Clusters 1-3; Nephrogenic Interstitium: Clusters

4-5; Proximal Tubule Interstitium: Clusters 6-8; Interstitium Medullary to Proximal Tubule (Outer Medulla): Cluster 9; Outer strip of inner

885 medulla Interstitium: Cluster 10; Papillary Interstitium: Clusters 10-12; Ureteric Interstitium: Cluster 13; Vascular Smooth Muscle:
886 Cluster 14; Pericyte: Cluster 15; Mesangium: Cluster 16; Indeterminate Signature: Cluster 17.

887

888 **Supplemental Table 2:** separate file

889

890 **Supplemental Table 2:** List of stromal gene against which mRNA in situ hybridizations were conducted, the image of the in situ, along
891 with the rank of the gene in the DEG list for each cluster. Yellow indicates the gene was ranked in the top 100 DEGs, while grey
892 indicates the gene was not present in the DEG list for that specific cluster.

893

894

895