Defining Strawberry Uniformity using 3D Imaging and Genetic Mapping

Bo Li^{1,2#*}, Helen M. Cockerton^{1*}, Abigail W. Johnson¹, Amanda Karlström¹, Eleftheria Stavridou¹, Greg Deakin¹, Richard J. Harrison^{1,3}

¹NIAB EMR, East Malling, Kent, ME19 6BJ, United Kingdom

²University of the West of England, Bristol, BS16 1QY, United Kingdom

³NIAB, Huntingdon Road, Cambridge, CB3 0LE, United Kingdom

Bo2.li@uwe.ac.uk

Helen.cockerton@emr.ac.uk

Abi.johnson@emr.ac.uk

Amanda.karlstrom@emr.ac.uk

Eleftheria.stavridou@emr.ac.uk

Greg.deakin@emr.ac.uk

Richard.harrison@niab.com

Corresponding author: bo2.li@uwe.ac.uk. Tel: +44(0)1173281117

* Authors contributed equally.

Abstract

Strawberry uniformity is a complex trait, influenced by multiple genetic and environmental components. To complicate matters further, the phenotypic assessment of strawberry uniformity is confounded by the difficulty of quantifying geometric parameters 'by eye' and variation between assessors. An in-depth genetic analysis of strawberry uniformity has not been undertaken to date, due to the lack of accurate and objective data. Nonetheless, uniformity remains one of the most important fruit quality selection criteria for the development of a new variety. In this study, a 3D-imaging approach was developed to characterise berry uniformity. We show that circularity of the maximum circumference had the closest predictive relationship with the manual uniformity score. Combining five or six automated metrics provided the best predictive model, indicating that human assessment of uniformity is highly complex. Furthermore, visual assessment of strawberry fruit quality in a multi-parental QTL mapping population has allowed the identification of genetic components controlling uniformity. A "regular shape" QTL was identified and found to be associated with three uniformity metrics. The QTL was present across a wide array of germplasm, indicating a strong candidate for marker-assisted breeding. A greater understanding of berry uniformity has been achieved through the study of the relative impact of automated metrics on human perceived uniformity. Furthermore, the comprehensive definition of strawberry uniformity using 3D imaging tools has allowed precision phenotyping, which has improved the accuracy of trait quantification. This tool has allowed us to illustrate the use of advanced image analysis towards the breeding of greater uniformity in strawberry.

Key Words

3D imaging, uniformity, achene, carpel-viability, breeding, fruit development

Introduction

Strawberries (*Fragaria x ananassa*) are not true fruits. The red fleshy pseudocarp of a strawberry is formed from a swollen flower base or receptacle. The true fruits are, in fact, the achenes which develop from a whorl of carpels and together form an aggregate-accessory fruit. The viability of both carpels and pollen play an important role in the resulting uniformity of berries¹. Carpel position, density and viability dictate the shape, size and uniformity of a strawberry. Indeed, strawberry breeders have selected for high carpel densities in order to produce larger fruits¹. Simple, classical studies

which remove all or part of the achenes from undeveloped pseudocarps has led to a cessation in the auxin “swelling signal” in the area beneath each achene and thus uneven fruit development². In a similar fashion to achene removal, uneven pollination of the carpels, or absence of achene development, are the main causes of uneven pseudocarps³. Uneven successful pollination can be caused by damage to flowers through high temperature, frost or precipitation¹. A late frost in spring could lead to carpel and other damage, resulting not only in malformation but also complete lack of strawberry development¹. Strawberry flowers have a variable proportion of viable carpels and anthers between flower orders, both within a plant and also between different cultivars^{4,5}. Indeed, primary fruit are more likely to be malformed due to the relatively lower quantities of viable anthers and pollen^{6,7}.

In spite of the environmental factors known to influence uniformity, literature has shown that strawberry uniformity still has a large genetic component and can be improved through breeding^{8,9}. Indeed, where breeders have selected for increased uniformity within and among berries, improvements in uniformity were observed over time⁸. Cultivars have been shown to differ in their susceptibility to misshapen fruit, indicating a significant genetic component controlling uniformity^{1,8,10}. For example, ‘Florida Elyana’ is susceptible to rain damage, disrupting carpel development and thus misshapen fruit leading to lower market value⁹, similarly ‘Camerosa’ has been noted as a cultivar which is particularly susceptible to misshapen fruit with ~4% of yields lost as a result of misshapes^{10,11}. By contrast, ‘Florida Radiance’ has high marketable yields and does not exhibit a high proportion of misshapen fruits⁹. Breeders can influence the proportion of uniform strawberries through selecting- be it directly or indirectly- for 1) even allocation of viable carpels across the receptacle within the flower 2) ready access to pollen within flowers and 3) high fertility of carpels ensuring even successful pollination.

Strawberry is an important fruit crop with a global market revenue of 21,171 million USD in 2015¹². Producing visually appealing strawberry fruit is one of the primary objectives in a strawberry breeding program¹³. Shape uniformity is an essential trait of strawberry fruits due to the direct association with product quality and value¹⁴. Increasing the uniformity of berries can increase the proportion of marketable fruit as berry irregularity is one of the primary imperfections leading to culling and reduced marketable yield⁸.

As there is no well-defined strawberry phenotyping guidelines for fruit uniformity, the current system at NIAB EMR relies on visual assessments, which are subjective and laborious. Unlike morphological traits such as length, volume and colour, which can be accurately measured manually in a low-throughput manner, uniformity assessment is

extremely subjective. As there is no quantitative method of generating phenotypic data for uniformity, the genetic determinants of strawberry uniformity are still unknown.

Computer vision has shown great potential to quantify external fruit quality and 2D imaging has been successfully implemented to measure the shape and size of fruits such as strawberries¹⁵, apples¹⁶, watermelon¹⁷, cherries¹⁸ and mangos¹⁹. Basic shape traits such as length, width, aspect ratio and volume, and more sophisticated traits such as elliptic Fourier descriptors²⁰ have been quantified and used to describe variation in fruit quality. 3D imaging has been successfully used for phenotyping the crop canopy^{21,22} and root architecture^{23,24}, and a 3D strawberry phenotyping platform has been explored in our previous study²⁵. With the 3D point cloud reconstructed based on the Structure from Motion (SfM) method²⁶, basic size-related parameters have been measured in three dimensions allowing volume estimation with high accuracy²⁷. Compared with shape and size evaluation, uniformity is a multi-dimensional trait, therefore it is not possible to quantify through 2D image analysis with a single viewing angle. The application of 3D image analysis for phenotyping the external qualities of fruit has not been sufficiently explored, and the basic, previously characterised, shape- and size-related parameters are not adequate for understanding uniformity.

Here the application of a 3D phenotyping platform allows us to investigate the genetic basis of strawberry uniformity. The 3D image analysis software leverages the previously developed platform²⁵ in order to define eight new external variables and investigate their importance on manual uniformity assessment. This method was applied to a multi-parental strawberry mapping population in order to quantify the genetic components underpinning strawberry uniformity.

Materials and methods

Plant material and experimental set-up

A multi-parental strawberry population was generated through crossing 26 diverse cultivars and breeding lines to create a population of 416 genotypes made up of 26 families each containing 16 individuals (Suppl. Figure 1). Progenitors were selected to represent diversity across multiple fruit quality traits. Twelve replicate runner plants were pinned down from each genotype into 9 cm square pots containing compost. Clonal plants were separated from parental plants and then placed in cold storage (-2 °C) until the start of the experiment. Plants were potted into 2 L pots containing coir and fertigated at 1kg L⁻¹ (rate: 10 seconds every 45 min) using Vitex Vitafeed (N:P:K, 176:36:255). Blocks were horizontal intersections across the polytunnels. Due to the large scale of the experiment, replicate blocks were set up at three week intervals. A Natupol Koppert bumble bee hive was added into each polytunnel to assist even

pollination. Strawberries were picked when ripe into egg boxes. Boxes were labeled with QR codes to assist tracking of genotypes. Strawberry uniformity was scored on a scale from 1 (irregular) to 9 (uniform) with extensive training provided for all assessors. Strawberry shape was allocated into 9 categories: globose, globose-conic, conic, long-conic, bi-conic, conic-wedge, wedge, square and miscellaneous. Manual uniformity scores were recorded in the field book app²⁸, the QR scanning feature allowed quick access to the correct entry form.

Genotyping and Linkage map

DNA was extracted for each genotype from unopened leaflets using the Qiagen DNeasy plant mini extraction kit. Genotyping was conducted using the Axiom[®] IStraw35 384HT array²⁹ (i35k). Crosslink was used to generate linkage maps- a program developed specifically for polyploid plant species³⁰. The map orders from 5 populations were combined to make the consensus map as detailed in the study of Harrison et al³¹. *Fragaria* × *ananassa* chromosome number is denoted by 1-7 and the sub-genome number is represented by A-D as specified in³².

3D reconstruction

The 3D imaging platform was a modified version of that developed by He et al.²⁵. Strawberry fruit were placed in the middle of a turntable, on a dark blue holder made by polymeric foam (38 mm x 19 mm x 19 mm; height, length and width). Unlike the previous study, a webcam (Logitech C920, Newark, CA, USA) was fitted at a height of 30 cm and horizontal distance of 25 cm away from the sample. QR codes on containers were scanned through the webcams allowing tracking of berries and automated labeling of image files. The imaging rig was placed inside a photography studio tent with constant LED illumination. The turntable rotated at a frequency of 50 seconds per full turn, and an image was captured every second. Six imaging platforms allowed concurrent imaging of replicate berries. The 3D reconstruction was implemented with Agisoft Photoscan (Agisoft, LLC, St. Petersburg, Russia), and in order to increase the accuracy and processing speed, all images were pre-processed by cropping to a smaller size (400 x 600 pixels). Background subtraction was achieved through arbitrary colour thresholding. The image processing software for webcam control and automated image pre-processing were written in C++ utilising the OpenCV Library^{25,33}.

Data processing pipeline of phenotypic traits extraction

Point cloud preprocessing

In the preprocessing stage (Fig. 1), each point cloud model was converted from the colour space of RGB (Red, Green and Blue) to HSV (Hue, Saturation and Value). Arbitrary thresholding on the hue channel was used to remove the noise introduced in the reconstruction stage. The clean point cloud was translated to the origin of the 3D

coordinate system based on the distance between the moment of the point cloud and the origin. By calculating the eigenvector associated with the largest eigenvalue of the coordinates of points, a rotation matrix could be derived to represent the main orientation of the point cloud, which can be used to rotate the point cloud with the main orientation aligned with the z-axis. After rotation, the arbitrary threshold was applied again on the hue channel in order to segment the strawberry body and blue holder from the whole point cloud. The height of the holder was obtained by calculating the difference between the maximum and minimum values of the holder point cloud on the z-axis. As the original coordinate system generated by Structure from Motion (SfM) method has an arbitrary scale, each point cloud model needed to be standardised by the height of the holder, so that the sizes of all point clouds are comparable.

Figure 1. Point cloud pre-processing for strawberry body extraction, translation to origin of xyz coordinate system and size standardisation.

Uniformity-related traits measurements

Eight uniformity-related traits were calculated from the point cloud data of strawberry body after preprocessing. These are:

Coefficient of variation (CV) of side view areas (CV_A) and the ratio between maximum and minimum side view area (Max_A/Min_A)

All side views should be identical in a perfectly uniform strawberry. In order to eliminate the heterogeneity introduced from the calyx and the holder, only the points within the middle 50% of the body height of each point cloud were retained for analysis (Fig. 2). In order to understand the heterogeneity of different side views of a point cloud, each point cloud was rotated along the z-axis by 3.6° for 99 rotations, and after each rotation, the

side view of the point cloud was projected onto the x-z plane in 2D (labelled in white). A convex hull was fitted to each projected image and the contour area was calculated. For area metrics, two traits were obtained; the CV of side view areas (CV_A) and the ratio between the maximum and minimum area (Max_A/Min_A). An ideal uniform strawberry will have a value of zero for CV_A and one for Max_A/Min_A.

CV of principal orientations (CV_D)

The major eigenvector indicating the main orientation was calculated by principal component analysis (PCA) for all 100 side view projected images, and the heterogeneity of the orientations of the projected images was quantified by calculating the CV of angles of the main orientation. Like CV_A, a perfectly uniform strawberry will have a value of zero for CV_D.

(a)

(b)

(c)

Figure 2. Side view of strawberry body for the CV measurement of the area and principal orientations. Convex hulls are outlined in blue, and blue and green arrows indicate the principal orientations (a). Extraction of example slice images horizontal to x-

y plane at the height of 20%, 40%, 60% and 80% of the total height. A minimum bounding box is fitted to each slice image (b). Sixteen patches of points labelled in different colours for curvature estimation (c).

Aspect ratio of the minimum bounding box (L/W)

A lateral slice image was obtained by identifying the intersection between the plane in parallel with the x-y plane and point cloud (Fig. 2b). Based on the values on the z-axis, 100 evenly spaced slice images were obtained. The slice image with the largest contour was obtained by calculating the contour area of the convex hulls for all slice images. The main orientation of the contour was indicated by the major eigenvector of the PCA and the minimum bounding box was fitted to the slice images. The ratio between the length and width of the largest bounding box was derived and the ratio should be one for a perfectly uniform fruit.

Circularity of the maximum circumference (CIR)

Visually, the circularities of the contours in horizontal slice images are high if the strawberry is uniform. Circularity (CIR) was calculated as previously described³⁴:

$$C = \frac{4\pi A}{p^2}$$

Where A and p are the area and perimeter of the convex hull respectively. For each point cloud, the circularity was calculated for the slice image with largest contour area.

Straightness of centre axis (STR)

The coordinates of the centroids for each horizontal slice image can be located by calculating the moment of the contour. The centroids can be connected as a straight line for a uniform strawberry. The centroids were calculated for all the slice images within the middle 80% of the body height, and the straightness of the central axis was characterised by:

$$STR = \frac{\sum_{i=1}^{N-1} d_i}{D}$$

Where N ($N = 80$) is the number of slice images used for the analysis, d_i is the Euclidean distance between neighbouring slice images, and D is the Euclidean distance between the centroids of the top and bottom slice images.

CV of curvature and the ratio between maximum and minimum curvatures (Max_C/Min_C)

The principal curvature can be calculated for each point in the point cloud, which describes how much the curve deviates from a straight line at this point. It can be imagined that the 3D curve can be sliced orthogonally around the direction of normal in

to 2D curve, and the maximum curvature k_1 and minimum curvatures k_2 are the two principal curvatures for the 3D curve³⁵. The average curvature k , which is defined as the mean value of the magnitudes of principal curvatures in the two main directions was applied to quantify the curvature for a given point.

As the curvature measurement is sensitive to noise, the point cloud surface of strawberry body was first smoothed by using Moving Least Squares (MLS) method³⁶, which could reconstruct a smooth surface from the noisy point cloud. Sixteen patches of the points were selected evenly from the points forming the largest slice in parallel with x-y plane (Fig. 2c). For each patch, the first half largest curvatures were averaged and used to represent the curvature of the patch. With the curvatures of all 16 patches, the CV of curvature (CV_C) and the ratio between maximum and minimum curvatures (Max_C/Min_C) were calculated.

Statistical analysis

Ordinal regression

Statistical analysis was performed using R (version 3.5.1) and the Genstat statistical package (Version 13.0, VSN International Ltd. England). Differences in uniformity traits within each shape groups were distinguished using ANOVA and Tukey post-hoc test. Pearson coefficients of correlation were calculated between all proposed uniformity-related traits. As the group labels are ordinal dependent variables, ordinal regression was used to evaluate the performances of all traits³⁷. Model fit was ascertained by using selection criterion values based on the Akaike Information Criterion (AIC) and the Bayesian Information Criterion (BIC). In general, a better model fit generates lower values for both AIC and BIC³⁸. In order to identify the optimal variable combination related to manual assessment, stepwise AIC and BIC methods were applied³⁹. The most significant variable was identified by comparing the criterion values of all models. Other variables were added successively and retained if the model fit was improved.

Genetic Analysis

The best linear unbiased estimate (BLUE) was calculated for all genotypes in order to correct for the influence of assessor, data and block. Linear mixed-effects models were generated for each phenotypic trait with and without covariates. Grand scores for each genotype were calculated using mixed models to account for significant covariates.

Composite interval mapping

Multiparental QTL mapping was conducted in R using package “mppR”⁴⁰. A permutation test determined the significance threshold⁴¹. A two-step QTL analysis was implemented: the selection of cofactors was achieved through Simple Interval Mapping

(SIM) proceeded by a multi-QTL model search using composite interval mapping (CIM)^{42,43}. As a multi-parent population CIM works on parent relationships. Therefore the 'CPEM0162' x 'Rumba' cross was removed as it is not directly related through the parental cultivar network. All other crosses were interrelated and formed a single network (Sup. Figure 1).

Results

Characterisation of uniformity-related traits

All the uniformity-related traits were calculated based on the point cloud, the mean values and the standard errors for each visual uniformity class are presented in Figure 3. ANOVA results showed that significant differences were observed between uniformity classes for all traits ($p < 0.001$). The Pearson's linear correlation coefficients were calculated between all traits, and strong correlations were found amongst Max_A/Min_A, L/W and CIR (Table 1).

Ordinal regression models were constructed for all variables and each variable independently. L/W was not significant due to the high correlation with other variables and CIR showed the best model fit with the lowest AIC and BIC values (Table 2). New variables were added sequentially to the model until no further improvement of the criterion value was observed. The AIC and BIC based stepwise selection methods showed inconsistent results (Table 3). The AIC based method showed the optimal criterion value with all variables except L/W and Max_C/Min_C, but BIC based method showed that STR could not improve the model fit.

The influence of Shape on uniformity

The shape of a strawberry influences the uniformity trait score. Bi-conic strawberries were seen to have high uniformity based on the area overlap measures (CV_A & Max_A/Min_A), L/W and CIR scores indicating bi-conic strawberries have consistently circular horizontal transects at the mid point. Whereas for curvature uniformity measures (CV_C & Max_C/Min_C) globose fruit are the most uniform and miscellaneous fruit the least (Data not shown). Both the manual uniformity score and CIR could discriminate miscellaneous shapes from the other shape categories (Figure 4).

QTL identification

A total of 28 QTL were found to be associated with uniformity traits (Table 4). Of which 25 were detected in more than one progenitor (Figure 5). Five focal SNP's, on chromosome 2B and 5D were found to represent more than one trait (Table 4, Figure 6). The same focal SNP AX.166521303 was identified as important region in

Max_A/Min_A, CV_A and CIR uniformity traits. Global adjusted R^2 values for linear models were between 5.07 and 32.15 indicating the proportion of variation explained by identified QTLs (Table 5). All uniformity traits apart from CV_A were significantly affected by date of picking. CV_D had the largest broad sense heritability score of 38.4 (Table 5).

Figure 3. Mean value and standard error of calculated uniformity-related traits by the newly developed 3D image analysis software against defined uniformity scale based on manual assessment.

Figure 4. CIR scores for each manually classified strawberry shape category. Letters denote significant differences between categories. Error bars are standard errors of the mean.

Table 1. Pearson's linear correlation coefficients among all uniformity-related traits. All values are significant at $p < 0.05$ level.

	CV_A	Max_A/Min_A	CV_D	L/W	CIR	STR	CV_C	Max_C/Min_C
CV_A	1.00							
Max_A/Min_A	0.61	1.00						
CV_D	0.13	0.29	1.00					
L/W	0.54	0.90	0.27	1.00				
CIR	-0.48	-0.85	-0.32	-0.85	1.00			
STR	0.12	0.26	0.06	0.26	-0.29	1.00		
CV_C	0.21	0.24	0.07	0.23	-0.25	0.15	1.00	

Max_C/Min_C 0.17 0.20 0.15 0.17 -0.27 0.10 0.57 1.00

Table 2. Summary of individual ordinal models and variable significance of ordinal model with all variables, towards prediction of manual assessment uniformity scores. * p < 0.05, ** p < 0.01, *** p < 0.001.

Model	LogLik	AIC	BIC	Signif. codes
CIR	-1691.82	3401.65	3445.35	***
Max_A/Min_A	-1707.44	3432.89	3476.85	***
LW	-1742.98	3503.96	3547.66	
CV_A	-1790.55	3599.09	3642.79	***
CV_C	-1824.20	3666.41	3710.11	***
Max_C/Min_C	-1839.63	3697.25	3740.95	*
CV_D	-1857.94	3733.87	3777.57	***
STR	-1861.85	3741.70	3785.39	*

Table 3. Model comparison values for uniformity metrics, towards prediction of manual assessment uniformity scores based on AIC and BIC.

Model	1	2	3	4	5	6	7	8
CIR	x	x	x	x	x	x	x	x
Max_A/Min_A		x	x	x	x	x	x	x
LW			x					
CV_A				x	x	x	x	x
CV_C					x	x	x	x
Max_C/Min_C						x		
CV_D							x	x
STR								x

AIC	3401.65	3374.40	3374.29	3362.57	3302.01	3299.53	3291.87	3288.40
BIC	3445.35	3422.95	3427.70	3415.97	3360.28	3362.65	3354.99	3356.38

Supplementary Table 4. Focal SNPs representing strawberry uniformity QTL. The position of QTL is reported in Mb as scaled to the vesca version 4 genome

Marker Names	Chromosome	Pos Mb	$\log^{10} p$ value	R^2	Trait
AX.89804099	3B	96.3	3.29	6.9	Visual Uniformity
AX.123357183	2D	63.1	4.07	3.81	Max_C/Min_C
AX.166513757	4B	239.6	3.03	5.05	Max_C/Min_C
AX.166523206	4D	272.0	3.89	4.03	Max_C/Min_C
AX.166514922	5D	1.8	3.02	9.9	Max_C/Min_C
AX.89886024	5D	36.8	3.32	2.87	Max_C/Min_C
AX.166525020	6B	34.4	3.49	7.59	Max_C/Min_C
AX.166526040	7B	47.5	3.07	2.71	Max_C/Min_C
AX.166521293	2B	60.7	3.74	6.41	CV_C
AX.166519032	2C	183.5	3.04	5.07	STR
AX.166521303	2B	63.1	3.29	9.41	CIR
AX.89791395	4B	130.4	3.17	2.71	CIR
AX.166509340	4C	241.7	3.00	2.37	CIR
AX.89890707	5C	212.2	3.11	1.86	CIR
AX.166526395	7C	228.6	3.81	3.95	CIR
AX.89873861	1A	139.7	3.73	5.27	L/W
AX.123357666	3B	303.6	3.35	3.42	L/W
AX.89829301	4D	182.3	3.03	2.06	L/W

AX.166524494	5B	103.6	3.25	3.83	L/W
AX.89791880	5D	123.7	3.52	1.46	L/W
AX.89792701	5D	182.0	3.61	10.64	L/W
AX.166503621	2C	266.2	3.89	2.19	CV_D
AX.166504095	3A	149.5	4.10	2.8	CV_D
AX.166525754	6C	74.7	3.25	3.58	CV_D
AX.166521303	2B	63.1	4.55	15.08	Max_A/Min_A
AX.89791880	5D	123.7	4.22	3.11	Max_A/Min_A
AX.166506309	5D	150.8	3.16	2.48	Max_A/Min_A
AX.166523943	5D	233.9	3.42	4.47	Max_A/Min_A
AX.166521303	2B	63.1	3.86	6.28	CV_A
AX.89788547	5D	40.9	3.61	5.8	CV_A

Table 5. Broad sense heritability scores for each automated uniformity trait, the influence of block and date of assessment on the trait measured. The number of QTL and the coefficient of determination associated with combined QTL.

Trait	Heritability	Significance of Block	Significance of Date	GxE	Number of QTL	R2	R2 adj
STR	21.70	***	***	***	1	5.07	4.63
CIR	22.00	***	***	***	5	21.2	19.3
CV_D	38.40	***	***	***	3	9.35	8.05
CV_A	19.95	NS	NS	***	2	12.13	11.29
CV_C	24.63	***	***	***	1	6.41	5.97
L/W	19.20	***	**	***	6	24.77	22.59
Max_C/Min_C	16.70	NS	*	NS	7	32.15	29.85

Max_A/Min_A	22.70	**	***	***	4	23.1	21.62
Uniformity	8.00	NS	***	***	1	6.9	6.46

	Man Uni	CV_A	Max_A/Min_A					CV_D			LAW					CIR					STR	CV_C	Max_C/Min_C							
	AX.89800099	AX.166511303	AX.89798547	AX.166511303	AX.166506309	AX.89791880	AX.166513943	AX.166508621	AX.166504095	AX.166525754	AX.89872861	AX.123317666	AX.89829301	AX.166524494	AX.89791880	AX.89792701	AX.166511303	AX.89791395	AX.166509340	AX.89890707	AX.166526395	AX.166519032	AX.166511293	AX.123317183	AX.166513757	AX.16651206	AX.166514022	AX.89886024	AX.166515020	AX.166526040
FA2311	1.26						-0.68											-0.52	-0.84	-0.49		1.01					NA		1.04	
FA2807			-0.72				-0.68			-1.56								-0.52	-0.84	-0.49	-0.74							-0.75		-0.80
FA0121							-0.68	NA										-0.52	-0.84	-0.49	-0.74	1.01								1.04
FA0193					0.65		-0.68											-0.52	-0.84	-0.49	-0.74	1.01								
FA0248					0.65		-0.68											-0.52	-0.84	-0.49	-0.74	1.01								
FA0307	1.26		-0.72				-0.68	1.18			1.11	-0.90	1.37					-0.52	-0.84	-0.49	-0.74	1.01				0.70	-0.75		-0.40	
FA0676							-0.68	2.32			2.23							-0.52	-0.84	-0.49	-0.74	1.01				0.70	-0.75		1.04	-0.40
FA0677		NA					-0.68		NA				-0.90					-0.52	-0.84	-0.49	-0.74	1.01								
FA0900				0.65		-1.35												-0.52	-0.84	-0.49	-0.74	1.01		-0.86						
FA1111																		-0.52	-0.84	-0.49	-0.74	1.01				0.70			2.07	-0.40
FA1211	1.26																	-0.52	-0.84	-0.49	-0.74	1.01				0.70			2.07	-0.40
FA1217		-1.03	-0.72	-1.73	0.65		-1.35				1.11					-2.66	1.35	-0.52	-0.84	-0.49	-0.74			-1.16	-0.86					-0.40
FA2654											1.11							-0.52	-0.84	-0.49	-0.74	1.01								
FA6969	1.26	-1.03	-0.72	-1.73			-0.68				1.11	1.80				-2.66	1.35	-0.52	-0.84	-0.49	-0.74	1.01		-1.16						-0.40
FA0508	1.26			-1.44			-0.68											-0.52	-0.84	-0.49	-1.47							-0.75	-0.40	
FA9521				-1.44			-0.68	NA										-1.04	-0.52	-0.74	-1.47				-0.86	0.70	-0.75		-0.40	
FA2807	-1.03			-1.73			-0.68					1.80					1.35	-0.52	-0.99	-1.47			-1.16	-0.86	0.70	-0.75		-0.40		
FA2807	1.26				0.65		-1.35											-0.52	-0.84	-0.49	-0.74	1.01			2.47					-0.40
FA0900	-1.03	-0.72	-1.73		0.65	-1.35				-1.56				NA	0.82			-0.52	-0.84	-0.49	-0.74	1.01		-1.16		0.70				-0.40
FA1711		-0.72			0.65	1.25		-1.35		-1.56								-0.52	-0.84	-0.49	-0.74	1.01		-0.86	2.47		1.72		1.04	
FA0900					0.65		-1.35											NA	-0.84	-0.49	-0.74	1.01								
FA0900		-1.03	-0.72	-1.73	NA		NA			-1.56	NA					NA		-0.52	-0.84	-0.49	-0.74	NA		-1.16		0.70		NA	1.04	NA
FA0752		-1.03	-0.72	-1.73	NA	1.25	-0.68			NA						0.82	NA	1.35	-0.52	-0.84	-0.99	-0.74	NA	-1.16		0.70		NA	1.04	NA
M.Centenary		-1.03	-0.72	-1.73	NA												NA	1.35	-0.52	-0.84	-0.99	-0.74	NA	-1.16		0.70		NA	1.04	NA
FA0907		-0.72						2.32				1.80					NA		-0.52	-0.84	-0.49	NA		-0.86		NA		NA		NA
Vibrant					NA											NA		-0.52	-0.84	-0.49	NA				NA		NA		NA	NA

Figure 5. Effect sizes associated with each QTL in each of the 26 progenitors; blue colour is associated with lower uniformity and red colour is associated with higher uniformity.

Figure 6. Location of QTL on the octoploid consensus map scaled to the *Fragaria vesca* 'version four' genome. Horizontal grey lines represent istraw 35k axiom array markers

Supplementary Figure 1. A network of crosses conducted to generate the multiparental mapping population used in this study. Cultivars are represented by circles, families are represented by lines.

(a)

Supplementary Figure 2. Manhattan plots of (a) Max_A/Min_A, (b) L/W and (c) CIR.

Discussion

We report for the first time a robust method to measure strawberry uniformity and apply this technique to generate genetic markers for uniformity traits. Several studies have attempted to quantify strawberry fruit shape using 2D images with neural networks⁴⁴, 3D imaging⁴⁵ and by machine learning¹⁵. However, none of these studies investigated berry uniformity. Unlike the aforementioned studies, who measure a relatively small number of genotypes intensively, we have implemented a high throughput imaging platform across a large population to facilitate genetic analysis of the trait. Although strawberry shape has received greater attention in the literature, berry uniformity is a more important trait for a breeder to improve (Personal communication, Abigail Johnson).

In current strawberry breeding practice, there is no widely accepted criteria for quantifying uniformity due to the difficulty of defining a multidimensional trait. Here, the manual strawberry uniformity scale has been designed by NIAB EMR breeders. As such, the absence of a straightforward definition, has meant that it has not been possible to study the genetic components controlling strawberry uniformity in the past. To overcome this, we have used 3D image analysis to define the parameters underlying a breeder's perception of strawberry uniformity. The original 3D strawberry phenotyping system²⁵ could accurately measure basic size-related traits. In this study, the point cloud analysis software was further developed to quantify strawberry uniformity through eight proposed metrics. By comparing with the manual scale, the image processing pipeline has demonstrated an objective method of characterising strawberry uniformity components.

Quantifying berry uniformity

Circularity of the maximum circumference (CIR) of strawberries showed the best predictive ability for manual uniformity scores based on the ordinal regression model fit, when studying individual variables alone. A completely misshapen fruit with a severely undulating fruit surface will score a value of 1 for manual assessments, and these completely misshapen fruits were the easiest category to identify by eye, as they were clearly distinct from regular shapes. A low CIR value appears to represent the undulating misshapen and "miscellaneous" fruit (Figure 4 & 6). Miscellaneous berries are the most undesirable fruit shape category therefore it is highly beneficial to select against them. When multiple traits are combined to describe uniformity, the best fitting model required the combination of CIR, CV_A and Max_A/Min_A, CV_D and CV_C. The five factors required for optimal model construction indicate that there are multiple uniformity components influencing the manual uniformity score.

Misshapen fruit QTL

One of the QTL represented by the focal marker AX.166521303 on chromosome 2B was found to be associated with CIR, this QTL was also associated with CV_A and Max_A/Min_A, each of which were found in the best fitting model used to describe the manual uniformity score. The focal SNP AX.166521303 was found to be present and significant in six progenitors and had an effect size of 9.14% on CIR. Therefore, this marker is a good candidate for marker assisted breeding in selection against completely mis-shapen and irregular strawberries. Furthermore, this work has highlighted a region of interest for further study to pinpoint the causative allele associated with reduced uniformity. Dissecting the contribution of genetic and environmental components believed to underpin strawberry uniformity; susceptibility to heat stress, carpel and pollen viability, achene position, size and distribution¹ may help to further elucidate the mechanism of uniformity segregating in the multiparental population.

Uniformity trait selection

The trait L/W shows little improvement on the overall combined trait model fit due to the high correlation with other traits including Max_A/Min_A and CIR, but it was still a good predictor of uniformity based on the model fit when studying individual variables alone. AIC and BIC based stepwise feature selection showed disagreement on the selection of the STR parameter. The difference between calculating AIC and BIC is that AIC does not account for the sample size, so when sample number is large, BIC applies larger penalty for complex models and leads to a simpler model⁴⁶. However, this study does not aim to identify the optimal feature combination to develop prediction model related to manual uniformity evaluation, but develop a new image based quantification to replace the manual scale, because the ground-truth data are subjective and as such any large bias can reduce the robustness of model development. Moreover, the manual scale cannot be considered a comprehensive assessment as the parameter STR cannot be visually evaluated by eye. However, it must be said that if a trait cannot be detected by the human eye, then it is not a valuable trait for a strawberry breeder to select upon.

Limitations of the system

The 3D point cloud analysis software is independent of the imaging acquisition system, and the uniformity-related traits can be extracted automatically in a high-throughput manner. However, the imaging collection throughput was 50 seconds per fruit and the 3D reconstruction has to be performed separately, which limits use to pre breeding experiments as opposed to use as a breeders tool. Due to the occlusion from the viewing angles, the strawberry nose cannot be fully reconstructed especially for globose shaped fruit, which decreases the accuracy of STR measurements and also limits the study on automated shape classification. To increase the throughput and accuracy of 3D phenotyping, it is necessary to further develop the hardware with multiple cameras to allow more viewing angles or a structured light based imaging system with a robotic arm, and also integrate the hardware driver with 3D reconstruction software. The current point cloud image analysis software was able to characterise many key external traits which are important for strawberry breeding, however, the measurement of other parameters such as achene density must be investigated through an improved phenotyping platform in future studies.

Genetic Control of Fruit Quality

Papers detailing strawberry fruit quality QTL report genetic alleles associated with multiple fruit quality traits including fresh weight, metabolites, external colour and firmness^{29,47,48}, however, there are currently no papers which report QTL associated with strawberry uniformity. Here, we provide a phenotyping platform which has

facilitated the assessment of the genetic components underlying strawberry uniformity for the first time. The use of a multi-parental population has allowed the study of a diverse set of germplasm and has ensured that resulting QTL to have a greater relevance for breeders when compared to alleles identified in bi-parental studies. Overall, 25 of the QTL were found to have an effect on uniformity in more than one of the 26 progenitors indicating that there has been limited linkage decay between the causal allele and marker, and that the relationships have been maintained across generations. Furthermore, the QTL on chromosome 2B was observed three times across different uniformity traits, such traits are only partially correlated and thus describe discrete components, as such this allele can be seen to play a major role in uniformity.

Genetic control of strawberry fruit shape

Unlike uniformity the mechanism controlling fruit shape has been studied extensively in the wild strawberry; *Fragaria vesca* and may act as a surrogate model for the cultivated octoploid strawberry *Fragaria x ananassa*. In *F. vesca*, fruit shape is primarily controlled by phytohormones^{49,50}. Auxin increases the width of fruit and by contrast gibberellic acid (GA) increases the length of a strawberry whereas Absciscic acid (ABA) down regulates both Auxin and GA and thus reduces fruit expansion^{49,50}. GA deficient Vesca mutants were found to have a “short” or globose berry shape, which, through the application of GA to the berry, could be restored to result in a “long” or long conic fruit shape⁵⁰.

It is clear that breeders wish to select for greater berry uniformity however the confounding relationship between shape and uniformity must also be considered. For example, square, wedge and wedge-conic strawberries may have high 2D symmetry but not 3D symmetry. UK breeders primarily aim to select for conic or long conic fruit whereas globose, square, wedge and miscellaneous berries are classified as undesirable and biconic, globose-conic and conic-wedge fruit are seen as acceptable shapes (Personal Communication, Abigail Johnson). Here we provide an objective measure (CIR) that can be used to discriminate the least desirable berries - miscellaneous or misshapen berries and select for regular fruit shapes.

Heritability of Uniformity

Broadsense heritability scores were between 16.70 and 38.40 for automated uniformity metrics indicating a greater genetic component than that associated with manual uniformity (8.00). These values indicate the proportion of variation segregating in the study population, however improvement in the heritability may also be caused, in part, by more accurate phenotypic measurements. In particular, high heritability was observed for CV_D which indicates the angle of a strawberry related to whorl of carpels

(Figure 4) is under strong genetic control. Date of picking was seen to have a significant impact on all uniformity metrics apart from CV_D which had a large genetic component. The high significance of date indicates the developmental environmental conditions has a significant impact on strawberry uniformity. Extreme temperatures were observed during the experiment which may have caused the significance of date. All traits apart from CV_C showed a significant genotype by environment interaction indicating that genotypes were responding differently to heat stress. Misshapen fruit have been found to have a greater proportion of small underdeveloped achenes following exposure to extreme temperatures during embryo development^{51,52}.

Here we provide a comprehensive dissection of the traits underlying strawberry uniformity and show that the visual perception of a strawberry can be represented by 5 metrics. The generation of an objective measure of uniformity has allowed the assessments of genetic components in a multi-parental breeding population. We show uniformity has a strong genetic component that can be improved by breeding and identify genetic components controlling uniformity that are present across a wide array of germplasm.

Availability of data, materials and methods

The software developed and datasets generated and analysed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

The authors acknowledge project partners Soloberry, Sainsburys, Botanicoir and Agrovista for their involvement and support of the project. The authors acknowledge Robert Vickerstaff for generating the octoploid consensus map as part of other projects and Dr Beatrice Denoyes, INRA and Dr Amparo Monfort, CRAG for granting the use of their informative markers in the production of the strawberry consensus linkage map. This work was supported by grants from the Biotechnology and Biological Sciences Research Council (BBSRC) BB/M01200X/2, BB/P005039/1 and Innovate UK project 101914.

Authors' contributions

BL – Analysed imaging data
 AJ, HMC, ES, BL, RJH - Conceived and designed experiments
 HMC – Conducted quantitative genetics analysis
 AJ, HMC, AK - Performed experiments
 AK - Performed genotyping

GD - Ordinal regression analysis

BL, HMC & RJH wrote the manuscript with contributions from all authors.

BL and HMC equally contributed to the manuscript.

Conflict of interests

On behalf of all authors, the corresponding author states that there is no conflict of interests regarding the publication of this work.

List of Abbreviations

CIM- composite interval mapping

CIR - Circularity

CV_A - Coefficient of Variation of side view areas

CV_C - Coefficient of Variation of curvatures

CV_D - Coefficient of Variation of principal orientations

i35k - Istraw35 Affymetrix chip

L/W - Aspect ratio of the minimum bounding box

Max_A/Min_A - ratio between maximum and minimum side view areas

Max_C/Min_C - ratio between maximum and minimum curvatures

MedR - Median number of roots

QTL- Quantitative Trait Loci

QR - Quick Response

RGB- Red Green Blue

SNP - Single Nucleotide Polymorphism

STR- Straightness of centre axis

Reference

1 Carew JG, Morretini M, Battey NH. Misshapen fruits in strawberry. *Small Fruits Review*. 2003; **2**: 37–50.

2 Nitsch JP. Growth and morphogenesis of the strawberry as related to Auxin. *American Journal of Botany*. 1950; **37**: 211.

3 Kronenberg HG, Braak JP, Zeilinga AE. Poor fruit setting in strawberries. II. *Euphytica*. 1959; **8**: 245–251.

4 Thompson PA. Environmental effects on pollination and receptacle development in the strawberry. *Journal of Horticultural Science*. 1971; **46**: 1–12.

- 628 5 Darrow GM. The importance of sex in the strawberry. *Journal of Heredity*. 1925; **16**:
629 193–204.
- 630 6 Darrow GM. The strawberry. *The Strawberry* 1966.
- 631 7 Gilbert C, Breen PJ. Low pollen production as a cause of fruit malformation in
632 strawberry. *J Am Soc Hortic Sci* 1987.
- 633 8 Whitaker VM, Hasing T, Chandler CK, Plotto A, Baldwin E. Historical trends in
634 strawberry fruit quality revealed by a Trial of University of Florida Cultivars and
635 Advanced Selections. *HortScience*. 2011; **46**: 553–557.
- 636 9 Chandler CK, Santos BM, Peres NA, Jouquand C, Plotto A. ‘Florida Elyana’
637 strawberry. *HortScience*. 2009; **44**: 1775–1776.
- 638 10 Zanin DS, Fagherazzi AF, dos Santos AM, Martins R, Kretschmar AA, Rufato L.
639 Agronomic performance of cultivars and advanced selections of strawberry in the
640 South Plateau of Santa Catarina State. *Revista Ceres*. 2019; **66**: 159–167.
- 641 11 Ariza MT, Soria C, Medina-Mínguez JJ, Martínez-Ferri E. Incidence of misshapen
642 fruits in strawberry plants grown under tunnels is affected by cultivar, planting date,
643 pollination, and low temperatures. *HortScience*. 2012; **47**: 1569–1573.
- 644 12 Xiong Y, From PJ, Isler V. Design and evaluation of a novel cable-driven gripper
645 with perception capabilities for strawberry picking robots. *2018 IEEE International
646 Conference on Robotics and Automation (ICRA)*. 2018.
647 doi:10.1109/icra.2018.8460705.
- 648 13 Faedi W, Mourgues F, Rosati C. Strawberry breeding and varieties: situation and
649 perspectives. *Acta Horticulturae*. 2002; 51–59.
- 650 14 Nielsen JA, Lovell PH. Value of morphological characters for cultivar identification
651 in strawberry (*Fragaria xananassa*). *New Zealand Journal of Crop and Horticultural
652 Science*. 2000; **28**: 89–96.
- 653 15 Ishikawa T, Hayashi A, Nagamatsu S *et al*. Classification of strawberry fruit shape
654 by machine learning. *ISPRS - International Archives of the Photogrammetry,
655 Remote Sensing and Spatial Information Sciences*. 2018; **XLII-2**: 463–470.
- 656 16 Mir JI, Ahmed N, Singh DB *et al*. Diversity evaluation of fruit quality of apple (*Malus
657 x domestica* Borkh.) germplasm through cluster and principal component analysis.
658 *Indian Journal of Plant Physiology*. 2017; **22**: 221–226.
- 659 17 Akodagali J, Balaji S. Computer Vision and Image Analysis based techniques for
660 automatic characterization of fruits a review. *International Journal of Computer
661 Applications*. 2012; **50**: 6–12.
- 662 18 Beyer M, Hahn R, Peschel S, Harz M, Knoche M. Analysing fruit shape in sweet

- 663 cherry (*Prunus avium* L.). *Scientia Horticulturae*. 2002; **96**: 139–150.
- 664 19 Naik S, Patel B, Pandey R. Shape, size and maturity features extraction with fuzzy
665 classifier for non-destructive mango (*Mangifera Indica* L., cv. Kesar) grading. 2015
666 *IEEE Technological Innovation in ICT for Agriculture and Rural Development*
667 (*TIAR*). 2015. doi:10.1109/tiar.2015.7358522.
- 668 20 Hiraoka Y, Kuramoto N. Identification of *Rhus succedanea* L. cultivars using Elliptic
669 Fourier Descriptors based on fruit shape. *Silvae Genetica*. 2004; **53**: 221–226.
- 670 21 Paproki A, Sirault X, Berry S, Furbank R, Fripp J. A novel mesh processing based
671 technique for 3D plant analysis. *BMC Plant Biol* 2012; **12**: 63.
- 672 22 Coupel-Ledru A, Pallas B, Delalande M *et al.* Multi-scale high-throughput
673 phenotyping of apple architectural and functional traits in orchard reveals genotypic
674 variability under contrasted watering regimes. *Hortic Res* 2019; **6**: 52.
- 675 23 Topp CN, Iyer-Pascuzzi AS, Anderson JT *et al.* 3D phenotyping and quantitative
676 trait locus mapping identify core regions of the rice genome controlling root
677 architecture. *Proc Natl Acad Sci U S A* 2013; **110**: E1695–704.
- 678 24 Cockerton, H.M., Li, B., Stavridou, E., Johnson, A., Karlström, A., Armitage, A.D.,
679 Martinez-Crucis, A., Arjona, L.G. and Harrison, R.J. Genetic and phenotypic
680 associations between root architecture, arbuscular mycorrhizal fungi colonisation
681 and low phosphate tolerance in strawberry (*Fragaria × ananassa*).
682 doi:10.21203/rs.2.18300/v1.
- 683 25 He JQ, Harrison RJ, Li B. A novel 3D imaging system for strawberry phenotyping.
684 *Plant Methods* 2017; **13**: 93.
- 685 26 Westoby MJ, Brasington J, Glasser NF, Hambrey MJ, Reynolds JM. ‘Structure-
686 from-Motion’ photogrammetry: A low-cost, effective tool for geoscience
687 applications. *Geomorphology*. 2012; **179**: 300–314.
- 688 27 Kovacs L, Eder M, Hollweck R *et al.* Comparison between breast volume
689 measurement using 3D surface imaging and classical techniques. *Breast* 2007; **16**:
690 137–145.
- 691 28 Rife TW, Poland JA. Field Book: An Open-Source Application for field data
692 collection on Android. *Crop Science*. 2014; **54**: 1624.
- 693 29 Verma S, Zurn JD, Salinas N *et al.* Clarifying sub-genomic positions of QTLs for
694 flowering habit and fruit quality in U.S. strawberry (*Fragaria x ananassa*) breeding
695 populations using pedigree-based QTL analysis. *Horticulture Research*. 2017; **4**.
696 doi:10.1038/hortres.2017.62.
- 697 30 Vickerstaff RJ, Harrison RJ. Crosslink: A fast, scriptable genetic mapper for
698 outcrossing species. *BioRxiv*, doi:10.1101/135277.

- 699 31 Cockerton HM, Vickerstaff RJ, Karlström A *et al.* Identification of powdery mildew
700 resistance QTL in strawberry (*Fragaria x ananassa*). *Theor Appl Genet* 2018; **131**:
701 1995–2007.
- 702 32 van Dijk T, Pagliarani G, Pikunova A *et al.* Genomic rearrangements and
703 signatures of breeding in the allo-octoploid strawberry as revealed through an allele
704 dose based SSR linkage map. *BMC Plant Biol* 2014; **14**: 55.
- 705 33 Laganiere R. OpenCV 3 Computer Vision Application Programming Cookbook.
706 Packt Publishing Ltd, 2017.
- 707 34 Podczeck F, Rahman SR, Newton JM. Evaluation of a standardised procedure to
708 assess the shape of pellets using image analysis. *Int J Pharm* 1999; **192**: 123–138.
- 709 35 Taubin G. Estimating the tensor of curvature of a surface from a polyhedral
710 approximation. *Proceedings of IEEE International Conference on Computer Vision*
711 1995. doi:10.1109/iccv.1995.466840.
- 712 36 Lancaster P, Salkauskas K. Surfaces generated by moving least squares methods.
713 *Mathematics of Computation*. 1981; **37**: 141–141.
- 714 37 Gutierrez PA, Perez-Ortiz M, Sanchez-Monedero J, Fernandez-Navarro F, Hervás-
715 Martínez C. Ordinal Regression methods: survey and experimental study. *IEEE*
716 *Transactions on Knowledge and Data Engineering*. 2016; **28**: 127–146.
- 717 38 Chakrabarti A, Ghosh JK. AIC, BIC and recent advances in model selection.
718 *Philosophy of Statistics*. 2011; 583–605.
- 719 39 Yamashita T, Yamashita K, Kamimura R. A stepwise AIC method for variable
720 selection in linear regression. *Communications in Statistics - Theory and Methods*.
721 2007; **36**: 2395–2403.
- 722 40 Garin V, Wimmer V, Mezrouk S, Malosetti M, van Eeuwijk F. How do the type of
723 QTL effect and the form of the residual term influence QTL detection in multi-parent
724 populations? A case study in the maize EU-NAM population. *Theor Appl Genet*
725 2017; **130**: 1753–1764.
- 726 41 Churchill GA, Doerge RW. Empirical threshold values for quantitative trait mapping.
727 *Genetics* 1994; **138**: 963–971.
- 728 42 Zeng ZB. Theoretical basis for separation of multiple linked gene effects in mapping
729 quantitative trait loci. *Proc Natl Acad Sci U S A* 1993; **90**: 10972–10976.
- 730 43 Zeng Z-B. Precision mapping of quantitative trait loci. *Genetics* 1994; **136**: 1457–
731 1468.
- 732 44 Oo LM, Aung NZ. A simple and efficient method for automatic strawberry shape
733 and size estimation and classification. *Biosystems Engineering*. 2018; **170**: 96–107.

- 734 45 Kochi N, Tanabata T, Hayashi A *et al.* A 3D shape-measuring system for assessing
735 strawberry fruits. *International Journal of Automation Technology*. 2018; **12**: 395–
736 404.
- 737 46 Aho K, Derryberry D, Peterson T. Model selection for ecologists: the worldviews of
738 AIC and BIC. *Ecology*. 2014; **95**: 631–636.
- 739 47 Castro P, Lewers KS. Identification of quantitative trait loci (QTL) for fruit-quality
740 traits and number of weeks of flowering in the cultivated strawberry. *Molecular*
741 *Breeding*. 2016; **36**. doi:10.1007/s11032-016-0559-7.
- 742 48 Lerceteau-Köhler E, Moing A, Guérin G *et al.* Genetic dissection of fruit quality
743 traits in the octoploid cultivated strawberry highlights the role of homoeo-QTL in
744 their control. *Theor Appl Genet* 2012; **124**: 1059–1077.
- 745 49 Liao X, Li M, Liu B *et al.* Interlinked regulatory loops of ABA catabolism and
746 biosynthesis coordinate fruit growth and ripening in woodland strawberry. *Proc Natl*
747 *Acad Sci U S A* 2018; **115**: E11542–E11550.
- 748 50 Wang S-M, Wei-jia LI, Yue-xue LIU, He LI, Yue MA, Zhang Z-H. Comparative
749 transcriptome analysis of shortened fruit mutant in woodland strawberry (*Fragaria*
750 *vesca*) using RNA-Seq. *Journal of Integrative Agriculture*. 2017; **16**: 828–844.
- 751 51 Ariza MT, Soria C, Medina JJ, Martínez-Ferri E. Fruit misshapen in strawberry
752 cultivars (*Fragaria xananassa*) is related to achenes functionality. *Annals of Applied*
753 *Biology*. 2011; **158**: 130–138.
- 754 52 Pipattanawong R, Yamane K, Fujishige N, Bang S-W, Yamaki Y. Effects of high
755 temperature on pollen quality, ovule fertilization and development of embryo and
756 achene in ‘Tochiotome’ strawberry. *Journal of the Japanese Society for*
757 *Horticultural Science*. 2009; **78**: 300–306.