The Genetic Architecture of Leaf Stable Carbon Isotope Composition in *Zea mays* and the Effect of Transpiration Efficiency on Elemental Accumulation

Crystal A. Sorgini^{*}, Lucas M. Roberts^{*}, Asaph B. Cousins[†], Ivan Baxter[‡], Anthony J. Studer^{*,§}

Affiliation:

^{*} Department of Crop Sciences, University of Illinois, Urbana, IL 61801, USA

[†] School of Biological Sciences, Washington State University, Pullman, Washington 99164, USA

[‡] Donald Danforth Plant Science Center, St. Louis, MO 63132, USA

[§] Corresponding Author

Running Title:

Genetic control of maize leaf $\delta^{13}\text{C}$

Key Words:

Zea mays, carbon isotopes, transpiration efficiency, specific leaf area, ionomics

Article Summary:

Quantitative genetics approaches were used to investigate the genetic architecture of leaf stable carbon isotope discrimination ($\delta^{13}\text{C}$) in maize. Developing a better understanding of leaf $\delta^{13}\text{C}$ could facilitate its use in breeding for reduced transpirational water loss. Several genomic regions were identified that contribute to the variation observed in leaf $\delta^{13}\text{C}$. Furthermore, contrary to what has been observed in other species, leaf $\delta^{13}\text{C}$ was not correlated with specific leaf area. Finally, a leaf ionomic analysis indicates that a reduction in transpiration, and thus mass flow, would not result in a decrease in nutrient accumulation.

Corresponding Author:

Anthony J. Studer

Department of Crop Sciences, University of Illinois

1201 West Gregory Drive, Edward R. Madigan Laboratory, #289

Urbana, IL 61801 USA

217-244-5469, astuder@illinois.edu

ABSTRACT

With increased demand on freshwater resources for agriculture, it is imperative that more water-use efficient crops are developed. Leaf stable carbon isotope composition, $\delta^{13}\text{C}$, is a proxy for transpiration efficiency and a possible tool for breeders, but the underlying mechanisms effecting $\delta^{13}\text{C}$ in C_4 plants are not known. It has been suggested that differences in specific leaf area, which potentially reflects variation in internal CO_2 diffusion, can impact leaf $\delta^{13}\text{C}$. However, at this point the relationship has not been tested in maize. Furthermore, although it is known that water movement is important for elemental uptake, it is not clear how manipulation of transpiration for increased water-use efficiency may impact nutrient accumulation. Here we characterize the underlying genetic architecture of leaf $\delta^{13}\text{C}$ and test its relationship to specific leaf area and the ionome in four biparental populations of maize. Five significant QTL for leaf $\delta^{13}\text{C}$ were identified, including both novel QTL as well as some that were identified previously in maize kernels. One of the QTL regions contains an Erecta-like gene, the ortholog of which has been shown to regulate transpiration efficiency and leaf $\delta^{13}\text{C}$ in *Arabidopsis*. Our data does not support a relationship between $\delta^{13}\text{C}$ and specific leaf area, and of the 19 elements analyzed, only a weak correlation between molybdenum and $\delta^{13}\text{C}$ was detected. Together these data begin to build a genetic understanding of leaf $\delta^{13}\text{C}$ in maize and suggest the potential to improve plant water use without significantly influencing elemental homeostasis.

INTRODUCTION

The impacts of global population growth and climate change on natural resources indicate that the future of food security will depend on increasing both the productivity and sustainability of agriculture systems (National Academies of Sciences, Engineering, and Medicine, 2018). Improving crop water-use efficiency (WUE) would ameliorate the effects of the increasing frequency and severity of droughts (Scheffield and Wood 2008; Chapman *et al.* 2012, Leakey 2019). Agronomic WUE can be defined as the amount of yield, whether grain or biomass, produced per the total amount of water utilized by the crop (Condon *et al.* 2004). Many factors can affect WUE including transpirational water loss through the stomatal pores on the leaf's surface. In C_3 plants the amount of carbon available for assimilation is limited by stomatal and mesophyll conductances to CO_2 (Flexas *et al.* 2016) and therefore correlated to the rate of transpiration. For example, yield was shown to be positively associated with cumulative transpiration in soybean (Purcel 2007), and higher net carbon assimilation was accompanied by higher transpiration in rice (Adachi *et al.* 2017). However, higher rates of biomass yield do not always correspond to higher transpiration rates in C_4 plants due to the evolution of the carbon concentrating mechanism. The uncoupling of CO_2 assimilation and transpiration has been demonstrated in field and greenhouse grown maize (Walker 1986, Kolbe *et al.* 2018a). Thus, there is the potential to increase transpiration efficiency, or carbon gain per amount of water transpired, without reducing productivity in C_4 species (Leakey, 2019). A large amount of variation is present in the transpiration rates of C_4 crop species, including sorghum (Hammer *et al.* 1997) and maize (Bunce 2010), suggesting that existing occurring alleles could be exploited for optimizing WUE.

Although increasing transpiration efficiency provides a strategy to avoid the negative effects of water limitation on plant growth and development (Passioura 1996, Chaves *et al.* 2002, Jaleel *et al.* 2009), there is the possibility of pleiotropic side effects given the fundamental requirement for water movement in plants. A potential impact of reducing transpiration could be a corresponding reduction in the uptake and mobilization of water-soluble nutrients. As water is absorbed by roots, nutrients in solution come in contact with the root surface in a process known as mass flow (Barber *et al.* 1963). Most nutrients are acquired by mass flow, although phosphorus is a notable exception that contacts the root through diffusion (Barber *et al.* 1962).

Reducing transpiration may also affect nutrient uptake facilitated by symbiosis with mycorrhizal fungi (Marschner and Dell 1994). Therefore, the manipulation of basic plant processes such as transpiration for improved WUE must also consider potential impacts on the availability of essential plant nutrients. Previous research has shown in the C₄ plant sorghum that total leaf mineral content is positively correlated to transpiration efficiency (Masle *et al.* 1992). While meta-analyses of high CO₂ grown plants with reduced transpiration have shown a drastic reduction in nutrient accumulation in C₃ crops (McGrath and Lobell 2013), sorghum showed no difference and maize had similar levels of zinc, protein, and phytate, but a decrease in iron accumulation (Meyers *et al.* 2014). Although part of the reduction in nutrient content can be explained by dilution, due to increased growth at high CO₂, this does not completely account for the observed reduction. An ionomics (high-throughput elemental profiling) approach has been used in maize to assess kernel nutrient content (Baxter *et al.* 2014). A similar ionomics approach in leaf tissue could be used to assess the effect of transpiration on nutrient uptake.

The difficulty and labor-intensive nature of accurately quantifying the amount of water that an individual plant transpires has been a major limitation to breeding for transpiration efficiency. This has resulted in the selection for drought tolerance rather than applying a direct selection for water use. One alternative method is the use of leaf stable carbon isotopes as a proxy for transpiration efficiency. The stable carbon isotope composition, $\delta^{13}\text{C}$, reflects the amount of ¹³C present in plant tissue relative to a standard (Keeling 1979). Enzymes in the process of carbon fixation discriminate differently against the heavier ¹³C atoms in a process known as fractionation (Farquhar *et al.* 1982, O'Leary 1988). It has been widely shown that stable carbon isotopes can be used as a proxy trait for quantifying a plant's transpiration efficiency in C₃ plants (Farquhar *et al.* 1989a, Farquhar *et al.* 1989b, Condon *et al.* 1990, Virgona *et al.* 1990, Condon *et al.* 1993, Barbour *et al.* 2010) and in C₄ plants (Henderson *et al.* 1998, von Cammerer *et al.* 2014, Ellsworth *et al.* 2017, Twohey III *et al.* 2019, Ellsworth *et al.* 2020). Studies have also shown that $\delta^{13}\text{C}$ can be influenced by environmental factors including light intensity and drought (reviewed in Cernusak *et al.* 2013). However, the genetic control of $\delta^{13}\text{C}$ remains unknown in C₄ species.

Kolbe *et al.* showed that $\delta^{13}\text{C}$ did not correlate with any of the photosynthetic enzymes previously posited to control for $\delta^{13}\text{C}$ variation (2018b). Additionally, a transcriptome analysis was unable to identify a clear candidate gene (Kolbe *et al.* 2018b). Quantitative genetic approaches have the potential to reveal the genetic control of $\delta^{13}\text{C}$ in C_4 species because genomic locations are tested for associations with the trait of interest, without *a priori* knowledge of the mechanism underlying the variation. Maize is ideal for use in mapping studies due to its high level of recombination and low linkage disequilibrium (Yu and Buckler 2006). Mapping methods have been successfully used for decades to identify genes controlling complex traits in maize, with evolving approaches to tackle more difficult traits (Wallace *et al.* 2014). In addition, maize is both a model organism with available populations and genomic data, and one of the three most important global crops contributing to 30% of the total calories consumed by humans (Shiferaw *et al.* 2011).

There have been several previous studies that used quantitative genetics to investigate $\delta^{13}\text{C}$ in C_3 species (Teulat *et al.* 2002, Masle *et al.* 2005, Rebetzke *et al.* 2008, Xu *et al.* 2009). In *Arabidopsis* the gene *ERECTA* was identified in a QTL study for isotopic discrimination and was found to alter transpiration efficiency by altering stomatal density (Masle *et al.* 2005). Genetic mapping of leaf $\delta^{13}\text{C}$ has also been performed in the C_4 species *Setaria viridis* (Feldman *et al.* 2018, Ellsworth *et al.* 2020) and kernel $\delta^{13}\text{C}$ has been mapped in the C_4 maize (Gresset *et al.* 2014, Avramova *et al.* 2019). Although the QTL found for C_4 species still require fine-mapping to identify the causative gene, no correlation was observed between kernel $\delta^{13}\text{C}$ and leaf $\delta^{13}\text{C}$ (Foley 2012). The lack of correlation may be the result of post-photosynthetic fractionation (Badeck *et al.* 2005), and therefore mapping QTL for $\delta^{13}\text{C}$ in leaves may reveal additional loci not found using kernels. In this manuscript, we focus on leaf $\delta^{13}\text{C}$ in maize and its association with leaf elemental composition. We also investigate variation in specific leaf area (SLA) and its potential relationship to leaf $\delta^{13}\text{C}$ by CO_2 diffusion. Characterization of the genetic architecture of leaf $\delta^{13}\text{C}$ will provide a better context for understanding what drives $\delta^{13}\text{C}$, which will allow breeders to utilize this trait in crop improvement.

MATERIALS AND METHODS

Plant Material

All experiments were planted at the University of Illinois Crop Sciences Research Farm, Urbana IL and were subject to natural conditions without supplemental irrigation. NAM RIL families CML103, CML333, NC358, and Tx303 and NAM founder parents (McMullen *et al.* 2009) are publicly available through the Maize Genetic Cooperative Stock Center. The NAM RIL families were planted in the summer of 2015 using an augmented incomplete block design. For this experiment fifteen kernels were planted in each 3.7 meter row with 0.8 meter spacing between rows and 0.9 meter alleys. The families were randomized together, with each block consisting of 20 lines and 2 checks (B73 and one of the other founder lines). Of the 880 plots, 10% were dedicated to checks with the common parent B73 appearing in 40 plots and each of the four founder lines appears in ten plots. All lines used for the GWAS experiment are publicly available through the USDA Germplasm Resources Information Network (GRIN). This experiment was planted on May 23rd 2016. Twenty kernels were planted in each 3.7 meter row with 0.8 meter spacing between rows and 0.9 meter alleys, and then thinned to 15 plants per row. A complete list of lines used can be found in Tables S1-S3.

Tissue Sampling

Samples for $\delta^{13}\text{C}$ analysis from the NAM RIL populations were collected six weeks after planting as follows. A rectangular piece of tissue approximately 7.5 cm X 5 cm was taken from the center of the leaf blade of the uppermost fully expanded leaf from four plants in each plot. Samples were placed in a coin envelope and dried at 65°C for at least 7 days. After drying four hole punches (each 0.058532 cm²) were taken and placed in a 6 mm x 4 mm tin capsules (OEA Laboratories # C11350.500P) for analysis using a Delta PlusXP (Washington State University) isotope ratio mass spectrometer. Leaf samples to measure specific leaf area (SLA) were collected from four plants in each of the plots (preferentially but not necessarily the same plants as were collected for $\delta^{13}\text{C}$) using a 1.6 cm diameter cork borer. Leaf discs were dried at 65°C for at least 7 days prior to weighing on an analytical balance (Model MS204S). Specific leaf area (SLA) was calculated as the area of a leaf disc divided by its dry weight. These same leaf discs were then used for ionomics analyses as described in Pauli *et al.* 2018. Leaf samples for $\delta^{13}\text{C}$ analysis from the GWAS panel were collected from the uppermost fully expanded leaf seven weeks after

planting using the hole punch method and processed as described previously (Twohey III *et al.* 2019). Due to the high level of diversity in this panel, some lines were flowering when samples were collected, which resulted in tissue being collected from the flag leaf. These samples were analyzed using a Costech instruments elemental combustion system and a Delta V Advantage isotope ratio mass spectrometer.

Statistical Analysis

All analyses were completed using custom scripts and statistical packages in R (R Core Team 2017).

Correlation analysis: Pearson correlations using phenotype mean values were calculated with `corr.test()` in R package ‘psych’ (Revelle 2018) using complete observations and Holm’s method (Holm 1979) to adjust p-values for multiple testing. The correlation matrix was visualized using `pairs.panel()` in the R package ‘psych’ (Revelle 2018).

Stepwise regression QTL mapping: The analysis was completed using NAM_phasedImputed_1cM_AllZeaGBSv2.3 dataset. The file contains fully imputed and phased genotypes for most of the RILs in the NAM population (Zhao *et al.* 2006; Lipka *et al.* 2015). This HapMap format file was converted to numeric format where 0 is the B73 homozygote reference, 1 is a heterozygote, and 2 is the homozygote alternative parent. Phenotypic means were regressed onto genotype. Lowest p-values from the ANOVA values of the linear model were recorded (i.e. `pvalues[i] = anova(lm(mypheno~geno[i,]))`). The previously identified marker was added to the model and re-run in a stepwise regression procedure. Significance thresholds were determined by 200 permutations and alpha was set at 0.05. All analysis was completed using custom scripts in R (R Core Team 2017). Results were then compared to composite interval QTL mapping completed in R package ‘r/QTL’ (Broman *et al.* 2003).

Joint linkage mapping: The analysis was completed using HapMapv2 (Chia *et al.* 2012). The genotypic dataset consisted of 836 markers were scored on 624 RILs from four biparental families with B73 as a common parent. The marker subset represented markers that could be placed unambiguously on the physical map. Unambiguous markers are defined by those

anchored in CDS positions of genes that have held consistent over genome versions verified by MaizeGDB cross reference tables. Missing data were imputed as previously described in Tain *et al.* (2011). Joint linkage models were constructed using custom script in R (R Core Team 2017) by a stepwise regression procedure. In general, we used linkage to test every marker across all four families to find the most significant QTL. The model has a family term and a marker:family term. The family term accounts for differences in mean phenotype between families. Inclusion of the marker:family term means that for each QTL we are assigning a separate effect to each family. The family term was included in the model and each of the 836 possible marker-by-family terms were assessed. Lowest p-values from the ANOVA values of the linear regression model were recorded (i.e. `JL_pvalues[i] = anova(lm(my_pheno~family+geno[,i]:family))`). All 836 marker-by-family terms were tested. SNP effects were nested in families to reflect the potential for unique QTL allele effects within each family. The lowest resulting *p*-value was recorded for each permutation. Significance thresholds were determined using 1000 permutations for each family independently and alpha was set at 0.05.

Genome wide association study: A subset of 413 of 503 diverse lines from Hirsch *et. al* 2014 that included the Wisconsin Diversity Set of Hansey *et al* 2011 was grown in 2016 and listed in Table S2. Hirsch *et. al* 2014 collected RNA from whole seedling tissue which was sequenced via IlluminaHiSeq and filtered to create a working set of 485,179 SNPs that is available at <https://datadryad.org/resource/doi:10.5061/dryad.r73c5>. The 413 lines were grown in 2016 and tissue was sampled when B73 was at the developmental stage V10. Isotopic analysis is described above. A genome wide association analysis was run using R package ‘GAPIT’ (Lipka *et al.* 2012) on leaf $\delta^{13}\text{C}$. Removal of SNPs with a minor allele frequency of less than 0.05 resulted in a subset of 438,222 SNPs being used in this analysis. A MLM model was used with model selection set to true to find the optimum number of principal components to account for population structure (Lipka *et. al* 2012). Significance threshold were calculated using the Bonferroni correction of familywise error rate. An alternative significance test was calculated using the Benjamini-Hochberg procedure for controlling the false discovery rate (Benjamini & Hochberg 1995).

Data Availability

Genotypic datasets were downloaded from Panzea CyVerse iPlant Data Storage Commons (<http://datacommons.cyverse.org/browse/iplant/home/shared/panzea>). All phenotypic datasets were quality controlled for complete technical replicates, outliers, and availability of genotypic data. A list of all genotypes used in each analysis is provided in Tables S1-S4 and have been uploaded to figshare. Briefly, the $\delta^{13}\text{C}$ analysis was completed with 640 RILs; including 156 CML103 RILs, 160 CML333 RILs, 159 NC358 RILs, and 165 Tx303 RILs (Table S1). The element analysis was completed using a total of 704 RILs; including 175 CML103 RILs, 181 CML333 RILs, 175 NC358 RILs, and 173 Tx303 RILs (Table S1). The SLA analysis used a total of 683 RILs; including 172 CML103 RILs, 176 CML333 RILs, 168 NC358 RILs, and 167 Tx303 RILs (Table S1). The Joint linkage analysis was completed using a total of 624 RILs; including 154 CML103 RILs, 159 CML333 RILs, 151 NC358 RILs, and 160 Tx303 RILs (Table S2). Table S3 lists the 413 lines used in the GWAS of leaf $\delta^{13}\text{C}$. Table S4 includes the QTL coordinates identified in the elemental QTL analyses. Figure S1 shows the distribution of leaf $\delta^{13}\text{C}$ for each of the NAM RIL families. Figure S2 presents the correlation matrix for the elemental analysis, and Figure S3 shows the chromosomes where significant QTL were identified for each element. Figure S4 is the LOD plot from the GWAS mapping of leaf $\delta^{13}\text{C}$.

RESULTS

Single family QTL mapping

Previous studies investigating leaf $\delta^{13}\text{C}$ in maize indicated that the NAM founder lines CML103, CML333, and Tx303 consistently contrast B73 with respect to leaf $\delta^{13}\text{C}$ (Kolbe *et al.* 2018b; Twohey III *et al.* 2019). The founder line NC358 had a moderate leaf $\delta^{13}\text{C}$ value (Kolbe *et al.* 2018b; Twohey III *et al.* 2019) and was also included in this study. The RIL families generated from these four founder lines were grown for linkage analysis. Consistent with previous studies, both the CML103 and CML333 parent lines had a significantly less negative leaf $\delta^{13}\text{C}$ than B73 ($p < 0.05$), when grown as replicated controls among the RILs. However, the Tx303 and NC358 parental lines were not found to be significantly different from B73. Transgressive segregation was observed in all four RIL families (Fig. S1).

Stepwise regression analyses found significant QTL for leaf $\delta^{13}\text{C}$ in the NAM RIL families CML103, CML333, and Tx303 but not in NC358 (Fig. 1A). Interestingly, none of these QTL were shared between RIL families in this analysis. The CML103 RIL family had two significant QTL, one on chromosome 5 at 211.7 Mb and another on chromosome 7 at 142.4 Mb. Combined these two QTL accounted for 21.36% of the total phenotypic variance (Table 1). The RIL family CML333 had one significant QTL on chromosome 3 at 183.9 Mb, which accounted for 8.37% of the total phenotypic variation explained (Table 1). Finally, the Tx303 RIL family had a significant QTL on chromosome 2 at 13.5 Mb explaining 9.48% of phenotypic variation (Table 1). No significant QTL for leaf $\delta^{13}\text{C}$ were identified in the NC358 RIL family.

Specific leaf area was used as a proxy trait to test for a relationship between leaf thickness and leaf $\delta^{13}\text{C}$. No significant correlation was observed between SLA and leaf $\delta^{13}\text{C}$ ($p = 0.304$). In addition to testing for a correlation with leaf $\delta^{13}\text{C}$, QTL mapping was performed for SLA to identify any possible overlaps with genomic regions identified for leaf $\delta^{13}\text{C}$. Mapping of SLA in the four RIL families identified two significant QTL. In the CML103 RIL family, a QTL was identified on chromosome 5 at 86.1 Mb and in the Tx303 RIL family a QTL on chromosome 9 at 107.8 Mb. Neither of the SLA QTL identified overlapped with QTL for leaf $\delta^{13}\text{C}$ (Fig. 1B).

To test a potential link between transpiration and nutrient uptake, an elemental analysis was performed on leaf samples from each of the four RIL families. Samples were analyzed for 19 different elements using an IPC-MS. A full correlation matrix shows that some elements are highly correlated with each other (Fig. S2), but no strong correlations ($r > \pm 0.7$) were identified with $\delta^{13}\text{C}$. However, there was a weak but significant correlation ($p = 6.745\text{E-}05$, $r = 0.18$) between leaf $\delta^{13}\text{C}$ and Mo (Fig. 2). Subsequent QTL mapping of the 19 element concentrations identified 28 QTL across 12 different elements (Fig. 3, Fig. S3). Significant QTL were found for B, Mg, P, S, K, Fe, Mn, Co, Cu, Rb, Sr, and Mo (Table S4). None of the elemental QTL overlapped with those found for leaf $\delta^{13}\text{C}$ or SLA. However, in some cases multiple elements had common QTL, such as Mg and Mn on chromosome 10 in the CML103 RIL family and Co and Cu on chromosome 3 in the NC358 RIL family. Additionally, common QTL for an element were found across families, as in the case of Mg in the NC358 and Tx303 RIL families.

Joint linkage QTL mapping

A joint linkage analysis was performed for leaf $\delta^{13}\text{C}$ to test whether any additional QTL would be identified by combining the four RIL families into a single analysis. The joint linkage analysis identified the same significant QTL for leaf $\delta^{13}\text{C}$ on chromosomes 2, 3, and 5 (Table 2) as in the single family stepwise regression analysis. Although the QTL on chromosome 7 was not found using the joint linkage approach, an additional significant QTL was identified on chromosome 1. Given that no significant QTL for leaf $\delta^{13}\text{C}$ were identified in the NC358 RIL family, we tested whether removing this family from the joint linkage analysis would change the outcome. When the joint linkage analysis was rerun excluding family NC358, the same four QTL were reidentified with decreased p -values, and the total phenotypic variation explained (R^2 value) increased in later steps of the model. However, no new QTL were identified with this approach.

Genome wide association study

Once significant QTL intervals were identified for leaf $\delta^{13}\text{C}$ using a biparental mapping strategy (Fig. 1), we performed a genome wide association study to try and narrow down the intervals to specific genic regions. The Wisconsin Diversity Panel was chosen because it represents a large portion of variation found within maize and has a robust publicly available 485,179 SNP set. A subset of 413 of the possible lines were chosen due to seed availability, and were grown in a single randomized block. No significant SNP associations with leaf $\delta^{13}\text{C}$ were identified (Fig. S4).

DISCUSSION

Leaf $\delta^{13}\text{C}$ has a moderately high heritability in maize (Twohey III *et al.* 2019), which facilitates the use of quantitative genetics approaches to pinpoint the genomic locations controlling this trait. Here we characterized the genetic control of $\delta^{13}\text{C}$ in maize using leaf tissue collected at vegetative stage V9-V10 to reflect the photosynthetic pool during active growth. We were able to identify several significant QTL for leaf $\delta^{13}\text{C}$ across three NAM RIL families using stepwise regression. Using these populations we were also successful in identifying QTL for SLA and 12

different elements. Contrary to our hypothesis, no significant correlation was observed between leaf $\delta^{13}\text{C}$ and SLA or elemental composition.

We strategically picked NAM RIL families based on the founder parents that had the largest differences in their $\delta^{13}\text{C}$ for single family and joint linkage analyses. However, we also included a parent which was not extremely different from B73. Interestingly, we were unable to identify significant QTL in the NC358 RIL family despite transgressive segregation. We interpret this result as an indication the NC358 contains only small effect QTL that were not detected in this study. Alternatively, NC358 leaf $\delta^{13}\text{C}$ may be more sensitive to the growing environment with a smaller genetic component. Twohey III *et al.* noted that while several maize lines were stable when tested in greenhouse and field environments, there were other lines that had highly variable isotopic signatures (2019). A large amount of environmental influence over this trait in some backgrounds would obscure the genetic contribution and our ability to detect significant QTL.

When we compared the regions identified here with regions previously mapped in *S. viridis* no obvious overlap was observed (Ellsworth *et al.*, 2020). However, of the QTL identified for kernel $\delta^{13}\text{C}$ in maize (Gresset *et al.* 2014, Avramova *et al.* 2019), our QTL for leaf $\delta^{13}\text{C}$ overlapped those on chromosomes 1, 3, and 7. This result demonstrates that some QTL for $\delta^{13}\text{C}$ are shared between tissues, and that these QTL are identified across several populations and environments. Therefore, while metabolic processes have the possibility of influencing the $\delta^{13}\text{C}$ as products are mobilized from source to sink tissues, our data would suggest that the initial source signature is maintained to a large degree in the kernel.

Although the QTL analyses presented here do not provide gene-level resolution, we were able to look for candidate genes within the intervals. The chromosome 5 QTL includes an Erecta-like gene (*er1*, GRMZM2G463904, 211.8 Mb). Unfortunately, stomatal density data were not collected on these populations, which would further support the role of *er1* in variation of leaf $\delta^{13}\text{C}$. This would be an interesting area of future research given that this gene was found to effect $\delta^{13}\text{C}$ in Arabidopsis by changing stomatal density (Masle *et al.* 2005). We also looked for genes that have been previously shown to directly influence transpiration efficiency in maize (reviewed in Leakey 2019). However, none of these were found to be located in our QTL intervals.

The linkage analyses using biparental mapping populations identified several significant QTL, but none of the single family QTL were independently identified in more than one family (Fig. 1). This result indicates that leaf $\delta^{13}\text{C}$ can be controlled by different factors depending on the genetic background. Furthermore, if in fact leaf $\delta^{13}\text{C}$ is controlled by many small effect QTL, this may explain why the GWAS did not identify any significant SNP associations with leaf $\delta^{13}\text{C}$. Identifying rare alleles with small to moderate effect size is a known weakness of the GWAS method (Bazakos *et al.* 2017). A better understanding of the mechanisms influencing leaf $\delta^{13}\text{C}$ would allow future analyses to move beyond single marker tests and instead look at SNPs in genes representing a particular pathway or process that could be collectively significant. This approach was successfully used to study maize lipid biosynthesis (Li *et al.* 2019).

The diffusion of CO_2 into mesophyll cells is a potential source of variation in leaf $\delta^{13}\text{C}$, which could be linked to stomatal density or leaf thickness. Previous work in maize has shown that stomatal density is not correlated with leaf $\delta^{13}\text{C}$ in a small diversity panel of maize (Foley 2012). SLA has not been linked to $\delta^{13}\text{C}$ in maize, but in rice $\delta^{13}\text{C}$ and SLA have shared QTL (This *et al.* 2010). In this study we were able to test SLA and $\delta^{13}\text{C}$ in four RIL families, and no correlation was observed. Likewise, a comparison of the QTL analyses showed no overlapping regions for SLA and those mapped for leaf $\delta^{13}\text{C}$. This result suggests two possibilities. First, it is possible that differences in SLA observed in these populations are not due to leaf thickness, but rather composition. Identifying the causative genes underlying the QTL would give insight into the mechanism. A second possibility is that leaf anatomical traits other than leaf thickness influence leaf $\delta^{13}\text{C}$. A variety of anatomical traits could affect $\delta^{13}\text{C}$ and would not be captured by measurement of SLA.

With our data we were able to indirectly test the relationship between nutrient uptake and transpiration. If reducing transpiration limits nutrient uptake, transpiration efficiency as a trait for increasing WUE would have limited application. The nineteen elements tested here were previously reported to have narrow-sense heritabilities ranging from 0.11 to 0.66 (Baxter 2014). The only element found to be significantly correlated to leaf $\delta^{13}\text{C}$ was Molybdenum. Molybdenum is required for several vital biological processes related to nitrogen and water

(Baxter 2008). Because molybdenum is a required cofactor for ABA synthesis, maize plants overexpressing molybdenum cofactor sulfuryase gene have increased drought tolerance (Lu *et al.* 2013). However, in our study we observed a positive correlation between leaf $\delta^{13}\text{C}$ and molybdenum, which is contrary to expectation given that an increase in $\delta^{13}\text{C}$ signifies a decrease in WUE. Overall, it is encouraging that the majority of elements sampled were not associated with $\delta^{13}\text{C}$. This suggests that breeding for leaf $\delta^{13}\text{C}$ as a means to reduce transpiration would be unlikely to result in plants with nutrient uptake deficiencies.

Although the main focus of this study was to investigate leaf $\delta^{13}\text{C}$ and its relationship to SLA and nutrient accumulation, the QTL mapping of the analyzed elements was an interesting biproduct. Mapping the leaf ionome of the four RIL families resulted in many significant QTL, including some overlapping intervals for different elements. Multi element QTLs are common, and are thought to be due to loci affecting processes such as the acidification of the rhizosphere or altering the permeability of the casparian strip (Baxter 2015). Interestingly there was not much overlap between the ionomic QTL identified here and a previous study on kernels (Table 4 Baxter *et al.* 2014). The only overlapping QTL was for Rb85 on chromosome 3. There are several possible causes for the limited overlap between these methods. The ionome is strongly influenced by genotype by environment interactions, with many of the QTL identified in previous studies being location specific (Asaro *et al.* 2016). Additionally, there could be differences between the leaf and grain ionome is due to differential mobilization of nutrients from vegetative tissues into kernels during grain fill.

Acknowledgements

We thank Aaron Slack, Robert Twohey III, and Mengqiao Han for technical assistance with growing and sampling the mapping populations. This work was supported by United States Department of Agriculture - Hatch, a United States Department of Agriculture - Agriculture and Food Research Initiative grant (2019-67013-29195), the United States Department of Agriculture - Agricultural Research Service (5070-21000-039-00D), as well as a Department of Energy, Office of Science, Office of Biological and Environmental Research grant (DE-SC0018277).

Tables

Table 1: $\delta^{13}\text{C}$ Single Family Stepwise Regression QTL

Step	Marker	Family	Chr.	Position (Mb)	<i>p</i> -value	TPVE* (%)	Effect Size	1LOD Interval (Mb)
1	824	CML103	5	211.7	7.03E-06	12.32	0.1286	211.5 - 212.9
2	1032	CML103	7	142.4	7.52E-05	21.36	-0.1064	141.2 - 149.7
1	470	CML333	3	183.9	2.07E-04	8.37	0.1137	178.6 - 195.7
1	251	Tx303	2	13.5	6.04E-05	9.48	-0.1457	13.5 - 15.2

* Total Percent of Variation Explained

Table 2: $\delta^{13}\text{C}$ Joint Linkage Mapping QTL

All Four Families								
Step	QTL	Family	Chr.	Position (Mb)	<i>p</i> -value	TPVE* (%)	Effect Size	1LOD Interval (Mb)
1	m0200	Tx303	2	13.8	2.82E-06	7.9	-0.144	12.6 - 15.8
2	m0677	CML103	5	211.2	1.98E-04	11.8	0.123	211.2 - 212.7
3	m0385	CML103	3	182.1	1.44E-04	15.24	0.115	180.0 - 195.3
4	m0132	Tx303	1	263.2	4.41E-05	18.19	-0.126	257.1 - 263.6
Excluding NC358								
Step	QTL	Family	Chr.	Position (Mb)	<i>p</i> -value	TPVE* (%)	Effect Size	1LOD Interval (Mb)
1	m0200	Tx303	2	13.8	5.70E-06	7.47	-0.144	12.6 - 15.9
2	m0677	CML103	5	211.2	2.86E-04	12.31	0.123	211.2 - 212.7
3	m0385	CML103	3	182.1	1.60E-04	16.45	0.115	180.0 - 195.3
4	m0132	Tx303	1	263.2	5.97E-05	20.05	-0.121	253.0 - 263.6

Figure Captions

Figure 1 $\delta^{13}\text{C}$ and SLA Single Family Stepwise Regression QTL Mapping. $\delta^{13}\text{C}$ QTL (A) were identified in NAM RIL families CML103 (black), CML333 (orange), and Tx303 (grey) but not in NC358 (blue). Specific leaf area (SLA) QTL (B) were identified in NAM RIL families CML103 (black) and Tx303 (grey). Significance thresholds (dashed horizontal line) were determined by 200 permutations and an alpha of 0.05.

Figure 2 Pearson's r Correlations. Correlations of mean phenotypic values using complete observations and Holm's method to adjust p -values for multiple testing. The heat map shows no strong correlations between $\delta^{13}\text{C}$ mean values and element mean values. $\delta^{13}\text{C}$ and Mo have a significant p -value ($p = 6.745\text{E-}05$, $r = 0.18$).

Figure 3 Element Single Family Stepwise Regression QTL Mapping. QTL mapping identified 28 QTL across 12 different elements. Significant QTL ($\alpha = 0.05$) for each element are plotted. QTL location is shown across the 10 maize chromosomes (cM) on the x-axis. Dashes indicate a significant QTL, with the NAM RIL family in which the QTL was found designated by color; CML103 (black), CML333 (orange), Tx303 (grey), NC358 (blue). All dashes are the same length for visibility.

Supplemental Figure 1 NAM RIL Transgressive Segregation. NAM RIL families CML103 (A), CML333 (B), NC358 (C), and Tx303 (D) were sorted by $\delta^{13}\text{C}$ and plotted. Parental lines are shown in red.

Supplemental Figure 2 Element and $\delta^{13}\text{C}$ Full Correlation Matrix. A full correlation matrix of element and $\delta^{13}\text{C}$ mean values is shown. The diagonal displays histograms of each dataset. Pearson's r is shown in the upper panel. Scatter plots and best fit line are shown in the lower panel.

Supplemental Figure 3 Element QTL Mapping by Chromosome. Significant element QTL are shown by maize chromosomes 1 through 10 on the x-axis (in cM). Each NAM RIL family is represented by a symbol; CML103 ($\circ$), CML333 ($\times$), NC358 ($\square$), and Tx303 (Δ). Each element

is designated by color. Significance thresholds (dashed horizontal line) were determined using 200 permutations, $\alpha=0.05$ for each QTL independently.

Supplemental Figure 4 Genome Wide Association Study for $\delta^{13}\text{C}$. Manhattan plot showing significance of SNPs derived from a mixed linear model using the Bayesian information criterion to select the optimal number of principal components. The significance threshold represents the Bonferroni correction of familywise error rate.

Literature Cited

- Adachi, S., K. Yoshikawa, U. Yamanouchi, T. Tanabata, J. Sun, T. *et al.*, 2017 Fine mapping of carbon assimilation rate 8, a quantitative trait locus for flag leaf nitrogen content, stomatal conductance and photosynthesis in rice. *Front. Plant Sci.*, 8: 60.
- Asaro, A., G. Ziegler, C. Ziyomo, O. A. Hoekenga, B. P. Dilkes, B.P. and Baxter, I., 2016 The interaction of genotype and environment determines variation in the maize kernel ionome. *G3: Genes, Genomes, Genetics*, 6: 4175-4183.
- Avramova, V., A. Meziane, E. Bauer, S. Blankenagel, S. Eggels *et al.*, 2019 Carbon isotope composition, water use efficiency, and drought sensitivity are controlled by a common genomic segment in maize. *Theor. Appl. Genet.* 132: 53-63.
- Badeck, F., G. Tcherkez, S. Nogues, C. Piel, and J. Ghashghaie, 2005 Post-photosynthetic fractionation of stable carbon isotopes between plant organs—a widespread phenomenon. *Rapid Comm. Mass Spect.* 19: 1381-1391.
- Barber, S.A. 1962 A diffusion and mass-flow concept of soil nutrient availability. *Soil Sci.* 93: 39–49.
- Barber, S. A., J. M. Walker, and E. H. Vasey, 1963 Mechanisms for movement of plant nutrients from soil and fertilizer to plant root. *J. Agric. Food Chem.* 11: 204-207.
- Barbour, M. M., C. R. Warren, G. D. Farquhar, G. Forrester, and H. Brown, 2010 Variability in mesophyll conductance between barley genotypes, and effects on transpiration efficiency and carbon isotope discrimination. *Plant Cell Environ.* 33: 1176-1185.
- Baxter, I., B. Muthukumar, H. C. Park, P. Buchner, B. Lahner *et al.*, 2008 Variation in molybdenum content across broadly distributed populations of *Arabidopsis thaliana* is controlled by a mitochondrial molybdenum transporter (MOT1). *PLoS Genet.* 4: e1000004.

Baxter, I. R., G. Ziegler, B. Lahner, M. V. Mickelbart, R. Foley *et al.*, 2014 Single-kernel ionomic profiles are highly heritable indicators of genetic and environmental influences on elemental accumulation in maize grain (*Zea mays*). PLoS ONE 9: e87628.

Baxter, I. 2015 Should we treat the ionome as a combination of individual elements, or should we be deriving novel combined traits? J. Exp. Bot. 66: 2127-2131.

Bazakos, C., Hanemian, M., Trontin, C., Jiménez-Gómez, J. M., and Loudet, O. 2017 New strategies and tools in quantitative genetics: how to go from the phenotype to the genotype. Ann. Rev. Plant Biol. 68: 435-455.

Broman, K. W., H. Wu, S. Sen, and G. A. Churchill, 2003 R/qtl: QTL mapping in experimental crosses. Bioinformatics 19: 889-890.

Bunce, J. A., 2010 Leaf transpiration efficiency of some drought-resistant maize lines. Crop Sci. 50: 1409-1413.

von Caemmerer, S., O. Ghannoum, J. J. L. Pengelly, and A. B. Cousins, 2014 Carbon isotope discrimination as a tool to explore C_4 photosynthesis. J. Exp. Bot. 65: 3459-3470.

Cernusak, L. A., N. Ubierna, K. Winter, J. A. M. Holtum, J. D. Marshall *et al.*, 2013 Environmental and physiological determinants of carbon isotope discrimination in terrestrial plants. New Phytol. 200: 950-965.

Chapman, S. C., S. Chakraborty, M. F. Dreccer, and S. M. Howden, 2012 Plant adaptation to climate change—opportunities and priorities in breeding. Crop Pasture Sci. 63: 251-268.

Chaves, M. M., J. S. Pereira, J. Maroco, M. L. Rodrigues, C. P. P. Ricardo *et al.*, 2002 How plants cope with water stress in the field? Photosynthesis and growth. Annals Bot. 89: 907-916.

- Chia, J., C. Song, P. J. Bradbury, D. Costich, N. De Leon *et al.*, 2012 Maize HapMap2 identifies extant variation from a genome in flux. *Nat. Genet.* 44: 803.
- Condon, A. G., G. D. Farquhar, and R. A. Richards, 1990 Genotypic variation in carbon isotope discrimination and transpiration efficiency in wheat. Leaf gas exchange and whole plant studies. *Funct. Plant Biol.* 17: 9-22.
- Condon, A. G., R. A. Richards, and G. D. Farquhar, 1993 Relationships between carbon isotope discrimination, water use efficiency and transpiration efficiency for dryland wheat. *Australian J. Agric. Res.* 44: 1693-1711.
- Condon, A. G., R. A. Richards, G. J. Rebetzke, and G. D. Farquhar, 2004 Breeding for high water-use efficiency. *J. Exp. Bot.* 55: 2447-2460.
- Ellsworth, P. Z., P. V. Ellsworth, and A. B. Cousins, 2017 Relationship of leaf oxygen and carbon isotopic composition with transpiration efficiency in the C₄ grasses *Setaria viridis* and *Setaria italica*. *J. Exp. Bot.* 68: 3513-3528.
- Ellsworth, P., M. Feldman, I. Baxter, and A. Cousins, 2020 A genetic link between whole-plant water use efficiency and leaf carbon isotope composition in the C₄ grass *Setaria*. *Plant J.* <https://doi.org/10.1111/tpj.14696>.
- Farquhar, G. D., J. R. Ehleringer, and K. T. Hubick, 1989a Carbon isotope discrimination and photosynthesis. *Annu. Rev. Plant Biol.* 40: 503-537.
- Farquhar, G. D., K. T. Hubick, A. G. Condon, and R. A. Richards, 1989b Carbon isotope fractionation and plant water-use efficiency, pp. 21-40 in *Stable isotopes in ecological research*, edited by P. W. Rundel, J. R. Ehleringer, and K. A. Nagy. Springer, New York.

Feldman, M. J., P. Z. Ellsworth, N. Fahlgren, M. A. Gehan, A. B. Cousins *et al.*, 2018
Components of water use efficiency have unique genetic signatures in the model C₄ grass
Setaria. *Plant Physiol.* 178: 699-715.

Flexas J., Díaz-Espejo A., Conesa M. A., Coopman R. E., Douthe C., *et al.* 2016 Mesophyll
conductance to CO₂ and Rubisco as targets for improving intrinsic water use efficiency in C₃
plants. *Plant, Cell & Environment.* 39: 965-82.

Gresset, S., P. Westermeier, S. Rademacher, M. Ouzunova, T. Presterl *et al.*, 2014 Stable carbon
isotope discrimination is under genetic control in the C₄ species maize with several genomic
regions influencing trait expression. *Plant Physiol.* 164: 131-143.

Hirsch, C. N., J. M. Foerster, J. M. Johnson, R. S. Sekhon, G. Muttoni *et al.* 2014 Insights into
the maize pan-genome and pan-transcriptome. *Plant Cell* 26: 121-135.

Hammer, G. L., G. D. Farquhar, and I. J. Broad, 1997 On the extent of genetic variation for
transpiration efficiency in sorghum. *Australian J. Agric. Res.* 48: 649-656.

Hansey, C. N., J. M. Johnson, R. S. Sekhon, S. M. Kaeppler, and N. de Leon, 2011 Genetic
diversity of a maize association population with restricted phenology. *Crop Sci.* 51: 704-715.

Henderson, S., S. Von Caemmerer, G. D. Farquhar, L. Wade, and G. Hammer, 1998 Correlation
between carbon isotope discrimination and transpiration efficiency in lines of the C₄ species
Sorghum bicolor in the glasshouse and the field. *Funct. Plant Biol.* 25: 111-123.

Holm, S., 1979 A simple sequentially rejective multiple test procedure. *Scandinavian J. Stat.* 6:
65-70.

Jaleel, C. A.I, P. Manivannan, A. Wahid, M. Farooq, H. J Al-Juburi *et al.* 2009 Drought stress in
plants: a review on morphological characteristics and pigments composition. *Int. J. Agric. Biol.*
11: 100-105.

578
579 Keeling, C. D., W. G. Mook, and P. P. Tans 1979 Recent trends in the $^{13}\text{C}/^{12}\text{C}$ ratio of
580 atmospheric carbon dioxide. *Nature* 277: 121-123.
581
582 Kolbe, A. R., T. P. Brutnell, A. B. Cousins, A. J. Studer, 2018a. Carbonic anhydrase mutants in
583 *Zea mays* have altered stomatal responses to environmental signals. *Plant Physiol.*, 177: 980-989.
584
585 Kolbe, A. R., A. J. Studer, and A. B. Cousins, 2018b Biochemical and transcriptomic analysis of
586 maize diversity to elucidate drivers of leaf carbon isotope composition. *Funct. Plant Biol.* 45:
587 489-500.
588
589 Leakey, A. D. B., J. N. Ferguson, C. P. Pignon, A. Wu, Z. Jin *et al.* 2019 Water use efficiency as
590 a constraint and target for improving the resilience and productivity of C_3 and C_4 crops. *Annu.*
591 *Rev. Plant Biol.* 70: 781-808.
592
593 Li, H., Thrash, A., Tang, J. D., He, L., Yan, J., and Warburton, M. L. 2019 Leveraging GWAS
594 data to identify metabolic pathways and networks involved in maize lipid biosynthesis. *Plant J.*,
595 98: 853-863.
596
597 Lipka, A. E., F. Tian, Q. Wang, J. Peiffer, M. Li *et al.* 2012 GAPIT: genome association and
598 prediction integrated tool. *Bioinformatics* 28: 2397-2399.
599
600 Lipka, A. E., C. B. Kandianis, M. E. Hudson, J. Yu, J. Drnevich *et al.* 2015 From association to
601 prediction: statistical methods for the dissection and selection of complex traits in plants. *Curr.*
602 *Opin. Plant Biol.* 24: 110-118.
603
604 Lu, Y., Y. Li, J. Zhang, Y. Xiao, Y. Yue *et al.* 2013 Overexpression of *Arabidopsis* molybdenum
605 cofactor sulfurase gene confers drought tolerance in maize (*Zea mays* L.). *PloS ONE* 8: e52126.
606
607 Marschner, H., and B. Dell, 1994 Nutrient uptake in mycorrhizal symbiosis. *Plant Soil* 159: 89-
608 102.

Masle, J., G. D. Farquhar, and S. C. Wong, 1992 Transpiration ratio and plant mineral content are related among genotypes of a range of species. *Funct. Plant Biol.* 19: 709-721.

Masle, J., Gilmore, S. R., and Farquhar, G. D., 2005. The ERECTA gene regulates plant transpiration efficiency in *Arabidopsis*. *Nature*, 436: 866.-870

McGrath, J. M., and Lobell, D. B. 2013 Reduction of transpiration and altered nutrient allocation contribute to nutrient decline of crops grown in elevated CO₂ concentrations. *Plant, Cell & Environment*, 36: 697-705.

McMullen, M. D., S. Kresovich, H. S. Villeda, P. Bradbury, H. Li *et al.* 2009 Genetic properties of the maize nested association mapping population. *Science* 325: 737-740.

Myers, S. S., Zanobetti, A., Kloog, I., Huybers, P., Leakey, A. D., *et al.* 2014 Increasing CO₂ threatens human nutrition. *Nature*, 510: 139-142.

National Academies of Sciences, Engineering, and Medicine. 2019 Science Breakthroughs to Advance Food and Agricultural Research by 2030. Washington, DC: Natl. Acad. Press.

O'Leary, M. H., 1988 Carbon isotopes in photosynthesis. *Bioscience* 38: 328-336.

Passioura, J. B., 1996 Drought and drought tolerance. *Plant Growth Reg.* 20: 79-83.

Pauli, D., G. Ziegler, M. Ren, M. A. Jenks, D. J. Hunsaker, *et al.* 2018 Multivariate analysis of the cotton seed ionome reveals a shared genetic architecture. *G3: Genes, Genomes, Genetics*, 8: 1147-1160.

Purcell, L. C., J. T. Edwards, and K. R. Brye, 2007 Soybean yield and biomass responses to cumulative transpiration: Questioning widely held beliefs. *Field Crops Res.* 101: 10-18.

640

641 R Core Team 2017 R: A language and environment for statistical computing. R Foundation for
642 Statistical Computing, Vienna, Austria. Available at: <https://www.R-project.org/>.

643

644 Rebetzke, G. J., A. G. Condon, G. D. Farquhar, R. Appels, and R. A. Richards, 2008
645 Quantitative trait loci for carbon isotope discrimination are repeatable across environments and
646 wheat mapping populations. *Theor. Appl. Genet.* 118: 123-137.

647

648 Revelle, W. R., 2017 psych: Procedures for personality and psychological research. Software,
649 <https://CRAN.R-project.org/package=psych>, version 1.8.4.

650

651 Sheffield, J., and E. F. Wood 2008 Projected changes in drought occurrence under future global
652 warming from multi-model, multi-scenario, IPCC AR4 simulations. *Climate Dynamics* 31: 79-
653 105.

654

655 Shiferaw, B., B. M. Prasanna, J. Hellin, and M. Bänziger, 2011 Crops that feed the world 6. Past
656 successes and future challenges to the role played by maize in global food security. *Food Sec.* 3:
657 307.

658

659 Tian, F., P. J. Bradbury, P. J. Brown, H. Hung, Q. Sun *et al.* 2011 Genome-wide association
660 study of leaf architecture in the maize nested association mapping population. *Nature Genet.* 43:
661 159.

662

663 Teulat, B., O. Merah, X. Sirault, C. Borries, R. Waugh *et al.* 2002 QTLs for grain carbon isotope
664 discrimination in field-grown barley. *Theor. Appl. Genet.* 106: 118-126.

665

666 This, D., J. Comstock, B. Courtois, Y. Xu, N. Ahmadi *et al.* 2010 Genetic analysis of water use
667 efficiency in rice (*Oryza sativa* L.) at the leaf level. *Rice* 3: 72.

668

669 Twohey III, R J., L. M. Roberts, and A. J. Studer 2019 Leaf stable carbon isotope composition
670 reflects transpiration efficiency in *Zea mays*. *Plant J.* 97: 475-484.

671
672 Virgona, J. M., K. T. Hubick, H. M. Rawson, G. D. Farquhar, and R. W. Downes, 1990
673 Genotypic variation in transpiration efficiency, carbon-isotope discrimination and carbon
674 allocation during early growth in sunflower. *Funct. Plant Biol.* 17: 207-214.
675
676 Walker, G. K., 1986 Transpiration efficiency of field-grown maize. *Field Crops Res.* 14: 29-38.
677
678 Wallace, J. G., S. J. Larsson, and E. S. Buckler, 2014 Entering the second century of maize
679 quantitative genetics. *Heredity* 112: 30.
680
681 Xu, Y., D. This, R. C. Pausch, W. M. Vonhof, J. R. Coburn *et al.* 2009 Leaf-level water use
682 efficiency determined by carbon isotope discrimination in rice seedlings: genetic variation
683 associated with population structure and QTL mapping. *Theor. Appl. Genet.* 118: 1065-1081.
684
685 Yu, J., and E. S. Buckler, 2006 Genetic association mapping and genome organization of maize.
686 *Curr. Opin. Biotechnol.* 17: 155-160.
687
688 Zhao, W., P. Canaran, R. Jurkuta, T. Fulton, J. Glaubitz *et al.* 2006 Panzea: a database and
689 resource for molecular and functional diversity in the maize genome. *Nucleic Acids Res.* 34:
690 D752-D757.

Figure 1:

Figure 1 $\delta^{13}\text{C}$ and SLA Single Family Stepwise Regression QTL Mapping. $\delta^{13}\text{C}$ QTL (A) were identified in NAM RIL families CML103 (black), CML333 (orange), and Tx303 (grey) but not in NC358 (blue). Specific leaf area (SLA) QTL (B) were identified in NAM RIL families CML103 (black) and Tx303 (grey). Significance thresholds (dashed horizontal line) were determined by 200 permutations and an alpha of 0.05.

Figure 2:

Figure 2 Pearson's r Correlations. Correlations of mean phenotypic values using complete observations and Holm's method to adjust p -values for multiple testing. The heat map shows no strong correlations between $\delta^{13}\text{C}$ mean values and element mean values. $\delta^{13}\text{C}$ and Mo have a significant p -value ($p = 6.745\text{E-}05$, $r = 0.18$).

Figure 3:

Figure 3 Element Single Family Stepwise Regression QTL Mapping. QTL mapping identified 28 QTL across 12 different elements. Significant QTL ($\alpha = 0.05$) for each element are plotted. QTL location is shown across the 10 maize chromosomes (cM) on the x-axis. Dashes indicate a significant QTL, with the NAM RIL family in which the QTL was found designated by color; CML103 (black), CML333 (orange), Tx303 (grey), NC358 (blue). All dashes are the same length for visibility.

Figure S1:

Supplemental Figure 1 NAM RIL Transgressive Segregation. NAM RIL families CML103 (A), CML333 (B), NC358 (C), and Tx303 (D) were sorted by $\delta^{13}\text{C}$ and plotted. Parental lines are shown in red.

Figure S2:

Supplemental Figure 2 Element and $\delta^{13}\text{C}$ Full Correlation Matrix. A full correlation matrix of element and $\delta^{13}\text{C}$ mean values is shown. The diagonal displays histograms of each dataset. Pearson's r is shown in the upper panel. Scatter plots and best fit line are shown in the lower panel.

Figure S3:

Supplemental Figure 3 Element QTL Mapping by Chromosome. Significant element QTL are shown by maize chromosomes 1 through 10 on the x-axis (in cM). Each NAM RIL family is represented by a symbol; CML103 ($\circ$), CML333 ($\times$), NC358 ($\square$), and Tx303 (Δ). Each element is designated by color. Significance thresholds (dashed horizontal line) were determined using 200 permutations, $\alpha=0.05$ for each QTL independently.

Figure S4:

Supplemental Figure 3 Element QTL Mapping by Chromosome. Significant element QTL are shown by maize chromosomes 1 through 10 on the x-axis (in cM). Each NAM RIL family is represented by a symbol; CML103 ($\circ$), CML333 ($\times$), NC358 ($\square$), and Tx303 (Δ). Each element is designated by color. Significance thresholds (dashed horizontal line) were determined using 200 permutations, $\alpha=0.05$ for each QTL independently.