Loss of *prdm1a* accelerates melanoma onset and progression

Ritsuko Iwanaga^{1,*}, Brittany T. Truong^{1,2,*}, Jessica Y. Hsu³, Karoline A. Lambert⁴, Rajesh Vyas⁷, David Orlicky⁵, Yiqun G. Shellman⁴, Aik-Choon Tan⁶, Craig Ceol⁷ and [Kristin Bruk Artinger](#)¹

¹Department of Craniofacial Biology, ²Human Medical Genetics & Genomics Graduate Program, ³Pharmacology Graduate Program, ⁴Department of Dermatology, ⁵Department of Pathology, ⁶Division of Medical Oncology, University of Colorado Denver Anschutz Medical Campus, Aurora, CO, USA

⁷Program in Molecular Medicine, University of Massachusetts Medical School, Worcester, MA, USA

*These two are joint first authors.

Correspondence:

Kristin Bruk Artinger, Department of Craniofacial Biology, University of Colorado Anschutz Medical Campus, 12801 E. 17th Ave., MS8120, Aurora, CO 80045.

Email: Kristin.arteringer@cuanschutz.edu

<https://orcid.org/0000-0002-3003-604>

Key words: melanoma, zebrafish, PRDM1, neural crest cells

Acknowledgements:

The State of Colorado, Tobacco Settlement Funds pilot grant

SDRC Gates Center Pilot Grant, University of Colorado P30AR057212-10

The National Institutes of Health P30CA046934, University of Colorado Cancer Center Support Grant

ABSTRACT

Melanoma is an aggressive, deadly skin cancer derived from melanocytes, a neural crest cell derivative. Melanoma cells mirror the developmental program of neural crest cells in that they exhibit the same gene expression patterns and utilize similar cellular mechanisms, including increased cell proliferation, EMT and migration. Here we studied the role of neural crest regulator PRDM1 in melanoma onset and progression. In development, *Prdm1a* functions to promote neural crest progenitor fate, and in melanoma, we found that *PRDM1* has reduced copy number and is recurrently deleted in both zebrafish and humans. When examining expression of neural crest and melanocyte development genes, we show that *sox10* progenitor expression is high in *prdm1a*^{-/-} mutants, while more differentiated melanocyte markers are reduced, suggesting that normally *Prdm1a* is required for differentiation. Data mining of human melanoma datasets indicate that high *PRDM1* expression in human melanoma is correlated with better patient survival and decreased *PRDM1* expression is common in metastatic tumors. When one copy of *prdm1a* is lost in the zebrafish melanoma model (Tg[mitfa:BRAF^{V600E}];p53^{-/-}; *prdm1a*^{+/-}), melanoma onset occurs more quickly, and the tumors that form have a larger area with increased expression of *sox10*. These data demonstrate a novel role for PRDM1 as a tumor suppressor in melanoma.

INTRODUCTION

Melanoma is an aggressive cancer that accounts for the majority of human skin cancer deaths. The 5-year survival rate of metastatic melanoma is only 23%¹. Early stage melanoma can be cured by surgical excision; however, only a few treatment options exist once the melanoma has metastasized. More than 50% of melanoma patients present with a somatic *BRAF* mutation², and up to 90% of these are from a single amino acid change from valine (V) to glutamic acid (E) at position 600 (*V600E*)³. As such, the high frequency of *BRAFV600E* mutations among melanoma patients has prompted the development of targeted therapies such as BRAF and MEK inhibitors that dampen BRAF/MEK pathway signaling⁴. Dabrafenib and trametinib were the first combined inhibitors to be approved by the Food and Drug Administration (FDA) for patients with advanced *BRAFV600E* melanoma^{5,6}. The combined inhibitors improve patient survival rates and have fewer adverse side effects compared to monotherapy treatments. About one third of melanoma patients experience disease recurrence or metastasis years after initial treatment^{7,8}. The acquirement of additional somatic mutations (i.e. point mutations, chromosomal rearrangements, copy number variations, etc.) that bypass the targets of BRAF and MEK inhibitors may account for some cases of incomplete response to the primary line of targeted therapy. However, other mechanisms of resistance or melanoma progression exist that have yet to be studied. For example, we previously showed that gene amplification of *SETDB1*, in conjunction with a *BRAFV600E* mutation, accelerates melanoma tumor formation in a zebrafish model by causing disruptions in transcriptional regulation, including a decrease in expression of *HOX* genes⁹. Identifying additional genes and pathways involved in melanoma initiation is critical, as it can lead to the development of future therapeutic and treatment options.

The cell origins of melanoma tumors are thought to arise from transformed melanocytes¹⁰, which are derived from neural crest cells, a multipotent population of cells that forms between the neural and non-neural ectoderm. During embryonic development, neural crest cells undergo an epithelial-to-mesenchymal transition and migrate to different regions of the developing embryo before giving rise to a variety of cell types, including peripheral nerves, chondrocytes, osteoblasts, and melanocytes. As neural crest cells are a highly migratory population of cells, it has been hypothesized that the same genes and pathways utilized by neural crest cells during development to aid in the trafficking of cells to their final destinations in the embryo can be later hijacked by cancer cells to enhance their proliferation and migration^{11,12}. Indeed, several publications have shown that there are distinct parallels in behaviors and gene expression patterns between melanoma cells and neural crest cells¹³. For instance, *Brn3a*, a neural crest transcription factor in the neuronal lineage, is frequently expressed in melanoma tumors and has been shown to accelerate tumor growth by promoting survival¹⁴. Interestingly, mouse melanoma cells can arise from a population of seemingly differentiated melanocytes that respond to environmental cues to undergo transcriptional reprogramming¹⁵. One cue is likely a neural crest growth factor, as melanoma cells express CD271 receptors on the cell surface¹⁶.

Apart from these studies that have implicated single neural crest-related genes in melanoma progression, there is strong evidence that a neural crest transcriptional program plays a causative role in melanoma progression. Genes such as *crestin*, *sox10*, *ednrb*, *tyr*, and *dct*, which are characteristic markers of neural crest progenitors and melanocytes, are highly enriched in zebrafish and mouse melanoma tumors^{17,18}. In zebrafish, *crestin* is normally downregulated at later stages in development after neural crest cells have fully differentiated¹⁹. Using a *crestin:EGFP* reporter in a zebrafish model, Kaufman et al. (2016) used live-cell imaging of

melanoma tumor formation and showed that a single melanocyte can initiate tumor growth by reactivating *crestin*²⁰. Moreover, *crestin:EGFP*-positive zebrafish scales, but not adjacent *crestin*-negative scales, exhibit gene signatures that were enriched with neural crest and melanoma genes, such as *mitfa* and *sox10*²⁰. This work suggests that reactivation of specific neural crest cell genes coincides with tumor initiation. It is evident that inappropriate activation of developmental genes can lead to cancer; however, it is not well-known whether expression of neural crest cell genes can also be protective.

Here, we identify neural crest regulator PRDM1 as a novel tumor suppressor in melanoma. *PRDM1* (zebrafish ortholog *prdm1a*) is a highly conserved transcription factor, whose structure consists of an N-terminal PR/SET domain, followed by a proline-rich domain, and five DNA-binding C₂H₂ zinc fingers. PRDM1 regulates transcription both by direct DNA binding²¹ and by forming complexes with chromatin-modifying proteins including histone methyltransferases Prmt5²² and G9a²³ and histone deacetylases HDAC1/2²⁴. *PRDM1* is a critical factor in neural crest cell development. It is highly expressed at the neural plate border in zebrafish^{25,26}, *Xenopus laevis*²⁷, and lamprey²⁸ and is required for Rohon-Beard sensory neuron and neural crest cell formation^{25,26}. Homozygous loss of *prdm1a* in developing zebrafish embryos results in a significant reduction of neural crest cell derivatives, including pigment cells and craniofacial cartilage²⁵, suggesting that it is required for neural crest differentiation. Similarly, PRDM1 has also been shown to be required for the differentiation of both adaxial cells into slow-twitch muscle cells²⁹, as well as retinal progenitors to photoreceptor cells^{30,31}. In addition to being important for cell fate specification during development, PRDM1 and other members of the PRDM family function as tumor suppressors for many cancers, including B, T, and natural killer cell lymphomas, colorectal cancer, and lung cancer³²⁻³⁵. In lymphoid

malignancies, genomic deletions, missense mutations, and premature truncations of *PRDM1* are frequently observed^{34,36,37}, causing inactivation of the protein and misregulation of lymphocyte differentiation and cell cycle progression. These studies provide strong evidence and support a role for PRDM1 as a tumor suppressor. In this study, we show that *Prdm1a* is required for melanocyte differentiation from neural crest cells during development and that decreased *PRDM1* expression is correlated with worse melanoma patient survival. We also report that loss of *PRDM1* in a zebrafish model leads to an acceleration of melanoma onset and progression.

MATERIALS & METHODS

Zebrafish

Zebrafish were maintained as previously described³⁸. The Institutional Animal Care and Use Committee (IACUC) at the University of Colorado Denver Anschutz Medical Campus (UCD-AMC) reviewed and approved all animal procedures. Wildtype strains include AB and EKK lines. Transgenic lines include *prdm1a*^{m805} (*nrd*)³⁹, referred to as *prdm1a*^{-/-}; *tp53*^{M214/M214}, referred to as *p53*^{-/-}; Tg(*mitfa*:BRAF^{V600E});*p53*^{-/-}^{17,40}, and Tg(*mitfa*:BRAF^{V600E});*p53*^{-/-}.

The standard length of the zebrafish was measured from the tip of the snout to the base of the tail (**Figure 2A**).

Microinjections were performed in *p53*^{-/-} and *prdm1a*^{+/-} zebrafish at the single-cell stage. The *mitfa*:BRAF^{V600E} vector was a gift from Richard M. White¹⁷. The vector was linearized with SalI and BssHI, purified, and resuspended in 100 mM KCl for a final concentration of 100 ng/mL. 500 pL were injected into each embryo.

Zebrafish were observed and scored weekly for hyperpigmentation, which indicated the presence of melanoma tumors. We compared tumor-free curves over time using a log-rank test

where $p < 0.05$ was statistically significant. The tumor area was measured using ImageJ and normalized to body length.

aCGH, JISTIC, and comparative analysis

Array comparative genomic hybridization (aCGH) was performed and analyzed as previously described⁴¹. Briefly, we custom designed an array with 398,426 unique probes covering 97% of the zebrafish genome (based on Zv9 assembly). Genomic DNA was extracted from zebrafish melanoma tumors (Tg[*mitfa*:BRAFF^{V600E}];p53-/-^{17,40}) and from a normal region of the same fish using the Qiagen DNeasy Blood and Tissue kit (Qiagen, Germantown, MD, #69504). The DNA was fragmented and hybridized to the array. Raw data was generated from scanned images with the Agilent Feature Extraction software (v10.7) and normalized using the Agilent Genomic workbench to identify copy number differences (Agilent, Santa Clara, CA). The JISTIC algorithm was used to calculate significantly altered regions, and peak calling was done using a q-value cut-off of 0.25. For human melanomas, copy number data was downloaded from Tumorscape^{42,43} and JISTIC analysis was conducted as described previously⁴¹. Genes from within peaks were pooled to define species-specific sets of recurrently deleted genes. Human orthologs of zebrafish genes were determined using Ensembl and BLAST.

Survival analysis of The Cancer Genome Atlas (TCGA) cutaneous melanoma data

Clinical and mutational data of 329 TCGA melanoma patients were obtained from the supplementary tables of the TCGA publication (<http://cancergenome.nih.gov/>). Normalized RNA-seq gene expression profiles (Level 3, RSEM value) for these patients were downloaded from the cBioPortal website (<http://www.cbioportal.org>). Patients were assigned to high (hi) and low (lo) groups based on above or below mean gene expression of *PRDM1*. In this study, we only focus on the 286 metastatic melanoma patients with outcome data. Kaplan-Meier survival

analysis was performed comparing hi and lo groups. A log-rank test of $p < 0.05$ was considered statistically significant.

Mining public gene expression data sets

Gene expression profiles of the Riker data set (GSE7553)⁴⁴ were downloaded from the Gene Expression Omnibus (GEO) (<http://www.ncbi.nlm.nih.gov/gds>). This data set contains 87 melanoma patients and was profiled by the Affymetrix Human Genome U133 Plus 2.0 microarray. Raw data was downloaded and normalized with the Robust Multiarray Average (RMA) method using the Affymetrix Power Tools. From this data set, there were 4 normal skin, 14 primary melanoma and 40 metastatic melanoma patients. Gene expression for *PRDMI* was compared across these patients, and a one-way ANOVA test was performed to determine the different expressions. Error bars represent mean $\pm$ SD.

Histology

Zebrafish samples were fixed in 4% paraformaldehyde and embedded in paraffin. The samples were then sectioned at a 5 μ M thickness and fixed to glass slides. H&E staining was performed by the Gates Center Regenerative Medicine Histology Core at UCD-AMC. Histology images were captured on an Olympus BX51 microscope equipped with a four megapixel Macrofire digital camera (Optronics, Goleta, CA) using the Picture Frame Application 2.3 (Optronics).

RNA extraction and real-time PCR (RT-qPCR)

Total RNA was isolated from dissected zebrafish tumors and zebrafish embryos (20 pooled embryos). The zebrafish samples were stored in TRIzol reagent (Invitrogen Corp., Carlsbad, CA, #15596018) and then purified with the Direct-zol RNA kit (Zymo Research, Irvine, CA). RNA (1 μ g) was converted to cDNA using the Verso cDNA Synthesis Kit (Thermo

Fisher Scientific, Waltham, MA). All procedures were conducted according to the manufacturers' suggestions.

Real-time PCR reactions (RT-qPCR) were performed in the Step One Plus Real-time PCR System (Thermo Fisher Scientific). Taqman primers for *PRDM1*, *prdm1a*, *sox10*, *dct*, *mitfa*, *mclr*, and *tyr* were purchased from Thermo Fisher Scientific. *rpl13a* was used as the zebrafish internal control. (*rpl13a*: probe 5'-/56-FAM/ATGAAGTTGGCTGGAAGTACCA/3IABlk_FQ/-3', forward 5'-CACACGCAAATTTGCCCTGCTTGG-3', and reverse 5'-GCCTGCTTAGTCAGCTTCATC-3'). Reactions were performed in three biological and technical replicates. Transcript abundance and relative fold change were quantified using the $2^{-\Delta\Delta Ct}$ method relative to control. Relative abundance was compared using an unpaired, independent t-test. Error bars represent the mean $\pm$ SD. Results were statistically significant when $p < 0.05$.

Western blotting

Western blotting was performed as previously described⁴⁵. 50 to 100 zebrafish embryos were pooled at each time point. Calcium-free Ginzberg Fish Ringer's solution (NaCl, KCl, NaHCO₃) was added to the embryos for deysolking. Samples were then washed in deysolking wash buffer (NaCl, Tris [pH 8.5], KCl, and CaCl₂) and lysed in lysis buffer using a syringe (0.1% glycerol, 0.01% SDS, 0.1M Tris-HCL [pH6.8]). Total protein concentration was determined with the Bio-Rad DC Protein assay (Bio-Rad, Hercules, CA). Proteins (20 μ g) were separated by SDS-PAGE (12%) and transferred to polyvinylidene difluoride membranes. Primary antibodies against PRDM1 (1:750 dilution, gift from Dr. Philip W. Ingham)⁴⁶ and total histone H3 (1:5000 dilution, Cell Signaling Technology, Danvers, MA, #9715) were used. The blots were incubated with secondary antibody, HRP-conjugated goat anti-rabbit (1:5000 dilution,

Sigma-Aldrich, St. Louis, MO, #A0545) for 2 hours. Chemiluminescent detection was performed with Luminata Classico Western HRP substrate (Millipore Sigma, Burlington, MA). Bio-Rad Chemidoc multiplex imaging system was used to detect and quantify the protein levels. Band intensity was quantified in ImageJ and normalized to the total histone H3 loading control. The experiment was performed at least twice.

Tyrosinase assay

The tyrosinase assay was performed as previously described with a few modifications^{47,48}. About 100 embryos at two days post fertilization (dpf) were dechorionated and sonicated in Pro-Prep protein extraction solution (iNtRON Biotechnology, Burlington, MA, #17081). 200 µg of total protein in 75 uL of lysis buffer was added to 75 µL of 1.0 mM of L-3,4-dihydroxyphenylalanine (L-DOPA) in a 96-well plate. The control well contained 75 µL of lysis buffer and 75 µL of 1.0 mM L-DOPA. After incubation at 37°C for 60 minutes, absorbance was read at 425 nm. The assay was performed twice with three technical replicates in each trial. Error bars represent the mean ± SD. The data were analyzed using a one-way ANOVA test followed by a multiple comparisons test where $p < 0.05$ was statistically significant.

RESULTS

PRDM1 is recurrently deleted in human and zebrafish melanoma and is required for melanocyte differentiation

Somatic gene mutations, including gene copy number variations, are frequently observed in cancer, including melanoma^{9,43,49,50}. We used a cross-species oncogenomics approach to identify recurrently deleted genes in both human and zebrafish melanomas. As previously described, we performed aCGH on Tg(*mitfa*:BRAF^{V600E});p53^{-/-} zebrafish to generate copy

number variant (CNV) profiles for zebrafish that develop melanomas (see below)^{17,40,41} (**Figure 1A**). Human data were obtained from Tumorscape, a dataset that contains copy-number analyses for over 3,000 human cancer specimens^{42,43}. The aCGH values and human data were analyzed using the JISTIC algorithm⁵¹, which then listed recurrently amplified and deleted genes. We identified 2,617 genes recurrently deleted in human melanomas and 1,515 genes in zebrafish melanomas. Of these recurrently deleted genes, 72 were deleted in both human and zebrafish melanomas (**Figure 1B**), and furthermore 34 of these genes were also identified as transcriptionally downregulated in a microarray dataset comparing expression in zebrafish melanoma tissue and normal melanocytes⁴¹. Among the 34 downregulated genes in zebrafish melanoma was the family of *prdm1* paralogs, *prdm1a-c* (**Figure 1C**). *prdm1a* was selected as a candidate gene because of its known role as a neural crest regulator²⁵. Paralogs *prdm1b* and *prdm1c* were also downregulated, but they were not recurrently deleted based on the JISTIC analysis (**Figure 1C**). Therefore, we did not explore these paralogs further.

PRDM1 is a known tumor suppressor in multiple cancers, including B and T-cell lymphoma^{36,52}, colorectal cancer⁵³, lung cancer⁵⁴, and glioma⁵⁵. Moreover, *Prdm1a* has been shown to play a role in neural crest differentiation in zebrafish^{25,39}. Homozygous loss of *prdm1a* results in reduced neural crest cell derivatives, including peripheral neurons and mature pigment cells (**Figure 2A-D**)³⁹. *prdm1a*^{-/-} mutants die at 6-7 dpf, and wildtype and heterozygotes are indistinguishable by phenotype (**Figure 2B-C'**). The standard length of the zebrafish from the tip of the snout to the base of the tail is not affected when one or both copies of *prdm1a* are mutated ($p>0.05$) (**Figure 2E-F**). *prdm1a* transcripts (**Figure 2G**) and protein levels (**Figure 2H-I**) are highest a 2 days post fertilization (dpf), when pigment cells are differentiating along body of the zebrafish⁵⁶. To better understand the role of *Prdm1a* in melanocyte development, we performed

RT-qPCR on whole wildtype and *prdm1a* ^{-/-} mutant embryos for a series of genes that mark early (*sox10*) and late melanocyte lineage (*dct*, *mclr*, *mitfa*, *tyr*)¹⁰ during a time course of development from 1-5 dpf (Schematic **Figure 3A**). When neural crest cells commit to the melanocyte lineage, they express different markers that indicate their stage of differentiation (**Figure 3A**). *sox10* is an early melanocyte stem cell marker and is required for melanocyte differentiation from neural crest cells in that it activates expression of *mitf*, *dct*, and *tyr*^{10,57}. Interestingly, in wildtype embryos, *sox10* expression is steady over time, but *prdm1a* ^{-/-} mutants exhibit a significant increase in expression at 2 and 4 dpf, corresponding to the first and second waves of melanocyte migration during zebrafish development⁵⁸ (p<0.0001) (**Figure 3B**). Expression of *dct*, *mitfa*, and *tyr*, markers of the melanoblast precursor state, decrease steadily over developmental time, while *mitfa* increases in wildtype embryos (**Figure 3C-F**). In contrast, the *prdm1a* ^{-/-} mutants consistently exhibit an overall decrease in the later melanocyte markers compared to wildtype embryos (p<0.05) (**Figure 3C-F**). The increase in *sox10*, along with the overall decrease in later melanocyte markers, suggests that the cells could be maintained as early melanocyte progenitors, unable to fully differentiate into mature, pigment producing melanocytes.

To further test whether loss of *prdm1a* affected proper melanogenesis, we tested for tyrosinase activity, the enzyme responsible for pigment production, in *prdm1a* ^{-/-} mutant, wildtype (EKK strain), and wildtype/heterozygote mixed embryos at 2 dpf. Pooled embryos were incubated with the tyrosinase substrate L-DOPA, and spectrophotometry was used to quantify pigment production. Compared to both the wildtype and wildtype/heterozygote mix, *prdm1a* ^{-/-} mutants had significantly decreased pigment production (p<0.01) and this loss of tyrosinase activity aligns with the levels of *prdm1a*, as lysates from a combination of wildtype and

prdm1a^{+/-} embryos exhibited an intermediate loss of tyrosine activity that was significantly different from that of both the wildtype and *prdm1a*^{-/-} lysates (**Figure 3G**). Together with our RT-qPCR results demonstrating that *prdm1a* mutants are deficient in the expression of late melanocyte markers, these data suggest that melanocyte progenitors require *Prdm1a* to terminally differentiate and produce pigment. Moreover, this suggests that even a partial loss of *prdm1a* affects melanogenesis, which is important to note given that our adult melanoma models are *prdm1a*^{+/-} heterozygotes. Together, the data indicate that *Prdm1a* is important for melanocyte development from neural crest cells. Given that there are key similarities between melanoma and neural crest cells and that *PRDM1* is already a known tumor suppressor, we determined whether it played a role in melanoma as well.

Low *PRDM1* expression is correlated with worse patient outcomes and metastatic melanoma in publicly available datasets.

To determine whether *PRDM1* expression was associated with human melanoma patient prognosis, we analyzed The Cancer Genome Atlas (TCGA) cutaneous melanoma data set and plotted the overall survival of patients with metastatic melanoma based on their expression of *PRDM1*. Lower *PRDM1* expression is correlated with a worse overall survival rate ($p < 0.001$) (**Figure 4A**). We also mined the Riker melanoma data set (GSE7553 Moffitt data) and observed a significant decrease in *PRDM1* expression in metastatic melanoma tumors compared to normal skin ($p < 0.001$) and primary melanoma tumors ($p < 0.05$) (**Figure 4B**).

Heterozygous loss of *prdm1a* in zebrafish increases melanoma onset and progression

To determine whether loss of *PRDM1* promotes melanoma *in vivo*, we used two zebrafish models, an injected and transgenic *mitfa*:BRAF^{V600E} line. In both models, the BRAF mutation is driven by a *mitfa* promoter to drive melanoma in melanocytes specifically, with an

additional *p53*^{-/-} mutant background that further accelerates melanoma onset^{17,40}. For the injected model, the oncogenic *mitfa*:BRAF^{V600E} construct was introduced into *p53*^{-/-} and *p53*^{-/-}; *prdm1a*^{+/-} zebrafish embryos at the single-cell stage^{17,40}. Again, homozygous *prdm1a*^{-/-} mutants die at 6-7 dpf. The construct is integrated randomly into the genome and is expressed transiently. We monitored the zebrafish and noted when hyperpigmentation occurred in concentrated areas of the body. This hyperpigmentation was indicative of the initiation of melanoma tumor formation. Due to transient expression of the *mitfa*:BRAF^{V600E} mutation in the injected model, tumor formation was slower than previously reported^{17,20,40}. The *p53*^{-/-} zebrafish formed tumors 12 months post-injection (n=12), while the *p53*^{-/-}; *prdm1a*^{+/-} zebrafish developed tumors significantly more quickly at 10 months post-injection (n=8) (p<0.05) (**Figure 4A**). Unfortunately, due to the *p53*^{-/-} mutant background, the zebrafish often developed malignant peripheral nerve sheath tumors (MPNST)⁵⁹ at about 10 months post-injection before melanoma formation. Therefore, these fish were excluded from our melanoma study.

To circumvent this issue, we established two stable transgenic lines, Tg(*mitfa*:BRAF^{V600E})*p53*^{-/-} and Tg(*mitfa*:BRAF^{V600E})*p53*^{-/-}; *prdm1a*^{+/-}, and monitored tumor formation. As expected, these tumors formed more quickly than in the injected model. By 17 weeks, about 50% of the zebrafish for both genotypes had formed tumors (**Figure 5B**). With one copy of *prdm1a* mutated, the Tg(*mitfa*:BRAF^{V600E})*p53*^{-/-}; *prdm1a*^{+/-} zebrafish (n=33) quickly surpassed their wildtype Tg(*mitfa*:BRAF^{V600E})*p53*^{-/-} siblings (n=18) and developed melanoma tumors (p<0.05) (**Figure 5B**). By week 19, more than 60% of the Tg(*mitfa*:BRAF^{V600E})*p53*^{-/-}; *prdm1a*^{+/-} zebrafish had visible tumors. In contrast, approximately 50% of the wildtype siblings were still tumor free, suggesting that loss of *prdm1a* accelerates tumor onset. Representative images of melanoma tumors are shown at 7 and 11 months for both genotypes (**Figure 5C**), and

tumor area was measured and normalized to the body length (**Figure 5D**). The tumor area of *Tg(mitfa:BRAF^{V600E});p53^{-/-};prdm1a^{+/-}* (n=7) was significantly larger than wildtype *Tg(mitfa:BRAF^{V600E});p53^{-/-}* tumors (n=5) (p<0.05) (**Figure 5C-D**). However, this may be because tumor onset occurred earlier in *Tg(mitfa:BRAF^{V600E});p53^{-/-};prdm1a^{+/-}* zebrafish (**Figure 5B**). Similar to the injected model, several transgenic fish developed MPNST tumors due to the *p53^{-/-}* mutant background⁵⁹. Therefore, the melanoma tumors on these fish were not large enough to measure and were not included in the quantification.

As previously described, Sox10 is a conserved marker of neural crest cells and potentially co-opted in melanoma progenitor cells in many vertebrate models^{20,60}, thus we compared its expression between dissected *Tg(mitfa:BRAF^{V600E});p53^{-/-};prdm1a^{+/-}* and *Tg(mitfa:BRAF^{V600E});p53^{-/-}* tumors. For each tumor, *sox10* expression was compared to normal, adjacent muscle tissue of the same individual. *sox10* expression varied but was elevated for both genotypes. On average, tumor samples from *Tg(mitfa:BRAF^{V600E});p53^{-/-};prdm1a^{+/-}* zebrafish had a threefold increase in *sox10* compared to their *Tg(mitfa:BRAF^{V600E});p53^{-/-}* siblings (p<0.01) (**Figure 5E**). This suggests that reduction of *prdm1a* expression causes an upregulation of *sox10* in both neural crest development and melanoma cells^{21,61}. In melanoma, we predict that the gain in *sox10* expression supports the ability of melanoma to grow and spread.

To examine the zebrafish tumors in more detail, we paired zebrafish with similarly sized tumors and then sectioned the samples for H&E staining. In both tumors, we observed invasion into the neighboring muscle tissue with variable levels of melanin across the tumors (n=3 for each genotype; **Figure 5F**). The melanoma in *Tg(mitfa:BRAF^{V600E});p53^{-/-};prdm1a^{+/-}* zebrafish have an overall larger tumor area (**Figure 5D**), with melanin observed in the neighboring tissue

(**Figure 5F**) whereas the Tg(*mitfa*:BRAF^{V600E});*p53*^{-/-} tumor remains somewhat more localized at the skin surface with less invasion into the surrounding skeletal muscle.

DISCUSSION

Melanoma cells often acquire features of neural crest cells during tumor formation by reactivating genes and pathways utilized by neural crest cells^{12,13}. The work presented in this study suggest that *Prdm1a*, a known neural crest regulator, is required for melanocyte differentiation from neural crest cells and that loss of *Prdm1a* accelerates melanoma onset and progression. During development, we have shown that *prdm1a*^{-/-} mutants have a reduction in neural crest derivatives, including mature pigment cells^{25,26}. Here, we demonstrate that loss of *prdm1a* leads to an increase in expression of *sox10*, an early melanocyte stem cell marker, and an overall decrease in later melanocyte markers. *prdm1a*^{-/-} mutants also have significantly reduced pigment production by tyrosinase. Together, these data suggest that *Prdm1a* is required for the differentiation of neural crest cells to mature melanocytes. It is likely that in the absence of *Prdm1a*, the cells are maintained as early melanocyte progenitors and thus function in supporting melanoma formation and progression. The case for *PRDM1* functioning as a tumor suppressor in melanoma is further supported by the fact that *PRDM1/prdm1a-c* copy number is significantly reduced in both human and zebrafish melanomas, and its expression is reduced in melanoma cells compared to normal melanocytes. Moreover, in human patients, low *PRDM1* expression is correlated with worse human patient survival. Finally, we demonstrate that heterozygous loss of *prdm1a* promotes melanoma tumorigenesis in a zebrafish model and results in an increase in *sox10* expression in the tumors. Thus, our data demonstrates a novel role for the neural crest regulator *PRDM1* as a melanoma tumor suppressor.

Melanoma is known to have increased somatic mutation rates compared to other cancers due to mutagenesis by UV exposure^{49,50}, with *BRAF*^{V600E} being the most common mutation³. Other mutations, such as *NRAS*, are common in large congenital nevi, which are often melanogenic and can be precursors to melanoma⁶². Interestingly when *NRAS*^{Q61K} is driven by a tyrosinase promoter, transgenic mice develop congenital naevi and melanoma and express the neural crest progenitor marker Sox10¹⁸, again iterating that Sox10 expression is required for malignant melanoma formation. Reducing one copy of Sox10 in this background reduces the number of congenital naevi and melanoma tumors by regulating cell proliferation¹⁸. This suggests that Sox10's function in development, to maintain neural crest stem cells, may be dysregulated during disease progression.

In our previous work and this current body of work, we have demonstrated that the developmental regulator, *Prdm1a*, is required for neural crest differentiation to mature melanocytes, which is consistent with its function in other cell types during development²⁹⁻³¹. We show that *prdm1a* may promote differentiation through regulation of *sox10*⁶¹, as suggested by the increase in its expression in *prdm1a*^{-/-} mutants. However, because the *prdm1a*^{-/-} mutants have visibly less pigmentation compared to wildtype embryos, this may have affected our interpretation of the melanocyte marker gene expression data. Moreover, just from their reduced pigmentation and reduced expression of late melanocyte markers, we cannot conclusively claim that *prdm1a*^{-/-} larvae in fact have less melanoblasts or melanocytes, as *prdm1a* loss may disrupt neural crest cells' ability to differentiate into these cell types or impair the proliferation of these neural crest derivatives. In future studies, we can address these issues by performing live-imaging experiments and/or lineage tracing experiments to parse out the exact effects of *prdm1a* loss on neural crest derivatives. However, it is interesting that reducing *prdm1a* in heterozygous

embryos also reduces tyrosinase activity, even when there is not a discernable difference in pigmentation. Therefore, our results suggest that *Prdm1a* is an important regulator of neural crest cell differentiation into melanocytes. Our data contribute to the growing literature regarding the parallels between neural crest cell development and melanoma tumor formation¹³.

Interestingly, *PRDM1* is known to act as a tumor suppressor in other cancers, particularly B, T, and natural killer cell lymphomas and lung cancer^{34,36,37,54}. We have now shown in a zebrafish model that heterozygous loss of *prdm1a* can accelerate melanoma onset, and it is likely acting as a tumor suppressor. We hypothesize that loss of *PRDM1* coordinates with mutations in *BRAF* and/or *TP53* to promote tumorigenesis. *PRDM1* deletions have been reported to be associated with loss of *TP53* in anaplastic large cell lymphoma patients³⁷, and this association likely contributed to the pathogenicity of the cancer. Our zebrafish models carried mutations in both *BRAF* and *TP53*. Additional mutations in *PRDM1* led to accelerated melanoma tumor onset and suggested a trend towards metastasis (data not shown), similar to what has been shown with *PRDM1* and lung cancer⁵⁴. Whether *PRDM1* coordinates with these or additional tumor suppressor/oncogenic genes during tumorigenesis must be explored further.

In conclusion, this study demonstrates a novel role for *PRDM1* as a tumor suppressor in *p53* mutated melanoma. We showed that even partial loss of *prdm1a* in zebrafish accelerates melanoma onset, and this can potentially be sourced to the impairment of melanocyte differentiation from neural crest cells during embryonic development observed upon *prdm1a* loss. Broadly, these results compound on accumulating evidence that neural crest cells and melanomas operate on overlapping gene regulatory networks and suggest that factors that regulate neural crest development are a source of therapeutic value for melanomas.

Author Contributions: R.I. and B.T.T. contributed equally to this work, performed, analyzed, and interpreted data, and performed the statistical analysis on the zebrafish; A-C.T. performed bioinformatic analysis; D.O. performed histological analysis and imaging. K.B.A. conceptualized the project and obtained funding, collaborating with Y.S.; B.T.T. and K.B.A. wrote the manuscript with the help of Y.S.; and R.I., J.H., C.C. and A-C.T. edited the manuscript.

Funding: This work is supported by pilot grants from the Skin Disease Research Center at the University of Colorado Anschutz Medical Campus P30AR057212-10, The State of Colorado, Tobacco Settlement funds pilot grants, and P30CA046934 to the University of Colorado Cancer Center.

Acknowledgments: We thank the Artinger and Shellman labs for experimental feedback; Morgan Singleton and UCD-AMC CLAC for zebrafish care; Virginia Ware and Aaron Clark for technical help with experiments; and Richard White for sharing microarray data, fish lines and advice.

Conflicts of Interest: We have no conflicts to declare.

FIGURES

Figure 1: PRDM1 is recurrently deleted in human and zebrafish melanoma. (A) Circos plot displaying gene copy number gains and losses in zebrafish melanomas as previously described⁴¹. JISTIC G-scores are displayed as pale red shading (amplifications [minimum = 0; maximum = 1,550]) and blue shading (deletions [minimum = 0; maximum = 2,150]). $-\log_{10}$ -transformed JISTIC Q-values with a cutoff of 0.6 (corresponding to an untransformed Q-value of 0.25) are shown as bold red lines (amplifications [minimum = 0; maximum = 11]) and bold blue (deletions [minimum = 0; maximum = 11]). Dotted circles represent the $-\log_{10}$ -transformed Q-value of 0 (center) and 11 (outer: amplification; inner: deletion). (B) Venn diagram of orthologous genes significantly deleted in human and zebrafish melanomas from 10,380 human-zebrafish gene pairs (hypergeometric test, p-value: 2.0E-16). (C) Genes downregulated in zebrafish melanomas as compared with melanocytes (analyzed from previously obtained microarray data set⁴¹). *prdm1a* (red dot) and its paralogs *prdm1b* and *prdm1c* (orange dots) are indicated.

Figure 2: *prdm1a* (A) Schematic showing how the standard length of the embryo was measured. Lateral images of wildtype (B), *prdm1a*^{+/-} (C) and *prdm1a*^{-/-} (D) at 2 dpf. Ventral images of wildtype (B'), *prdm1a*^{+/-} (C') and *prdm1a*^{-/-} (D') at 4 dpf. Quantification of standard length at 2 (E) and 4 (F) dpf. Statistical analysis was performed with a one-way ANOVA. (G) RT-qPCR was performed on 20 pooled embryos at each time point (n=3 replicate experiments) for *prdm1a*. Proteins were resolved by SDS-PAGE and western blotting for *prdm1a* and total H3 (H) was performed. (I) Quantification of band intensity from the western blot (G). Relative intensity was normalized to total histone H3 for each sample. Abbreviations: dpf, days post fertilization

Figure 3: *prdm1a* is required for melanocyte differentiation and pigment production. (A)

Melanocyte lineage from neural crest to mature melanocytes and their characteristic markers.

Whole wildtype embryos were pooled for RNA isolation and protein extraction over developmental time. (B-F) Whole wildtype and *prdm1a*^{-/-} mutant embryos were pooled for RNA isolation over developmental time. 20 whole embryos were pooled for each genotype and at each time point (n=3 replicate experiments). RT-qPCR was performed for *sox10* (B), *dct* (C), *mitfa* (D), *mc1r* (E), and *tyr* (F) over developmental time. Expression was compared using an unpaired, independent t-test. (J) 100 embryos were pooled and incubated with L-DOPA for a tyrosinase functional assay. Pigment production was measured by spectrometry in whole wildtype, wildtype/heterozygote mix, and *prdm1a*^{-/-} mutant embryos. Statistical analysis was performed with a one-way ANOVA followed by a multiple comparisons test. Error bars represent mean $\pm$ SD. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Abbreviations: dpf, days post fertilization

Figure 4: Low *PRDM1* expression in humans is correlated with worse patient outcomes and metastatic melanoma. (A)

Overall survival of patients with metastatic melanoma compared to *PRDM1* expression obtained from The Cancer Genome Atlas (TCGA). Statistical analysis was performed with a log-rank test. (B) *PRDM1* expression in normal skin, primary melanoma tumors and metastatic melanoma tumors obtained from the Riker melanoma data set. Statistical analysis was performed using a one-way ANOVA. Error bars represent mean $\pm$ SD. *p<0.05, **p<0.01, ***p<0.001

Figure 5: Loss of *prdm1a* in zebrafish accelerates tumor onset. (A) Tumor free curve from p53^{-/-} (n=12) and p53^{-/-}; *prdm1a*^{+/-} (n=8) zebrafish injected with BRAF^{V600E} construct to induce melanoma. (B) Tumor free curve from Tg(*mitfa*:BRAF^{V600E}); p53^{-/-} (n=18) and Tg(*mitfa*:BRAF^{V600E}); p53^{-/-}; *prdm1a*^{+/-} (n=33) zebrafish. Statistical analysis for tumor free curves was performed with a log-rank test. (C) Representative images of melanoma tumors from Tg(*mitfa*:BRAF^{V600E}); p53^{-/-} and Tg(*mitfa*:BRAF^{V600E}); p53^{-/-}; *prdm1a*^{+/-} zebrafish at 7 and 11 months. (D) Tumor area was measured in five Tg(*mitfa*:BRAF^{V600E}); p53^{-/-} tumors and seven Tg(*mitfa*:BRAF^{V600E}); p53^{-/-}; *prdm1a*^{+/-} zebrafish tumors. The area was then divided by the body length to normalize. Relative tumor area was compared using an unpaired, Mann-Whitney U test. Error bars represent median and range. (E) RNA was isolated from dissected tumors and RT-qPCR was performed for *sox10*. Expression was compared using an unpaired, Mann-Whitney U test. Error bars represent mean ± SD. *p<0.05. (F) H&E staining of size-matched tumors for Tg(*mitfa*:BRAF^{V600E}); p53^{-/-}; *prdm1a*^{+/-} (7 months) and Tg(*mitfa*:BRAF^{V600E}); p53^{-/-} (9 months) zebrafish compared to normal skin. Upper panels show 40X magnification. The lower panels are zoomed in to 200X magnification.

REFERENCES

1. *Cancer Facts & Figures*. American Cancer Society, 2019.
2. Ascierto, P., Kirkwood, J., Grob, J.-J., et al., *The role of BRAF V600 mutation in melanoma*. Journal of Translational Medicine, 2012. **10**(85): p. 1-9.
3. Bradish, J.R. and Cheng, L., *Molecular pathology of malignant melanoma: changing the clinical practice paradigm toward a personalized approach*. Hum Pathol, 2014. **45**(7): p. 1315-26.
4. Mackiewicz, J. and Mackiewicz, A., *BRAF and MEK inhibitors in the era of immunotherapy in melanoma patients*. Contemp Oncol (Pozn), 2018. **22**(1A): p. 68-72.
5. Robert, C., Karaszewska, B., Schachter, J., et al., *Improved overall survival in melanoma with combined dabrafenib and trametinib*. N Engl J Med, 2015. **372**(1): p. 30-9.
6. Eroglu, Z. and Ribas, A., *Combination therapy with BRAF and MEK inhibitors for melanoma: latest evidence and place in therapy*. Ther Adv Med Oncol, 2016. **8**(1): p. 48-56.
7. Tas, F. and Erturk, K., *Relapse patterns in patients with local and regional cutaneous melanoma*. Clin Transl Oncol, 2019. **21**(4): p. 412-419.
8. Soong, S., Harrison, R., McCarthy, W., et al., *Factors affecting survival following local, regional, or distant recurrence from localized melanoma*. J Surg Oncol, 1998. **67**(4): p. 228-233.
9. Ceol, C.J., Houvras, Y., Jane-Valbuena, J., et al., *The histone methyltransferase SETDB1 is recurrently amplified in melanoma and accelerates its onset*. Nature, 2011. **471**(7339): p. 513-7.
10. White, R.M. and Zon, L.I., *Melanocytes in development, regeneration, and cancer*. Cell Stem Cell, 2008. **3**(3): p. 242-52.
11. Powell, D.R., Blasky, A.J., Britt, S.G., et al., *Riding the crest of the wave: parallels between the neural crest and cancer in epithelial-to-mesenchymal transition and migration*. Wiley Interdiscip Rev Syst Biol Med, 2013. **5**(4): p. 511-22.
12. Maguire, L.H., Thomas, A.R., and Goldstein, A.M., *Tumors of the neural crest: Common themes in development and cancer*. Developmental Dynamics, 2015. **244**: p. 311-322.
13. Shakhova, O., *Neural crest stem cells in melanoma development*. Curr Opin Oncol, 2014. **26**(2): p. 215-21.
14. Hohenauer, T., Berking, C., Schmidt, A., et al., *The neural crest transcription factor Brn3a is expressed in melanoma and required for cell cycle progression and survival*. EMBO Mol Med, 2013. **5**(6): p. 919-34.
15. Kohler, C., Nittner, D., Rambow, F., et al., *Mouse Cutaneous Melanoma Induced by Mutant BRaf Arises from Expansion and Dedifferentiation of Mature Pigmented Melanocytes*. Cell Stem Cell, 2017. **21**(5): p. 679-693 e6.
16. Boiko, A.D., Razorenova, O.V., van de Rijn, M., et al., *Human melanoma-initiating cells express neural crest nerve growth factor receptor CD271*. Nature, 2010. **466**(7302): p. 133-7.
17. White, R.M., Cech, J., Ratanasirintraooot, S., et al., *DHODH modulates transcriptional elongation in the neural crest and melanoma*. Nature, 2011. **471**(7339): p. 518-22.
18. Shakhova, O., Zingg, D., Schaefer, S.M., et al., *Sox10 promotes the formation and maintenance of giant congenital naevi and melanoma*. Nat Cell Biol, 2012. **14**(8): p. 882-90.

19. Luo, R., An, M., Arduini, B.L., et al., *Specific pan-neural crest expression of zebrafish Crestin throughout embryonic development*. Dev Dyn, 2001. **220**(2): p. 169-74.
20. Kaufman, C.K., Mosimann, C., Fan, Z.P., et al., *A zebrafish melanoma model reveals emergence of neural crest identity during melanoma initiation*. Science, 2016. **351**(6272): p. aad2197.
21. Powell, D.R., Hernandez-Lagunas, L., LaMonica, K., et al., *Prdm1a directly activates foxd3 and tfap2a during zebrafish neural crest specification*. Development, 2013. **140**(16): p. 3445-55.
22. Ancelin, K., Lange, U.C., Hajkova, P., et al., *Blimp1 associates with Prmt5 and directs histone arginine methylation in mouse germ cells*. Nat Cell Biol, 2006. **8**(6): p. 623-30.
23. Gyory, I., Wu, J., Fejer, G., et al., *PRDI-BF1 recruits the histone H3 methyltransferase G9a in transcriptional silencing*. Nat Immunol, 2004. **5**(3): p. 299-308.
24. Yu, J., Angelin-Duclos, C., Greenwood, J., et al., *Transcriptional Repression by Blimp-1 (PRDI-BF1) Involves Recruitment of Histone Deacetylase*. Molecular and Cellular Biology, 2000. **20**(7): p. 2592-2603.
25. Hernandez-Lagunas, L., Choi, I.F., Kaji, T., et al., *Zebrafish narrowminded disrupts the transcription factor prdm1 and is required for neural crest and sensory neuron specification*. Dev Biol, 2005. **278**(2): p. 347-57.
26. Roy, S. and Ng, T., *Blimp-1 specifies neural crest and sensory neuron progenitors in the zebrafish embryo*. Curr Biol, 2004. **14**: p. 1772-1777.
27. Rossi, C.C., Hernandez-Lagunas, L., Zhang, C., et al., *Rohon-Beard sensory neurons are induced by BMP4 expressing non-neural ectoderm in Xenopus laevis*. Dev Biol, 2008. **314**(2): p. 351-61.
28. Nikitina, N., Tong, L., and Bronner, M.E., *Ancestral network module regulating prdm1 expression in the lamprey neural plate border*. Dev Dyn, 2011. **240**(10): p. 2265-71.
29. Roy, S., Wolff, C., and Ingham, P.W., *The u-boot mutation identifies a Hedgehog-regulated myogenic switch for fiber-type diversification in the zebrafish embryo*. Genes & Development, 2001. **15**: p. 1563-1576.
30. Brzezinski, J.A., Lamba, D.A., and Reh, T.A., *Blimp1 controls photoreceptor versus bipolar cell fate choice during retinal development*. Development, 2010. **137**(4): p. 619-29.
31. Katoh, K., Omori, Y., Onishi, A., et al., *Blimp1 suppresses Chx10 expression in differentiating retinal photoreceptor precursors to ensure proper photoreceptor development*. J Neurosci, 2010. **30**(19): p. 6515-26.
32. Mzoughi, S., Tan, Y.X., Low, D., et al., *The role of PRDMs in cancer: one family, two sides*. Curr Opin Genet Dev, 2016. **36**: p. 83-91.
33. Hayashi, K., Yoshida, K., and Matsui, Y., *A histone H3 methyltransferase controls epigenetic events required for meiotic prophase*. Nature, 2005. **438**(7066): p. 374-8.
34. Kucuk, C., Iqbal, J., Hu, X., et al., *PRDM1 is a tumor suppressor gene in natural killer cell malignancies*. Proc Natl Acad Sci U S A, 2011. **108**(50): p. 20119-24.
35. Boi, M., Zucca, E., Inghirami, G., et al., *PRDM1/BLIMP1: a tumor suppressor gene in B and T cell lymphomas*. Leuk Lymphoma, 2015. **56**(5): p. 1223-8.
36. Mandelbaum, J., Bhagat, G., Tang, H., et al., *BLIMP1 is a tumor suppressor gene frequently disrupted in activated B cell-like diffuse large B cell lymphoma*. Cancer Cell, 2010. **18**(6): p. 568-79.

37. Boi, M., Rinaldi, A., Kwee, I., et al., *PRDM1/BLIMP1 is commonly inactivated in anaplastic large T-cell lymphoma*. Blood, 2013. **122**(15): p. 2683-93.
38. Westerfield, M., *The Zebrafish Book; A Guide for the Laboratory Use of Zebrafish (Brachydanio rerio)*. 5th Edition. University of Oregon Press, 2007.
39. Artinger, K.B., Chitnis, A.B., Mercola, M., et al., *Zebrafish narrowminded suggests a genetic link between formation of neural crest and primary sensory neurons*. Development, 1999. **126**(18): p. 3969-79.
40. Patton, E.E., Widlund, H.R., Kutok, J.L., et al., *BRAF mutations are sufficient to promote nevi formation and cooperate with p53 in the genesis of melanoma*. Curr Biol, 2005. **15**(3): p. 249-54.
41. Venkatesan, A.M., Vyas, R., Gramann, A.K., et al., *Ligand-activated BMP signaling inhibits cell differentiation and death to promote melanoma*. J Clin Invest, 2018. **128**(1): p. 294-308.
42. Lin, W.M., Baker, A.C., Beroukhi, R., et al., *Modeling genomic diversity and tumor dependency in malignant melanoma*. Cancer Res, 2008. **68**(3): p. 664-73.
43. Beroukhi, R., Mermel, C.H., Porter, D., et al., *The landscape of somatic copy-number alteration across human cancers*. Nature, 2010. **463**(7283): p. 899-905.
44. Riker, A.I., Enkemann, S.A., Fodstad, O., et al., *The gene expression profiles of primary and metastatic melanoma yields a transition point of tumor progression and metastasis*. BMC Med Genomics, 2008. **1**(13): p. 1-16.
45. Shull, L., Sen, R., Menzel, J., et al., *The conserved and divergent role of Prdm3 and Prdm16 in zebrafish and mouse craniofacial development*. bioRxiv, 2019.
46. von Hofsten, J., Elworthy, S., Gilchrist, M.J., et al., *Prdm1- and Sox6-mediated transcriptional repression specifies muscle fibre type in the zebrafish embryo*. EMBO Rep, 2008. **9**(7): p. 683-9.
47. Choi, T.Y., Kim, J.H., Ko, D.H., et al., *Zebrafish as a new model for phenotype-based screening of melanogenic regulatory compounds*. Pigment Cell Res, 2007. **20**(2): p. 120-7.
48. Cha, S.H., Ko, S.C., Kim, D., et al., *Screening of marine algae for potential tyrosinase inhibitor: those inhibitors reduced tyrosinase activity and melanin synthesis in zebrafish*. J Dermatol, 2011. **38**(4): p. 354-63.
49. Hodis, E., Watson, I.R., Kryukov, G.V., et al., *A landscape of driver mutations in melanoma*. Cell, 2012. **150**(2): p. 251-63.
50. Pleasance, E.D., Cheetham, R.K., Stephens, P.J., et al., *A comprehensive catalogue of somatic mutations from a human cancer genome*. Nature, 2010. **463**(7278): p. 191-6.
51. Sanchez-Garcia, F., Akavia, U.D., Mozes, E., et al., *JISTIC: identification of significant targets in cancer*. BMC Bioinformatics, 2010. **11**(189).
52. Calado, D.P., Zhang, B., Srinivasan, L., et al., *Constitutive canonical NF-kappaB activation cooperates with disruption of BLIMP1 in the pathogenesis of activated B cell-like diffuse large cell lymphoma*. Cancer Cell, 2010. **18**(6): p. 580-9.
53. Liu, C., Banister, C.E., Weige, C.C., et al., *PRDM1 silences stem cell-related genes and inhibits proliferation of human colon tumor organoids*. Proc Natl Acad Sci U S A, 2018. **115**(22): p. E5066-E5075.
54. Zhu, Z., Wang, H., Wei, Y., et al., *Downregulation of PRDM1 promotes cellular invasion and lung cancer metastasis*. Tumour Biol, 2017. **39**(4): p. 1010428317695929.

55. Wang, X., Wang, K., Han, L., et al., *PRDM1 is directly targeted by miR-30a-5p and modulates the Wnt/beta-catenin pathway in a Dkk1-dependent manner during glioma growth*. Cancer Lett, 2013. **331**(2): p. 211-9.
56. Kimmel, C.B., Ballard, W.W., Kimmel, S.R., et al., *Stages of Embryonic Development of the Zebrafish*. Developmental Dynamics, 1995. **203**: p. 253-310.
57. Harris, M.L., Baxter, L.L., Loftus, S.K., et al., *Sox proteins in melanocyte development and melanoma*. Pigment Cell Melanoma Res, 2010. **23**(4): p. 496-513.
58. Milos, N. and Dingle, A.D., *Dynamics of pigment pattern formation in the zebrafish, Brachydanio rerio. I. Establishment and regulation of the lateral line melanophore stripe during the first eight days of development*. Journal of Experimental Zoology, 1978. **205**(2): p. 205-216.
59. Berghmans, S., Murphey, R.D., Wienholds, E., et al., *tp53 mutant zebrafish develop malignant peripheral nerve sheath tumors*. Proc Natl Acad Sci U S A, 2005. **102**(2): p. 407-12.
60. Mohamed, A., Gonzalez, R.S., Lawson, D., et al., *SOX10 Expression in malignant melanoma, carcinoma, and normal tissues*. Appl Immunohistochem Mol Morphol, 2013. **21**(6): p. 506-510.
61. Olesnick, E., Hernandez-Lagunas, L., and Artinger, K.B., *prdm1a Regulates sox10 and islet1 in the development of neural crest and Rohon-Beard sensory neurons*. Genesis, 2010. **48**(11): p. 656-66.
62. Bauer, J., Curtin, J.A., Pinkel, D., et al., *Congenital melanocytic nevi frequently harbor NRAS mutations but no BRAF mutations*. J Invest Dermatol, 2007. **127**(1): p. 179-82.

URLs:

<http://www.cbioportal.org/>
<http://cancergenome.nih.gov/>
<http://www.ncbi.nlm.nih.gov/gds/>

A

E

F

G

H

I

Melanocyte lineage

B

C

D

E

F

G