Novel genes required for surface-associated motility in

Acinetobacter baumannii

Ulrike Blaschke ^{a,b,*}, Evelyn Skiebe ^a and Gottfried Wilharm ^{a,c,*}

^a Robert Koch-Institute, Project group P2, Burgstr. 37, D-38855 Wernigerode, Germany

* Corresponding addresses: ulrikeblaschke@googlemail.com; WilharmG@rki.de

^b ORCID: 0000-0002-3496-5447

^c ORCID: 0000-0002-1771-6799

Keywords

Acinetobacter baumannii, surface-associated motility, transposon mutant library, pellicle biofilm, antibiotic resistance, *Galleria mellonella*

Funding

This project was funded by the Deutsche Forschungsgemeinschaft (DFG) within FOR 2251 (WI 3272/3-1 and WI 3272/3-2).

Abstract

Acinetobacter baumannii is an opportunistic and increasingly multi-drug resistant human pathogen rated as a critical priority 1 pathogen for the development of new antibiotics by the WHO in 2017. Despite the lack of flagella, *A. baumannii* can move along wet surfaces in 2 different ways: via twitching motility and surface-associated motility. While twitching motility is known to depend on type IV pili, the mechanism of surface-associated motility is poorly understood. In this study we established a library of 30 *A. baumannii* ATCC 17978 mutants that displayed deficiency in surface-associated motility. By making use of natural competence we also introduced these mutations into strain 29D2 to differentiate strain-specific versus species-specific effects of mutations. Mutated genes were associated with purine/pyrimidine/folate biosynthesis (e.g. *purH*, *purF*, *purM*, *purE*), alarmone/stress metabolism (e.g. Ap4A hydrolase), RNA modification/regulation (e.g. methionyl-tRNA synthetase), outer membrane proteins (e.g. *ompA*), and genes involved in natural competence (*comEC*). All tested mutants originally identified as motility-deficient in strain ATCC 17978 also displayed a motility-deficient phenotype in 29D2. By contrast, further comparative characterization of the mutant sets of both strains regarding pellicle biofilm formation, antibiotic resistance, and virulence in the *Galleria mellonella* infection model revealed numerous strain-specific mutant phenotypes. Our studies highlight the need for comparative analyses to characterize gene functions in *A. baumannii* and for further studies on the mechanisms underlying surface-associated motility.

Introduction

Acinetobacter baumannii is a Gram-negative and strictly aerobic coccobacillus [1,2]. Being an opportunistic human pathogen [3], *A. baumannii* is associated with nosocomial diseases including soft tissue, bloodstream, and urinary tract infections as well as pneumonia [2]. Worldwide, about 9% of culture-positive infections found in intensive care units arise from *Acinetobacter spp.* [4]. Increased multi-drug resistance in *A. baumannii* has become problematic in recent years [5,6]. A global surveillance study found that 44% of 18,741 collected isolates were multi-drug resistant. During the study period the proportion of multi-drug resistant *A. baumannii* isolates increased from 23% in 2004 to 63% in 2014 [7]. As a consequence of rising multi-drug resistance, *A. baumannii* was rated as one of the critical priority 1 pathogens for the development of new antibiotics by the WHO in 2017 [8]. Drug resistance and environmental persistence have enabled *A. baumannii* to successfully establish in the hospital environment. Some clinical isolates can survive 100 days or more under dry conditions [9-13]. An important factor for the interaction of *A. baumannii* with biotic or abiotic surfaces is the formation of biofilms, a feature that is associated with an increased tolerance to desiccation stress [14].

A connection between *A. baumannii* virulence and motility has been shown in the *Caenorhabditis elegans* infection model where hypermotility resulted in increased virulence [15]. Although *A. baumannii* does not produce flagella, it is capable of moving in two different ways: via twitching motility and surface-associated motility. For *A. baumannii*, twitching motility has been shown to depend on type IV pili (T4P) [16,17] which drive the bacteria via retraction of attached T4P [18-25]. Inactivation of the putative T4P retraction ATPase *pilT* reduces twitching motility [26-28,11] but does not abolish surface-associated motility [26,16]. Surface-associated motility in *A.*

baumannii occurs at the surface of semi-dry media and is independent of T4P [26,29]. Surface-associated motility is poorly understood mechanistically, but was demonstrated to be controlled by quorum sensing [26], light [30], and iron availability [31,32]. Also, the synthesis of 1,3-diaminopropane (DAP) [33] and lipopolysaccharide (LPS) production [32] were shown to contribute to surface-associated motility of *A. baumannii*. Several genes have been identified which contribute to *A. baumannii*'s capacity for surface-associated motility [26,32], including a ribonuclease T2 family protein [34] and the superoxide dismutase SodB [35]. A recent study revealed the regulatory control of surface-associated motility and biofilm formation by a cyclic-di-GMP signaling network in *A. baumannii* strain ATCC 17978 [36]. Interestingly, studies on phase-variable phenotypes in *A. baumannii* strain AB5075 showed that “opaque phase” bacterial colonies had improved surface-associated motility [37,38]. A correlation between pellicle biofilm formation and surface-associated motility has been described in *A. baumannii* [39]. Given the fact that many *A. baumannii* clinical isolates exhibit surface-associated motility, it could be an important trait associated with infection [33,28,26].

To investigate the mechanisms underlying surface-associated motility, we utilized a previously generated transposon mutant library of ATCC 17978 [33] which we screened for a surface-associated motility-deficient phenotype. The motility-deficient mutations were found to affect purine/pyrimidine/folate biosynthesis, alarmone/stress metabolism, RNA modification/regulation, outer membrane proteins, and DNA modification. We characterized these mutants with respect to growth, pellicle biofilm formation, antibiotic resistance, and virulence in the *Galleria mellonella* infection model. To facilitate distinguishing between strain-specific and species-specific traits

some mutations were also introduced into the naturally competent *A. baumannii* strain 29D2 [40].

Materials and Methods

Bacterial strains and culture conditions

A. baumannii strain ATCC 17978 was purchased from LGC Promochem. The *A. baumannii* strain 29D2 was isolated from a white stork [40] and is naturally competent [41]. All strains were grown at 37°C in Luria-Bertani (LB) broth or on LB agar, and mutants were supplemented with 50 µg/mL of kanamycin. All strains used in this work are listed in supplementary Table S1. Single colonies were used as inoculum for overnight cultures or motility plates. Neither strain ATCC 17978 nor strain 29D2 exhibited phase variation [38,37,42].

Bacterial transformation and generation of an *A. baumannii* mutant library

ATCC 17978 transposon mutants were generated using the EZ-Tn5™ <KAN-2> Insertion Kit (Epicentre Biotechnologies) as previously described [33]. Transformation of the transposome complex into ATCC 17978 was performed by electroporation [43]. 29D2 mutants were generated by making use of the strain's ability for natural competence. The transforming DNA was isolated from the ATCC 17978 mutants described above. A suspension of DNA-accepting bacteria was generated by resuspending a few colonies in 100 µL of sterile PBS. The bacterial suspension was then mixed with equal volumes of the transforming DNA (~400 ng/µL). This mixture was stabbed into motility agar plates 10 times, pipetting 2 µL of the mixture with each

stabbing [16]. The motility plates were incubated for 18 h at 37°C. After incubation, the bacteria were flushed off the motility plates with 1 mL of sterile PBS and 100 µL was plated on selective agar plates (50 g/mL of kanamycin). After sub-culturing of selected colonies transformation was confirmed by PCR.

Identification of transposon insertion sites by single-primer PCR

To identify the transposon insertion sites of ATCC 17978 motility mutants, single-primer PCR was performed as described previously [33] using one of the following primers: FP-2Kana 5'-CTTCCCGACAACGCAGACCG-3'; FP-3Kana 5'-GAGTTGAAGGATCAGATCACGC-3'; RP-2Kana 5'-CCCTTGTATTACTGTTTATGTAAGC-3'; RP-3Kana 5'-CGCGGCCTCGAGCAAGACG-3'; Tn5-Kana-For4 5'-GTTTTCTCCTTCATTACAGAAACG-3'; and Tn5-Kana-Rev4 5'-CCCATACAATCGATAGATTGTCTG-3'. Transposon insertions of all mutants (ATCC 17978 and 29D2) were confirmed by PCR using primers for the EZ-Tn5™ <KAN-2> kanamycin cassette (Suppl. Fig. S1), which are specified in the manufacturer's instructions, and appropriate gene target site primers (Suppl. Table S2; Suppl. Figs. S2 and S3).

Surface-associated motility

Motility assays were performed as described previously [33]. A single bacterial colony from a nutrient agar plate (Oxoid) or selective agar plates (supplemented with 50 µg/mL of kanamycin for the mutants) of either wildtype (ATCC 17978 and 29D2) or mutants was lifted with a pipette tip and transferred to the surface of a motility plate

(0.5% agarose). Plates were incubated for 16 h at 37°C. The diameter of the surface motility spreading zone was measured and quadruplicates were statistically analyzed.

Bacterial growth curves

Growth curves were determined by growing overnight cultures at 37°C in LB medium (supplemented with 50 µg/mL of kanamycin for the mutants). Overnight cultures were adjusted to 1 OD (600 nm) in LB medium. In 250 mL baffled flasks, 50 mL of LB medium (without antibiotics) was inoculated with 1 mL of the OD-adjusted inoculum. The cultures were incubated at 37°C for 9 h with shaking at 160 rpm. OD measurements at 600 nm were performed every hour by sampling 100 µL of every culture. For each strain, data obtained from 3 independent cultures grown on the same day were averaged and represented by the mean ± SD.

Infection in the *Galleria mellonella* caterpillar

For *G. mellonella* caterpillar infection, bacteria were grown in LB medium overnight at 37°C (50 µg/mL of kanamycin was added to mutant strains). Cultures were diluted 1:50 in LB medium and incubated for another 4 h at 37°C. Bacteria were pelleted for 5 min at 7500 rpm at room temperature (RT) and the supernatant was discarded. Bacteria were resuspended in 500 µL sterile PBS, adjusted to an OD₆₀₀ nm of 1.0 and diluted 1:10 in sterile PBS. 5 µL of this dilution, corresponding to 3 x 10⁵ colony-forming units (CFUs), was injected into the last right proleg of *G. mellonella* caterpillars (purchased from TZ-TERRARISTIK, Germany, and BioSystems Technology TruLarv, UK). As a control, caterpillars were injected with 5 µL of sterile

PBS. Three independent experiments were performed with groups of 16 caterpillars for every bacterial strain and control. The caterpillars were incubated at 37°C for 5 days and checked daily for vitality. Experiments with more than 2 dead caterpillars within 5 days in the control group were not considered valid. CFUs were determined by serial dilutions, plated on nutrient agar, and colonies were counted after incubation at 37°C for 18 h. For each strain, data obtained from 3 independent experiments were averaged and represented by the mean $\pm$ SD.

Determination of susceptibility to antibiotics

For the minimal inhibitory concentration (MIC) tests, bacteria were grown in LB medium overnight at 37°C (to mutant strains 50 μ g/mL of kanamycin was added). Cultures were diluted 1:50 in LB medium and incubated (without antibiotics) another 4 h at 37°C. Agar plates were flushed with 2 mL of each culture and E-test strips (Liofilchem, Italy) were deposited on nutrient agar plates. MICs were determined after incubation for 16 h at 37°C. Three independent experiments were performed and statistical significance was tested by the Student's *t* test (2-tailed, unpaired).

Pellicle biofilm assays

A. baumannii strains were grown in LB medium overnight at 37°C (50 μ g/mL of kanamycin was added to mutant strains). The cultures were adjusted to an OD_{600 nm} of 1.0 and 3 mL of LB medium (without antibiotics) was inoculated with 15 μ L of OD-adjusted culture. Samples were incubated at RT for at least 3 days. The LB medium was removed using a thin cannula and the biofilm (sticking to the tube wall) was stained with a 0.5% crystal violet solution (w/v in Aqua Bidest) for 20 min. The crystal

violet was removed and the biofilm was washed twice with 4 mL Aqua Bidest. The biofilm was scrubbed and flushed off the tube walls with a pipet tip and 96% alcohol solution. The absorption at 550 nm was determined. Samples which showed an OD > 1.0 were diluted 1:10 with 96% ethanol for measurement. For each strain 3 independent experiments were performed and statistical significance was analyzed by the Student's *t* test (2-tailed, unpaired).

Microscopy

The bacterial strains ATCC 17978, ATCC 17978 *ompA::Km*, 29D2, and 29D2 *ompA::Km* were grown for 16 h at 37°C under constant shaking. One µL of each bacterial overnight culture was pipetted on a glass slide and analyzed under the bright field microscope (200 times magnification).

Statistical analysis

All experiments were performed at least 3 times. Comparison between groups was performed using GraphPad Prism 7 with Student's *t* test (2-tailed, unpaired). P-values less than 0.05 were considered to be statistically significant.

Results

Surface-associated motility

Approximately 2,000 transposon mutants of ATCC 17978 were screened for surface-associated motility phenotypes and 30 were identified with motility defects. Previous studies were limited to the characterization of mutations in single strains. Here, to

provide a comparative study, we introduced at least one mutation of every gene function category into 29D2 to get insight into strain-specific and species-specific traits.

To this end, surface-associated motility was analyzed on 0.5% agarose plates. The diameter ($\emptyset$) of the surface motility spreading zone of 3 independent experiments was measured and analyzed (Fig. 1A and Suppl. Table S3). All selected motility-deficient mutants of ATCC 17978 exhibited at least a 7-fold reduction of the spreading zone. Subsequently, DNA isolated from these transposon mutants was used to generate mutants in 29D2. All 29D2 mutants displayed a motility-deficient phenotype compared to the wildtype strain (Fig. 1B). Note that the surface-associated motility spreading zone of the wildtype ATCC 17978 (mean $\emptyset$ of 78 mm) was more than twice as large as that of the 29D2 wildtype strain (mean $\emptyset$ of 30 mm). Most ATCC 17978 mutants showed a 16-fold reduced surface-associated motility compared to the wildtype strain (Fig 2A), whereas the *a1s_0806* (encoding an aminotransferase) mutant lacked almost any measurable surface-associated motility (mean $\emptyset$ of 1 mm). 3 mutants, *purH::Km* (mean $\emptyset$ 10.25 mm), *1970::Km* (mean $\emptyset$ 8.75 mm), and *3297::Km* (mean $\emptyset$ 11 mm), showed 10-fold reduced surface-associated motility. Most 29D2 mutants displayed a 4-fold reduction in their surface-associated motility. The most pronounced reduction in motility appeared in mutants *purH::Km* (mean $\emptyset$ 5 mm), *purF::Km* (mean $\emptyset$ 3.75 mm), and *ddc::Km* (mean $\emptyset$ 4.25 mm). The mutant *purM::Km* (mean $\emptyset$ of about 16 mm) had the lowest reduction in surface-associated motility.

To summarize, all mutations initially identified in ATCC 17978 that conferred motility defects were also found to cause motility-deficient phenotypes when introduced into the orthologous genes of 29D2.

Pellicle biofilm formation

The formation of pellicles, a specific form of biofilm, occurs at the air-liquid interface and is distinct from submerged biofilms [39,44,45]. A correlation between surface-associated motility and pellicle biofilm formation has been described for *A. baumannii* [39]. We examined the ability of our motility-deficient mutants to form pellicles. Pellicle biofilms were incubated 3 days, stained with a 0.5% crystal violet solution, and analyzed by OD measurements (Suppl. Table S4). Pellicle-biofilm formation in wildtype ATCC 17978 was measured to be about 8.6 at OD₅₅₀ nm (Fig. 2A). A broad spread between low and high pellicle-producing mutants was visible, ranging between a 1.8-fold increase to more than a 25-fold decrease. For 15 of 30 mutants less than 67% of the wildtype-specific pellicle biomass was quantified (Table 1 and Fig. 2A). In the mutants *carB::Km*, *0414::Km*, and *prpF::Km* a pellicle biomass less than 8% compared to the wildtype biomass was measured. This dramatic decrease was not observed by inactivation of the orthologous gene in the 29D2 background. In ATCC 17978, 8 mutants (*purH::Km*, *purM::Km*, *purE::Km*, *0530::Km*, *3297::Km*, *galE::Km*, *0806::Km*, and *1055::Km*) were able to produce more pellicle biomass compared to the wildtype strain, of which the mutants *0530::Km*, *galE::Km*, and *0806::Km* produced 50-80% more pellicle biomass compared to wildtype (Fig. 2A). 29D2 mutants only displayed small changes in pellicle biofilm formation compared to wildtype, with a range of the mutants' pellicle biomass production from a 1.3-fold increase to a 2.1-fold decrease. 13 of 21 tested 29D2 mutants did not display any significant change in their pellicle biofilm formation compared to the parental strain (Table 2). Deficiencies could be observed in the following 6 mutant strains: *carB::Km*, *1970::Km*, *ddc::Km*, *dat::Km*, *gidA::Km*, and *3026::Km*, which produced less than

70% of the 29D2 wildtype-specific pellicle biomass. Only the following 2 mutants produced significantly more pellicle biomass (Fig. 2B) compared to the wildtype strain: *purH::Km* (30%) and *galE::Km* (38%).

In summary, the ATCC 17978 parental strain produced more pellicle biofilms compared to 29D2. Conspicuous changes in biofilm formation could mainly be observed among ATCC 17978 mutants. Concordance of pellicle formation phenotypes between the mutants of both strains was limited suggesting that strain-specific traits that are independent of surface-associated motility influence pellicle biomass production.

Bacterial growth

The ability of motility-deficient mutants to grow as a planktonic culture under aeration was assayed. Growth curves and data for all tested strains are provided in supplementary Fig. S4 (ATCC 17978 mutants), supplementary Fig. S5 (29D2 mutants), and Table S5. For 17978, 22 of 30 tested mutant strains exhibited significant growth defects compared to the parental strain (Table 1). The most striking growth defects (Fig. 3A) were observed in the mutants defective in purine biosynthesis (*purH::Km*, *purF::Km*, *purM::Km*, and *purE::Km*), pyrimidine biosynthesis (*carB::Km*), and diaminopropane biosynthesis (*ddc::Km* and *dat::Km*). Only 8 of 30 tested mutant strains were able to grow without any defect compared to the parental strain (Table 1). By testing the 29D2 mutant strains we observed 13 of 21 strains with notable planktonic growth defects (Table 2). Within this group most striking defects were observed with mutations associated with purine biosynthesis (*purH::Km*, *purF::Km*, *purM::Km*, and *purE::Km*), pyrimidine biosynthesis (*carB::Km*), folate

biosynthesis (*1566::Km*), and diaminopropane biosynthesis (*ddc::Km* and *dat::Km*). Additionally, *galE::Km*, *comEC::Km*, and *prpF::Km* mutants displayed strong growth deficiencies (Fig. 3B). The mutant *ompA::Km* showed growth comparable to the parental strain for up to 4 h, reached a growth maximum of 2.5 ± 0.28 OD_{600 nm} after 5 h, but then slowly collapsed to 1.36 ± 0.73 after 9 h. No growth defects were observed in 8 of 21 tested mutants (Table 2).

In summary, we found that genes involved in purine/pyrimidine and diaminopropane biosynthesis, oxidative stress, and propionate catabolism were crucial for growth of ATCC 17978 and 29D2 in LB medium.

***G. mellonella* caterpillar infection**

To gain insight into a possible correlation between motility and virulence we made use of the *G. mellonella* infection model. Caterpillars were infected with 3×10^5 CFU of different *A. baumannii* strains and the death of larvae was monitored over a time period of 5 days. *G. mellonella* infection of ATCC 17978 wildtype and mutant strains is shown in supplementary Fig. S6 and levels of significance for 5 days post-infection are presented in Table 1. A detailed listing of p-values for every monitored timepoint is provided in supplementary Table S6. After 24 h post-infection with the 17978 wildtype strain about 60% of larvae were dead. This number increased to over 80% of dead larvae after 5 days post-infection. 15 of 30 tested mutant strains displayed a significant attenuation in *G. mellonella* infection (Table 1). Another 4 mutant strains (*purE::Km*, *1624::Km*, *rpmG::Km*, and *ddc::Km*) showed some attenuation but this was not significant. The remaining 11 mutant strains did not display attenuation (Suppl. Fig. S6 and Table 1). Most pronounced attenuation was observed in strains

carB::Km, *metG::Km*, *ompA::Km*, and *galE::Km* (Fig. 4A). These results suggest an important role for these genes in *A. baumannii* virulence. However, to exclude the possibility that attenuation could be due to decreased planktonic growth, we compared the caterpillar infection results to our bacterial growth data (Fig. 3, Suppl. Fig. S4, and Fig. S5). Among the above mentioned mutants, only the *galE::Km* mutant was not significantly affected in growth. Overall, we found that for 11 of 15 significantly attenuated mutant strains the caterpillar infection data could possibly be influenced by decreased growth rates (Table 1 and Fig. 3).

The *G. mellonella* infection with 29D2 wildtype and mutant strains data is shown in supplementary Fig. S7 and significance levels (for 5 days post-infection) are shown in Table 2. A detailed listing of p-values for every monitored timepoint is provided in supplementary Table S7. 11 of 21 29D2 mutants were significantly attenuated in the *G. mellonella* infection model (Suppl. Fig S7 and Table 2). Within this group the most pronounced attenuation was observed in strains *carB::Km*, *ompA::Km*, *galE::Km*, and *comEC::Km* (Fig. 4B). The mutant strains *purE::Km*, *gidA::Km*, and *0806::Km* showed some attenuation at 5 days post-infection but p-values ranged between 0.060 – 0.067 (Suppl. Table S7). Interestingly, 8 of 11 significantly attenuated mutant strains (*purH::Km*, *1566::Km*, *carB::Km*, *ompA::Km*, *ddc::Km*, *comEC::Km*, *galE::Km*, and *2761::Km*) manifested a growth deficiency compared to the parental strain (Suppl. Fig. S5 and Table 2).

In summary, concordant infection traits were observed for 12 mutants of both strains including mutants affected in purine/pyrimidine/folate biosynthesis. Among these 12 strains, most significant attenuation was observed for *carB::Km*, *ompA::Km*, and *galE::Km*.

As a control, the CFUs were determined from the OD-adjusted bacterial cultures used for the infection experiments. To this end, OD-adjusted cultures were serially diluted and plated on nutrient agar. Colonies were counted after incubation for 18 h at 37°C. Interestingly, for both ATCC 17978 *ompA::Km* and 29D2 *ompA::Km* mutants we observed 1-2 log scale lower CFU numbers compared to the OD-adjusted suspension (data not shown). Sticking of cells to the tube wall during the dilution process was minimized by using low-binding tubes (Eppendorf). When growing both *ompA::Km* mutants on agar plates we observed a very sticky colony texture upon touching with a glass rod. Based on these findings we examined the cell morphology of *ompA::Km* mutant strains under the microscope. A distinct cell elongation or chain formation of both *ompA::Km* mutant strains compared to their parental strains was observed (Supplementary Fig. S8).

MIC Determination

We aimed to elucidate the correlation between motility-deficient mutants and their sensitivity to the bactericidal antibiotics ampicillin and imipenem as well as to the bacteriostatic antibiotic tetracycline. For ATCC 17978, 18 of 30 mutants displayed a significant resistance to ampicillin compared to the parental strain. The highest MIC values were obtained in mutant strains *0414::Km* (4-fold increase compared to the parental strain), *3026::Km* (4-fold increase), and *1566::Km* (3.7-fold increase). The only mutant strain which showed decreased resistance (0.7-fold decrease) to ampicillin was *aamA::Km* (Table 3). By contrast, a significantly increased sensitivity to imipenem was observed in six (*purE::Km*, *carB::Km*, *1624::Km*, *rpmG::Km*, *1970::Km*, and *aamA::Km*) of the tested strains (Table 1). Furthermore, a significantly

increased resistance to imipenem was observed in 4 of the tested mutant strains (*ompA::Km*, *3297::Km*, *0806::Km*, and *3026::Km*). For tetracycline, we found 13 of 30 mutants to be significantly more sensitive compared to the parental strain. Only 2 of 30 mutant strains, *purF::Km* and *galE::Km*, displayed significantly increased resistance to tetracycline (Table 1). Next, we analyzed all 29D2 mutant strains with respect to the MIC values for ampicillin, imipenem, and tetracycline. A significantly increased sensitivity to ampicillin was observed in 15 of 21 tested mutant strains compared to wildtype (Table 2). The only strain with significantly increased resistance to ampicillin was *3026::Km*, with a 1.7-fold increased MIC value (Table 3). Another mutant strain with a 1.5-fold increased ampicillin MIC value, although not significant, was *aamA::Km*. Similar effects were observed for imipenem (Table 3). Here, strains *3026::Km* and *aamA::Km* displayed significant resistance compared to the parental strain (Table 2). Increased sensitivity to imipenem was observed in 8 of 21 tested mutants. For the MIC values of tetracycline, we found the 3 mutant strains *1566::Km* (3.6-fold decrease), *comEC::Km* (5.4-fold decrease), and *prpF::Km* (6.5-fold decrease) to be significantly more susceptible compared to the parental strain (Table 3). Only one mutant, *dat::Km*, was significantly more resistant to tetracycline with a 1.8-fold increase (Table 3).

In conclusion, mutants from the 29D2 background predominantly showed increased sensitivity to all tested antibiotics. By contrast, many mutants of ATCC 17978 showed increased resistance to ampicillin, but increased sensitivity to imipenem and tetracycline.

Discussion

Recently, *A. baumannii* was demonstrated to exhibit motility on semi-dry plates, with agar concentrations between 0.2-0.4%, and motility was dependent on the type of agar that was used. Bacterial surface spreading was shown not to depend on type IV pili [26]. Here, we characterized 30 genes involved in *A. baumannii* surface-associated motility with respect to bacterial growth, pellicle biofilm formation, virulence, and antibiotic resistance. We discuss motility-deficient mutants with regards to their known/putative gene function in the bacterial cell (Fig. 5).

Genes involved in purine/pyrimidine/folate biosynthesis

In our study we identified 4 proteins involved in purine (*pur*) biosynthesis to be essential for *A. baumannii* surface-associated motility: PurH, PurF, PurM, and PurE (Tables 1 and 2). The decrease in surface motility of these mutants ranged from a 25-fold decrease in ATCC 17978 to a 2.1-fold decrease in 29D2 compared to their parental strains (Suppl. Table S3). In *A. nosocomialis* strain M2, EZ::Tn insertion in gene *purK* (*a1s_2963*) has been previously described to result in a 70% reduction in surface motility compared to the parental strain [26]. Mutations in the genes *purD*, *purF*, *purH*, *purL*, and *purM* abolished K⁺-dependent colony spreading in *Bacillus subtilis* [46]. The *pur* genes were also demonstrated to be essential for biofilm formation in *Bacillus cereus* (*purH* and *purD* [47]; *purA* [48]; *purA*, *purC*, and *purL* [49]). Interestingly, our study revealed no defective role of *pur* genes in pellicle biofilm formation. In contrast, mutations *purH::Km*, *purM::Km*, and *purE::Km* in 17978 and *purH::Km* in 29D2 produced significantly more pellicle biomass than their parental strains (Tables 1 and 2). A pellicle proteome study in 17978 showed that the *pur*

proteins were differentially expressed/accumulated under planktonic (PurH, PurF, and PurA), 1-day pellicle (PurM and PurB), and 4-day pellicle (PurD) growth conditions [44].

In addition to the motility deficiency, we found all tested *pur* mutants to display bacterial growth defects in LB media (Suppl. Figs. S4 and S5). For various bacteria, *pur* genes were identified to be required for bacterial growth in human serum including *Enterococcus faecium* (*purL*, *purH*, and *purD*) [50], *E. coli* (*purA*, *C*, *D*, *E*, *F*, *H*, *K*, *L*, and *M*), *S. enterica* (*purA*, *B*, *C*, *D*, *E*, *F*, *G*, and *H*), and *B. anthracis* (*purE* and *purK*) [51]. Due to the fact that all *pur* mutants displayed bacterial growth defects we expected these mutants to be attenuated in the *G. mellonella* infection, but we only found the 2 mutants ATCC 17978 *purM::Km* and 29D2 *purH::Km* had significant attenuation (Fig. 4). Purine biosynthesis mutants (*purF*, *purD*, and *purL*) in *Burkholderia cenocepacia* were also found to be attenuated in the *G. mellonella* infection model as well as in *C. elegans* and *D. melanogaster* infections [52]. *De novo* purine biosynthesis has also been shown to be required for virulence in ATCC 17978 (*purF*, *purD*, *purN*, *purL*, *purM*, *purK*, *purE*, *purC*, *purP*, and *purO*) in the mouse lung [53], and in several other bacteria such as *Streptococcus pneumoniae* (*purE*, *purK*, *purC*, and *purL*) [54], *Bacillus anthracis* (*purH*) [55], and *Pasteurella multocida* (*purF*) [56].

The *pur* mutants were tested for antibiotic sensitivity/resistance against ampicillin, imipenem, and tetracycline. For ampicillin MIC's we obtained very contrary results. The *pur* mutants of 29D2 showed increased sensitivity to ampicillin, but, except for *purH::Km*, the *pur* mutants of ATCC 17978 were significantly more resistant compared to their parental strain (Table 3). In *S. aureus* defective rifampicin persistence was shown for *purB*, *purF*, *purH*, and *purM* [57].

429 The *A. baumannii* gene *a1s_2687* encodes the large subunit (*carB*) of
 430 carbamoylphosphate synthase and is arranged with the small subunit (*carA*) in the
 431 *carAB* operon, which is required for the *de novo* synthesis of arginine and pyrimidines
 432 (reviewed in [58]), and in turn pyrimidines are known to be involved in biofilm
 433 formation in *P. aeruginosa* [59] and *E. coli* [60]. In *A. baumannii*, inactivation of *carB*
 434 caused a significantly decreased persistence in a mouse pneumonia model [53]. The
 435 contribution of *carB* to *A. baumannii* virulence was confirmed by our results, showing
 436 significant attenuation in both ATCC 17978 *carB::Km* and 29D2 *carB::Km* mutant
 437 strains (Fig. 4). Additionally, in a *P. aeruginosa* competition study against *B. cepacia*,
 438 *K. pneumoniae*, and *S. aureus*, the *carB* gene and hence uracil/pyrimidine
 439 biosynthesis was identified to be essential [61]. Inactivation of *carB* in ATCC 17978
 440 and 29D2 resulted in the greatest phenotypic alterations in planktonic growth, pellicle
 441 biofilm formation, and *G. mellonella* caterpillar infection of all tested mutants (Tables
 442 1 and 2). Interestingly, similar observations were also made for the
 443 gammaproteobacterium *Xanthomonas citri* subsp. *citri*. In that study, the knockout of
 444 *carB* resulted in a 70% decrease in biofilm formation, an extensive reduction in
 445 swimming motility, and alterations in bacterial growth [62]. CarB was also found to be
 446 required for growth of *E. coli* in human serum [51]. A motility-deficient phenotype was
 447 identified for the gene *a1s_1566* (putative 6-pyruvoyl-tetrahydropterin synthase),
 448 involved in folate biosynthesis and thus crucial for biosynthesis of purines and
 449 deoxythymidine monophosphate (Fig. 1). Here we observed an involvement in
 450 virulence, bacterial growth, and pellicle biofilm formation (Tables 1 and 2). Taken
 451 together, these findings suggest that purine and pyrimidine genes contribute to
 452 important bacterial processes like motility, bacterial growth, pellicle biofilm formation,

and virulence not only in *Acinetobacter* but also in well studied genera like *Bacillus* and *Salmonella*.

Genes involved in alarmone/damage metabolism

The *A. baumannii* genes *a1s_0414* and *a1s_1624* encode for an Ap5A pyrophosphatase and an Ap4A hydrolase (ApaH-like), respectively, and are proposed to be involved in depletion of putative alarmones/signaling molecules [63,64] and/or damage metabolites [65,66]. Recent work suggests that dinucleoside polyphosphates can be used by RNA polymerases to initiate transcription and to act as 5'-RNA caps that may stabilize RNA, while ApaA-like hydrolases are able to remove these caps [67]. The Ap4A hydrolase knockout mutant *a1s_1624::Km* seems to play a role in *A. baumannii* surface motility and planktonic growth (Table 1). An *E. coli* Ap4A hydrolase (*apaH*) knockout mutant was previously associated with decreased motility [68]. In *Salmonella enterica* adhesion and invasion capacity into epithelial cells was reduced for the Δ *apaH* mutant [69]. Additionally, the *a1s_1624::Km* mutant exhibited increased sensitivity to imipenem (Table 1). Increased sensitivity of Δ *apaH* mutants against kanamycin and streptomycin was also shown for ATCC 17978, *E. coli*, and *P. aeruginosa* [70], and decreased sensitivity of Δ *apaH* mutants, in the form of persister cells, was found in *E. coli* [71]. Although we could not observe a significant impact of *a1s_1624::Km* on pellicle formation, others have shown A1S_1624 to be overproduced in pellicle cells in ATCC 17978 [44], and Ap4A metabolism impacts biofilm formation in *Pseudomonas fluorescens* [72]. The Ap5A pyrophosphatase knockout in ATCC 17978 resulted in a significant reduction in pellicle biofilm formation, significant attenuation of *G.*

mellonella infection, and resistance against ampicillin (Table 1). Interestingly, in the corresponding *a1s_0414::Km* mutant of 29D2, only surface-associated motility was affected (Table 2).

In general, Ap4A and Ap5A are thought to be synthesized by aminoacyl-tRNA synthetases in the absence of tRNAs during amino acid activation. This process requires ATP and a cognate amino acid [73,74]. Providing a possible link, we found a methionyl-tRNA synthetase in our surface motility-deficient library (discussed below).

Genes involved in RNA modification/regulation

We found 3 genes involved in the regulation and/or modification of RNAs: *metG* (methionyl-tRNA synthetase, *a1s_0778*), *rpmG* (50S ribosomal protein L33, *a1s_0447*), and *gidA* (glucose-inhibited division protein A, a tRNA modification enzyme, *a1s_2182*). The deficiency in motility of *gidA* mutants has been described mainly for swarming motility in bacteria like *Bacillus cereus* [48], *Serratia* species SCBI [75], and *Pseudomonas syringae* [76]. In the present study, a *gidA* null allele in strains ATCC 17978 and 29D2 resulted in small decreases in their planktonic growth (Suppl. Fig. S4 and S5). Contrary results for Δ *gidA* bacterial growth has been reported (reviewed in [77]). Interestingly, proteomic analysis of *A. baumannii* planktonic and biofilm growth identified *GidA* only under biofilm growth conditions [78], while several studies reported the negative effect of *gidA* mutants on biofilm formation in different bacteria [79,80]. In the present study we also saw a significant reduction of the pellicle-biofilm formation in both *gidA::Km* mutants (Fig. 2). An essential role of *gidA* in pellicle-biofilm formation was also shown in *Bacillus cereus* [48]. While several *GidA*-associated virulence effects have been reported (reviewed

in [77]) we did not see significant attenuation in the *G. mellonella* infection model (Suppl. Figs. S6 and S7).

In contrast, the knockout of *metG* (*a1s_0778*) was associated with a significant attenuation in the *G. mellonella* infection model (Fig. 4). Similarly, involvement of *metG* in *A. baumannii* virulence was also shown in a mouse pneumonia model [53]. The *metG::Km* mutant revealed a significantly reduced ability to form pellicles (Fig. 2). Moreover, MetG was found to be more abundant in *A. baumannii* pellicle cells than in planktonic cells [44]. In our study we found the *metG::Km* mutant to be more sensitive to tetracycline (Table 1), which agrees with observations of amino acid substitutions of MetG associated with increased antibiotic tolerance in *Burkholderia thailandensis* [81] and *E. coli* [82,83].

We observed increased sensitivity of the *rpmG::Km* mutant to imipenem and tetracycline (Table 1). This data is in line with a study which showed that a mitomycin C resistance phenotype was associated with RpmG overproduction in *E. coli* [84]

Genes involved in oxidative stress

The ATCC 17978 gene *a1S_3366* is predicted to encode a gamma-glutamate-cysteine ligase (*gshA*) which is required to synthesize glutathione (GSH), an antioxidant molecule that protects cells against oxidative stress [85,86]. Different studies observed a decrease in swarming [87], swimming [87,88], and twitching motility [88] of the *P. aeruginosa* $\Delta gshA$ mutant compared to the parental strain. Contrary results were found for the ability of *P. aeruginosa* $\Delta gshA$ to form biofilms (increased in [88] and decreased in [87]). We did not find any changes in pellicle biofilm production compared to the parental strains for both of our *gshA* mutants (Fig.

2). In *Acinetobacter baylyi* the knockout of *gshA* increased sensitivity to metronidazole and ciprofloxacin [89]. We observed an enhanced sensitivity to ampicillin for the 29D2 *gshA::Km* mutant, but the ATCC 17978 *gshA::Km* mutant showed a resistant phenotype (Table 3). Attenuation in *G. mellonella* infection was observed for the ATCC 17978 *gshA* mutant strain, which agrees with other studies describing *gshA* mutants to be attenuated in *C. elegans* infection (*P. aeruginosa* [90]) and a murine infection model (*Salmonella* [91]).

The *A. baumannii* gene *a1s_0530* encodes for a rhodanese domain-containing protein, a putative sulfurtransferase, supposed to be involved in oxidative stress detoxification and sulfur metabolism [92-95]. The only knockout-related phenotype, besides surface-associated motility-deficiency, that we observed in ATCC 17978 was a significant increase in pellicle biofilm production (Fig. 2A). Finally, the involvement of oxidative stress response proteins in air-liquid pellicles has been described recently in a proteomic study of ATCC 17978 [44].

Outer membrane proteins

The gene *a1s_3297* encodes a putative outer membrane protein and *a1s_1970* encodes a putative membrane-associated Zn-dependent protease (RseP). Here we show that both genes are involved in *A. baumannii* virulence, pellicle formation, and antimicrobial resistance (Tables 1 and 2).

We found OmpA to be involved in *A. baumannii* surface-associated motility, which has been described for the *A. nosocomialis* strain M2 by Clemmer *et al.* [26]. Several studies have reported the involvement of OmpA in biofilm formation [96-98] and OmpA, along with other outer membrane proteins, was observed to accumulate in *A.*

baumannii pellicle cells compared to planktonic cells [44]. We found the *ompA* knockout associated with a significant decrease in pellicle biofilm formation in ATCC 17978 but not in 29D2 (Fig. 2). For *A. baumannii* and a number of other pathogens, OmpA has been identified as a virulence factor and its importance in bacterial pathogenicity has been shown recently (reviewed in [99]). For example, the loss of OmpA impaired virulence of *A. baumannii* in *C. elegans* [100] and *Klebsiella pneumoniae* virulence in *G. mellonella* [101]. In our study, the knockout of *ompA* in both tested strains significantly decreased the mutant's ability to kill *G. mellonella* caterpillars (Fig. 4). We observed 1-2 log scale lower CFU numbers (for both mutant strains) compared to the OD-adjusted suspension which was used to infect the caterpillars. This observation led us to examine the mutant's cell morphology by microscopy. We found the *ompA::Km* mutants exhibiting filamentous cell phenotypes in contrast to the parental strains (Suppl. Fig. S8). Filamentous cell morphologies are known to provide bacterial survival advantages, e.g. protection against phagocytosis, resistance against antibiotics, and enhanced response to environmental cues like quorum sensing [102]. In other bacteria, the loss of outer membrane proteins, like the Tol-Pal system or OmpA-like proteins, resulted in reduced membrane integrity and alterations in cell division [103,104]. OmpA is involved in the ability of *A. baumannii* to grow and persist in human serum [11,105] and in the adherence and invasion of epithelial cells [106]. A resistance phenotype was only observed for the ATCC 17978 *ompA::Km* mutant strain. This mutant showed a 2-fold increase in MIC for imipenem compared to the parental strain (Table 3), correlating with the published finding that the *A. baumannii* OmpA C-terminus is important for resistance to antibiotics including imipenem [107].

Genes involved in 1,3-diaminopropane biosynthesis

As previously shown, mutations in the genes *dat* and *ddc* resulted in a dramatic reduction in surface-associated motility, but can be restored by supplementation with 120 μ M DAP [33]. In the present study, we observed motility deficiency for these genes in 29D2 (Table 2). We also gained new insight into the pleiotropic effects of the *dat::Km* and *ddc::Km* mutants, such as a significant decrease in pellicle biofilm formation. This observation might represent species-specific traits as we see this effect in both tested strains (ATCC 17978 and 29D2), whereas we see contradictory results for the MIC assays (Table 3).

Genes involved in lipopeptide synthesis/export

The genes *a1s_0113* and *a1s_0116* are involved in the synthesis and export of a lipopeptide and are part of an operon consisting of 8 genes [26,108]. The knockout of *a1s_0113* (acyl-CoA dehydrogenase) in *A. nosocomialis* clinical isolate M2 resulted in a significant surface motility defect [26], which correlates with our observation in ATCC 17978 (Fig. 1A). We found both mutants to show similar pleiotropic effects in ATCC 17978 (Table 1). Additionally, other genes of this operon have been reported to be necessary for motility (*a1s_0112* and *a1s_0115* [39]), pellicle biofilm formation (*a1s_0112* and *a1s_0115* [39]; *a1s_0114* [108]), and biofilm formation on abiotic surfaces (*a1s_0114* [108,109]). A pellicle proteome analysis in ATCC 17978 found the proteins A1S_0112-A1S_0118, with the exception of A1S_0114, to accumulate in the pellicle [44]. Since the gene *a1s_0116* encodes an RND superfamily transporter, it may thus play a role in multi-drug resistance. Deletion of *a1s_0116* in ATCC 17978 resulted in significantly increased ampicillin resistance compared to the parental

strain whereas no differences were observed for testing with imipenem and tetracycline (Table 3). A transcriptomic study on imipenem-resistant ATCC 17978 cells showed decreased expression of genes from the *a1s_0112-a1s_0119* cluster [110]. Clemmer *et al.* speculated that the lipopeptide synthesized from the *a1s_0112-a1s_0119* operon may act as a surfactant to promote motility, but they could not detect any surfactant activity in *A. nosocomialis* culture supernatants [26]. While we could not show a significant effect of *a1s_0113* or *a1s_0116* inactivation on virulence in *G. mellonella*, significant attenuation was observed in the same model for an *a1s_0114* mutant [108]. No essential role of any of the *a1s_0112-a1s_0119* genes in virulence was also found for strain AB5075 [111]. In conclusion, our data confirm findings by other groups [26,39,109,108,44] indicating that genes of the *a1s_0112-a1s_0119* operon are essential for surface motility and pellicle biofilm formation in *A. baumannii*.

Genes involved in DNA modification/repair/uptake

We found 4 genes in our library to be involved in DNA modification, uptake, and recombination. The gene *a1s_0222*, designated as *aamA*, encodes a Type II N6-adenine DNA methyltransferase [112,113]. Methylation is important for the regulation of various physiological processes [114,115]. We speculate the phenotype of both *aamA* mutants to represent strain-specific traits (Tables 1 and 2). In bacteria DNA methylation is the most studied epigenetic mechanism and the *E. coli* Dam protein is the most prominent orphan DNA adenine methyltransferase [116]. For other bacteria like *S. enterica*, *Y. enterocolitica*, and *K. pneumoniae* different phenotypes of *dam*

mutants and *dam* overexpression were shown to affect motility, virulence, and other traits (reviewed in [117]).

The *A. baumannii* gene *a1s_2334* encodes an S-adenosyl-L-homocysteine hydrolase (*sahH*), which takes part in the recycling of S-adenosyl-L-methionine (SAM). Here we show that inactivation of *sahH* in *A. baumannii* leads to pleiotropic effects such as strong motility deficiency, a significant attenuation in *G. mellonella* caterpillar infection, and increased antibiotic resistance (Table 1). Furthermore, we found the Holliday junction helicase subunit A (*ruvA*/ *a1s_2587*) to be important for *A. baumannii* surface-associated motility, pellicle biofilm formation, and antibiotic resistance in ATCC 17978 (Table 1).

We identified the gene *a1s_2610* in our mutant library screening. Designated as *comEC*, this gene is involved in DNA uptake and incorporation of exogenous DNA into the genome. Phenotypically, a linkage between motility and natural transformation competence was shown in that *A. baumannii* can take up DNA while moving along wet surfaces [16] and its transformability is influenced by motility-determining parameters such as agarose concentration [41]. Genetically this interrelationship was illustrated by abolished twitching motility and natural transformation competence of *comEC* knockout mutants in *A. baumannii* strains 07-095 and 07-102, and a defect in surface-associated motility was ascribed for the ATCC 17978 *comEC::Km* mutant [16]. Deficiency in twitching motility has also been shown for $\Delta comEC$ in *Thermus thermophilus* [118]. Our results confirmed surface-associated motility deficiency in the 29D2 *comEC::Km* mutant strain (Fig. 2). Deficiency in twitching motility was also shown for $\Delta comEC$ in *Thermus thermophilus* [118]. A striking attenuation in *G. mellonella* caterpillar infection for the *comEC::Km* mutants in both 17978 and 29D2 was observed (Fig. 4), similar to attenuation of

comEC::Km mutant derivatives of *A. baumannii* strains DSM 30011, 07-102, and 07-095 [16]. In *Listeria monocytogenes*, *comEC* was demonstrated to be involved in phagosomal escape, intracellular growth, and virulence [119]. However, *com* genes have been reported to be involved in bacterial biofilm formation [120], which we could not confirm for our *comEC::Km* mutant strains (Fig. 2).

Other genes

The gene *a1s_0065* encodes a UDP-glucose 4-epimerase (*galE*) and is predicted to play a role in capsule and lipopolysaccharide biosynthesis [121]. Capsules are important virulence factors in *A. baumannii* [122]. In this study, a knockout of *galE* resulted in a reduced motility phenotype in ATCC 17978 and 29D2 (Fig. 1). The involvement of lipopolysaccharides in *Acinetobacter* surface motility has recently been shown for the gene *rmIB* which is part of the O-antigen in Gram-negative bacteria [26]. A proteomic study of *A. baumannii* revealed GalE to be only expressed in biofilm growth mode [78]. Additionally, multiple studies have revealed that UDP-glucose 4-epimerases play a role in biofilm formation, including in *Sinorhizobium meliloti* [123], *Vibrio cholerae* [124], *Bacillus subtilis* [125], and *Thermus thermophiles* [126]. The knockout of *galE* resulted in a significant increase in pellicle biomass production for both mutants compared to their parental strains (Fig. 2). Similar observations were also made for a *galE* mutant in *Haemophilus parasuis* [127] and *Porphyromonas gingivalis* [128]. Moreover, other proteins necessary for the catabolism of D-galactose (Leloir pathway), GalM and GalU, were found to be upregulated in *A. baumannii* biofilms [129]. Infection of *G. mellonella* caterpillars with the *galE::Km* mutants resulted in a significant attenuation. The caterpillar survival rate

5 days post infection was 98.9% for the ATCC 17978 *galE::Km* mutant and 95.8% for 29D2 *galE::Km* (Fig. 4). Similarly, a significant decrease in persistence in a mouse pneumonia model of *A. baumannii* was previously reported for the *a1s_0065* mutant [53]. Several other studies demonstrated UDP-glucose 4-epimerases to be involved in virulence/pathogenesis, for example in *Bacillus anthracis* [130], *Streptococcus iniae* [131], and the plant-pathogenic fungus *Leptosphaeria maculans* [132]. In our study, we observed a resistance phenotype for the ATCC 17978 *galE::Km* mutant to ampicillin and tetracycline. By contrast, the 29D2 *galE::Km* mutant showed significant sensitivity to ampicillin compared to the parental strain (Table 3). The *galE* mutant in *Porphyromonas gingivalis* was shown to be significantly more susceptible to benzylpenicillin, oxacillin, cefotaxime, imipenem, and vancomycin compared to the wildtype [128] and involvement of UDP-glucose 4-epimerases in antibiotic resistance/sensitivity was reported for several other bacteria [133-135].

The gene *a1s_0806* encodes for an adenosylmethionine-8-amino-7-oxononanoate aminotransferase (*bioA*), belonging to the acetyl ornithine aminotransferase family, which is part of the pyridoxal phosphate-dependent aspartate aminotransferase superfamily. BioA is part of the biotin biosynthesis pathway and biotin is essential for cell metabolism in prokaryotes and eukaryotes, and only bacteria and plants can synthesize biotin *de novo* [136-138]. Inactivation of *a1s_0806* in ATCC 17978 resulted in the strongest surface motility defect (Fig. 1) and the greatest pellicle biomass production of all tested mutants (Fig. 2). In contrast, the mutant's ability to kill *G. mellonella* caterpillars was not significantly affected (Suppl. Figs. S6 and S7). Other studies in *M. tuberculosis* have demonstrated *bioA* to be essential for establishment of infection and persistence in mice [139].

A knockout of gene *a1s_1055*, encoding a LysM peptidoglycan-binding domain-containing lytic transglycosylase, resulted in a significantly increased pellicle biomass production. Similar to our findings, mutation of lytic transglycosylase (A1S_3027) in *A. nosocomialis* strain M2 was found to exhibit surface motility deficiency [26]. A1S_1055 seems to play a role in *A. baumannii* virulence, since mutants of both parental backgrounds led to attenuation in the *G. mellonella* infection assay (Fig. 4).

The gene *a1s_2761* encodes for a 2-methylnaconitate cis-trans isomerase (PrpF), involved in the 2-methylcitric acid cycle and propionate catabolism. The inactivation of *prpF* in 17978 resulted in one of the most reduced pellicle formation phenotypes of all tested mutant strains (Fig. 2), which correlates with the previous finding that PrpF accumulated in mature 4-days pellicles in 17978 [44]. The mutant's virulence capacity was not significantly affected (Suppl. Fig. S6). In contrast, the 29D2 *prpF::Km* mutant strain displayed a significant attenuation in the *G. mellonella* infection model (Fig. 4) but did not affect pellicle biofilm formation (Fig. 2).

The *A. baumannii* gene *a1s_3026* is predicted to encode a secreted ribonuclease T2 family protein (RNase T2-family). In the *A. baumannii* strain 98-37-09 a deficiency in surface motility for the *a1s_3026* knockout was previously reported [34]. Additionally, Jacobs *et al.* indicated that the *a1s_3026* mutant showed reduced colonization on abiotic surfaces like glass, polystyrene, and stainless steel, and that the *a1s_3026* knockout was shown to be associated with decreased expression of genes involved in motility and biofilm formation [34]. Despite the deficiency in surface-associated motility we observed a significant decrease in pellicle biofilm formation (Tables 1 and 2). Interestingly, A1S_3026 was shown to be involved in *A. baumannii* colistin resistance [140] and both *3026::Km* mutant strains exhibited elevated resistance values for ampicillin and imipenem (Table 3).

The *A. baumannii* gene *a1s_3129* encodes for a succinylarginine dihydrolase (*astB*) and is involved in the arginine succinyltransferase (AST) pathway [141]. In a mouse pneumonia model of *A. baumannii*, the *astB* insertion caused a significant decrease in persistence [53]. However, we did not observe a significant attenuation in the *G. mellonella* infection model (Table 1). Furthermore, we observed a significant reduction of pellicle formation in the *astB::Km* mutant (Fig. 2A). This is in line with an accumulation of AstB in ATCC 17978 pellicle cells described previously [44].

***G. mellonella* caterpillar infection**

The *G. mellonella* caterpillar, which is an established insect model system for bacterial infections [142], was used to study virulence traits of the motility-deficient mutants. A study on virulence and resistance to antibiotic and environmental stress analyzed 250,000 *A. baumannii* AB5075 transposon mutants for growth within *G. mellonella* larvae, and TnSeq experiments identified 300 genes essential for growth [111]. When comparing with these results, we could not identify concordant genes in our library, but we found that main categories of genes do match. For example we found *galE* to be essential and in AB5075 numerous genes involved in structure and function of the cell envelopment were found to be required for growth in *G. mellonella* [111]. Conversely, for example, the *gidA::Km* mutant was not attenuated in *G. mellonella* infection in our study (Suppl. Fig. S6 and Fig. S7), but was stated to be essential for growth of AB5075 in *G. mellonella* [111]. It is known that AB5075 is more virulent than ATCC 17978 [143,111], therefore, comparative studies are needed to unravel strain-specific and species-specific traits.

Limitations

While our study highlights the need for comparative studies of specific mutant phenotypes in different strains to distinguish strain-specific from species-specific traits, it is clear that the two strains studied in detail here do not provide a sufficient basis to deduce such insight. Such comparative studies in combination with genome-based analyses may pave the way for the identification of species-specific traits and, ultimately, novel target sites.

The use of marker-based mutagenesis and naturally competent strains to efficiently generate sets of mutants in different strains has its shortcomings as recombination events are not necessarily limited to the site of the marker gene. Apart from homology-based recombination events, transfer of mobile genetic elements and even illegitimate recombination events may occur [144,145] and could corrupt the mutants' phenotypes. A few of the mutations described in this study have been partially characterized previously using additional strains (*ddc*, *dat*, *comEC*, and *aamA* for example [16,33,112]). However, repetitive construction of the same mutants did not lead to significant phenotype variation arguing against a high frequency of such corrupting side-effects. The ability to rapidly reconstruct mutants is an advantage of this method which can allow for confirmation of mutant phenotypes.

Complementation experiments and site-specific deletion mutagenesis would exclude polar effects of the transposon insertions and help to verify the contribution of each gene. In support of the specificity of our findings, we found many groups of related mutants (e.g. purine and pyrimidine biosynthesis) and identified multiple linkages to motility mutants described in other organisms.

We could not achieve a saturated mutant library which indicates that surface-associated motility is probably under control of additional genes yet to be discovered. Our attempts were limited by the poor transformability of ATCC 17978 with the transposome complex, and by the transposition not being completely unbiased so that we obtained several insertion events repeatedly and independently.

Conclusion

In this study we made use of a previously generated *A. baumannii* ATCC 17978 EZ-Tn5™ <KAN-2> transposon mutant library [33] to screen for surface-associated motility-deficient mutants. We identified 30 genes involved in surface motility. All tested mutants originally identified as motility-deficient in strain ATCC 17978 also displayed a motility-deficient phenotype in the *A. baumannii* white stork isolate 29D2. Some of these genes have already been linked to motility in *A. baumannii* (e.g. *comEC*, *a1s_0113*, and *a1s_0116*) or other bacteria (e.g. *carB* and *gidA*), but some of our findings represent new insights into requirements for surface-associated motility. Furthermore, we analyzed these mutants with respect to bacterial growth, pellicle biofilm formation, virulence in *G. mellonella* infection, and antibiotic resistance and used the naturally competent strain 29D2 to indicate whether the mutations showed strain-specific or species-specific traits. In summary, we can state that mutations in genes involved in purine/pyrimidine/folate biosynthesis are essential for all tested categories. Mutants that targeted RNA modification/regulation seem to mainly play a role in motility and pellicle formation. The discovery of novel genes required for surface-associated motility in *A. baumannii* demonstrates that more work is required to further define its genetic basis.

790

791 **Conflict of Interest**

792 The authors declare that they have no conflict of interest.

793

Figures with legends

Fig. 1. ATCC 17978 mutants (A) and 29D2 mutants (B) deficient in surface-associated motility. Wildtypes and mutants of strains ATCC 17978 and 29D2 were inoculated on motility plates. Plates were incubated for 16 h at 37°C. The diameter ($\emptyset$) of the surface-associated motility spreading zone was measured and triplicates were statistically analyzed. All mutants of strains ATCC 17978 (A) and 29D2 (B) displayed a significant motility deficiency compared to their respective parental strain (p-value ≤ 0.05).

Fig. 2. Pellicle biofilm formation of ATCC 17978 wildtype and mutants (A) and 29D2 wildtype and mutants (B). A. *baumannii* pellicle biofilms developed within 3 days of incubation and were stained with a 0.5% crystal violet solution. The biofilm was scrubbed and flushed off the tube walls and the absorption was determined at 550 nm. For each strain 3 independent experiments were performed and statistical significance was analyzed by the Student's *t* test (2-tailed, unpaired). Significance as indicated: *, p -value ≤ 0.05 ; **, p -value ≤ 0.01 ; ***, p -value ≤ 0.001 ; ****, p -value ≤ 0.0001 .

Fig. 3. Growth deficiency of mutant strains from ATCC 17978 (A) and 29D2 (B) mutant libraries. OD-adjusted bacterial cultures were grown for 9 h at 37°C under constant shaking. Every hour cultures were measured at an OD of 600 nm. For each strain data obtained from 3 independent cultures grown on the same day were averaged and represented by the mean $\pm$ SD. Growth defects compared to wildtype ATCC 17978 were observed for mutants that are involved in purine, pyrimidine, and diaminopropane biosynthesis (A). In strain 29D2, growth defects were observed for mutants involved in purine/pyrimidine/folate and diaminopropane biosynthesis, and for mutants *galE::Km*, *ompA::Km*, and *prpF::Km* (B). See Supplementary Figs. S4 and S5 for growth curves of all strains described in this study.

Fig. 4. Attenuation of *A. baumannii* ATCC 17978 mutants (A) and 29D2 mutants (B) in the *G. mellonella* caterpillar infection model. Caterpillars were infected with 3×10^5 CFU of *A. baumannii* strains as indicated. Sterile PBS (black lines) was used as a control. Three independent experiments were performed with groups of 16 caterpillars for every bacterial strain and control. Data obtained from 3 independent experiments were averaged and represented by the mean $\pm$ SD. In strain ATCC 17978, 15 of 30 mutants showed a significant attenuation at 5 days post-infection (see Table 1 for p-values) compared to the wildtype strain (A). In strain 29D2, 11 of 21 mutants were attenuated (see Table 2 for p-values) in the *G. mellonella* infection model (B). See Supplementary Figs. S6 and S7 for infection data of all strains described in this study.

Fig. 5 Genes inactivated in *A. baumannii* ATCC 17978 mutants with a surface-associated motility defect and their known/predicted/putative function in the bacterial cell. A common color code indicates that mutants belong to the same functional, processual, and/or structural category. OM, outer membrane; IM, inner membrane; Ap4A, diadenosine tetraphosphate; Ap5A, diadenosine pentaphosphate; SAM, S-adenosyl-L-methionine; SAH, S-adenosylhomocysteine; Ado, adenosine; Hcy, homocysteine; Met, methionine; ASA, L-aspartate 4-semialdehyde; DAB, L-2,4-diaminobutanoate; 2-MMC, 2-methylcitric acid cycle.

References

1. Baumann P (1968) Isolation of *Acinetobacter* from soil and water. *J Bacteriol* 96 (1):39-42
2. Peleg AY, Seifert H, Paterson DL (2008) *Acinetobacter baumannii*: emergence of a successful pathogen. *Clin Microbiol Rev* 21 (3):538-582. doi:10.1128/CMR.00058-07
3. Harding CM, Hennon SW, Feldman MF (2018) Uncovering the mechanisms of *Acinetobacter baumannii* virulence. *Nat Rev Microbiol* 16 (2):91-102. doi:10.1038/nrmicro.2017.148
4. Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, Moreno R, Lipman J, Gomersall C, Sakr Y, Reinhart K, Investigators ELGo (2009) International study of the prevalence and outcomes of infection in intensive care units. *JAMA* 302 (21):2323-2329. doi:10.1001/jama.2009.1754
5. Landman D, Bratu S, Kochar S, Panwar M, Trehan M, Doymaz M, Quale J (2007) Evolution of antimicrobial resistance among *Pseudomonas aeruginosa*, *Acinetobacter baumannii* and *Klebsiella pneumoniae* in Brooklyn, NY. *J Antimicrob Chemother* 60 (1):78-82. doi:10.1093/jac/dkm129
6. Tognim MC, Andrade SS, Silbert S, Gales AC, Jones RN, Sader HS (2004) Resistance trends of *Acinetobacter* spp. in Latin America and characterization of international dissemination of multi-drug resistant strains: five-year report of the SENTRY Antimicrobial Surveillance Program. *Int J Infect Dis* 8 (5):284-291. doi:10.1016/j.ijid.2003.11.009
7. Giammanco A, Cala C, Fasciana T, Dowzicky MJ (2017) Global Assessment of the Activity of Tigecycline against Multidrug-Resistant Gram-Negative Pathogens between 2004 and 2014 as Part of the Tigecycline Evaluation and Surveillance Trial. *mSphere* 2 (1). doi:10.1128/mSphere.00310-16
8. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, Pulcini C, Kahlmeter G, Kluytmans J, Carmeli Y, Ouellette M, Outtersen K, Patel J, Cavaleri M, Cox EM, Houchens CR, Grayson ML, Hansen P, Singh N, Theuretzbacher U, Magrini N, Group WHOPPLW (2018) Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet Infect Dis* 18 (3):318-327. doi:10.1016/S1473-3099(17)30753-3
9. Roca I, Espinal P, Vila-Farres X, Vila J (2012) The *Acinetobacter baumannii* Oxymoron: Commensal Hospital Dweller Turned Pan-Drug-Resistant Menace. *Front Microbiol* 3:148. doi:10.3389/fmicb.2012.00148
10. Giannouli M, Antunes LC, Marchetti V, Triassi M, Visca P, Zarrilli R (2013) Virulence-related traits of epidemic *Acinetobacter baumannii* strains belonging to the international clonal lineages I-III and to the emerging genotypes ST25 and ST78. *BMC Infect Dis* 13:282. doi:10.1186/1471-2334-13-282

11. Antunes LC, Imperi F, Carattoli A, Visca P (2011) Deciphering the multifactorial nature of *Acinetobacter baumannii* pathogenicity. PLoS One 6 (8):e22674. doi:10.1371/journal.pone.0022674
12. Jawad A, Seifert H, Snelling AM, Heritage J, Hawkey PM (1998) Survival of *Acinetobacter baumannii* on dry surfaces: comparison of outbreak and sporadic isolates. J Clin Microbiol 36 (7):1938-1941
13. Wagenvoort JH, Joosten EJ (2002) An outbreak *Acinetobacter baumannii* that mimics MRSA in its environmental longevity. J Hosp Infect 52 (3):226-227. doi:10.1053/jhin.2001.1294
14. Greene C, Vadlamudi G, Newton D, Foxman B, Xi C (2016) The influence of biofilm formation and multidrug resistance on environmental survival of clinical and environmental isolates of *Acinetobacter baumannii*. Am J Infect Control 44 (5):e65-71. doi:10.1016/j.ajic.2015.12.012
15. Eijkelkamp BA, Stroeher UH, Hassan KA, Elbourne LD, Paulsen IT, Brown MH (2013) H-NS plays a role in expression of *Acinetobacter baumannii* virulence features. Infect Immun 81 (7):2574-2583. doi:10.1128/IAI.00065-13
16. Wilharm G, Piesker J, Laue M, Skiebe E (2013) DNA uptake by the nosocomial pathogen *Acinetobacter baumannii* occurs during movement along wet surfaces. J Bacteriol 195 (18):4146-4153. doi:10.1128/JB.00754-13
17. Harding CM, Tracy EN, Carruthers MD, Rather PN, Actis LA, Munson RS, Jr. (2013) *Acinetobacter baumannii* strain M2 produces type IV pili which play a role in natural transformation and twitching motility but not surface-associated motility. MBio 4 (4). doi:10.1128/mBio.00360-13
18. Harshey RM (2003) Bacterial motility on a surface: many ways to a common goal. Annu Rev Microbiol 57:249-273. doi:10.1146/annurev.micro.57.030502.091014
19. Jarrell KF, McBride MJ (2008) The surprisingly diverse ways that prokaryotes move. Nat Rev Microbiol 6 (6):466-476. doi:10.1038/nrmicro1900
20. Mattick JS (2002) Type IV pili and twitching motility. Annu Rev Microbiol 56:289-314. doi:10.1146/annurev.micro.56.012302.160938
21. Merz AJ, So M, Sheetz MP (2000) Pilus retraction powers bacterial twitching motility. Nature 407 (6800):98-102. doi:10.1038/35024105
22. Skerker JM, Berg HC (2001) Direct observation of extension and retraction of type IV pili. Proc Natl Acad Sci U S A 98 (12):6901-6904. doi:10.1073/pnas.121171698
23. Wall D, Kaiser D (1999) Type IV pili and cell motility. Mol Microbiol 32 (1):1-10. doi:10.1046/j.1365-2958.1999.01339.x
24. Henrichsen J (1984) Not gliding but twitching motility of *Acinetobacter calcoaceticus*. J Clin Pathol 37 (1):102-103. doi:10.1136/jcp.37.1.102-b

25. McBride MJ (2010) Shining a light on an opportunistic pathogen. *J Bacteriol* 192 (24):6325-6326.
doi:10.1128/JB.01141-10
26. Clemmer KM, Bonomo RA, Rather PN (2011) Genetic analysis of surface motility in *Acinetobacter baumannii*. *Microbiology* 157 (Pt 9):2534-2544. doi:10.1099/mic.0.049791-0
27. Henrichsen J, Blom J (1975) Correlation between twitching motility and possession of polar fimbriae in *Acinetobacter calcoaceticus*. *Acta Pathol Microbiol Scand B* 83 (2):103-115.
doi:10.1111/j.1699-0463.1975.tb00078.x
28. Eijkelkamp BA, Stroehrer UH, Hassan KA, Papadimitriou MS, Paulsen IT, Brown MH (2011) Adherence and motility characteristics of clinical *Acinetobacter baumannii* isolates. *FEMS Microbiol Lett* 323 (1):44-51. doi:10.1111/j.1574-6968.2011.02362.x
29. Barker J, Maxted H (1975) Observations on the growth and movement of *Acinetobacter* on semi-solid media. *J Med Microbiol* 8 (3):443-446. doi:10.1099/00222615-8-3-443
30. Mussi MA, Gaddy JA, Cabruja M, Arivett BA, Viale AM, Rasia R, Actis LA (2010) The opportunistic human pathogen *Acinetobacter baumannii* senses and responds to light. *J Bacteriol* 192 (24):6336-6345. doi:10.1128/JB.00917-10
31. Eijkelkamp BA, Hassan KA, Paulsen IT, Brown MH (2011) Investigation of the human pathogen *Acinetobacter baumannii* under iron limiting conditions. *BMC Genomics* 12:126.
doi:10.1186/1471-2164-12-126
32. McQueary CN, Kirkup BC, Si Y, Barlow M, Actis LA, Craft DW, Zurawski DV (2012) Extracellular stress and lipopolysaccharide modulate *Acinetobacter baumannii* surface-associated motility. *J Microbiol* 50 (3):434-443. doi:10.1007/s12275-012-1555-1
33. Skieba E, de Berardinis V, Morczinek P, Kerrinnes T, Faber F, Lepka D, Hammer B, Zimmermann O, Ziesing S, Wichelhaus TA, Hunfeld KP, Borgmann S, Grobner S, Higgins PG, Seifert H, Busse HJ, Witte W, Pfeifer Y, Wilharm G (2012) Surface-associated motility, a common trait of clinical isolates of *Acinetobacter baumannii*, depends on 1,3-diaminopropane. *Int J Med Microbiol* 302 (3):117-128. doi:10.1016/j.ijmm.2012.03.003
34. Jacobs AC, Blanchard CE, Catherman SC, Dunman PM, Murata Y (2014) An ribonuclease T2 family protein modulates *Acinetobacter baumannii* abiotic surface colonization. *PLoS One* 9 (1):e85729. doi:10.1371/journal.pone.0085729
35. Heindorf M, Kadari M, Heider C, Skieba E, Wilharm G (2014) Impact of *Acinetobacter baumannii* superoxide dismutase on motility, virulence, oxidative stress resistance and susceptibility to antibiotics. *PLoS One* 9 (7):e101033. doi:10.1371/journal.pone.0101033

36. Ahmad I, Nygren E, Khalid F, Myint SL, Uhlin BE (2020) A Cyclic-di-GMP signalling network regulates biofilm formation and surface associated motility of *Acinetobacter baumannii* 17978. *Sci Rep* 10 (1):1991. doi:10.1038/s41598-020-58522-5
37. Tipton KA, Rather PN (2017) An ompR-envZ Two-Component System Ortholog Regulates Phase Variation, Osmotic Tolerance, Motility, and Virulence in *Acinetobacter baumannii* Strain AB5075. *J Bacteriol* 199 (3). doi:10.1128/JB.00705-16
38. Tipton KA, Dimitrova D, Rather PN (2015) Phase-Variable Control of Multiple Phenotypes in *Acinetobacter baumannii* Strain AB5075. *J Bacteriol* 197 (15):2593-2599. doi:10.1128/JB.00188-15
39. Giles SK, Stroehrer UH, Eijkelkamp BA, Brown MH (2015) Identification of genes essential for pellicle formation in *Acinetobacter baumannii*. *BMC Microbiol* 15:116. doi:10.1186/s12866-015-0440-6
40. Wilharm G, Skiebe E, Higgins PG, Poppel MT, Blaschke U, Leser S, Heider C, Heindorf M, Brauner P, Jackel U, Bohland K, Cuny C, Lopinska A, Kaminski P, Kasprzak M, Bochenski M, Ciebiera O, Tobolka M, Zolnierowicz KM, Siekiera J, Seifert H, Gagne S, Salcedo SP, Kaatz M, Layer F, Bender JK, Fuchs S, Semmler T, Pfeifer Y, Jerzak L (2017) Relatedness of wildlife and livestock avian isolates of the nosocomial pathogen *Acinetobacter baumannii* to lineages spread in hospitals worldwide. *Environ Microbiol* 19 (10):4349-4364. doi:10.1111/1462-2920.13931
41. Godeux AS, Lupo A, Haenni M, Guette-Marquet S, Wilharm G, Laaberki MH, Charpentier X (2018) Fluorescence-Based Detection of Natural Transformation in Drug-Resistant *Acinetobacter baumannii*. *J Bacteriol* 200 (19). doi:10.1128/JB.00181-18
42. Ahmad I, Karah N, Nadeem A, Wai SN, Uhlin BE (2019) Analysis of colony phase variation switch in *Acinetobacter baumannii* clinical isolates. *PLoS One* 14 (1):e0210082. doi:10.1371/journal.pone.0210082
43. Choi KH, Kumar A, Schweizer HP (2006) A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA fragment transfer between chromosomes and plasmid transformation. *J Microbiol Methods* 64 (3):391-397. doi:10.1016/j.mimet.2005.06.001
44. Kentache T, Ben Abdelkrim A, Jouenne T, De E, Hardouin J (2017) Global Dynamic Proteome Study of a Pellicle-forming *Acinetobacter baumannii* Strain. *Mol Cell Proteomics* 16 (1):100-112. doi:10.1074/mcp.M116.061044
45. Nait Chabane Y, Marti S, Rihouey C, Alexandre S, Hardouin J, Lesouhaitier O, Vila J, Kaplan JB, Jouenne T, De E (2014) Characterisation of pellicles formed by *Acinetobacter baumannii* at the air-liquid interface. *PLoS One* 9 (10):e111660. doi:10.1371/journal.pone.0111660

46. Kinsinger RF, Kearns DB, Hale M, Fall R (2005) Genetic requirements for potassium ion-dependent colony spreading in *Bacillus subtilis*. *J Bacteriol* 187 (24):8462-8469. doi:10.1128/JB.187.24.8462-8469.2005
47. Yan F, Yu Y, Gozzi K, Chen Y, Guo JH, Chai Y (2017) Genome-Wide Investigation of Biofilm Formation in *Bacillus cereus*. *Appl Environ Microbiol* 83 (13). doi:10.1128/AEM.00561-17
48. Okshevsky M, Louw MG, Lamela EO, Nilsson M, Tolker-Nielsen T, Meyer RL (2018) A transposon mutant library of *Bacillus cereus* ATCC 10987 reveals novel genes required for biofilm formation and implicates motility as an important factor for pellicle-biofilm formation. *Microbiologyopen* 7 (2):e00552. doi:10.1002/mbo3.552
49. Vilain S, Pretorius JM, Theron J, Brozel VS (2009) DNA as an adhesin: *Bacillus cereus* requires extracellular DNA to form biofilms. *Appl Environ Microbiol* 75 (9):2861-2868. doi:10.1128/AEM.01317-08
50. Zhang X, de Maat V, Guzman Prieto AM, Prajsnar TK, Bayjanov JR, de Been M, Rogers MRC, Bonten MJM, Mesnage S, Willems RJL, van Schaik W (2017) RNA-seq and Tn-seq reveal fitness determinants of vancomycin-resistant *Enterococcus faecium* during growth in human serum. *BMC Genomics* 18 (1):893. doi:10.1186/s12864-017-4299-9
51. Samant S, Lee H, Ghassemi M, Chen J, Cook JL, Mankin AS, Neyfakh AA (2008) Nucleotide biosynthesis is critical for growth of bacteria in human blood. *PLoS Pathog* 4 (2):e37. doi:10.1371/journal.ppat.0040037
52. Schwager S, Agnoli K, Kothe M, Feldmann F, Givskov M, Carlier A, Eberl L (2013) Identification of *Burkholderia cenocepacia* strain H111 virulence factors using nonmammalian infection hosts. *Infect Immun* 81 (1):143-153. doi:10.1128/IAI.00768-12
53. Wang N, Ozer EA, Mandel MJ, Hauser AR (2014) Genome-wide identification of *Acinetobacter baumannii* genes necessary for persistence in the lung. *MBio* 5 (3):e01163-01114. doi:10.1128/mBio.01163-14
54. Polissi A, Pontiggia A, Feger G, Altieri M, Mottl H, Ferrari L, Simon D (1998) Large-scale identification of virulence genes from *Streptococcus pneumoniae*. *Infect Immun* 66 (12):5620-5629
55. Jenkins A, Cote C, Twenhafel N, Merkel T, Bozue J, Welkos S (2011) Role of purine biosynthesis in *Bacillus anthracis* pathogenesis and virulence. *Infect Immun* 79 (1):153-166. doi:10.1128/IAI.00925-10
56. Fuller TE, Kennedy MJ, Lowery DE (2000) Identification of *Pasteurella multocida* virulence genes in a septicemic mouse model using signature-tagged mutagenesis. *Microb Pathog* 29 (1):25-38. doi:10.1006/mpat.2000.0365

57. Yee R, Cui P, Shi W, Feng J, Zhang Y (2015) Genetic Screen Reveals the Role of Purine Metabolism in *Staphylococcus aureus* Persistence to Rifampicin. *Antibiotics (Basel)* 4 (4):627-642. doi:10.3390/antibiotics4040627
58. Charlier D, Nguyen Le Minh P, Roovers M (2018) Regulation of carbamoylphosphate synthesis in *Escherichia coli*: an amazing metabolite at the crossroad of arginine and pyrimidine biosynthesis. *Amino Acids* 50 (12):1647-1661. doi:10.1007/s00726-018-2654-z
59. Ueda A, Attila C, Whiteley M, Wood TK (2009) Uracil influences quorum sensing and biofilm formation in *Pseudomonas aeruginosa* and fluorouracil is an antagonist. *Microb Biotechnol* 2 (1):62-74. doi:10.1111/j.1751-7915.2008.00060.x
60. Garavaglia M, Rossi E, Landini P (2012) The pyrimidine nucleotide biosynthetic pathway modulates production of biofilm determinants in *Escherichia coli*. *PLoS One* 7 (2):e31252. doi:10.1371/journal.pone.0031252
61. Beaume M, Kohler T, Fontana T, Tognon M, Renzoni A, van Delden C (2015) Metabolic pathways of *Pseudomonas aeruginosa* involved in competition with respiratory bacterial pathogens. *Front Microbiol* 6:321. doi:10.3389/fmicb.2015.00321
62. Zhuo T, Rou W, Song X, Guo J, Fan X, Kamau GG, Zou H (2015) Molecular study on the *carAB* operon reveals that *carB* gene is required for swimming and biofilm formation in *Xanthomonas citri* subsp. *citri*. *BMC Microbiol* 15:225. doi:10.1186/s12866-015-0555-9
63. Kisselev LL, Justesen J, Wolfson AD, Frolova LY (1998) Diadenosine oligophosphates (Ap(n)A), a novel class of signalling molecules? *FEBS Lett* 427 (2):157-163. doi:10.1016/s0014-5793(98)00420-7
64. Bochner BR, Lee PC, Wilson SW, Cutler CW, Ames BN (1984) AppppA and Related Adenylylated Nucleotides Are Synthesized as a Consequence of Oxidation Stress. *Cell* 37 (1):225-232. doi:10.1016/0092-8674(84)90318-0
65. Linster CL, Van Schaftingen E, Hanson AD (2013) Metabolite damage and its repair or pre-emption. *Nat Chem Biol* 9 (2):72-80. doi:10.1038/nchembio.1141
66. Despotovic D, Brandis A, Savidor A, Levin Y, Fumagalli L, Tawfik DS (2017) Diadenosine tetraphosphate (Ap4A) - an *E. coli* alarmone or a damage metabolite? *FEBS J* 284 (14):2194-2215. doi:10.1111/febs.14113
67. Hudecek O, Benoni R, Reyes-Gutierrez PE, Culka M, Sanderova H, Hubalek M, Rulisek L, Cvacka J, Krasny L, Cahova H (2020) Dinucleoside polyphosphates act as 5'-RNA caps in bacteria. *Nat Commun* 11 (1):1052. doi:10.1038/s41467-020-14896-8

68. Farr SB, Arnosti DN, Chamberlin MJ, Ames BN (1989) An apaH mutation causes AppppA to accumulate and affects motility and catabolite repression in Escherichia coli. Proc Natl Acad Sci U S A 86 (13):5010-5014. doi:10.1073/pnas.86.13.5010
69. Ismail TM, Hart CA, McLennan AG (2003) Regulation of dinucleoside polyphosphate pools by the YgdP and ApaH hydrolases is essential for the ability of Salmonella enterica serovar typhimurium to invade cultured mammalian cells. J Biol Chem 278 (35):32602-32607. doi:10.1074/jbc.M305994200
70. Ji X, Zou J, Peng H, Stolle AS, Xie R, Zhang H, Peng B, Mekalanos JJ, Zheng J (2019) Alarmone Ap4A is elevated by aminoglycoside antibiotics and enhances their bactericidal activity. Proc Natl Acad Sci U S A 116 (19):9578-9585. doi:10.1073/pnas.1822026116
71. Hansen S, Lewis K, Vulic M (2008) Role of global regulators and nucleotide metabolism in antibiotic tolerance in Escherichia coli. Antimicrob Agents Chemother 52 (8):2718-2726. doi:10.1128/AAC.00144-08
72. Monds RD, Newell PD, Wagner JC, Schwartzman JA, Lu W, Rabinowitz JD, O'Toole GA (2010) Diadenosine tetraphosphate (Ap4A) metabolism impacts biofilm formation by Pseudomonas fluorescens via modulation of c-di-GMP-dependent pathways. J Bacteriol 192 (12):3011-3023. doi:10.1128/JB.01571-09
73. Goerlich O, Foeckler R, Holler E (1982) Mechanism of synthesis of adenosine(5')tetraphospho(5')adenosine (AppppA) by aminoacyl-tRNA synthetases. Eur J Biochem 126 (1):135-142. doi:10.1111/j.1432-1033.1982.tb06757.x
74. Fraga H, Fontes R (2011) Enzymatic synthesis of mono and dinucleoside polyphosphates. Biochim Biophys Acta 1810 (12):1195-1204. doi:10.1016/j.bbagen.2011.09.010
75. Petersen LM, Tisa LS (2014) Molecular characterization of protease activity in Serratia sp. strain SCBI and its importance in cytotoxicity and virulence. J Bacteriol 196 (22):3923-3936. doi:10.1128/JB.01908-14
76. Kinscherf TG, Willis DK (2002) Global regulation by gidA in Pseudomonas syringae. J Bacteriol 184 (8):2281-2286. doi:10.1128/jb.184.8.2281-2286.2002
77. Shippy DC, Fadl AA (2015) RNA modification enzymes encoded by the gid operon: Implications in biology and virulence of bacteria. Microb Pathog 89:100-107. doi:10.1016/j.micpath.2015.09.008
78. Shin JH, Lee HW, Kim SM, Kim J (2009) Proteomic analysis of Acinetobacter baumannii in biofilm and planktonic growth mode. J Microbiol 47 (6):728-735. doi:10.1007/s12275-009-0158-y

79. Zhang W, Zhao Z, Zhang B, Wu XG, Ren ZG, Zhang LQ (2014) Posttranscriptional regulation of 2,4-diacetylphloroglucinol production by GidA and TrmE in *Pseudomonas fluorescens* 2P24. *Appl Environ Microbiol* 80 (13):3972-3981. doi:10.1128/AEM.00455-14
80. Li D, Shibata Y, Takeshita T, Yamashita Y (2014) A novel gene involved in the survival of *Streptococcus mutans* under stress conditions. *Appl Environ Microbiol* 80 (1):97-103. doi:10.1128/AEM.02549-13
81. Yi H, Lee H, Cho KH, Kim HS (2018) Mutations in MetG (methionyl-tRNA synthetase) and TrmD [tRNA (guanine-N1)-methyltransferase] conferring meropenem tolerance in *Burkholderia thailandensis*. *J Antimicrob Chemother* 73 (2):332-338. doi:10.1093/jac/dkx378
82. Brauner A, Fridman O, Gefen O, Balaban NQ (2016) Distinguishing between resistance, tolerance and persistence to antibiotic treatment. *Nat Rev Microbiol* 14 (5):320-330. doi:10.1038/nrmicro.2016.34
83. Fridman O, Goldberg A, Ronin I, Shores N, Balaban NQ (2014) Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. *Nature* 513 (7518):418-421. doi:10.1038/nature13469
84. Bolt EL, Jenkins T, Russo VM, Ahmed S, Cavey J, Cass SD (2015) Identification of *Escherichia coli* ygaQ and rpmG as novel mitomycin C resistance factors implicated in DNA repair. *Biosci Rep* 36 (1):e00290. doi:10.1042/BSR20150249
85. Smirnova GV, Oktyabrsky ON (2005) Glutathione in bacteria. *Biochemistry (Mosc)* 70 (11):1199-1211. doi:10.1007/s10541-005-0248-3
86. Jamieson DJ (1998) Oxidative stress responses of the yeast *Saccharomyces cerevisiae*. *Yeast* 14 (16):1511-1527. doi:10.1002/(SICI)1097-0061(199812)14:16<1511::AID-YEA356>3.0.CO;2-S
87. Van Laar TA, Esani S, Birges TJ, Hazen B, Thomas JM, Rawat M (2018) *Pseudomonas aeruginosa* gshA Mutant Is Defective in Biofilm Formation, Swarming, and Pyocyanin Production. *mSphere* 3 (2). doi:10.1128/mSphere.00155-18
88. Wongsaroj L, Saninjuk K, Romsang A, Duang-Nkern J, Trinachartvanit W, Vattanaviboon P, Mongkolsuk S (2018) *Pseudomonas aeruginosa* glutathione biosynthesis genes play multiple roles in stress protection, bacterial virulence and biofilm formation. *PLoS One* 13 (10):e0205815. doi:10.1371/journal.pone.0205815
89. Gomez MJ, Neyfakh AA (2006) Genes involved in intrinsic antibiotic resistance of *Acinetobacter baylyi*. *Antimicrob Agents Chemother* 50 (11):3562-3567. doi:10.1128/AAC.00579-06
90. Feinbaum RL, Urbach JM, Liberati NT, Djonovic S, Adonizio A, Carvunis AR, Ausubel FM (2012) Genome-wide identification of *Pseudomonas aeruginosa* virulence-related genes using a

1113 Caenorhabditis elegans infection model. PLoS Pathog 8 (7):e1002813.
 1114 doi:10.1371/journal.ppat.1002813

1115 91. Song M, Husain M, Jones-Carson J, Liu L, Henard CA, Vazquez-Torres A (2013) Low-molecular-
 1116 weight thiol-dependent antioxidant and antinitrosative defences in Salmonella pathogenesis.
 1117 Mol Microbiol 87 (3):609-622. doi:10.1111/mmi.12119

1118 92. Westley J (1981) Thiosulfate: cyanide sulfurtransferase (rhodanese). Methods Enzymol 77:285-
 1119 291. doi:10.1016/s0076-6879(81)77039-3

1120 93. Donadio S, Shafiee A, Hutchinson CR (1990) Disruption of a rhodaneselike gene results in cysteine
 1121 auxotrophy in Saccharopolyspora erythraea. J Bacteriol 172 (1):350-360.
 1122 doi:10.1128/jb.172.1.350-360.1990

1123 94. Pagani S, Bonomi F, Cerletti P (1984) Enzymic synthesis of the iron-sulfur cluster of spinach
 1124 ferredoxin. Eur J Biochem 142 (2):361-366. doi:10.1111/j.1432-1033.1984.tb08295.x

1125 95. Bonomi F, Pagani S, Kurtz DM, Jr. (1985) Enzymic synthesis of the 4Fe-4S clusters of Clostridium
 1126 pasteurianum ferredoxin. Eur J Biochem 148 (1):67-73. doi:10.1111/j.1432-1033.1985.tb08808.x

1127 96. Navidifar T, Amin M, Rashno M (2019) Effects of sub-inhibitory concentrations of meropenem
 1128 and tigecycline on the expression of genes regulating pili, efflux pumps and virulence factors
 1129 involved in biofilm formation by Acinetobacter baumannii. Infect Drug Resist 12:1099-1111.
 1130 doi:10.2147/IDR.S199993

1131 97. Yang CH, Su PW, Moi SH, Chuang LY (2019) Biofilm Formation in Acinetobacter Baumannii:
 1132 Genotype-Phenotype Correlation. Molecules 24 (10). doi:10.3390/molecules24101849

1133 98. Gaddy JA, Tomaras AP, Actis LA (2009) The Acinetobacter baumannii 19606 OmpA protein plays a
 1134 role in biofilm formation on abiotic surfaces and in the interaction of this pathogen with
 1135 eukaryotic cells. Infect Immun 77 (8):3150-3160. doi:10.1128/IAI.00096-09

1136 99. Confer AW, Ayalew S (2013) The OmpA family of proteins: roles in bacterial pathogenesis and
 1137 immunity. Vet Microbiol 163 (3-4):207-222. doi:10.1016/j.vetmic.2012.08.019

1138 100. Skerniskyte J, Karazijaite E, Deschamps J, Krasauskas R, Briandet R, Suziedeliene E (2019) The
 1139 Mutation of Conservative Asp268 Residue in the Peptidoglycan-Associated Domain of the OmpA
 1140 Protein Affects Multiple Acinetobacter baumannii Virulence Characteristics. Molecules 24 (10).
 1141 doi:10.3390/molecules24101972

1142 101. Insua JL, Llobet E, Moranta D, Perez-Gutierrez C, Tomas A, Garmendia J, Bengoechea JA (2013)
 1143 Modeling Klebsiella pneumoniae pathogenesis by infection of the wax moth Galleria mellonella.
 1144 Infect Immun 81 (10):3552-3565. doi:10.1128/IAI.00391-13

1145 102. Justice SS, Hunstad DA, Cegelski L, Hultgren SJ (2008) Morphological plasticity as a bacterial
 1146 survival strategy. Nat Rev Microbiol 6 (2):162-168. doi:10.1038/nrmicro1820

- 1147 103. Robertson GT, Case ED, Dobbs N, Ingle C, Balaban M, Celli J, Norgard MV (2014)
1148 FTT0831c/FTL_0325 contributes to *Francisella tularensis* cell division, maintenance of cell shape,
1149 and structural integrity. *Infect Immun* 82 (7):2935-2948. doi:10.1128/IAI.00102-14
- 1150 104. Egan AJF (2018) Bacterial outer membrane constriction. *Mol Microbiol* 107 (6):676-687.
1151 doi:10.1111/mmi.13908
- 1152 105. Kim SW, Choi CH, Moon DC, Jin JS, Lee JH, Shin JH, Kim JM, Lee YC, Seol SY, Cho DT, Lee JC (2009)
1153 Serum resistance of *Acinetobacter baumannii* through the binding of factor H to outer
1154 membrane proteins. *FEMS Microbiol Lett* 301 (2):224-231. doi:10.1111/j.1574-
1155 6968.2009.01820.x
- 1156 106. Choi CH, Lee JS, Lee YC, Park TI, Lee JC (2008) *Acinetobacter baumannii* invades epithelial cells
1157 and outer membrane protein A mediates interactions with epithelial cells. *BMC Microbiol* 8:216.
1158 doi:10.1186/1471-2180-8-216
- 1159 107. Kwon HI, Kim S, Oh MH, Na SH, Kim YJ, Jeon YH, Lee JC (2017) Outer membrane protein A
1160 contributes to antimicrobial resistance of *Acinetobacter baumannii* through the OmpA-like
1161 domain. *J Antimicrob Chemother* 72 (11):3012-3015. doi:10.1093/jac/dkx257
- 1162 108. Rumbo-Feal S, Perez A, Ramelot TA, Alvarez-Fraga L, Vallejo JA, Beceiro A, Ohneck EJ, Arivett BA,
1163 Merino M, Fiester SE, Kennedy MA, Actis LA, Bou G, Poza M (2017) Contribution of the A.
1164 *baumannii* A1S_0114 Gene to the Interaction with Eukaryotic Cells and Virulence. *Front Cell*
1165 *Infect Microbiol* 7:108. doi:10.3389/fcimb.2017.00108
- 1166 109. Rumbo-Feal S, Gomez MJ, Gayoso C, Alvarez-Fraga L, Cabral MP, Aransay AM, Rodriguez-
1167 Ezpeleta N, Fullaondo A, Valle J, Tomas M, Bou G, Poza M (2013) Whole transcriptome analysis
1168 of *Acinetobacter baumannii* assessed by RNA-sequencing reveals different mRNA expression
1169 profiles in biofilm compared to planktonic cells. *PLoS One* 8 (8):e72968.
1170 doi:10.1371/journal.pone.0072968
- 1171 110. Chang KC, Kuo HY, Tang CY, Chang CW, Lu CW, Liu CC, Lin HR, Chen KH, Liou ML (2014)
1172 Transcriptome profiling in imipenem-selected *Acinetobacter baumannii*. *BMC Genomics* 15:815.
1173 doi:10.1186/1471-2164-15-815
- 1174 111. Gebhardt MJ, Gallagher LA, Jacobson RK, Usacheva EA, Peterson LR, Zurawski DV, Shuman HA
1175 (2015) Joint Transcriptional Control of Virulence and Resistance to Antibiotic and Environmental
1176 Stress in *Acinetobacter baumannii*. *MBio* 6 (6):e01660-01615. doi:10.1128/mBio.01660-15
- 1177 112. Blaschke U, Suwono B, Zafari S, Ebersberger I, Skiebe E, Jeffries CM, Svergun DI, Wilharm G
1178 (2018) Recombinant production of A1S_0222 from *Acinetobacter baumannii* ATCC 17978 and
1179 confirmation of its DNA-(adenine N6)-methyltransferase activity. *Protein Expr Purif* 151:78-85.
1180 doi:10.1016/j.pep.2018.06.009

113. Roberts RJ, Vincze T, Posfai J, Macelis D (2015) REBASE-a database for DNA restriction and modification: enzymes, genes and genomes. *Nucleic Acids Research* 43 (D1):D298-D299. doi:10.1093/nar/gku1046
114. Wion D, Casadesus J (2006) N6-methyl-adenine: an epigenetic signal for DNA-protein interactions. *Nat Rev Microbiol* 4 (3):183-192. doi:10.1038/nrmicro1350
115. Adhikari S, Curtis PD (2016) DNA methyltransferases and epigenetic regulation in bacteria. *FEMS Microbiol Rev* 40 (5):575-591. doi:10.1093/femsre/fuw023
116. Marinus MG, Morris NR (1973) Isolation of deoxyribonucleic acid methylase mutants of *Escherichia coli* K-12. *J Bacteriol* 114 (3):1143-1150
117. Collier J (2009) Epigenetic regulation of the bacterial cell cycle. *Curr Opin Microbiol* 12 (6):722-729. doi:10.1016/j.mib.2009.08.005
118. Salzer R, Kern T, Joos F, Averhoff B (2016) The *Thermus thermophilus* comEA/comEC operon is associated with DNA binding and regulation of the DNA translocator and type IV pili. *Environ Microbiol* 18 (1):65-74. doi:10.1111/1462-2920.12820
119. Rabinovich L, Sigal N, Borovok I, Nir-Paz R, Herskovits AA (2012) Prophage excision activates *Listeria* competence genes that promote phagosomal escape and virulence. *Cell* 150 (4):792-802. doi:10.1016/j.cell.2012.06.036
120. Yoshida A, Kuramitsu HK (2002) Multiple *Streptococcus mutans* Genes Are Involved in Biofilm Formation. *Appl Environ Microbiol* 68 (12):6283-6291. doi:10.1128/aem.68.12.6283-6291.2002
121. Lees-Miller RG, Iwashiki JA, Scott NE, Seper A, Vinogradov E, Schild S, Feldman MF (2013) A common pathway for O-linked protein-glycosylation and synthesis of capsule in *Acinetobacter baumannii*. *Mol Microbiol* 89 (5):816-830. doi:10.1111/mmi.12300
122. Russo TA, Luke NR, Beanan JM, Olson R, Saubaran SL, MacDonald U, Schultz LW, Umland TC, Campagnari AA (2010) The K1 capsular polysaccharide of *Acinetobacter baumannii* strain 307-0294 is a major virulence factor. *Infect Immun* 78 (9):3993-4000. doi:10.1128/IAI.00366-10
123. Schaper S, Wendt H, Bamberger J, Sieber V, Schmid J, Becker A (2019) A Bifunctional UDP-Sugar 4-Epimerase Supports Biosynthesis of Multiple Cell Surface Polysaccharides in *Sinorhizobium meliloti*. *J Bacteriol* 201 (10). doi:10.1128/JB.00801-18
124. Nesper J, Lauriano CM, Klose KE, Kapfhammer D, Kraiss A, Reidl J (2001) Characterization of *Vibrio cholerae* O1 El tor galU and galE mutants: influence on lipopolysaccharide structure, colonization, and biofilm formation. *Infect Immun* 69 (1):435-445. doi:10.1128/IAI.69.1.435-445.2001

125. Chai Y, Beauregard PB, Vlamakis H, Losick R, Kolter R (2012) Galactose metabolism plays a crucial role in biofilm formation by *Bacillus subtilis*. *MBio* 3 (4):e00184-00112. doi:10.1128/mBio.00184-12
126. Niou YK, Wu WL, Lin LC, Yu MS, Shu HY, Yang HH, Lin GH (2009) Role of galE on biofilm formation by *Thermus* spp. *Biochem Biophys Res Commun* 390 (2):313-318. doi:10.1016/j.bbrc.2009.09.120
127. Zou Y, Feng S, Xu C, Zhang B, Zhou S, Zhang L, He X, Li J, Yang Z, Liao M (2013) The role of galU and galE of *Haemophilus parasuis* SC096 in serum resistance and biofilm formation. *Vet Microbiol* 162 (1):278-284. doi:10.1016/j.vetmic.2012.08.006
128. Nakao R, Senpuku H, Watanabe H (2006) *Porphyromonas gingivalis* galE is involved in lipopolysaccharide O-antigen synthesis and biofilm formation. *Infect Immun* 74 (11):6145-6153. doi:10.1128/IAI.00261-06
129. Cabral MP, Soares NC, Aranda J, Parreira JR, Rumbo C, Poza M, Valle J, Calamia V, Lasa I, Bou G (2011) Proteomic and functional analyses reveal a unique lifestyle for *Acinetobacter baumannii* biofilms and a key role for histidine metabolism. *J Proteome Res* 10 (8):3399-3417. doi:10.1021/pr101299j
130. Chateau A, Lunderberg JM, Oh SY, Abshire T, Friedlander A, Quinn CP, Missiakas DM, Schneewind O (2018) Galactosylation of the Secondary Cell Wall Polysaccharide of *Bacillus anthracis* and Its Contribution to Anthrax Pathogenesis. *J Bacteriol* 200 (5). doi:10.1128/JB.00562-17
131. Zeng Y, He Y, Wang KY, Wang J, Zeng YK, Chen YX, Chen D, Geng Y, OuYang P (2016) cpsJ gene of *Streptococcus iniae* is involved in capsular polysaccharide synthesis and virulence. *Antonie Van Leeuwenhoek* 109 (11):1483-1492. doi:10.1007/s10482-016-0750-1
132. Remy E, Meyer M, Blaise F, Simon UK, Kuhn D, Balesdent MH, Rouxel T (2009) A key enzyme of the Leloir pathway is involved in pathogenicity of *Leptosphaeria maculans* toward oilseed rape. *Mol Plant Microbe Interact* 22 (6):725-736. doi:10.1094/MPMI-22-6-0725
133. Anton DN (1995) Resistance to mecillinam produced by the co-operative action of mutations affecting lipopolysaccharide, spoT, and cya or crp genes of *Salmonella typhimurium*. *Mol Microbiol* 16 (3):587-595. doi:10.1111/j.1365-2958.1995.tb02421.x
134. Hickman FW, Rhoden DL, Esaias AO, Baron LS, Brenner DJ, Farmer JJ, 3rd (1982) Evaluation of two *Salmonella typhi* strains with reduced virulence for use in teaching and proficiency testing. *J Clin Microbiol* 15 (6):1085-1091

135. Carlson-Banning KM, Chou A, Liu Z, Hamill RJ, Song Y, Zechiedrich L (2013) Toward repurposing
ciclopirox as an antibiotic against drug-resistant *Acinetobacter baumannii*, *Escherichia coli*, and
Klebsiella pneumoniae. *PLoS One* 8 (7):e69646. doi:10.1371/journal.pone.0069646
136. Dey S, Lane JM, Lee RE, Rubin EJ, Sacchettini JC (2010) Structural characterization of the
Mycobacterium tuberculosis biotin biosynthesis enzymes 7,8-diaminopelargonic acid synthase
and dethiobiotin synthetase. *Biochemistry* 49 (31):6746-6760. doi:10.1021/bi902097j
137. Salaemae W, Azhar A, Booker GW, Polyak SW (2011) Biotin biosynthesis in *Mycobacterium*
tuberculosis: physiology, biochemistry and molecular intervention. *Protein Cell* 2 (9):691-695.
doi:10.1007/s13238-011-1100-8
138. Said HM (2009) Cell and molecular aspects of human intestinal biotin absorption. *J Nutr* 139
(1):158-162. doi:10.3945/jn.108.092023
139. Woong Park S, Klotzsche M, Wilson DJ, Boshoff HI, Eoh H, Manjunatha U, Blumenthal A, Rhee K,
Barry CE, 3rd, Aldrich CC, Ehrt S, Schnappinger D (2011) Evaluating the sensitivity of
Mycobacterium tuberculosis to biotin deprivation using regulated gene expression. *PLoS Pathog*
7 (9):e1002264. doi:10.1371/journal.ppat.1002264
140. Mu X, Wang N, Li X, Shi K, Zhou Z, Yu Y, Hua X (2016) The Effect of Colistin Resistance-Associated
Mutations on the Fitness of *Acinetobacter baumannii*. *Front Microbiol* 7:1715.
doi:10.3389/fmicb.2016.01715
141. Shirai H, Mizuguchi K (2003) Prediction of the structure and function of AstA and AstB, the first
two enzymes of the arginine succinyltransferase pathway of arginine catabolism. *FEBS Lett* 555
(3):505-510. doi:10.1016/s0014-5793(03)01314-0
142. Peleg AY, Jara S, Monga D, Eliopoulos GM, Moellering RC, Jr., Mylonakis E (2009) *Galleria*
mellonella as a model system to study *Acinetobacter baumannii* pathogenesis and therapeutics.
Antimicrob Agents Chemother 53 (6):2605-2609. doi:10.1128/AAC.01533-08
143. Jacobs AC, Thompson MG, Black CC, Kessler JL, Clark LP, McQueary CN, Gancz HY, Corey BW,
Moon JK, Si Y, Owen MT, Hallock JD, Kwak YI, Summers A, Li CZ, Rasko DA, Penwell WF, Honnold
CL, Wise MC, Waterman PE, Lesho EP, Stewart RL, Actis LA, Palys TJ, Craft DW, Zurawski DV
(2014) AB5075, a Highly Virulent Isolate of *Acinetobacter baumannii*, as a Model Strain for the
Evaluation of Pathogenesis and Antimicrobial Treatments. *MBio* 5 (3):e01076-01014.
doi:10.1128/mBio.01076-14
144. Harms K, Lunnan A, Hultner N, Mourier T, Vinner L, Andam CP, Marttinen P, Fridholm H, Hansen
AJ, Hanage WP, Nielsen KM, Willerslev E, Johnsen PJ (2016) Substitutions of short heterologous
DNA segments of intragenomic or extragenomic origins produce clustered genomic
polymorphisms. *Proc Natl Acad Sci U S A* 113 (52):15066-15071. doi:10.1073/pnas.1615819114

1279 145. Domingues S, Harms K, Fricke WF, Johnsen PJ, da Silva GJ, Nielsen KM (2012) Natural
1280 transformation facilitates transfer of transposons, integrons and gene cassettes between
1281 bacterial species. PLoS Pathog 8 (8):e1002837. doi:10.1371/journal.ppat.1002837
1282

1 **Table 1. Summary of experimental results on genes involved in surface-associated motility in *A. baumannii* ATCC 17978**

Locus tag in ATCC 17978	Annotation/gene name	Predicted function	Motility deficiency ^[a]	Growth deficiency ^[b]	Attenuation in <i>Galleria mellonella</i> infection ^[c]	Pellicle biofilm formation ^[d]	MIC values ^[e]		
							Ampicillin	Imipenem	Tetracycline

Purine/pyrimidine/folate biosynthesis

A1S_2187	<i>purH</i>	phosphoribosylaminoimidazolecarboxamide formyltransferase (purine synthesis)	+	Y	N.s.	↑**	N.s.	N.s.	N.s.
A1S_2251	<i>purF</i>	amidophosphoribosyltransferase (purine synthesis)	++	Y	N.s.	N.s.	R ***	N.s.	R *
A1S_2605	<i>purM</i>	phosphoribosylaminoimidazole synthetase (purine synthesis)	+++	Y	*	↑*	R ***	N.s.	S *
A1S_2964	<i>purE</i>	phosphoribosylaminoimidazole	+++	Y	N.s.	↑***	R ***	S **	S *

A1S_1566		carboxylase mutase subunit (purine synthesis)	+++	Y	**	↓ ****	R ****	N.s.	N.s.
		6-pyruvoyl-tetrahydropterin synthase (folate biosynthesis)							
A1S_2687	<i>carB</i>	carbamoylphosphate synthase subunit (pyrimidine synthesis)	++	Y	****	↓ ****	N.s.	S ***	S **

Alarmones/ stress metabolism

A1S_0414		Ap5A pyrophosphatase	++	Y	*	↓ ****	R ****	N.s.	N.s.
A1S_1624		Ap4A hydrolase	++	Y	N.s.	N.s.	N.s.	S ***	N.s.

RNA modification/ regulation

A1S_0447	<i>rpmG</i>	50S ribosomal protein L33	+++	Y	N.s.	N.s.	N.s.	S *	S **
A1S_0778	<i>metG</i>	methionyl-tRNA synthetase	+++	Y	****	↓ ****	N.s.	N.s.	S ***
A1S_2182	<i>gidA</i>	Glucose-inhibited division protein A, FAD-binding protein	++	Y	N.s.	↓ ****	R *	N.s.	N.s.

Oxidative stress

A1S_0530		rhodanese domain-containing protein, sulfurtransferase	++	N	N.s.	↑****	N.s.	N.s.	N.s.
A1S_3366	<i>gshA</i>	gamma-glutamate-cysteine ligase	++	Y	*	N.s.	R ***	N.s.	N.s.

Outer membrane proteins

A1S_1970		outer membrane protein (Omp85 family)	++	Y	**	↓**	R ****	S *	N.s.
A1S_2840	<i>ompA</i>	outer membrane protein	++	N	****	↓****	N.s.	R ***	S *
A1S_3297		putative outer membrane protein	+	Y	**	↑**	R ***	R *	N.s.

1,3-diaminopropane biosynthesis

A1S_2453	<i>ddc</i>	L-2,4-diaminobutyrate decarboxylase, biosynthesis of 1,3-diaminopropane (DAP)	+++	Y	N.s.	↓****	N.s.	N.s.	S ***
A1S_2454	<i>dat</i>	L-2,4-diaminobutyrate:2-	+++	Y	*	↓****	N.s.	N.s.	S ***

		ketoglutarate 4-aminotransferase, biosynthesis of 1,3-diaminopropane (DAP)							

Lipopeptide synthesis/export

A1S_0113		acyl-CoA dehydrogenase (putative lipoprotein biosynthesis)	++	N	N.s.	↓**	R *	N.s.	N.s.
A1S_0116		RND superfamily transporter (Efflux pump)	++	N	N.s.	↓**	R ***	N.s.	N.s.

DNA modification/repair/uptake

A1S_0222	<i>aamA</i>	adenine-specific methyltransferase	++	Y	***	↓****	N.s.	S ***	S **
A1S_2334	<i>sahH</i>	S-adenosyl-L-homocysteine hydrolase	+++	Y	*	N.s.	R ***	N.s.	N.s.
A1S_2587	<i>ruvA</i>	holliday junction helicase subunit A	++	Y	N.s.	↓****	R ***	N.s.	S **
A1S_2610	<i>comEC</i>	competence factor	++	N	**	N.s.	N.s.	N.s.	N.s.

Others

A1S_0065	<i>galE</i>	UDP-glucose 4-epimerase, Lipopolysaccharide biosynthesis	++	N	****	↑****	R ***	N.s.	R *
A1S_0806		adenosylmethionine-8-amino-7-oxononanoate aminotransferase	+++	Y	N.s.	↑****	R ***	R ****	N.s.
A1S_1055		soluble lytic murein transglycosylase	++	N	**	↑**	N.s.	N.s.	S *
A1S_2761	<i>prpF</i>	2-methylaconitate isomerase	+++	Y	N.s.	↓****	R *	N.s.	S **
A1S_3026		Hyp. Secreted Ribonuclease T2 (predicted secretion signal)	++	N	N.s.	N.s.	R ****	R ***	N.s.
A1S_3129	<i>astB</i>	succinylarginine dihydrolase	+++	Y	N.s.	↓****	R ***	N.s.	S **

4 ^[a] Compared to ATCC 17978 wild type strain (WT); diameter of the spreading zone was measured: '+++', 0-4 mm; '++', < 4-9 mm; '+', < 9 mm

5 ^[b] Comparison of bacterial growth curves. Y, growth deficiency compared to WT was observed; N, no growth deficiency was observed

6 ^[c] Compared to ATCC 17978 WT; unpaired t-test was performed after 5 days p.i.: N.s., not significant; *, p-value ≤ 0.05; **, p-value ≤ 0.01; ***, p-value ≤ 0.001;

7 ****, p-value ≤ 0.0001

8 ^[d] Compared to ATCC 17978 WT; unpaired t-test was performed: N.s., not significant; *, p-value ≤ 0.05; **, p-value ≤ 0.01; ***, p-value ≤ 0.001; ****, p-value ≤
9 0.0001; ↑, more than ATCC 17978 WT; ↓, less than ATCC 17978 WT

10 ^[e] Compared to ATCC 17978 WT; unpaired t-test was performed: N.s., not significant; *, p-value ≤ 0.05; **, p-value ≤ 0.01; ***, p-value ≤ 0.001; ****, p-value ≤
11 0.0001; R, resistant; S, susceptible

Table 2. Summary of experimental results on genes involved in surface-associated motility in *A. baumannii* 29D2. A dark grey background indicates concordance to results obtained for strain ATCC 17978

Locus tag in ATCC 17978	Annotation/gene name	Motility deficiency ^[a]	Growth deficiency ^[b]	Attenuation in <i>Galleria mellonella</i> infection ^[c]	Pellicle biofilm formation ^[d]	MIC values ^[e]		
						Ampicillin	Imipenem	Tetracycline

Purine/pyrimidine/folate biosynthesis

A1S_2187	<i>purH</i>	++	Y	*	↑*	S *	N.s.	N.s.
A1S_2251	<i>purF</i>	++	Y	N.s.	N.s.	S **	S *	N.s.
A1S_2605	<i>purM</i>	+	Y	N.s.	N.s.	S *	N.s.	N.s.
A1S_2964	<i>purE</i>	+	Y	N.s.	N.s.	S **	N.s.	N.s.
A1S_1566		+	Y	***	N.s.	S *	N.s.	S *
A1S_2687	<i>carB</i>	+	Y	****	↓**	S **	S **	N.s.

Alarmones/stress metabolism

A1S_0414		+	N	N.s.	N.s.	N.s.	N.s.	N.s.

RNA modification/ regulation

A1S_2182	<i>gidA</i>	+	N	N.s.	↓*	S *	N.s.	N.s.

Oxidative stress

A1S_3366	<i>gshA</i>	+	Y	N.s.	N.s.	S *	N.s.	N.s.

Outer membrane proteins

A1S_1970		+	N	N.s.	↓**	N.s.	N.s.	N.s.
A1S_2840	<i>ompA</i>	+	Y	****	N.s.	S *	S **	N.s.

1,3-diaminopropane biosynthesis

A1S_2453	<i>ddc</i>	++	Y	*	↓**	S **	S *	N.s.
A1S_2454	<i>dat</i>	+	Y	N.s.	↓*	S **	N.s.	R *

Lipopeptide synthesis/export

A1S_0116		+	N	N.s.	N.s.	N.s.	N.s.	N.s.

DNA modification/repair/uptake

A1S_0222	<i>aamA</i>	+	N	*	N.s.	N.s.	R **	N.s.
A1S_2610	<i>comEC</i>	+	Y	****	N.s.	S **	S **	S *

Others

A1S_0065	<i>galE</i>	++	Y	****	↑*	S *	N.s.	N.s.
A1S_0806		+	N	N.s.	N.s.	S *	S ****	N.s.
A1S_1055		+	N	**	N.s.	N.s.	S *	N.s.
A1S_2761	<i>prpF</i>	+	Y	**	N.s.	S *	S *	S *
A1S_3026		+	N	**	↓*	R *	R ***	N.s.

^[a] Compared to 29D2 wild type strain (WT); diameter of the spreading zone was measured: '+++', 0-3 mm; '++', < 3-6 mm; '+', < 6 mm

^[b] Comparison of bacterial growth curves. Y, growth deficiency compared to WT was observed; N, no growth deficiency was observed

^[c] Compared to 29D2 WT; unpaired t-test was performed after 5 days p.i.: n.s., not significant; *, p-value ≤ 0.05 ; **, p-value ≤ 0.01 ; ***, p-value ≤ 0.001 ; ****, p-value ≤ 0.0001

^[d] Compared to 29D2 WT; unpaired t-test was performed: n.s., not significant; *, p-value ≤ 0.05 ; **, p-value ≤ 0.01 ; ***, p-value ≤ 0.001 ; ****, p-value ≤ 0.0001 ; ↑, more than 29D2 WT; ↓, less than 29D2 WT

^[e] Compared to 29D2 WT; unpaired t-test was performed: N.s., not significant; *, p-value ≤ 0.05 ; **, p-value ≤ 0.01 ; ***, p-value ≤ 0.001 ; ****, p-value ≤ 0.0001 ; R, resistant; S, susceptible

Table 3. Minimal inhibitory concentration (MIC) of ampicillin, tetracycline and imipenem determined from ATCC 17978 wildtype/mutants and 29D2 wildtype/mutants. A dark grey background indicates that MIC values of mutant strains are significantly elevated compared to the wildtype while a light grey background indicates increased susceptibility

Locus tag	Gene name	Ampicillin ^a		Imipenem ^a		Tetracycline ^a	
		ATCC 17978	29D2	ATCC 17978	29D2	ATCC 17978	29D2
Wildtype		25.3	36.5	0.23	0.25	2.1	3.25
A1S_2187	<i>purH</i>	32	18.6	0.23	0.21	1.6	1.5
A1S_2251	<i>purF</i>	48	16.0	0.25	0.16	3.0	2.3
A1S_2605	<i>purM</i>	53.3	17.3	0.21	0.18	1.0	2.3
A1S_2964	<i>purE</i>	53.3	10.6	0.13	0.21	1.5	2.0
A1S_1566		96	18.6	0.23	0.23	1.8	0.9
A1S_2687	<i>carB</i>	32	3.0	0.10	0.03	0.9	2.0
A1S_0414		106.6	32.0	0.23	0.25	1.6	2.0
A1S_1624		32	-	0.125	-	2.1	-
A1S_0447	<i>rpmG</i>	29.3	-	0.16	-	1.0	-
A1S_0778	<i>metG</i>	32	-	0.23	-	0.46	-
A1S_0530		24	-	0.19	-	2.0	-
A1S_3366	<i>gshA</i>	85.3	26.6	0.29	0.23	1.6	2.0
A1S_1970		64	32.0	0.16	0.25	2.5	1.8

A1S_2840	<i>ompA</i>	29.3	13.3	0.46	0.13	1.5	3.0
A1S_3297		58.6	-	0.33	-	2.0	-
A1S_2453	<i>ddc</i>	32	5.0	0.46	0.14	0.38	4.6
A1S_2454	<i>dat</i>	26.6	5.3	0.25	0.23	0.29	6.0
A1S_0113		42.6	-	0.25	-	2.0	-
A1S_0116		85.3	29.3	0.21	0.33	2.6	1.5
A1S_0222	<i>aamA</i>	18.6	42.6	0.125	0.38	0.9	2.6
A1S_0065	<i>galE</i>	53.3	13.3	0.23	0.18	3.0	2.0
A1S_0806		53.3	13.3	0.38	0.023	2.0	1.6
A1S_1055		26.6	21.3	0.29	0.14	1.3	1.6
A1S_2182	<i>gidA</i>	37.3	24.0	0.19	0.25	3.0	4.0
A1S_2334	<i>sahH</i>	74.6	-	0.18	-	1.8	-
A1S_2587	<i>ruvA</i>	53.3	-	0.19	-	1.0	-
A1S_2610	<i>comEC</i>	29.3	5.3	0.23	0.10	1.6	0.6
A1S_2761	<i>prpF</i>	42.6	14.6	0.25	0.16	0.8	0.5
A1S_3026		106.6	64.0	0.42	0.46	2.6	1.5
A1S_3129	<i>astB</i>	74.6	-	0.23	-	1.1	-

^a Averaged MIC values in [µg/mL] determined from three independent experiments. '-' indicates 'not tested'

9 **Table 4. Links between the genes identified in this study and the literature**

Locus tag in ATCC 17978	Annotation/ gene name	Known relationship in other bacteria
<i>Purine/pyrimidine/folate biosynthesis</i>		
A1S_2187	<i>purH</i>	biofilm formation in <i>Bacillus cereus</i> [48]; virulence in <i>B. anthracis</i> [54]; K ⁺ -dependent colony spreading in <i>Bacillus subtilis</i> [45]; <i>Enterococcus faecium</i> growth in human serum [49]; defects in rifampicin persistence in <i>S. aureus</i> [56]
A1S_2251	<i>purF</i>	virulence in <i>A. baumannii</i> [52]; K ⁺ -dependent colony spreading in <i>Bacillus subtilis</i> [45]; virulence in <i>Pasteurella multocida</i> [55]; virulence of <i>Burkholderia cenocepacia</i> in <i>G. mellonella</i> , <i>C. elegans</i> , <i>D. melanogaster</i> infection [51]; defects in rifampicin persistence in <i>S. aureus</i> [56]
A1S_2605	<i>purM</i>	virulence in <i>A. baumannii</i> [52]; K ⁺ -dependent colony spreading in <i>Bacillus subtilis</i> [45]; defects in rifampicin persistence in <i>S. aureus</i> [56]; pellicles in <i>A. baumannii</i> ATCC 17978 [43]
A1S_2964	<i>purE</i>	virulence in <i>S. pneumoniae</i> [53] and <i>A. baumannii</i> [52]; motility (<i>purK</i>) in <i>A. nosocomialis</i> [26]; pellicles in <i>A. baumannii</i> ATCC 17978 (PurB, PurD) [43]
A1S_1566		-
A1S_2687	<i>carB</i>	swimming motility and biofilm formation in <i>Xanthomonas citri</i> subsp. <i>citri</i> [61]; virulence in <i>A. baumannii</i> [52]; growth of <i>E. coli</i> in human serum [50]

Alarmons/ stress metabolite

A1S_0414		-
A1S_1624		motility in <i>E. coli</i> [66]; pellicles in <i>A. baumannii</i> ATCC 17978 [43]; biofilm formation in <i>Pseudomonas fluorescens</i> [70]; virulence in <i>Salmonella enterica</i> [67]; antibiotic susceptibility in <i>E. coli</i> , <i>A. baumannii</i> and <i>P. aeruginosa</i> [68]; antibiotic tolerance [69]

RNA modification/regulation

A1S_0447	<i>rpmG</i>	mitomycin C resistance in <i>E. coli</i> [82]
A1S_0778	<i>metG</i>	virulence in <i>A. baumannii</i> [52]; antibiotic tolerance in <i>Burkholderia thailandensis</i> [79] and <i>E. coli</i> [80,81]; pellicles in <i>A. baumannii</i> ATCC 17978 [43]
A1S_2182	<i>gidA</i>	review <i>gid</i> operon, virulence, motility, biofilm formation, antibiotic resistance, bacterial growth [75]; swarming motility, pellicle biofilm in <i>Bacillus cereus</i> [47]; swarming motility in <i>Serratia</i> species SCBI [73] and <i>Pseudomonas syringae</i> [74]; proteomic analysis in <i>A. baumannii</i> [76]; biofilm formation in <i>Pseudomonas fluorescens</i> [77] and <i>Streptococcus mutans</i> [78]

Oxidative stress

A1S_0530		virulence in <i>Salmonella Typhimurium</i> [144]; thioredoxin involved in <i>A. baumannii</i> virulence [145]; general oxidative stress response genes involved in pellicles in <i>A. baumannii</i> ATCC 17978 [43]
A1S_3366	<i>gshA</i>	swarming and swimming motility in <i>P. aeruginosa</i> , decrease in biofilm formation [85]; swimming and twitching motility in <i>P.</i>

		<i>aeruginosa</i> , increase in biofilm formation [86]; sensitivity to metronidazole and ciprofloxacin in <i>A. baylyi</i> [87]; <i>P. aeruginosa</i> <i>gshA</i> mutant attenuated in <i>C.elegans</i> infection [88]; <i>Salmonella gshA</i> mutant attenuated in murine model [89]

Outer membrane proteins

A1S_1970		-
A1S_2840	<i>ompA</i>	<i>A. nosocomialis</i> surface-associated motility [26]; biofilm formation [94-96]; bacterial pathogenicity – Review [97]; virulence [11,103,104]; <i>A. baumannii</i> virulence in <i>C. elegans</i> [98]; <i>K. pneumoniae</i> virulence in <i>G. mellonella</i> [99]; antibiotic resistance [105]; pellicles in <i>A. baumannii</i> ATCC 17978 [43]
A1S_3297		general outer membrane proteins involved in pellicles in <i>A. baumannii</i> ATCC 17978 [43]

1,3-diaminopropane biosynthesis

A1S_2453	<i>ddc</i>	surface-associated motility and virulence in <i>A. baumannii</i> [33]
A1S_2454	<i>dat</i>	

Lipopeptide synthesis/ export

A1S_0113		temperature dependent antibiotic resistance and surface motility in <i>A. baumannii</i> [146]; surface-associated motility in <i>A. baumannii</i> and <i>A. nosocomialis</i> [26,38]; pellicle biofilm formation in <i>A. baumannii</i> [38]; biofilm formation on abiotic surfaces in <i>A. baumannii</i> [107,106]; pellicles in <i>A. baumannii</i> ATCC 17978 [43]; imipenem-selected <i>A. baumannii</i> [108]
A1S_0116		

DNA modification/repair/uptake

A1S_0222	<i>aamA</i>	protein purification of <i>A. baumannii</i> AamA [110]; review of phenotypes caused by <i>dam</i>

		mutants or <i>dam</i> overexpression [115]
A1S_2334	<i>sahH</i>	biofilm formation [147]
A1S_2587	<i>ruvA</i>	-
A1S_2610	<i>comEC</i>	surface motility, twitching motility, virulence in <i>A. baumannii</i> [16]; twitching motility in <i>Thermus thermophilus</i> [116]; virulence and growth in <i>L. monocytogenes</i> [117]; biofilm formation [118]

Other

A1S_0065	<i>galE</i>	virulence in <i>A. baumannii</i> [52], <i>Bacillus anthracis</i> [128], <i>Streptococcus iniae</i> [129], <i>Leptosphaeria maculans</i> [130]; biofilm formation in <i>Sinorhizobium meliloti</i> [121], <i>Vibrio cholerae</i> [122], <i>Bacillus subtilis</i> [123], <i>Thermus thermophilus</i> [124], <i>Haemophilus parasuis</i> [125], <i>Porphyromonas gingivalis</i> [126], <i>A. baumannii</i> [76]; antibiotic resistance/susceptibility in <i>Porphyromonas gingivalis</i> [126], <i>Salmonella typhimurium</i> [131], <i>Salmonella typhi</i> [132]; <i>A. baumannii</i> biofilm (<i>galU</i> , <i>galM</i>) [127]; surface motility of <i>A. nosocomialis</i> (<i>rmlB</i>) [26]
A1S_0806		survival, growth and virulence of mycobacteria <i>bioA</i> [137,148-150]
A1S_1055		lytic transglycosylase (A1S_3027) in <i>A. nosocomialis</i> motility [26]
A1S_2761	<i>prpF</i>	pellicles in <i>A. baumannii</i> ATCC 17978 [43]
A1S_3026		reviews of T2 Family Ribonucleases [151,152]; abiotic surface colonization in <i>A. baumannii</i> [34]; colistin resistance in <i>A. baumannii</i> [138]
A1S_3129	<i>astB</i>	virulence in <i>A. baumannii</i> [52]; pellicles in <i>A. baumannii</i> ATCC 17978 [43]