Structural colours reflect individual quality: a meta-analysis

Thomas E. White^{1,2}

¹School of Life and Environmental Sciences, The University of Sydney, Sydney, Australia 2106

² Corresponding author.

E-mail: thomas.white@sydney.edu.au

Keywords: condition dependent, honest signal, communication, iridescence, sexual selection

Word count (excluding abstract): 2489

Number of figures: 1

Number of tables: 1

Supplementary material: Supplementary figures and table

Abstract

Ornamental colouration often communicates salient information to mates, and theory predicts covariance between signal expression and individual quality. This has borne out among pigment-based signals, but the potential for ‘honesty’ in structural colouration is unresolved. Here I synthesised the available evidence to test this prediction via meta-analysis and found that, overall, the expression of structurally coloured sexual signals is positively associated with individual quality. The effects varied by measure of quality, however, with body condition and immune function reliably encoded across taxa, but not age nor parasite resistance. The relationship was apparent for both the colour and brightness of signals and was slightly stronger for iridescent ornaments. These results suggest diverse pathways to the encoding and exchange of information among structural colours, while highlighting outstanding questions as to the development, visual ecology, and evolution of this striking adornment.

Introduction

Colour is a ubiquitous channel of communication in nature and is showcased at an extreme in the service of mate choice [1,2]. A central hypothesis in evolutionary biology is that sexual selection has driven the elaboration of ornamental colouration into reliable indicators of individual quality [3], with empirical tests guided by indicator and handicap models [4,5]. These models argue that conspicuous displays are selectively favoured because their production is differentially costly (handicap) and/or constrained (index) between individuals of varying quality, and so encode honest information to potential mates. A prediction common to honesty-based models is that signals should show heightened condition-dependent expression, and the most robust support to date among ornamental colouration is found in carotenoid-based signals [6,7]. As pigments that cannot be synthesised *de novo*, all carotenoids must ultimately be acquired via diet before being incorporated into signals directly or following bioconversion. This offers ample opportunity for selection to favour mechanistic links between foraging, metabolic performance, and sexual signal expression (that is, the combined perceptual features of hue, saturation, and brightness), which is now well established, at least among birds [8,9]. Relative to our knowledge of pigment-based colouration, however, the potential for structural colours to signal individual quality remains both understudied and poorly resolved.

Unlike pigments, which are selectively absorbent, structural colours result from the selective reflectance of light by nano-structured tissues [10,11]. Accumulating evidence also suggests that the development of these structures is driven by self-assembly — such as the phase separation of keratin and cytoplasm in nascent feather barbs [12-14] — rather than the active (and 'expensive') cellular processes that underlie some pigmentary colour production [8]. Three general arguments have been articulated around their potential for honesty among structural

colouration in sexual signalling. One is that if sufficient material is required to produce nano-architectures then it will establish a trade-off with other physiological needs that may be differentially met among individuals of varying 'quality', as consistent with a handicap explanation [4,6]. A non-exclusive alternative builds on the observation that features of the macro-scale expression of signals relies on the precision with which the underlying structures are organised [15,16]. If individuals heritably vary in their capacity to achieve such organisation as a result of, for example, physiological constraints on signal production, or the behavioural acquisition of stable developmental environments, then signals may serve as an index of underlying genetic quality [5]. Finally, the lack of obvious ecologically relevant material to trade-off against during signal construction, together with the self-assembly inherent in structural colours noted above, has motivated arguments against any general expectations for condition dependence *sensu lato* [12]. Though experimental work is able to partition these hypotheses in some contexts [17], most empirical studies to date have focused on the overarching question of honesty by examining the predicted covariance between fitness-related traits and signal expression. This has provided valuable insight into the central question, but diversity in signal designs, measures of 'quality', and taxonomy have presented a challenge for qualitative synthesis. Modern quantitative methods, however, provide robust tools for controlling for and capitalising on such variation (e.g via mixed-effects models and meta-regression), and so can offer substantive answers to longstanding questions [18].

Here I used phylogenetically controlled meta-analysis and meta-regression to examine whether structural colour signals encode salient information on individual quality. Specifically, I synthesised estimates of correlations between measures of individual quality and signal

expression to test the prediction of condition dependence, before examining methodological and theoretically derived mediators of effect-size variation among studies.

Methods

Literature search and study selection

I conducted a systematic literature search using *Web of Knowledge* and *Scopus* databases for publications up to September 2019, using the query ((colour OR color OR pigment) AND signal AND (quality OR condition OR condition dependent OR condition dependence OR ornament) OR honest*), as well as searching the references of included texts. This produced 3482 unique studies, from which 41 were ultimately suitable for quantitative synthesis following the screening of titles and abstracts (where n = 3430 were excluded for clear irrelevance), and full texts (see Fig. S1 for PRISMA statement). I used the R package ‘revtools’ v0.4.1 for title and abstract screening [19]. I included all experimental and observational studies that quantified the relationship between intersexual structural colour signal expression (via the measurement of hue, saturation, or brightness, or a composite thereof) and any one of age, body condition (size, size-corrected mass, or growth rate), immune function (oxidative damage, PHA response, circulating CORT or testosterone) or parasite resistance as a measure of individual quality. I excluded studies that conflated the structural and pigmentary contributions to signal expression during measurement or manipulation, only studied sexually immature juveniles, focused exclusively on intrasexual signalling, used human-subjective assessments of colouration (such as colour swatches or viewer rankings), or which did not provide adequate data in the form of raw effect sizes, or summary statistics or figures from which effect sizes might be estimated.

Effect size calculation

I used the correlation coefficient, Pearson's r , transformed to Fisher's z (given its preferable normalizing and variance-stabilizing qualities) as the effect size describing the relationship between colouration and individual quality for meta-analysis. These effects were extracted directly from text or figures, using the R package 'metadigitise' v1.0 [20], where possible ($n = 102$), or was otherwise converted from available test statistics or summary data ($n = 84$).

Meta-analyses

I ran both phylogenetic multi-level meta-analytic (intercept-only, MLM) and multi-level meta-regression (MLMR) models, using the package 'metafor' v2.1-0 [21] in R v3.5.2 [22]. Almost all studies reported multiple effects through the estimation of several colour metrics or multiple measures of individual quality, so I included both a study- and observation-level random effect in all models. From my MLM model I estimated a meta-analytic mean (i.e., intercept) effect size, which describes the overall support for the honesty of structural colour signals. I accounted for phylogenetic non-independence between effect sizes in all models by estimating relationships among species using the Open Tree of Life database [23], accessed via the R package 'rotl' v3.0.10 [24]. Given the resulting tree topology, I estimated a correlation matrix from branch lengths derived using Grafen's method [25] assuming node heights raised to the power of 0.5. Though this does not account for evolutionary divergence, it grants an approximate estimate of relatedness by accounting for phylogenetic topology (Fig. S2).

I then used separate MLMR models to examine the effects of moderators, both theoretical and methodological, which may be expected to alter the strength of the signal/quality

relationship. These included the measure of individual quality used—body condition, age, immune function, or parasite resistance (as defined above)—since ‘quality’ is multivariate (discussed below). There is a suite of metrics available for measuring colour, though they typically center on quantifying the perceptually relevant features of hue (the unique colour), saturation (spectral purity), and brightness, or a composite thereof [26]. I therefore classified every measure as such in order to test which, if any, signal features contain salient information on mate quality. In broad terms, the greater nano-structural organisation and/or material required to generate more saturated and (to a lesser extent) brighter signals predicts a positive correlation between these features and individual quality. Signal hue, by contrast, is a directionless measure in the sense that there is no clear biophysical reason for predicting consistent among-individual shifts toward longer or shorter wavelengths as a function of individual quality across taxa, and so I ignored the sign of correlations for estimates of hue alone. I also tested the effect of signal iridescence (i.e. the presence/absence of iridescent colouration), which I coded according to information presented in-text or via secondary sources (including personal observation). The rationale was twofold. For one, all iridescence arises from coherent light-scattering [27]. All things being equal, coherent light-scatterers demand a level of architectural organisation beyond that of incoherent scatterers (i.e. white colours) and possibly non-iridescent colours too, and so offers an indirect test of the hypothesised link between the demands of nano-scale organisation and signal honesty [14,16]. Second, iridescence is an inherently temporal feature of visual communication which may provide an additional or alternate conduit of information to potential mates, above and beyond that which is possible using non-iridescent signals (17,28,29; though this possibility remains unexplored directly). In both cases the prediction is a stronger correlation between colouration and quality among iridescent, as opposed to non-iridescent, ornaments.

Finally, and following the above, I secondarily examined the effects of both quality measures and colour metrics within each of the four taxonomic classes represented across the literature. I focused on these two moderators alone because potential taxonomic variation in the mechanistic links between colouration and individual quality are most likely to manifest via these moderators, and because the limited available data precludes the testing of all moderators on a per-class basis (note that even within these moderators, estimates were not possible across all taxonomic groups).

With respect to methodology I considered study type, given my inclusion of both experimental and observational studies, as well as the sex of focal animals. I also coded whether studies included measurements of non-sexual traits as controls in tests of *heightened* condition-dependence (see discussion). The prediction being that that studies including non-sexual controls may report larger effects than those without, given that many traits will scale with condition to some extent [30]. Note that both signal iridescence and the presence of controls were coded as binary (0/1) for simplicity in testing their respective predictions.

Publication bias

I explored evidence for publication bias by visually inspecting funnel plots of effect sizes versus standard errors (Fig. S3) and using an Egger's test on an intercept-only MLMA that included the random effects described above [31].

Data availability

All data and code are available via GitHub (https://github.com/EaSElab-18/ms_metacol) and are persistently archived through Zenodo (<https://dx.doi.org/10.5281/zenodo.3718617>).

Results

The final dataset comprised 186 effect sizes, across 28 species, from 41 studies [6, 17, 32-71]. Of those 186 effects, 117 were drawn from birds, 22 from insects, 28 from reptiles, and 11 from arachnids (Table S1; Fig. S2). As predicted, I found a positive overall correlation (i.e. meta-analytic mean effect) between individual quality and structural colour signal expression ($Z = 0.1573$, 95% CI = 0.084 to 0.231; Fig. 1; Table 1). The heterogeneity of effect sizes — that is, the variation in effect size estimates after accounting for sampling error — was high ($I^2 = 80.42\%$, 95% CI = 77.26 to 83.01), as is typical of meta-analytic data in ecology and evolutionary biology [72]. A small amount of heterogeneity was explained by among-study effects ($I^2 = 14.21\%$, 95% CI = 8.97 to 20.20), and only a very weak phylogenetic signal was evident ($I^2 = 2.17\%$, 95% CI = 1.18 to 3.47).

Of the measures of quality considered, body condition and immune function were reliably positively correlated with structural colour expression, while age and parasite resistance were not (see Table 1 for all corresponding numerical results henceforth). This varied slightly across taxa, however, with a robust effect of condition on colouration apparent across all groups, while effects of age and parasite resistance were apparent among insects and bird, respectively (table S2; though these estimates are based on limited within- and between-taxa samples). Both the hue and brightness of signals were similarly informative channels on-whole, while chromaticity was not consistently associated with individual quality across taxa (though this varied by taxa; table S2), nor was any correlation apparent when composite measures of colouration were used. Iridescent signals were subject to slightly stronger positive correlations than non-iridescent signals across all measures of condition. Signal honesty was apparent among males only though the weak, borderline effect and much smaller sample among females ($n =$

29/186 versus 146/186 for males) suggests a male bias in the literature similar to that in related fields [73], which may have partly driven this outcome. Experimental studies tended to report marginally stronger correlations than observational assays, which most likely reflects slightly exaggerated experimental manipulations of condition relative to natural variation [30]. Finally, the majority of studies ($n = 36$) did not include measurements of non-sexual control traits in tests of *heightened* condition dependence, though I found no clear difference in effect-size estimates between those that did and did not.

Publication bias

Visual inspection of the funnel plot showed little asymmetry (Fig. S3), as supported by non-significant Egger's tests ($t_{184} = -0.5535$, $p = 0.5806$), which suggests a minimal influence of missing data on effect size estimates.

Discussion

Ornamental colouration may be a reliable conduits of information on mate quality, though evidence for the predicted covariance between signal expression and mate quality among structural, as opposed to pigmentary, signals is equivocal. Here I found meta-analytic support for this link in the form of a positive correlation between structural colour expression and individual quality (Fig. 1), consistent with honesty-based models of sexual signal evolution [4,5]. The strength of the overall correlation, though moderate [74], was commensurate with meta-analytic estimates from pigment-based sexual signals [8,9,75], and suggests that structural colouration may similarly serve a reliable indicator of individual quality.

Quality is a multivariate feature of individuals, and this is reflected in the effect-size variation between measures. Both condition (as narrowly defined above), and proxy measures of immune system integrity were on-average positively correlated with signal expression across all taxa in which those relationships have been examined. This is consistent with experimental work showing that body mass and immune function are responsive to ecologically salient stressors, with consequences for colour production. Among birds, for example, disease and dietary stress produce abnormalities in the keratin barbules that contribute to colouration [16,76,77], while in butterflies the organisation of wing-scale architectures is disrupted by nutritional and environmental stress during pupal (hence, wing-structure) development [36,78]. In contrast, neither age nor parasite resistance were consistently informative of mate quality, though this varied slightly across taxa. These latter measures are often predicated on, or susceptible to, the mechanical degradation of structures post-development. Thus, the inherently heightened variability of sexual signals combined with parasite-induced damage (ectoparasite, in particular) and/or accumulated wear with age, combined with varied mechanisms of colour production of across animal classes, may compound to render the signals less accurate predictors on balance [59,79,80]. Curiously, the near inverse relationship was recently identified in a meta-analysis of carotenoid-based signalling. Weaver et al. [8] examined correlations across similar categories of quality as those used here but found no consistent relationship between signals and either of body condition or immune function. Given the fundamental optical and developmental differences between structural and pigmentary colour production (described above) the potential exists for each to signal unique aspects of individual mate quality, as is suggested by the totality of this work. This has also been directly supported by limited empirical work [65] and may hold

more broadly as an explanation for the often-integrated use of structural and pigmentary mechanisms in sexual colouration.

Colour is often assumed to be the central conduit of information exchange given its relative stability under variable natural illumination [81,82], though my results suggest both the colour and brightness of signals are similarly informative, considering the evidence to date (Fig. 1; Table S2). Furthermore, I identified slightly stronger condition dependence among iridescent, as opposed to non-iridescent, patches. While the underlying architecture varies across taxa, all iridescent colouration arises from coherent light interference and so may demand a level of architectural organisation beyond that of incoherent scattering [11,27], as well as non-iridescent coherent scattering (though evidence for the latter possibility is limited; [14]). Iridescence also introduces temporal structure to signals since the colour appearance depends on the precise arrangement of signals, viewers, and light sources. These combined features may render iridescent colouration particularly suitable as bearers of information [29] and so contribute to the ubiquity of the phenomenon [83,84]. Though only indirectly considered here, as few studies quantify between-individual variation in iridescence itself, this idea has found more immediate support via condition-dependent variation in signal angularity [17], and a predictive relationship between iridescence and mating success [85]. Empirically unravelling the function and perceptual significance of iridescence in the context of sexual signalling—where the effect is seen at its most extreme—remains an active challenge [28]. More generally, these results affirm the view that the extended spectral and temporal repertoire available to structural colours may facilitate the exploration of distinct ‘signalling niches’, with tangible evolutionary consequences [1,54].

By integrating the development of signal structure and fitness-related traits, structural colours may serve as informative signals during mate choice. A holistic understanding, however, awaits progress on several fronts. Most significant is the inclusion of appropriate non-sexual controls. Given that many traits will scale with overall condition, the ultimate evidence for handicap models lies in the demonstration of *heightened* condition-dependence among sexual traits. Though I found no clear difference in effect size estimates between studies with and without such controls the small sample size was limiting, and moreover represents a conceptual limitation that remains pervasive [30]. Partitioning indicator and handicap models of signal evolution and understanding the nature of direct and/or indirect benefits being signalled, are key challenges which requires both experimental and quantitative-genetic study [17]. The development of structural colours during ontogeny is also a central front for progress, and studies among invertebrates (which offer benefits in terms of tractability) would be invaluable in complementing the excellent work accumulating on birds [12-14]. Finally, signalling ecology should remain front-of-mind as accumulating evidence, consistent with that presented here, continues to highlight the inherent spatio-temporal complexity of signals and visual systems [86-88]. This offers exciting opportunities for integrative studies of signal development, production, and perception, which will fuel a richer view of this pervasive adornment of the natural world.

Acknowledgments

I'm grateful to Daniel Noble and Alistair Senior for advice on meta-analysis, and Elizabeth Mulvenna and Cormac White for their endless support. I appreciate the efforts of three anonymous reviewers whose thoughtful suggestions considerably improved this study.

282 **Funding**

283 None to report.

284

285

286

287

288

289

290

291

292

293

294

295

296

References

1. Maia R, Rubenstein DR, Shawkey MD. 2013 Key ornamental innovations facilitate diversification in an avian radiation. *Proceedings of the National Academy of Sciences* **110**, 10687–10692.
2. Dalrymple RL, Flores-Moreno H, Kemp DJ, White TE, Laffan SW, Hemmings FA, Hitchcock TD, Moles AT. 2018 Abiotic and biotic predictors of macroecological patterns in bird and butterfly coloration. *Ecological Monographs* **88**, 204–224.
3. Andersson MB. 1994 *Sexual selection*. Princeton, NJ: Princeton University Press.
4. Zahavi A. 1975 Mate selection—a selection for a handicap. *Journal of Theoretical Biology* **53**, 205–214.
5. Maynard Smith J, Harper D. 2003 *Animal signals*. Oxford University Press.
6. Keyser AJ, Hill GE. 1999 Condition-dependent variation in the blue-ultraviolet coloration of a structurally based plumage ornament. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **266**, 771–777.
7. Kemp DJ. 2008 Resource-mediated condition dependence in sexually dichromatic butterfly wing coloration. *Evolution* **62**, 2346–2358.
8. Weaver RJ, Santos ES, Tucker AM, Wilson AE, Hill GE. 2018 Carotenoid metabolism strengthens the link between feather coloration and individual quality. *Nature communications* **9**, 73.
9. Simons MJ, Cohen AA, Verhulst S. 2012 What does carotenoid-dependent coloration tell? Plasma carotenoid level signals immunocompetence and oxidative stress state in birds—a meta-analysis. *PLoS One* **7**, e43088.
10. Vukusic P, Sambles J. 2003 Photonic structures in biology. *Nature* **424**, 852–5.
11. Johnsen S. 2012 *The optics of life: A biologist's guide to light in nature*. Princeton University Press.

- 322 12. Prum RO, Dufresne ER, Quinn T, Waters K. 2009 Development of colour-producing β -
323 keratin nanostructures in avian feather barbs. *Journal of the Royal Society Interface* **6**, S253–
324 S265.
- 325 13. Dufresne ER, Noh H, Saranathan V, Mochrie SG, Cao H, Prum RO. 2009 Self-assembly of
326 amorphous biophotonic nanostructures by phase separation. *Soft Matter* **5**, 1792–1795.
- 327 14. Maia R, Macedo RHF, Shawkey MD. 2011 Nanostructural self-assembly of iridescent
328 feather barbules through depletion attraction of melanosomes during keratinization. *Journal of*
329 *the Royal Society Interface* (doi:[10.1098/rsif.2011.0456](https://doi.org/10.1098/rsif.2011.0456))
- 330 15. Ghiradella HT, Butler MW. 2009 Many variations on a few themes: A broader look at
331 development of iridescent scales (and feathers). *Journal of the Royal Society Interface*, rsif–
332 2008.
- 333 16. Shawkey MD, Estes AM, Siefferman LM, Hill GE. 2003 Nanostructure predicts intraspecific
334 variation in ultraviolet-blue plumage colour. *Proc R Soc B* **270**, 1455–1460.
335 (doi:[10.1098/rspb.2003.2390](https://doi.org/10.1098/rspb.2003.2390))
- 336 17. Kemp DJ, Rutowski RL. 2007 Condition dependence, quantitative genetics, and the potential
337 signal content of iridescent ultraviolet butterfly coloration. *Evolution* **61**, 168–183.
- 338 18. Koricheva J, Gurevitch J, Mengersen K. 2013 *Handbook of meta-analysis in ecology and*
339 *evolution*. Princeton University Press.
- 340 19. Westgate MJ. 2019 Revtools: An r package to support article screening for evidence
341 synthesis. *Research synthesis methods*
- 342 20. Pick JL, Nakagawa S, Noble DW. 2019 Reproducible, flexible and high-throughput data
343 extraction from primary literature: The metaDigitise r package. *Methods in Ecology and*
344 *Evolution* **10**, 426–431.
- 345 21. Viechtbauer W. 2010 Conducting meta-analyses in r with the metafor package. *Journal of*
346 *Statistical Software* **36**, 1–48.

22. R Core Team. 2018 *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. See <https://www.R-project.org/>.
23. Hinchliff CE *et al.* 2015 Synthesis of phylogeny and taxonomy into a comprehensive tree of life. *Proceedings of the National Academy of Sciences* **112**, 12764–12769.
24. Michonneau F, Brown JW, Winter DJ. 2016 Rotl: An r package to interact with the open tree of life data. *Methods in Ecology and Evolution* **7**, 1476–1481.
25. Grafen A. 1989 The phylogenetic regression. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* **326**, 119–157.
26. Maia R, White TE. 2018 Comparing colors using visual models. *Behavioral Ecology* **29**, 649–659.
27. Prum RO. 2006 Anatomy, physics, and evolution of structural colors. *Bird coloration* **1**, 295–353.
28. White TE. 2018 Illuminating the evolution of iridescence. *Trends in Ecology & Evolution* **33**, 374–375.
29. Rutowski RL, Merry JW, Morehouse NI, Yturralde K, Taylor-Taft L, Gaalema D, Kemp DJ, Papke RS. 2007 Iridescent ultraviolet signal in the orange sulphur butterfly (*colias eurytheme*): Spatial, temporal and spectral properties. *Biological Journal of the Linnean Society* **90**, 349–364.
30. Cotton S, Fowler K, Pomiankowski A. 2004 Do sexual ornaments demonstrate heightened condition-dependent expression as predicted by the handicap hypothesis? *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**, 771–783.
31. Egger M, Smith GD, Schneider M, Minder C. 1997 Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**, 629–634.
32. D’Alba L, Van Hemert C, Handel CM, Shawkey MD. 2011 A natural experiment on the condition-dependence of achromatic plumage reflectance in black-capped chickadees. *PLoS One* **6**.

33. Taylor LA, Clark DL, McGraw KJ. 2011 Condition dependence of male display coloration in a jumping spider (*habronattus pyrrithrix*). *Behavioral Ecology and Sociobiology* **65**, 1133–1146.
34. Barry KL, White TE, Rathnayake DN, Fabricant SA, Herberstein ME. 2015 Sexual signals for the colour-blind: Cryptic female mantids signal quality through brightness. *Functional Ecology* **29**, 531–539.
35. Peters A, Kurvers R, Roberts ML, Delhey K. 2011 No evidence for general condition-dependence of structural plumage colour in blue tits: An experiment. *Journal of evolutionary biology* **24**, 976–987.
36. Kemp DJ. 2008 Resource-mediated condition dependence in sexually dichromatic butterfly wing coloration. *Evolution: international journal of organic evolution* **62**, 2346–2358.
37. Megía-Palma R, Martínez J, Merino S. 2016 A structural colour ornament correlates positively with parasite load and body condition in an insular lizard species. *The Science of Nature* **103**, 52.
38. Siefferman L, Hill GE, Dobson FS. 2005 Ornamental plumage coloration and condition are dependent on age in eastern bluebirds *sialia sialis*. *Journal of Avian Biology* **36**, 428–435.
39. Perrier C, Lope F de, Møller AP, Ninni P. 2002 Structural coloration and sexual selection in the barn swallow *hirundo rustica*. *Behavioral Ecology* **13**, 728–736.
40. Doucet SM. 2002 Structural plumage coloration, male body size, and condition in the blue-black grassquit. *The Condor* **104**, 30–38.
41. Smiseth PT, Örnborg J, Andersson S, Amundsen T. 2001 Is male plumage reflectance correlated with paternal care in bluethroats? *Behavioral Ecology* **12**, 164–170.
42. Keyser AJ, Hill GE. 2000 Structurally based plumage coloration is an honest signal of quality in male blue grosbeaks. *Behavioral Ecology* **11**, 202–209.
43. Schull Q, Dobson FS, Stier A, Robin J-P, Bize P, Viblanc VA. 2016 Beak color dynamically signals changes in fasting status and parasite loads in king penguins. *Behavioral Ecology*, arw091.

- 398 44. Henderson LJ, Heidinger BJ, Evans NP, Arnold KE. 2013 Ultraviolet crown coloration in
399 female blue tits predicts reproductive success and baseline corticosterone. *Behavioral Ecology*
400 **24**, 1299–1305.
- 401 45. Zirpoli JA, Black J, Gabriel P. 2013 Parasites and plumage in steller’s jays: An experimental
402 field test of the parasite-mediated handicap hypothesis. *Ethology Ecology & Evolution* **25**, 103–
403 116.
- 404 46. Freeman-Gallant CR, Taff CC, Morin DF, Dunn PO, Whittingham LA, Tsang SM. 2010
405 Sexual selection, multiple male ornaments, and age-and condition-dependent signaling in the
406 common yellowthroat. *Evolution: International Journal of Organic Evolution* **64**, 1007–1017.
- 407 47. Nicolaus M, Le Bohec C, Nolan PM, Gauthier-Clerc M, Le Maho Y, Komdeur J, Jouventin
408 P. 2007 Ornamental colors reveal age in the king penguin. *Polar Biology* **31**, 53–61.
- 409 48. Quinard A, Cézilly F, Motreuil S, Rossi J-M, Biard C. 2017 Reduced sexual dichromatism,
410 mutual ornamentation, and individual quality in the monogamous zenaida dove *zenaida aurita*.
411 *Journal of Avian Biology* **48**, 489–501.
- 412 49. Molnár O, Bajer K, Mészáros B, Török J, Herczeg G. 2013 Negative correlation between
413 nuptial throat colour and blood parasite load in male european green lizards supports the
414 hamilton–zuk hypothesis. *Naturwissenschaften* **100**, 551–558.
- 415 50. Grindstaff JL, Lovern MB, Burtka JL, Hallmark-Sharber A. 2012 Structural coloration
416 signals condition, parental investment, and circulating hormone levels in eastern bluebirds (*sialia*
417 *sialis*). *Journal of Comparative Physiology A* **198**, 625–637.
- 418 51. Bajer K, Molnar O, Török J, Herczeg G. 2012 Temperature, but not available energy, affects
419 the expression of a sexually selected ultraviolet (uv) colour trait in male european green lizards.
420 *PLoS One* **7**, e34359.
- 421 52. Roberts ML, Ras E, Peters A. 2009 Testosterone increases uv reflectance of sexually selected
422 crown plumage in male blue tits. *Behavioral Ecology* **20**, 535–541.

- 423 53. Silva N, Avilés JM, Danchin E, Parejo D. 2008 Informative content of multiple plumage-
424 coloured traits in female and male european rollers. *Behavioral Ecology and Sociobiology* **62**,
425 1969.
- 426 54. Bitton P-P, Dawson RD. 2008 Age-related differences in plumage characteristics of male tree
427 swallows tachycineta bicolor: Hue and brightness signal different aspects of individual quality.
428 *Journal of Avian Biology* **39**, 446–452.
- 429 55. Balenger SL, Scott Johnson L, Brubaker JL, Ostlind E. 2007 Parental effort in relation to
430 structural plumage coloration in the mountain bluebird (sialia currucoides). *Ethology* **113**, 838–
431 846.
- 432 56. Siitari H, Alatalo RV, Halme P, Buchanan KL, Kilpimaa J. 2007 Color signals in the black
433 grouse (tetrao tetrix): Signal properties and their condition dependency. *The American Naturalist*
434 **169**, S81–S92.
- 435 57. Mougeot F, Arroyo BE. 2006 Ultraviolet reflectance by the cere of raptors. *Biology Letters* **2**,
436 173–176.
- 437 58. Kemp DJ, Vukusic P, Rutowski R. 2006 Stress-mediated covariance between nano-structural
438 architecture and ultraviolet butterfly coloration. *Functional Ecology* **20**, 282–289.
- 439 59. Kemp DJ. 2006 Heightened phenotypic variation and age-based fading of ultraviolet
440 butterfly wing coloration. *Evolutionary Ecology Research* **8**, 515–527.
- 441 60. Peters A, Delhey K, Goymann W, Kempenaers B. 2006 Age-dependent association between
442 testosterone and crown uv coloration in male blue tits (parus caeruleus). *Behavioral Ecology and*
443 *Sociobiology* **59**, 666–673.
- 444 61. Mougeot F, Redpath SM, Leckie F. 2005 Ultra-violet reflectance of male and female red
445 grouse, lagopus lagopus scoticus, sexual ornaments reflect nematode parasite intensity. *Journal*
446 *of Avian Biology* **36**, 203–209.
- 447 62. Hill GE, Doucet SM, Buchholz R. 2005 The effect of coccidial infection on iridescent
448 plumage coloration in wild turkeys. *Animal Behaviour* **69**, 387–394.

- 449 63. Doucet SM, Montgomerie R. 2003 Multiple sexual ornaments in satin bowerbirds:
450 Ultraviolet plumage and bowers signal different aspects of male quality. *Behavioral Ecology* **14**,
451 503–509.
- 452 64. Hahn S, Peter H-U. 2003 Sex and age dependency of wing patch size in brown skuas. *Emu*
453 **103**, 37–41.
- 454 65. McGraw KJ, Mackillop EA, Dale J, Hauber ME. 2002 Different colors reveal different
455 information: How nutritional stress affects the expression of melanin-and structurally based
456 ornamental plumage. *Journal of Experimental Biology* **205**, 3747–3755.
- 457 66. Molnár O, Bajer K, Török J, Herczeg G. 2012 Individual quality and nuptial throat colour in
458 male e uropean green lizards. *Journal of Zoology* **287**, 233–239.
- 459 67. Griggio M, Zanollo V, Hoi H. 2010 UV plumage color is an honest signal of quality in male
460 budgerigars. *Ecological Research* **25**, 77–82.
- 461 68. Viblanc VA *et al.* 2016 Mutually honest? Physiological ‘qualities’ signalled by colour
462 ornaments in monomorphic king penguins. *Biological Journal of the Linnean Society* **118**, 200–
463 214.
- 464 69. Lim ML, Li D. 2006 Effects of age and feeding history on structure-based uv ornaments of a
465 jumping spider (araneae: Salticidae). *Proceedings of the Royal Society B: Biological Sciences*
466 **274**, 569–575.
- 467 70. Bitton P-P, Dawson RD, Ochs CL. 2008 Plumage characteristics, reproductive investment
468 and assortative mating in tree swallows tachycineta bicolor. *Behavioral Ecology and*
469 *Sociobiology* **62**, 1543–1550.
- 470 71. Beck ML, Hopkins WA, Hawley DM. 2015 Relationships among plumage coloration, blood
471 selenium concentrations and immune responses of adult and nestling tree swallows. *Journal of*
472 *Experimental Biology* **218**, 3415–3424.
- 473 72. Senior AM, Grueber CE, Kamiya T, Lagisz M, O’dwyer K, Santos ES, Nakagawa S. 2016
474 Heterogeneity in ecological and evolutionary meta-analyses: Its magnitude and implications.
475 *Ecology* **97**, 3293–3299.

- 476 73. Ah-King M, Barron AB, Herberstein ME. 2014 Genital evolution: Why are females still
477 understudied? *PLoS Biology* **12**, e1001851.
- 478 74. Cohen J. 1998 *Statistical power analysis for the behavioral sciences*. Routledge.
- 479 75. Griffith SC, Parker TH, Olson VA. 2006 Melanin- versus carotenoid-based sexual signals: Is
480 the difference really so black and red? *Animal Behaviour* **71**, 749–763.
481 (doi:[10.1016/j.anbehav.2005.07.016](https://doi.org/10.1016/j.anbehav.2005.07.016))
- 482 76. Murphy ME, King JR, Lu J. 1988 Malnutrition during the postnuptial molt of white-crowned
483 sparrows: Feather growth and quality. *Canadian Journal of Zoology* **66**, 1403–1413.
- 484 77. Negro Juan Jose, Bildstein KL, Bird DM. 1994 Effects of food deprivation and handling
485 stress on fault-bar formation in nestling american kestrels (*falco sparverius*). *ARDEA* **82**, 263–
486 267.
- 487 78. Kemp DJ, Vukusic P, Rutowski RL. 2006 Stress-mediated covariance between nano-
488 structural architecture and ultraviolet butterfly coloration. *Functional Ecology* **20**, 282–289.
489 (doi:[10.1111/j.1365-2435.01100.x](https://doi.org/10.1111/j.1365-2435.01100.x))
- 490 79. Cotton S, Fowler K, Pomiankowski A. 2004 Condition dependence of sexual ornament size
491 and variation in the stalk-eyed fly *cyrtodiopsis dalmanni* (diptera: Diopsidae). *Evolution* **58**,
492 1038–1046.
- 493 80. Pomiankowski A, Møller AP. 1995 A resolution of the lek paradox. *Proceedings of the Royal*
494 *Society of London. Series B: Biological Sciences* **260**, 21–29.
- 495 81. Gegenfurtner KR, Kiper DC. 2003 Color vision. *Annual Review of Neuroscience* **26**.
- 496 82. Rubin JM, Richards W. 1982 Color vision and image intensities: When are changes material?
497 *Biological Cybernetics* **45**, 215–226.
- 498 83. Seago AE, Brady P, Vigneron J-P, Schultz TD. 2009 Gold bugs and beyond: A review of
499 iridescence and structural colour mechanisms in beetles (coleoptera). *Journal of the Royal*
500 *Society Interface* **6**, S165–S184. (doi:[10.1098/rsif.2008.0354.focus](https://doi.org/10.1098/rsif.2008.0354.focus))

84. White TE, Macedonia J, Birch D, Dawes J, Kemp DJ. 2012. The nanoanatomical basis of sexual dimorphism in iridescent butterfly colouration. *Australian Journal of Zoology*, 60(2), 101-107.
85. Loyau A, Gomez D, Moureau BT, Thery M, Hart NS, Saint Jalme M, Bennett ATD, Sorci G. 2007 Iridescent structurally based coloration of eyespots correlates with mating success in the peacock. *Behavioral Ecology* **18**, 1123–1131.
86. Hogan BG, Stoddard MC. 2018 Synchronization of speed, sound and iridescent color in a hummingbird aerial courtship dive. *Nature Communications* **9**.
87. White TE, Zeil J, Kemp DJ. 2015 Signal design and courtship presentation coincide for highly biased delivery of an iridescent butterfly mating signal. *Evolution* **69**, 14–25.
88. Foster DH, Amano K, Nascimento SM. 2001 How temporal cues can aid colour constancy. *Color Research & Application* **26**, S180–S185.

Table 1: Full parameter estimates from MLM and MLMR models of the mediators of the correlation between structural colour signal expression and individual quality. Shown are sample sizes, mean Fisher's z values and lower and upper 95% confidence intervals, and heterogeneity. Estimates whose 95% confidence intervals do not overlap zero are indicated in bold. Note that iridescence and the inclusion of controls are coded as binary (0/1), and so represent a test of difference in effect-sizes between their counterpart categories (see main text).

Model	n	Mean (Zr)	Lower CI	Upper CI	I ² (%)
Overall (intercept-only)	186	0.157	0.084	0.231	80.42
Quality measure					79.96
age	37	0.015	-0.119	0.148	
body condition	102	0.190	0.099	0.282	
immune function	11	0.353	0.126	0.580	
parasite resistance	36	0.114	-0.023	0.252	
Component					80.32
hue	50	0.224	0.123	0.345	
saturation	57	0.076	-0.029	0.181	
brightness	60	0.171	0.079	0.264	
composite	19	0.120	-0.056	0.296	
Sex					80.78
female	29	0.131	-0.004	0.267	
male	146	0.161	0.080	0.241	
not distinguished	11	0.183	-0.046	0.413	
Study type					80.39
experimental	52	0.216	0.104	0.329	
observational	134	0.128	0.058	0.199	
Optics					82.64
iridescent (vs not)	67	0.150	0.009	0.291	
Control					
included (vs not)	28	0.071	-0.117	0.260	83.28

Figure 1: Forest plot of the mediators of the correlation between structural colour signal expression and individual quality. Shown are Pearson's correlations back transformed from Fisher's z , with 95% confidence intervals about the mean. Sample sizes are displayed on the right. 'Composite' refers to measures that conflate hue, saturation, and brightness (such as PCA), while 'not distinguished' refers to studies in which the sex of focal animals was either not specified, or males and females were pooled. Note that iridescence and the inclusion of controls are coded as binary (0/1), and so represent a test of difference in effect-sizes between their counterpart categories (see main text).

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581