Reovirus core proteins $\lambda 1$ and $\sigma 2$ promote stability of disassembly intermediates and influence early replication events

Stephanie Gummersheimer^a and Pranav Danthi^{lab}

^aDepartment of Biology, Indiana University, Bloomington, Indiana, USA

Running Title: Reovirus core proteins influence entry and replication

^bCorresponding author: Mailing address: Department of Biology, Indiana University 212 S. Hawthorne Drive, Bloomington IN 47405. Phone: (812) 856-2449 Fax: (812) 856-5710. E-mail: pdanthi@indiana.edu

ABSTRACT

The capsids of mammalian reovirus contain two concentric protein shells, the core and the outer capsid. The outer capsid is comprised of $\mu 1$ - $\sigma 3$ heterohexamers which surround the core. The core is comprised of $\lambda 1$ decamers held in place by $\sigma 2$. After entry into the endosome, $\sigma 3$ is proteolytically degraded and $\mu 1$ is cleaved and exposed to form ISVPs. ISVPs undergo further conformational changes to form ISVP*s, resulting in the release of $\mu 1$ peptides which facilitate the penetration of the endosomal membrane to release transcriptionally active core particles into the cytoplasm. Previous work has identified regions or specific residues within reovirus outer capsid that impact the efficiency of cell entry. We examined the functions of the core proteins $\lambda 1$ and $\sigma 2$. We generated a reovirus T3D reassortant that carries strain T1L derived $\sigma 2$ and $\lambda 1$ proteins (T3D/T1L L3S2). This virus displays a lower ISVP stability and therefore converts to ISVP*s more readily. To identify the basis for lability of T3D/T1L L3S2, we screened for hyper-stable mutants of T3D/T1L L3S2 and identified three point mutations in $\mu 1$ that stabilize ISVPs. Two of these mutations are located in the C-terminal ϕ region of $\mu 1$, which has not previously been implicated in controlling ISVP stability. Independent from compromised ISVP stability, we also found that T3D/T1L L3S2 launches replication more efficiently and produces higher yields in infected cells. In addition to identifying a new role for the core proteins in disassembly events, these data highlight that core proteins may influence multiple stages of infection.

IMPORTANCE

Protein shells of viruses (capsids) have evolved to undergo specific changes to ensure the timely delivery of genetic material to host cells. The 2-layer capsid of reovirus

provides a model system to study the interactions between capsid proteins and the changes they undergo during entry. We tested a virus in which the core proteins were derived from a different strain than the outer capsid. We found that this mismatched virus was less stable and completed conformational changes required for entry prematurely. Capsid stability was restored by introduction of specific changes to the outer capsid, indicating that an optimal fit between inner and outer shells maintains capsid function. Separate from this property, mismatch between these protein layers also impacted the capacity of virus to initiate infection and produce progeny. This study reveals new insights into the roles of capsid proteins and their multiple functions during viral replication.

INTRODUCTION

In order to successfully launch replication, a virus must protect, transport and deliver its genome into the host cell. The viral capsid is a complex mechanical container with the primary function of assembling around the viral genome in one host cell, exiting that host cell and releasing the genome in another host cell. Therefore, the capsid has two demands that appear to be in direct conflict with one another. It must be stable enough to protect and transport the viral genome, but it also must be dynamic, poised to react to the right environment at the right time to perform functions during entry and to allow for replication of the genome. Viruses have evolved capsid proteins that are capable of these dynamic mechanical interactions in a number of diverse and fascinating ways. While some capsids are made up of only a single type of capsid protein, others have more complex combinations of proteins. One family of viruses, *Reoviridae*, have double or even triple layered capsids.

The *Reoviridae* family of viruses is made up of non-enveloped virions with capsids that are composed of 1 to 3 concentric protein shells that surround 9 to 12 ds RNA genome segments (1–3). While the outer layers of the multilayered capsids are proteolytically processed and undergo conformational changes during entry, the innermost capsid or core which contains the genome, remains intact throughout the remainder of replication (4). The core is a complex molecular machine that is capable of producing fully capped and functional RNA transcripts for translation by the host machinery (5, 6). The capsid proteins must, therefore, be capable of multiple functions in addition to their protective and structural roles. The outer capsid proteins must be poised to undergo the conformational changes required to release the cores and allow

them to be transcriptionally active. While studies of mammalian reovirus have provided many insights into how outer capsid proteins regulate and mediate entry events that lead to these conformational changes, the role of the core proteins in cell entry events remains unclear.

The capsids of mammalian reovirus are made up of two concentric protein shells, the core and the outer capsid (4). The outer capsid is primarily made up of $\mu 1$ - $\sigma 3$ heterohexamers (7) (Fig 1). These surround the core which is comprised of $\lambda 1$ decamers held in place by $\sigma 2$ (8). Turrets, made up of $\lambda 2$ pentamers, protrude from the core at the five-fold axis of symmetry. The $\sigma 1$ attachment protein forms trimers that are anchored in the $\lambda 2$ turret (8, 9). After entry, $\sigma 3$ is proteolytically degraded within the endosome and $\mu 1$ is cleaved into $\mu 1\delta$ and $\mu 1\phi$ fragments (10, 11). These $\mu 1$ fragments remain particle associated at this stage and the particle is referred to as an infectious subviral particle or ISVP (12–14). Further conformational changes within the endosome result in cleavage of $\mu 1\delta$ to form $\mu 1N$ and δ (15, 16). These particles are no longer infectious and are referred to as ISVP*. Release of $\mu 1$ peptides ($\mu 1N$ and ϕ) results in penetration of the endosomal membrane and the deposition of the transcriptionally active cores into the cytoplasm (13, 17, 18).

Previous work in identifying interactions and proteins involved in reovirus disassembly and cell entry have focused on the outer capsid proteins. Here, we focused on the structural functions of the core proteins $\lambda 1$ and $\sigma 2$. We characterized a reassortant virus containing the $\lambda 1$ encoding gene segment L3 and the $\sigma 2$ encoding gene segment S2 (T3D/T1L L3S2). We found that this new virus displays a higher ISVP-to-ISVP* conversion efficiency or lower ISVP stability. We identified three point

mutations in $\mu 1$ that increase the stability of T3D/T1L L3S2. Two of these mutations are located in the ϕ region of $\mu 1$, which has not previously been implicated in maintaining ISVP stability or controlling ISVP-to-ISVP* conversion. Additionally, this reassortant virus has increased growth resulting from higher transcription levels and, subsequently, higher protein production. This role does not appear to be related to the enhanced ISVP-to-ISVP* conversion of this virus. These results provide insights into understanding the structural functions of the core proteins and how their interactions may influence disassembly and early replication steps during infection.

RESULTS

ISVP to ISVP* conversion efficiency of T3D is altered by the T1L core proteins $\lambda 1$ and $\sigma 2$. Single gene reassortants between prototype reovirus strains T1L and T3D, which contain mismatches between outer capsid proteins, display altered capsid stability and efficiency of disassembly intermediate formation (19–21). To determine whether the core proteins $\lambda 1$ and $\sigma 2$ play a role in ISVP-to-ISVP* conversion efficiency, we characterized the properties of T3D/T1L L3S2. This virus contains the $\lambda 1$ -encoding gene segment L3 and the $\sigma 2$ -encoding gene segment S2 from T1L in an otherwise T3D genetic background. The resulting virus therefore contains major core proteins that do not match the outer capsid. The $\lambda 1$ protein from T3D and T1L are 99.3% identical (9 amino acid differences out of 1,275) and the $\sigma 2$ proteins are 98.8% identical (5 amino acid differences out of 418) (22–24). While $\sigma 2$ interacts with the outer capsid $\mu 1$ trimer, $\lambda 1$ is not known to interact with any outer capsid proteins (25).

To test overall virion stability, we incubated T3D and T3D/T1L L3S2 over a series of elevated temperatures and tested the protease stability of the major viral capsid proteins. Such an approach has been used previously to test stability of virions and viral entry intermediates (19). Based on the similarity in their protease sensitivity profiles, we think that virions of T3D/T1L L3S2 do not display significant changes in stability in comparison to the parent strain, T3D (Fig 2A). To determine if the mismatches in this reassortant virus alter stability of ISVPs or efficiency of ISVP-to-ISVP* conversion, we generated ISVPs of T3D and T3D/T1L L3S2 and incubated them over a gradient of increasing temperatures. ISVP* conversion was determined as a measure of trypsin sensitivity of the $\mu 1$ δ fragment (26). In comparison to the parent strain T3D, T3D/T1L L3 ISVPs have significantly reduced stability or enhanced ISVP-to-ISVP* conversion *in vitro* (Fig 2B).

Conversion to ISVP* results in loss of outer capsid proteins that are essential for entry (27, 28). As a consequence, ISVPs that are heated to temperatures that result in ISVP* conversion have significantly reduced titers. Measuring thermal stability of ISVP infectivity, therefore, is an alternate method to evaluate the efficiency of ISVP* formation. T3D/T1L L3S2 and T3D ISVPs were heated to 40°C and loss of infectivity was measured by plaque assay. Consistent with previous work (20), at 40°C, most of the T3D ISVPs did not convert to ISVP* and the change in titer was minimal (Fig 2C). However, heating T3D/T1L L3S2 ISVPs to 40°C resulted in ISVP* conversion and a significant loss in infectivity (Fig 2C). These data indicate that the reovirus core proteins play a role in stability of the ISVP, a function previously not attributed to the major core proteins $\lambda 1$ and $\sigma 2$.

Enhanced ISVP-to-ISVP* conversion is not due to interactions with RNA.

In addition to its structural functions, $\lambda 1$ plays a variety of other roles during infection. One of those functions involves interaction with RNA (29, 30). If $\lambda 1$ -RNA interactions affect stability of the particles, it may explain why viruses with different core proteins and, consequently, the potential for different RNA interaction properties, may have altered stability. To rule out this possibility, we tested the ISVP-to-ISVP* conversion efficiency of genome-containing particles in comparison to genome-deficient particles of T3D. This comparison would allow us to uncover the contribution of viral genomic RNA to ISVP stability. ISVP-to-ISVP* conversion was again determined as a measure of trypsin sensitivity of $\mu 1 \delta$. Stability of ISVPs of genome-containing particles and genome-deficient particles was not significantly different which suggests that $\lambda 1$ -RNA interactions do not affect ISVP-to-ISVP* conversion. Thus, the ISVP-to-ISVP* phenotype of T3D/T1L L3S2 is likely not related to differences in RNA interactions (Fig 3).

Isolation of heat resistant mutants in T3D/T1L L3S2.

Because ISVP particles that have converted to ISVP* are less infectious, it is possible to select for and isolate variants with mutations that render ISVPs more stable and thus still retain infectivity after exposure to heat (28, 31, 32). Mapping such mutations could reveal key interactions between viral structural proteins that contribute to maintaining ISVP stability. In order to better understand the basis for the lower stability of this reassortant, we sought to identify such mutations in T3D/T1L L3S2 (Fig 4A). ISVPs of T3D/T1L L3S2 were heated to 40°C (a temperature at which they display significantly lower infectivity than that of similarly treated wild-type ISVPs derived from T3D) and subjected

to plaque assay. Resulting plaques were isolated as potential heat resistant mutants. To confirm the heat resistance of these isolates, ISVPs of each isolate were again incubated at 40°C and loss of infectivity in comparison to ISVPs incubated at 4°C was determined by plaque assay. Of the 20 isolates tested, 7 were determined to have little to no loss in infectivity. These isolates were considered to be heat resistant (HR) (Fig 4B). For each of these 7 isolates, genome segments encoding $\lambda 1$, $\sigma 2$, and $\mu 1$ (L3, S2 and M2 respectively) were sequenced. We reasoned that these genome segments will bear mutations because T3D/T1L L3S2 differs from T3D in the properties of $\lambda 1$ and $\sigma 2$, and because $\mu 1$ has been previously implicated in controlling stability of ISVPs (28, 33–35). None of the isolates contained mutations in L3 or S2. Four isolates were identified with mutations in $\mu 1$. HR16 had a mutation at amino acid 459 (lysine to glutamic acid) which is located in the δ fragment of $\mu 1$. HR2, HR15 and HR17 had mutations in the ϕ fragment of $\mu 1$. The mutation in HR 15 was at amino acid 607 (proline to glutamine) and HR2 and HR17 had the same mutation at amino acid 615 (alanine to threonine) (Fig 4 C, D, E). Notably, the K459E mutation has been previously identified as a stabilizing mutation (31). However, mutations in $\mu 1$ ϕ that contribute to ISVP stability or ISVP-to-ISVP* conversion efficiency have not been previously identified.

Mutations in $\mu 1$ stabilize T3D/T1L L3S2 ISVPs. Because we did not sequence the entire genome of HR viruses, it remains possible that mutations in genome segments other than L3, S2 and M2 influence the thermal stability of ISVPs generated from the second-site revertants. To evaluate the stabilizing effect of the identified mutations on T3D/T1L L3S2 ISVPs, each mutation was introduced individually into a T3D/T1L L3S2 background. Each new mutant virus was tested for ISVP-to-ISVP* conversion efficiency.

ISVPs of each virus were generated and incubated over a gradient of temperatures. ISVP* conversion was determined as a measure of trypsin sensitivity of the $\mu 1$ δ fragment (21). In comparison to T3D/T1L L3S2, each of the ISVPs with $\mu 1$ mutations had increased stability, suggesting that these amino acid residues in $\mu 1$ play important roles in ISVP-to-ISVP* conversion (Fig 5A). To verify these results, the infectivity of ISVPs of T3D/T1L L3S2 and each of the $\mu 1$ mutants at 4°C and 40°C was compared by plaque assay. Consistent with the results seen in Fig 2C, T3D/T1L L3S2 ISVPs experience a loss of infectivity at this temperature. In contrast, introduction of $\mu 1$ changes identified in heat resistant mutants into T3D/T1L L3S2 result in ISVP particles that display greater stability (Fig 5B). These data indicate that mutations in $\mu 1$ are sufficient to restore wild-type like ISVP-to-ISVP* conversion efficiency and thermal stability to T3D/T1L L3S2.

Mutations in $\mu 1$ also affect ISVP-to-ISVP* conversion in wild-type T3D. The $\mu 1$ mutations that restored thermal stability of ISVPs and normal ISVP-to-ISVP* conversion efficiency of T3D/T1L L3S2 are not in a position to contact proteins that make up the core (25, 36). Thus, it seems unlikely that these mutations stabilize the capsid by directly altering core-outer capsid interactions. One possibility is that the changes in $\mu 1$ simply stabilize the capsid by strengthening interactions between $\mu 1$ monomers or between $\mu 1$ trimers. If so, the $\mu 1$ mutations would be expected to further stabilize ISVPs of T3D, which contains different L3 and S2 alleles. To test this idea, we also generated viruses containing one of each of the three $\mu 1$ changes found in the HR viruses in a wild-type T3D background. As before, ISVPs of each virus were generated and incubated over a gradient of temperatures. ISVP* conversion was again determined as

a measure of trypsin sensitivity of the $\mu 1$ δ fragment. In comparison to wild-type T3D, each mutant underwent ISVP-to-ISVP* conversion much less efficiently (Fig 6A). To verify these results, each virus was tested for loss of infectivity. ISVPs of T3D along with each mutant were heated to 49°C and loss in infectivity was measured by plaque assay. The 49°C temperature was determined empirically as the lowest temperature at which T3D ISVPs exhibit a loss in infectivity (data not shown). Consistent with the data in Fig 6A, at this temperature each of the $\mu 1$ mutants displayed no loss of infectivity when compared to wild-type T3D. These data suggest that the mutations identified in $\mu 1$ are generally stabilizing mutations and not directly related to the effects of the mismatched core proteins.

Mismatches between the core and outer capsid proteins affect viral replication

and transcription. To determine if the differences in the ISVP-to-ISVP* conversion efficiency of T3D and T3D/T1L L3S2 are relevant during a viral infection, we next examined viral growth via following infection of cells at an MOI of 0.1 PFU/cell. Virus titer at 24 h following infection was determined by plaque assay and viral yield was calculated as an increase in titer from 0 h post infection (which measured virus adsorbed to cells at the start of infection). We observed that infection with T3D/T1L L3S2 resulted in an increased yield in comparison to T3D (Fig 7A). To determine if this growth phenotype correlates with increased ISVP-to-ISVP* conversion efficiency, we tested the impact of introducing a representative $\mu 1$ mutation, A615T in these viruses. While growth of T3D was significantly higher than growth of T3D M2 A615T, growth of T3D/T1L L3S2 was not significantly different from T3D/T1L L3S2 M2A615T (Fig 7A). As the M2 A615T mutation in T3D/T1L L3S2 restores the stability of its ISVPs, these data

indicate that differences in replication efficiency of T3D and T3D/T1L L3S2 is not a consequence of the capacity of the reassortant virus to more easily convert to ISVP*. These data also suggest that T1L derived L3 and S2 genome segments can influence the properties of T3D in multiple independent ways.

The higher replication potential of T3D/T1L L3S2 could be related to the capacity of this virus to produce more viral gene products with faster kinetics or to a greater extent. To test this idea, we next examined viral gene expression early during infection using immunoblots. Cells were infected with ISVPs at an equal MOI and harvested at 0 or 10h post infection. Expression of the μ 1 protein was assessed as a representative. While expression of μ 1 was visible at 10 h post infection in all samples, T3D/T1L L3S2 displayed significantly higher protein expression than WT T3D (Fig 7B). To test the impact of ISVP-to-ISVP* conversion phenotypes on protein expression, we also examined T3D M2 A615T and T3D/T1L L3S2 M2 A615T. While both viruses displayed lower protein levels than T3D/T1L L3S2, the mutant in the T3D background had significantly lower μ 1 expression than the mutant in the T3D/T1L L3S2 suggesting that the ISVP-to-ISVP* conversion phenotype alone is not responsible for the increase in protein expression. To determine whether the greater level of viral protein expression is a consequence of a higher level of viral mRNA, we measured transcription of viral S1 mRNA early during infection using RT-qPCR. Cells were infected with ISVPs at an equal MOI and harvested at 6 hours post infection. At this timepoint, T3D/T1L L3S2 showed significantly greater reovirus S1 transcripts compared to T3D, indicating that the increase in protein production is likely due to enhanced transcription by the reassortant

virus (Fig 7C). These data indicate that swapping core proteins between T1L and T3D in a reassortant virus also influence post-entry viral replication events.

DISCUSSION

To test the role of the inner core proteins on disassembly and early entry events we generated a reassortant virus with major core proteins from T1L and all other proteins (including the outer capsid) from T3D (T3D/T1L L3S2). T3D/T1L L3S2 undergoes ISVP-to-ISVP* conversion much more efficiently. We identified mutations in $\mu 1$ that stabilize ISVPs. T3D/T1L L3S2 also demonstrated increased growth kinetics compared to the parental T3D strain. Surprisingly, the more efficient growth of T3D/T1L L3S2 was not related to its capacity to undergo more efficient ISVP-to-ISVP* transition. Instead, increased growth of T3D/T1L L3S2 relates to more rapid protein and mRNA production early in infection. These data suggest that alteration in properties of core proteins can impact the function of the outer capsid proteins in cell entry. Additionally, properties of core proteins can also influence the enzymatic functions of the capsid that are required to establish efficient infection of host cells.

The reovirus core is a T=1 icosahedron and it is surrounded by the outer capsid (T=13). The outer capsid is made up of 200 heterohexamers of $\mu 1$ and $\sigma 3$ proteins that cover the core (25, 36). It is penetrated by $\lambda 2$ pentameric turrets at each five-fold axis of symmetry. Trimers of the $\sigma 1$ attachment protein are situated inside these $\lambda 2$ turrets (8, 25). The core is made up of 120 copies of $\lambda 1$ arranged in asymmetric pairs of pentamers to form decamers. Twelve such decamers make up the core. The $\sigma 2$ protein

clamps onto $\lambda 1$ at 3 different sites within an asymmetric unit resulting in 150 copies of $\sigma 2$ stabilizing the core shell (8). At each five-fold axis of symmetry, channels penetrate the shell via the $\lambda 2$ pentameric turrets. It is at each of these channels that the $\lambda 3$ polymerase interacts with $\lambda 1$ and is thought to interact with the polymerase co-factor $\mu 2$ (5, 37, 38). There are no known contacts between $\lambda 1$ and the outer capsid. A majority of contacts between the core and the outer capsid occur between $\mu 1$ and $\sigma 2$. These interactions involve the bottom surface of $\mu 1$ and the top surface of $\sigma 2$ (25). The “hub and spoke” structure formed by the C-terminal 33 residues of $\mu 1$ which is thought to be important for stabilizing the $\mu 1$ lattice also makes contact with $\lambda 2$ and/or $\sigma 2$ (25). This interaction is thought to be stabilized in part by the $\mu 1$ residues 51-62 which form flexible loops (25). It is possible that the polymorphic differences between $\sigma 2$ proteins of T1L and T3D influence interaction with $\mu 1$ sufficiently enough such that disassembly is altered. $\lambda 1$ can also impact interaction of the core with $\mu 1$, thereby changing disassembly. However, because $\mu 1$ does not contact $\lambda 1$, this effect would occur indirectly if the structure or conformation of $\lambda 2$ or $\sigma 2$, two proteins that do interact with $\mu 1$, is altered due to differences in $\lambda 1$ residues. The relative contribution of differences in properties of $\sigma 2$ and $\lambda 1$ and the potential subtle differences in structure remain the focus of our ongoing work.

Differences in entry efficiency between different serotypes and laboratory strains of reovirus along with genetic approaches have been used to study how conformational changes and cleavage events required for entry are regulated. Panels of reassortant viruses link differences in ISVP-to-ISVP* conversion to $\mu 1$ (28). The autocatalytic cleavage of $\mu 1$ and efficiency of ISVP* formation have both been linked to distinct

portions of the δ fragment of $\mu 1$ (33). Multiple other studies have linked efficiency of entry related disassembly events to the δ fragment of $\mu 1$ (31, 33, 35, 36, 39). The $\mu 1N$ fragment is released from the particle during ISVP* conversion and the released fragment is thought to function in a positive feedback loop to further drive ISVP* conversion (40). The ISVP* promoting activity of $\mu 1N$ is most efficient in presence of membranes, likely because membrane associated $\mu 1N$ recruits ISVP-like particles (41, 42). Cleavage of ϕ , which is required for its release from particles during ISVP* conversion is also required for efficient interaction of ISVPs with membranes (42). However, precisely how ϕ functions in this step is not known. Here, we have identified two mutations in the ϕ fragment of $\mu 1$ that influence ISVP-to-ISVP* conversion. These mutations are the first indication that the ϕ fragment may also be involved in ISVP-to-ISVP* conversion efficiency. Because our ISVP-to-ISVP* reactions were performed in absence of membranes, the mutations are unlikely to influence ISVP-to-ISVP* conversion by affecting particle-membrane interaction. Thus, the precise mechanism by which the identified ϕ residues influence ISVP* remains unclear. While the majority of $\mu 1$ - $\sigma 3$ interactions occur in the jelly-roll domains (residues 306-514) of $\mu 1$, both of the mutations identified are in a region proposed to form the cradle for the base of $\sigma 3$ (36). This region has not been shown to interact with any core proteins or with neighboring $\mu 1$ monomers (36). Therefore, it is unlikely that these mutations stabilize ISVPs by strengthening inter- or intra- $\mu 1$ trimer interactions or by interacting with core proteins. The identified mutations are in a region of $\mu 1$ that may unfold in order to accommodate $\mu 1N$ release which is a necessary step of ISVP-to-ISVP* conversion (25, 36). Thus, one likely explanation is that ϕ properties affect the release of $\mu 1N$.

The L3 gene segment encoded $\lambda 1$ protein is thought of first as a structural protein that makes up the inner core of reovirus. However, as is the case with most viruses, proteins are capable of playing multiple roles during infection. In addition to its structural roles $\lambda 1$ has RNA helicase activity, phosphohydrolase activity and is known to interact with RNA (30). The role of each of these functions during infection is currently unknown. The polymerase $\lambda 3$ interacts with $\lambda 1$ on the inside of the shell at each five-fold axis (8, 43). While the position of the transcription cofactor protein $\mu 2$ within the capsid is not known, it is possible that it also interacts with $\lambda 1$. Thus, the interaction of $\lambda 1$ with these encapsidated enzymes could alter viral transcription efficiency. Reovirus serotypes T1L and T3D replicate with different efficiency in some cell lines, with T1L replicating to a higher extent and with faster kinetics (44). Reassortant analyses have partially linked this phenotype to the T1L derived $\lambda 1$ -encoding L3 gene segment (44). Our studies indicate that the enhanced infection efficiency of T3D/T1L L3S2 is not related to its greater propensity for ISVP-to-ISVP* transition. Instead, we propose that enhanced efficiency of infection is a result of differences in the activity of $\lambda 1$ itself or its impact on the activity of the transcriptional machinery comprised of $\lambda 3$ and $\mu 2$. While the role of $\sigma 2$ as a structural protein is well established, additional roles in replication have not been confidently identified. It has weak interactions with dsRNA and reassortant studies have linked $\sigma 2$ with increased induction and sensitivity to interferon in some cell types (45, 46). Additional studies are needed with monoreassortants bearing only S2 and L3 gene segments in the T3D background to precisely ascertain the basis for the enhanced replicative efficiency of T3D/T1L L3S2.

Recent studies from our laboratory have revealed that reassortant viruses display phenotypes that are unexpected and extend beyond the known function of the protein. First, we found that, even though the primary function of the M2 encoded protein $\mu 1$ is in membrane penetration, an M2 reassortant virus displays greater attachment to host cells (47). Second, we found that though the S1 encoded $\sigma 1$ protein is the attachment factor, an S1 reassortant impacts the stability of the $\mu 1$ layer with which it makes no interactions (20). Our current study presented here reveals that core proteins, previously only thought to have a structural role in packaging the genomic material, influence cell entry events regulated by the outer capsid. Until the advent of reverse genetics, reassortant analyses have been used as the main strategy to assign function to proteins of segmented viruses. This approach has also been useful to determine the genetic basis of viral disease. While this approach has been invaluable, we think our work suggests that the structure-function explanation of some phenotypes reported for reovirus and possibly other members of the *Reoviridae* family may be more complicated than previously appreciated.

MATERIALS AND METHODS

Cells and viruses. Spinner adapted murine L929 (L) cells were grown at 37°C in Joklik's minimal essential medium (Lonza) supplemented with 5% fetal bovine serum (FBS) (Life Technologies), 2 mM L-glutamine (Invitrogen), 100 U/ml penicillin (Invitrogen), 100 μ g/ml streptomycin (Invitrogen), and 25 ng/ml amphotericin B (Sigma-Aldrich). All virus strains used in this study were derived from reovirus type 3 Dearing (T3D) and reovirus type 1 Lang (T1L) and were generated by plasmid-based reverse

genetics (48). Mutations within the T3D M2 gene were generated by QuikChange site-directed mutagenesis (Agilent Technologies). Primer sequences are available upon request.

Virus propagation and purification. All wild-type and mutant viruses used in this study were propagated and purified as previously described (49, 50). Briefly, plaques isolated from plasmid based reverse genetics were propagated successively in T-25, T75 and T-175 flasks to generate P0, P1 and P2 virus stocks respectively. To generate purified virus, L cells infected with P2 reovirus stocks were lysed by sonication. Virus particles were extracted from the lysates using Vertrel-XF specialty fluid (Dupont) (51). The extracted particles were layered onto 1.2- to 1.4-g/cm³ CsCl step gradients. The gradients were then centrifuged at 187,000 × *g* for 4 h at 4°C. Bands corresponding to purified virus particles (~1.36 g/cm³) (52) were isolated and dialyzed into virus storage buffer (10 mM Tris, pH 7.4, 15 mM MgCl₂, and 150 mM NaCl). Following dialysis, the particle concentration was determined by measuring the optical density of the purified virus stocks at 260 nm (OD₂₆₀) (1 unit at OD₂₆₀ is equal to 2.1 × 10¹² particles/ml)

Generation of ISVPs. Purified virions of the indicated virus strains (2 × 10¹² particles/ml or 4 × 10¹² particles/ml) were digested with 200 µg/ml TLCK (*N*α-*p*-tosyl-L-lysine chloromethyl ketone)-treated chymotrypsin (Worthington Biochemical) in a total volume of 100 µl for 1 hour at 32°C. After 1 h, the reaction mixtures were incubated for 20 min on ice and quenched by the addition of 1 mM phenylmethylsulfonyl fluoride (Sigma-Aldrich). The generation of ISVPs was confirmed by SDS-PAGE and Coomassie brilliant blue staining.

Analysis of ISVP-ISVP* conversion. ISVPs (2×10^{12} particles/ml) of the indicated viral strains were divided into aliquots of equivalent volumes and heated at the indicated temperatures for 20 min. The reaction mixtures were cooled on ice and then digested with 0.10 mg/ml trypsin (Sigma-Aldrich) for 30 min on ice. Following addition of the SDS-PAGE loading dye, the samples subjected to SDS-PAGE analysis. For analysis by quantitative infectivity assay, P2 stocks or purified virus stocks of the indicated viruses were diluted 1:10 in virion storage buffer (10 mM Tris, pH 7.4, 15 mM $MgCl_2$, and 150 mM NaCl). 200 μ g/mL TLCK (*N* α -*p*-tosyl-L-lysine chloromethyl ketone)-treated chymotrypsin (Worthington Biochemical) was added to each sample. Samples were heated to 37°C for 30 min. The reaction was quenched by the addition of 1 mM phenylmethylsulfonyl fluoride (Sigma-Aldrich) and cooled on ice for 10 min. The reactions were divided in equivalent volumes and incubated at 4°C or 40°C for 20 min. Reactions were used to initiate infection of L929 cells and infectivity was determined by plaque assay. The change in infectivity at a given temperature (*T*) was calculated using the following formula: $\log_{10}(\text{PFU/ml})_T - \log_{10}(\text{PFU/ml})_{4^\circ\text{C}}$.

Analysis of virion stability. Virions (2×10^{12} particles/ml) of the indicated viral strains were divided into aliquots of equivalent volumes and heated at the indicated temperatures for 20 min. The reaction mixtures were cooled on ice and then digested with 0.10 mg/ml trypsin (Sigma-Aldrich) for 30 min on ice. Following addition of SDS loading dye, the samples were subjected to analysis by SDS-PAGE.

Isolation and verification of heat resistant (HR) mutants. ISVPs of purified T3D/T1L L3S2 were generated and subsequently heated to 40°C for 20 min. Resulting reactions were diluted in phosphate-buffered saline (PBS) supplemented with 2 mM $MgCl_2$ and

subjected to plaque assay. Heat resistant mutants were selected by plaque purification and propagated in L cells to obtain P0 viral stocks. P0 stocks were diluted 1:10 in virion storage buffer (10 mM Tris, pH 7.4, 15 mM MgCl₂, and 150 mM NaCl). 200µg/mL TLCK (*N*α-*p*-tosyl-L-lysine chloromethyl ketone)-treated chymotrypsin (Worthington Biochemical, Lakewood, NJ) was added to each sample. Samples were heated to 37°C for 30 min. The reaction was quenched by the addition of 1 mM phenylmethylsulfonyl fluoride (Sigma-Aldrich) and cooled on ice for 10 min. The reactions were divided in equivalent volumes and incubated at 4°C or 40°C for 20 min. Reactions were used to initiate infection of L929 cells and infectivity was determined by plaque assay. The change in infectivity at a given temperature (*T*) was calculated using the following formula: $\log_{10}(\text{PFU/ml})_T - \log_{10}(\text{PFU/ml})_{4^\circ\text{C}}$.

Plaque titration. Plaque assays were conducted in spinner-adapted L929 cells plated in 6-well plates (Greiner Bio-One). Cells were adsorbed with dilutions of virus in phosphate-buffered saline (PBS). Cells were overlaid with a molten mixture comprised of 1× medium 199 and 1% Bacto agar supplemented with 10 µg/ml chymotrypsin. Five days following infection, the monolayers were fixed by addition of 4% formaldehyde solution in PBS and incubated overnight. The agar overlay was peeled off, and the monolayers were stained with 1% crystal violet stain in 5% ethanol for 5 h at room temperature. The monolayers were washed with water. Virus titer was quantified by manual counting of plaques.

Analysis of protein levels by immunoblotting. The samples were whole-cell lysates of infected cells prepared using radioimmunoprecipitation assay (RIPA) lysis buffer (50 mM NaCl, 1 mM EDTA at pH 8, 50 mM Tris at pH 7.5, 1% Triton X-100, 1% sodium

deoxycholate, 0.1% SDS) supplemented with protease inhibitor cocktail (Roche) and 500 μ M PMSF, and they were resolved on 10% SDS-PAGE gels and transferred to nitrocellulose membranes. For immunoblotting using polyclonal rabbit antireovirus serum, the membranes were blocked with 5% milk in Tris-buffered saline (TBS) at room temperature for 1 h. Following blocking, rabbit anti-reovirus serum (1:1,000) or anti-PSTAIR was incubated with the membrane in appropriate blocking buffer at room temperature for 1 h. The membranes were washed with TBS supplemented with 0.1% Tween 20 (TBS-T) twice for 15 min and then incubated with Alexa Fluor-conjugated anti-rabbit IgG or anti-mouse IgG in blocking buffer. Following three washes, membranes were scanned using an Odyssey infrared imager (LI-COR), and intensities of bands were quantified using Image Studio Lite software (LI-COR).

Analysis of mRNA levels by RT-qPCR. RNA was extracted from infected cells, at various times after infection, using a total RNA minikit (Bio-Rad). For RT-qPCR, 0.5 to 2 μ g of RNA was reverse transcribed with the high-capacity cDNA RT kit (Applied Biosystems), using random hexamers for amplification of cellular and viral genes. Undiluted cDNA was subjected to PCR using SYBR Select Master Mix (Applied Biosystems) and primers specific for T3D S1 and GAPDH. Fold increases in gene expression with respect to control samples (indicated in figure legend) were measured using the $\Delta\Delta C_T$ method (53). Calculations for determining $\Delta\Delta C_T$ values and relative levels of gene expression were performed as follows:

$$\text{fold increase in viral gene expression} = 2_{[-(\Delta\Delta C_T)]}$$

Statistical analyses. The reported values represent the means of three independent biological replicates. The error bars indicate standard deviations (SD). *P* values were calculated using Student's *t* test (two-tailed; unequal variance assumed).

Modeling. Molecular graphics were created and analysis were performed with the UCSF Chimera package (54).

FIGURE LEGENDS

Figure 1. Schematic representation of reovirus capsid proteins.

Figure 2. T3D/T1L L3S2 exhibits increased efficiency of ISVP to ISVP* conversion *in vitro*. (A) T3D and T3D/T1L L3S2 virions (2×10^{12} particles/ml) were divided into aliquots of equal volume and incubated at either 4°C or over a range of temperatures (65-85°C) for 5 min. The reactions were chilled on ice and digested with 0.10mg/ml trypsin for 30 min. Following addition of loading dye, the samples were subjected to SDS-PAGE analysis. The positions of major capsid proteins are shown. $\mu 1$ runs as $\mu 1C$ (15). (B) ISVPs (2×10^{11} particles/ml) of T3D or T3D/T1L L3S2 were divided into aliquots of equivalent volume and incubated either at 4°C or over a range of temperatures (22-42°C) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The gels shown are representative of at least 3 independent experiments. The position of major capsid proteins is shown. $\mu 1$ runs as $\mu 1C$. (C) ISVPs generated from P2 stocks of the indicated virus strain were divided into aliquots of equivalent volume and incubated at either 4°C or 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. The data are plotted as mean loss of infectivity for three

independent samples in comparison to samples incubated at 4°C. Error bars indicate SD. *, P<0.05 as determined by student's t-test in comparison to T3D.

Figure 3. Increased ISVP to ISVP* efficiency in T3D/T1L L3S2 is not due to altered interactions with viral RNA. ISVPs (2×10^{11} particles/ml) derived from genome-containing or genome-deficient particles of strain T3D were divided into aliquots of equivalent volume and incubated at either 4°C or over a range of temperatures (22-40°C) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The position of major capsid proteins is shown. $\mu 1$ runs as $\mu 1C$.

Figure 4. Selection of viruses with mutations that confer stability to T3D/T1L L3S2 ISVPs. (A) Diagram depicting the process for selecting for mutants with reduced ISVP-ISVP* conversion efficiency of T3D/T1L L3S2. ISVPs of T3D/T1L L3S2 were incubated at 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. Viruses from resulting plaques were isolated and propagated to generate P0 stocks. Heat resistance of these putative heat resistant (HR) plaque isolates was confirmed by measuring the thermal stability of ISVPs incubated at 4°C or 40°C using a plaque assay. Mutants that were confirmed as heat resistant were sequenced. (B) ISVPs generated from P0 stocks were incubated at either 4°C or 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. ND, Not detectable. (C, D) Top (left) and side (right) views of $\mu 1$ trimer (C) and monomer (D) are shown. Position of mutations identified in HR viruses are shown in green. $\mu 1$ cleavage fragments are colored as indicated (E) with one $\mu 1$ monomer shown with darker colors.

Figure 5. Mutations in $\mu 1$ restore stability. (A) ISVPs (2×10^{11} particles/ml) of T3D/T1L L3S2 with the indicated M2 mutations were divided into aliquots of equivalent volume and incubated at either 4°C or over a range of temperatures (22-42°C) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The gels shown are representative of at least 3 independent experiments. The position of major capsid proteins is shown. $\mu 1$ runs as $\mu 1C$. (B) ISVPs generated from P2 stocks of the indicated virus strain were divided into aliquots of equivalent volume and incubated at either 4°C or 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. The data are plotted as mean loss of infectivity for three independent samples in comparison to samples incubated at 4°C. Error bars indicate SD. **, $P < 0.01$, ***, $P < 0.001$ as determined by student's t-test in comparison to T3D/T1L L3S2.

Figure 6. Mutations in $\mu 1$ hyperstabilize T3D. (A) ISVPs (2×10^{11} particles/ml) of T3D and T3D with the indicated M2 mutations were divided into aliquots of equivalent volume and incubated at either 4°C or over a range of temperatures (32-46°C) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The gels shown are representative of at least 3 independent experiments. The position of major capsid proteins is shown. $\mu 1$ runs as $\mu 1C$. (B) ISVPs generated from purified virions were divided into aliquots of equivalent volume and incubated at either 4°C or 42°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. The data are plotted as mean loss of infectivity for three independent samples in

comparison to samples incubated at 4°C. Error bars indicate SD. ***, $P < 0.001$, **, $P < 0.01$ as determined by student's T-test in comparison to T3D.

Figure 7. T3D/T1L L3S2 affects viral replication. (A) L cell monolayers were infected with T3D or T3D/T1L L3S2 or with the indicated mutant viruses at an MOI of 0.1 PFU/cell. At 0h and 24h post infection, the infected cells were lysed and the viral yield was quantified by plaque assay. Error bars indicate SD. *, $P < 0.05$, ***, $P < 0.001$ as determined by student's t-test in comparison to T3D. (B) L cell monolayers were infected with the indicated viruses at an MOI of 10 PFU/cell. At 10h post infection, the cells were lysed and protein production was determined by immunoblotting. Protein quantification of 3 replicates normalized to PSTAIR is shown. Error bars indicate SD. **, $P < 0.01$, ***, $P < 0.001$ as determined by student's t-test in comparison to T3D. (C) L cell monolayers were infected with the indicated viruses at an MOI of 10 PFU/cell. At the indicated times post infection, the cells were lysed and total RNA was isolated. cDNA was generated using primers for T3D S1 and GAPDH. mRNA production was measured by qPCR. Data shown are represented as fold change compared to mock infected samples and normalized to GAPDH. Error bars indicate SD. ***, $P < 0.001$ as determined by student's t-test in comparison to T3D

REFERENCES

1. SABIN AB. 1959. Reoviruses. A new group of respiratory and enteric viruses formerly classified as ECHO type 10 is described. *Science* 130:1387–1389.
2. Gomatos PJ, Tamm I. 1963. THE SECONDARY STRUCTURE OF REOVIRUS RNA. *Proc Natl Acad Sci U S A* 49:707–714.
3. Bellamy AR, Shapiro L, August JT, Joklik WK. 1967. Studies on reovirus RNA. I. Characterization of reovirus genome RNA. *J Mol Biol* 29:1–17.
4. Dermody TS, Parker JSL, Sherry B. 2013. Orthoreoviruses, p 1304–1346. *Fields virology*. Lippincott Williams & Wilkins, Philadelphia, PA.
5. Gillies S, Bullivant S, Bellamy AR. 1971. Viral RNA Polymerases: Electron Microscopy of Reovirus Reaction Cores. *Science* (80-) 174:694 LP – 696.
6. Skehel JJ, Joklik WK. 1969. Studies on the in vitro transcription of reovirus RNA catalyzed by reovirus cores. *Virology* 39:822–31.
7. Liemann S, Chandran K, Baker TS, Nibert ML, Harrison SC. 2002. Structure of the reovirus membrane-penetration protein, $\mu 1$, in a complex with its protector protein, $\sigma 3$. *Cell* 108:283–295.
8. Reinisch KM, Nibert ML, Harrison SC. 2000. Structure of the reovirus core at 3.6 Å resolution. *Nature* 404:960–967.
9. Mao ZX, Joklik WK. 1991. Isolation and enzymatic characterization of protein lambda 2, the reovirus guanylyltransferase. *Virology* 185:377–386.
10. Baer GS, Dermody TS. 1997. Mutations in reovirus outer-capsid protein sigma3

selected during persistent infections of L cells confer resistance to protease inhibitor E64. J Virol 71:4921–4928.

11. Silverstein SC, Astell C, Levin DH, Schonberg M, Acs G. 1972. The mechanisms of reovirus uncoating and gene activation in vivo. Virology 47:797–806.

12. Dryden KA, Wang G, Yeager M, Nibert ML, Coombs KM, Furlong DB, Fields BN, Baker TS. 1993. Early steps in reovirus infection are associated with dramatic changes in supramolecular structure and protein conformation: analysis of virions and subviral particles by cryoelectron microscopy and image reconstruction. J Cell Biol 122:1023–1041.

13. Nibert ML, Fields BN. 1992. A carboxy-terminal fragment of protein mu 1/mu 1C is present in infectious subviral particles of mammalian reoviruses and is proposed to have a role in penetration. J Virol 66:6408–6418.

14. Borsa J, Sargent MD, Lievaart PA, Copps TP. 1981. Reovirus: evidence for a second step in the intracellular uncoating and transcriptase activation process. Virology 111:191–200.

15. Nibert ML, Odegard AL, Agosto MA, Chandran K, Schiff LA. 2005. Putative autocleavage of reovirus mu1 protein in concert with outer-capsid disassembly and activation for membrane permeabilization. J Mol Biol 345:461–474.

16. Odegard AL, Chandran K, Zhang X, Parker JSL, Baker TS, Nibert ML. 2004. Putative autocleavage of outer capsid protein micro1, allowing release of myristoylated peptide micro1N during particle uncoating, is critical for cell entry by reovirus. J Virol 78:8732–8745.

17. Zhang L, Chandran K, Nibert ML, Harrison SC. 2006. Reovirus μ 1 structural rearrangements that mediate membrane penetration. *J Virol* 2006/09/27. 80:12367–12376.
18. Chandran K, Farsetta DL, Nibert ML. 2002. Strategy for nonenveloped virus entry: a hydrophobic conformer of the reovirus membrane penetration protein μ 1 mediates membrane disruption. *J Virol* 76:9920–33.
19. Snyder AJ, Wang JC-Y, Danthi P. 2019. Components of the Reovirus Capsid Differentially Contribute to Stability. *J Virol* 93:e01894-18.
20. Thete D, Danthi P. 2018. Protein mismatches caused by reassortment influence functions of the reovirus capsid. *bioRxiv*.
21. Snyder AJ, Wang JC-Y, Danthi P. 2018. The reovirus μ 1 protein contributes to the environmental stability of virions. *bioRxiv*.
22. Harrison SJ, Farsetta DL, Kim J, Noble S, Broering TJ, Nibert ML. 1999. Mammalian reovirus L3 gene sequences and evidence for a distinct amino-terminal region of the λ 1 protein. *Virology* 258:54–64.
23. Xu W, Coombs KM. 2009. Conserved structure/function of the orthoreovirus major core proteins. *Virus Res* 144:44–57.
24. Chapell JD, Goral MI, Rodgers SE, dePamphilis CW, Dermody TS. 1994. Sequence diversity within the reovirus S2 gene: reovirus genes reassort in nature, and their termini are predicted to form a panhandle motif. *J Virol* 68:750–756.
25. Zhang X, Ji Y, Zhang L, Harrison SC, Marinescu DC, Nibert ML, Baker TS. 2005.

Features of reovirus outer capsid protein mu1 revealed by electron cryomicroscopy and image reconstruction of the virion at 7.0 Angstrom resolution. Structure 13:1545–1557.

26. Snyder AJ, Danthi P. 2018. Infectious Subviral Particle to Membrane Penetration Active Particle (ISVP-to-ISVP*) Conversion Assay for Mammalian Orthoreovirus. Bio-protocol 8:e2700.

27. Snyder AJ, Danthi P. 2015. Lipid Membranes Facilitate Conformational Changes Required for Reovirus Cell Entry. J Virol 90:2628–2638.

28. Middleton JK, Severson TF, Chandran K, Gillian AL, Yin J, Nibert ML. 2002. Thermostability of reovirus disassembly intermediates (ISVPs) correlates with genetic, biochemical, and thermodynamic properties of major surface protein mu1. J Virol 76:1051–61.

29. Lemay G, Danis C. 1994. Reovirus λ1 protein: Affinity for double-stranded nucleic acids by a small amino-terminal region of the protein independent from the zinc finger motif. J Gen Virol 75:3261–3266.

30. Bisailon M, Bergeron J, Lemay G. 1997. Characterization of the nucleoside triphosphate phosphohydrolase and helicase activities of the reovirus λ1 protein. J Biol Chem 272:18298–18303.

31. Middleton JK, Agosto MA, Severson TF, Yin J, Nibert ML. 2007. Thermostabilizing mutations in reovirus outer-capsid protein mu1 selected by heat inactivation of infectious subviral particles. Virology 2007/01/17. 361:412–425.

32. Agosto MA, Middleton JK, Freimont EC, Yin J, Nibert ML. 2007. Thermolabilizing Pseudoreversions in Reovirus Outer-Capsid Protein $\mu 1$ Rescue the Entry Defect Conferred by a Thermostabilizing Mutation. *J Virol* 81:7400 LP – 7409.
33. Sarkar P, Danthi P. 2010. Determinants of strain-specific differences in efficiency of reovirus entry. *J Virol* 84:12723–12732.
34. Drayna D, Fields BN. 1982. Biochemical studies on the mechanism of chemical and physical inactivation of reovirus. *J Gen Virol* 63 (Pt 1):161–170.
35. Wessner DR, Fields BN. 1993. Isolation and genetic characterization of ethanol-resistant reovirus mutants. *J Virol* 67:2442–2447.
36. Liemann S, Chandran K, Baker TS, Nibert ML, Harrison SC. 2002. Structure of the reovirus membrane-penetration protein, $\mu 1$, in a complex with its protector protein, $\sigma 3$. *Cell* 108:283–295.
37. Farsetta DL, Chandran K, Nibert ML. 2000. Transcriptional activities of reovirus RNA polymerase in recoated cores. Initiation and elongation are regulated by separate mechanisms. *J Biol Chem* 275:39693–39701.
38. McDonald SM, Tao YJ, Patton JT. 2009. The ins and outs of four-tunneled Reoviridae RNA-dependent RNA polymerases. *Curr Opin Struct Biol* 2009/11/14. 19:775–782.
39. Hooper JW, Fields BN. 1996. Role of the $\mu 1$ protein in reovirus stability and capacity to cause chromium release from host cells. *J Virol* 70:459–467.
40. Agosto MA, Myers KS, Ivanovic T, Nibert ML. 2008. A positive-feedback

mechanism promotes reovirus particle conversion to the intermediate associated with membrane penetration. Proc Natl Acad Sci U S A 105:10571–6.

41. Snyder AJ, Danthi P. 2016. Lipids Cooperate with the Reovirus Membrane Penetration Peptide to Facilitate Particle Uncoating. J Biol Chem 2016/11/15. 291:26773–26785.

42. Ivanovic T, Agosto MA, Zhang L, Chandran K, Harrison SC, Nibert ML. 2008. Peptides released from reovirus outer capsid form membrane pores that recruit virus particles. EMBO J 27:1289–98.

43. Zhang X, Walker SB, Chipman PR, Nibert ML, Baker TS. 2003. Reovirus polymerase $\lambda 3$ localized by cryo-electron microscopy of virions at a resolution of 7.6 Å. Nat Struct Mol Biol 10:1011–1018.

44. Simon EJ, Howells MA, Stuart JD, Boehme KW. 2017. Serotype-specific killing of large cell carcinoma cells by reovirus. Viruses 9.

45. Dermody TS, Schiff LA, Nibert ML, Coombs KM, Fields BN. 1991. The S2 gene nucleotide sequences of prototype strains of the three reovirus serotypes: characterization of reovirus core protein sigma 2. J Virol 65:5721–5731.

46. Sherry B, Torres J, Blum MA. 1998. Reovirus induction of and sensitivity to beta interferon in cardiac myocyte cultures correlate with induction of myocarditis and are determined by viral core proteins. J Virol 72:1314–23.

47. Thete D, Danthi P. 2015. Conformational changes required for reovirus cell entry are sensitive to pH. Virology 483:291–301.

48. Boehme KW, Ikizler M, Kobayashi T, Dermody TS. 2011. Reverse genetics for mammalian reovirus. *Methods* 55:109–113.
49. Sarkar P, Danthi P. 2010. Determinants of strain-specific differences in efficiency of reovirus entry. *J Virol* 2010/10/13. 84:12723–12732.
50. Furlong DB, Nibert ML, Fields BN. 1988. Sigma 1 protein of mammalian reoviruses extends from the surfaces of viral particles. *J Virol* 62:246–256.
51. Mendez II, Hermann LL, Hazelton PR, Coombs KM. 2000. A comparative analysis of freon substitutes in the purification of reovirus and calicivirus. *J Virol Methods* 90:59–67.
52. Smith RE, Zweerink HJ, Joklik WK. 1969. Polypeptide components of virions, top component and cores of reovirus type 3. *Virology* 39:791–810.
53. Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 3:1101–1108.
54. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* 25:1605–1612.

Figure 1. Schematic representation of reovirus capsid proteins.

Figure 2. T3D/T1L L3S2 exhibits increased efficiency of ISVP to ISVP* conversion in vitro. (A) T3D and T3D/T1L L3S2 virions (2×10^{12} particles/ml) were divided into aliquots of equal volume and incubated at either 4°C or over a range of temperatures ($65\text{--}85^{\circ}\text{C}$) for 5 min. The reactions were chilled on ice and digested with 0.10mg/ml trypsin for 30 min. Following addition of loading dye, the samples were subjected to SDS-PAGE analysis. The positions of major capsid proteins are shown. $\mu 1$ runs as $\mu 1\text{C}$ (15). (B) ISVPs (2×10^{11} particles/ml) of T3D or T3D/T1L L3S2 were divided into aliquots of equivalent volume and incubated either at 4°C or over a range of temperatures ($22\text{--}42^{\circ}\text{C}$) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The gels shown are representative of at least 3 independent experiments. The position of major capsid proteins is shown. $\mu 1$ runs as $\mu 1\text{C}$. (C) ISVPs generated from P2 stocks of the indicated virus strain were divided into aliquots of equivalent volume and incubated at either 4°C or 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. The data are plotted as mean loss of infectivity for three independent samples in comparison to samples incubated at 4°C . Error bars indicate SD. *, $P < 0.05$ as determined by student's t-test in comparison to T3D.

Figure 3. Increased ISVP to ISVP* efficiency in T3D/T1L L3S2 is not due to altered interactions with viral RNA. ISVPs (2×10^{11} particles/ml) derived from genome-containing or genome-deficient particles of strain T3D were divided into aliquots of equivalent volume and incubated at either 4°C or over a range of temperatures (22-40°C) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The position of major capsid proteins is shown. μ 1 runs as μ 1C.

Figure 4. Selection of viruses with mutations that confer stability to T3D/T1L L3S2 ISVPs. (A) Diagram depicting the process for selecting for mutants with reduced ISVP-ISVP* conversion efficiency of T3D/T1L L3S2. ISVPs of T3D/T1L L3S2 were incubated at 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. Viruses from resulting plaques were isolated and propagated to generate P0 stocks. Heat resistance of these putative heat resistant (HR) plaque isolates was confirmed by measuring the thermal stability of ISVPs incubated at 4°C or 40°C using a plaque assay. Mutants that were confirmed as heat resistant were sequenced. (B) ISVPs generated from P0 stocks were incubated at either 4°C or 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. ND, Not detectable. (C, D) Top (left) and side (right) views of μ 1 trimer (C) and monomer (D) are shown. Position of mutations identified in HR viruses are shown in green. μ 1 cleavage fragments are colored as indicated (E) with one μ 1 monomer shown with darker colors.

A

B

Figure 5. Mutations in $\mu 1$ restore stability of T3D/T1L L3S2. (A) ISVPs (2×10^{11} particles/ml) of T3D/T1L L3S2 with the indicated M2 mutations were divided into aliquots of equivalent volume and incubated at either 4°C or over a range of temperatures ($22\text{--}42^{\circ}\text{C}$) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The gels shown are representative of at least 3 independent experiments. The position of major capsid proteins is shown. $\mu 1$ runs as $\mu 1\text{C}$. (B) ISVPs generated from P2 stocks of the indicated virus strain were divided into aliquots of equivalent volume and incubated at either 4°C or 40°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. The data are plotted as mean loss of infectivity for three independent samples in comparison to samples incubated at 4°C . Error bars indicate SD. **, $P < 0.01$, ***, $P < 0.001$ as determined by student's t-test in comparison to T3D/T1L L3S2.

Figure 6. Mutations in $\mu 1$ hyperstabilize T3D. (A) ISVPs (2×10^{11} particles/ml) of T3D and T3D with the indicated M2 mutations were divided into aliquots of equivalent volume and incubated at either 4°C or over a range of temperatures (32–46°C) for 20 min. The reactions were chilled on ice and digested with 0.10 mg/ml trypsin for 30 min. Following addition of loading dye the samples were subjected to SDS-PAGE analysis. The gels shown are representative of at least 3 independent experiments. The position of major capsid proteins is shown. $\mu 1$ runs as $\mu 1C$. (B) ISVPs generated from purified virions were divided into aliquots of equivalent volume and incubated at either 4°C or 42°C for 20 min. Reactions were then diluted in PBS and subjected to plaque assay. The data are plotted as mean loss of infectivity for three independent samples in comparison to samples incubated at 4°C. Error bars indicate SD. ***, $P < 0.001$, **, $P < 0.01$ as determined by student's t-test in comparison to T3D/T1L L3S2.

Figure 7. T3D/T1L L3S2 affects viral replication. (A) L cell monolayers were infected with T3D or T3D/T1L L3S2 or with the indicated mutant viruses at an MOI of 0.1 PFU/cell. At 0h and 24h post infection, the infected cells were lysed and the viral yield was quantified by plaque assay. Error bars indicate SD. *, $P < 0.05$, ***, $P < 0.001$ as determined by student's t-test in comparison to T3D. (B) L cell monolayers were infected with the indicated viruses at an MOI of 10 PFU/cell. At 10h post infection, the cells were lysed and protein production was determined by immunoblotting. Protein quantification of 3 replicates normalized to PSTAIR is shown. Error bars indicate SD. **, $P < 0.01$, ***, $P < 0.001$ as determined by student's t-test in comparison to T3D. (C) L cell monolayers were infected with the indicated viruses at an MOI of 10 PFU/cell. At the indicated times post infection, the cells were lysed and total RNA was isolated. cDNA was generated using primers for T3D S1 and GAPDH. mRNA production was measured by qPCR. Data shown are represented as fold change compared to mock infected samples and normalized to GAPDH. Error bars indicate SD. ***, $P < 0.001$ as determined by student's t-test in comparison to T3D.