1 Title Page

2 Original article

3 Angiotensin II induces apoptosis in human induced pluripotent stem cell-
4 derived cardiomyocytes

5
6 Qiang Gao¹, Ping Wang², Zhiming Wu³, Hailong Qiu¹, Bin Lin⁴, Jimei
7 Chen¹, Jianzheng Cen^{1*} and Jian Zhuang^{1*}

8

9 1 Department of Cardiac Surgery, Guangdong Cardiovascular Institute, Guangdong
10 Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou,
11 Guangdong 510100, China

12 2 School of Medical Imaging, Tianjin Medical University, Tianjin 300203, China

13 3 Department of Urology, Sun Yat-sen University Cancer Center, State Key Laboratory
14 of Oncology in South China, Collaborative Innovation Center for Cancer Medicine,
15 Guangzhou 510060, China

16 4 Guangdong Beating Origin Regenerative Medicine Co. Ltd., Foshan, Guangdong
17 528231, China

18 *Correspondence to: Jian Zhuang. Department of Cardiac Surgery, Guangdong
19 Cardiovascular Institute, Guangdong Provincial People's Hospital, Guangdong
20 Academy of Medical Sciences, Guangzhou, Guangdong 510100, China. Tel: +86
21 83827812-76210. Fax: +86 83827812. E-mail: zhuangjian2345@163.com; Jianzheng
22 Cen. Department of Cardiac Surgery, Guangdong Cardiovascular Institute, Guangdong

23 Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou,
 24 Guangdong 510100, China. Tel: +86 83827812-76210. Fax: +86 83827812. E-mail:
 25 jamesofsolar@163.com.

26 **Running Title: Angiotensin II induces apoptosis in iPSC-CM**

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

Abstract

Background: The renin-angiotensin system (RAS) functions fundamentally to regulate the pathological process of cardiovascular diseases, such as heart failure and hypertension. As the major effector in RAS, angiotensin II activates angiotensin II receptors to initiate the downstream pathways, which lead to the phenotypes including apoptosis, hypertrophy, and cardiac remodeling. Human induced pluripotent stem cell-derived cardiomyocytes (iPSC-CM) are being applied as a promising platform for personalized medicine to heart diseases. However, whether angiotensin II induces apoptosis in iPSC-CM is still obscure, which raises an uncertainty about the clinical applications of iPSC-CM.

Methods: We treated iPSC-CM with angiotensin II at eight concentrations (0 nM, 1 nM, 10 nM, 100 nM, 1 μ M, 10 μ M, 100 μ M and 1 mM) and four incubation durations (24 hours, 48 hours, 6 days and 10 days), then PrestoBlue reagent and a apoptosis marker were used to examine the viability and apoptosis status of cardiomyocytes from each group. The expression levels of some apoptosis and proliferation related genes were also analyzed.

Results: High concentration angiotensin II with a long-term treatment caused apoptosis and cell viability drop-off in iPSC-CM. Specifically, under a 10-day treatment with 1 mM angiotensin II, the viability of iPSC-CM was reduced by an average of 41% ($p=2.073E-08$), and the percentage

of apoptotic cells was 2.74 times higher than the controls averagely ($p=6.248E-12$). The data mining of previous RNA-seq data revealed that angiotensin II receptor type I was the major receptor in iPSC-CM.

Conclusions: For the first time, our data confirmed the apoptotic effect of angiotensin II to iPSC-CM. The angiotensin II concentrations and exposure time for apoptosis induction were depicted in our study, which provided supports to iPSC-CM as an *in vitro* model for cardiovascular disease study.

Keywords: Angiotensin II; concentration; iPSC-CM; apoptosis; cell viability.

Introduction

Heart failure (HF) is the most popular chronic disease among the elderly in developed countries (1), and its prevalence and incidence is increasing in the recent years (2-4). Approximately 26 million individuals are suffering HF worldwide (5), in which the survival rate within 5 years of admission was less than 53% (6-8). The chronic heart failure causes a tremendous economic burden with an estimation of \$108 billion per year globally (9). As a comprehensive medical condition, HF is an end stage of many cardiovascular diseases, such as ischaemic heart disease, chronic obstructive pulmonary disease, hypertensive heart disease, and rheumatic heart disease (10).

Like other multifactorial conditions, progression of HF is modified by the genetic diversity of affected individuals, which makes routine HF treatment difficult to achieve satisfying benefits (4,11). It is an urgent mission to establish an *in vitro* system to evaluate the effects of different medicine to each HF patient. The advances of cell reprogramming technology (especially induced pluripotent stem cell (iPSC) technology) (12,13), cardiomyocyte differentiation (14,15) and purification (16,17) protocols provide us powerful tools to generate cardiomyocytes from patients' somatic cells. These iPSC-derived cardiomyocytes (iPSC-CM) could mimic the cardiac phenotypes from the donors without ethics issues. To date, iPSC-CM were used to investigate long QT syndrome (18-20), arrhythmia (21,22), hypertrophic cardiomyopathy (23), dilated cardiomyopathy (24,25), mitochondrial cardiomyopathy (26), Duchenne muscular dystrophy (27), catecholaminergic polymorphic ventricular tachycardia (28), and other cardiovascular diseases.

On the other side, due to the high-throughput advantages, iPSC-CM were more and more applied to pharmacologic and cardiotoxicity testing (29-35). There medications, including aldosterone antagonist, angiotensin-converting enzyme inhibitors, angiotensin II receptor blockers, angiotensin receptor-neprilysin inhibitors, and beta-blockers (36,37), are available for testing in the patients' iPSC-CM. Notably, many of these drugs target to the angiotensin pathway. Indeed, angiotensin II (Ang II) plays a crucial role

in the pathological mechanism of HF (38-40). It was reported that Ang II caused apoptosis of cardiomyocytes in rats (41-43), vascular fibrosis and damage (44). In fact, Ang II was an apoptosis inducer of many kinds of cells, such as vascular smooth muscle cells (45), human endothelial cells (46), coronary artery endothelial cells (47), rat glomerular epithelial cells (48), alveolar epithelial cells (49), intestinal epithelial cells (50), renal proximal tubular cells (51,52), mesangial cells (53,54), and podocytes (55). However, there are no studies reporting that Ang II induced apoptosis in iPSC-CM. Instead, several reports showed that Ang II could induce hypertrophic phenotypes in iPSC-CM (56,57). Since the response to Ang II is a critical criterion for drug testing, it is necessary to figure out whether Ang II induces apoptosis in iPSC-CM.

In this research, we performed short-term and long-term treatments of Ang II to iPSC-CM. We presented that only long-term treatments with high concentration Ang II caused apoptosis in iPSC-CM, and Ang II triggered the downstream signaling cascades mainly through angiotensin II receptor type 1 (AT1).

Results

Cell viability was significantly reduced in iPSC-CM under long-term treatments of high concentration Ang II

To determine the pro-apoptotic effects by Ang II at various concentrations

133 and incubation time, 8 concentrations (0 nM (untreated), 1 nM, 10 nM, 100
134 nM, 1 μ M, 10 μ M, 100 μ M and 1 mM) and 4 treatment durations (24 hours,
135 48 hours, 6 days and 10 days) were tested in iPSC-CM. Here, 1 nM to 1
136 μ M were defined as “low concentrations”, while 10 μ M, 100 μ M and 1
137 mM were defined as “high concentrations”; similarly, 24 hours and 48
138 hours were defined as “short terms”, while 6 days and 10 days were defined
139 as “long terms”. Human iPS cell line DYR0100 (ATCC) was differentiated
140 to cardiomyocytes by using the protocols from previous reports (14,16).
141 After metabolic selections (16), iPSC-CM with > 98% purity were seeded
142 to 96-well plates for recovery. On differentiation day 30, iPSC-CM were
143 applied to Ang II treatments. Under short-term treatments (24 hours and 48
144 hours) of Ang II, there were no significant differences between control
145 iPSC-CM and Ang II-treated iPSC-CM in cell viability, no matter with low
146 or high concentrations (Figure 1a). Conversely, after 6-day treatments, 100
147 μ M and 1 mM Ang II significantly reduced the viability of iPSC-CM (Fold
148 change of cell viability, 100 μ M vs. untreated control: 0.763, $p=8.085E-07$;
149 1 mM vs. untreated control: 0.762, $p=7.306E-07$); after 10-day treatments,
150 10 μ M, 100 μ M and 1 mM Ang II caused ~21%, 36% and 41% drop-off in
151 cell viability, respectively (Fold change of cell viability, 10 μ M vs.
152 untreated control: 0.795, $p=0.034$; 100 μ M vs. untreated control: 0.638,
153 $p=4.885E-07$; 1 mM vs. untreated control: 0.590, $p=2.073E-08$). Ang II at
154 other concentrations seemed to slightly decrease the cell viability without

statistical significance (Figure 1b). Intriguingly, we observed that iPSC-CM under Ang II treatment for a short term showed a light increase in viability, which was inconsistent with the data from other types of cells.

Long-term incubation of high concentration Ang II induced apoptosis in iPSC-CM

Based on the results of cell viability test, the viability of iPSC-CM only declined under long-term treatments of Ang II. Therefore, to investigate the apoptotic rates, we performed apoptosis analysis in iPSC-CM after 10-day treatments of Ang II. iPSC-CM under 10-day incubations were collected and stained by Annexin V conjugated with Alexa Fluor 488, then analyzed by fluorescence-activated cell sorting (FACS). As Figure 2 shown, the apoptotic rate of untreated iPSC-CM was about 4.1%, whereas about 12.7% and 15.6% of iPSC-CM were apoptotic after 100 μ M and 1 mM Ang II treatments, respectively (Figure 2a), which indicated that the apoptotic cells under 100 μ M Ang II treatment were averagely 3.08 times as many as untreated group ($p=2.029E-09$), while the apoptotic cells under 1 mM Ang II treatment were averagely 3.74 times as many as untreated group ($p=6.248E-12$) (Figure 2b). There were no significant differences between treatments with other concentrations and controls. TUNEL assay was also performed to verify the apoptosis analysis. iPSC-CM were treated with 1 nM, 1 μ M, 100 μ M and 1 mM Ang II for 10 days then applied to

TUNEL assay. The TUNEL positive cells were significantly increased in 100 μ M and 1 mM Ang II-treated iPSC-CM. Compared to the untreated iPSC-CM, 100 μ M and 1 mM Ang II-treated iPSC-CM exhibited 1.63 ($p=7.144E-08$) and 2.32-fold ($p=1.298E-15$) increase in apoptotic rate, respectively (Figure 2c, d). 1 nM and 1 μ M Ang II treatments did not show differences compared with the control set.

We next evaluated the expression levels of apoptosis and proliferation related genes in iPSC-CM, which were treated with Ang II at different concentrations (1 nM, 1 μ M and 1 mM) for 2 days and 10 days (Figure 3). In the iPSC-CM under a 2-day Ang II treatment, 1 nM and 1 μ M Ang II did not change the expression levels of *BCL2*, *TP53* and *GADD45A*, while 1 mM Ang II led to 1.25-fold increase in *BAX* ($p=0.0004$) and down-regulation of *BCL2* ($p=1.432E-06$). The expression of *MKI67* was decreased after Ang II treatment at all the three concentrations. On the other hand, after the 10-day incubation with 1 mM Ang II, the expression of *BAX*, *TP53*, *GADD45A* and *MKI67* was significantly up-regulated (with 2.25-, 1.81-, 1.83-, and 2.53-fold increase, respectively; P values were 0.0001, 0.0006, 0.0030, 5.348E-07, respectively), while the expression of *BCL2* dropped by 70% ($p=8.583E-06$). Contrary to the 2-day treatment, 10-day treatment elevated the expression of *MKI67* at all the three concentrations. Encoding a pro-apoptotic protein, *BAX* functions as an apoptotic activator in the apoptosis pathway (58). In contrast, *BCL2* was

an anti-apoptotic molecule that promoted cell survival (59) and blocked programmed cell death (60). *TP53* and *GADD45A* were cell cycle-regulated genes, which were also reported to mediate apoptosis in cells (61,62). *MKI67* and *PCNA* were markers of cell proliferation (63). It is worth noting that 1 mM Ang II already caused up-regulation of *BAX* and down-regulation of *BCL2* in a short-term treatment, and with the increase of incubation time, the *BAX/BCL2* ratio, which was considered as an apoptotic index, was elevated sharply. Besides, 1 mM Ang II increased the expression of *TP53* and *GADD45A* as well, indicating that the cardiomyocytes were in the apoptotic status.

AT1, not AT2 was expressed in iPSC-CM

There are two types of angiotensin II receptors: AT1 and AT2. Basically, AT1 was implicated in the major effects of Ang II, such as proliferation, hypertrophy, and fibrosis, while AT2 acted to antagonize the effects of AT1 (64,65). Generally, it was considered that Ang II caused apoptosis and cardiac remodeling through binding to AT2 (66,67), however, it was reported that AT1 was also associated with Ang II-mediated apoptosis (68-70). To investigate which receptor was involved in the Ang II-mediated apoptosis in iPSC-CM, we first examined the expression levels of the renin-angiotensin system (RAS)-related genes (*AGTR1*, *AGTR2*, *ACE*, *REN*, and *AGT*) by re-analyzing the expression profiling data of cells from

different time points of cardiac differentiation (GSE116574 and GSE137920) (71,72). In the data set GSE116574, *AGTR1* started to be expressed from differentiation day 9, whereas the transcripts of *AGTR2* were too low to be considered undetectable (Figure 4a). In the data set GSE137920, the transcripts of *AGTR1* could be detected on differentiation day 14, however, *AGTR2* was not expressed in the cells from all the time points (Figure 4b). We performed reverse transcription PCR to detect the expression levels of *AGTR1* and *AGTR2* in iPSC-CM on differentiation day 30. In agreement with the previous reports, we found that *AGTR1* was expressed in the iPSC-CM, while *AGTR2* was hardly detectable (data not shown). According to these observations, we proposed that Ang II induced apoptosis in the iPSC-CM through binding to AT1.

Discussion

The renin-angiotensin system is the key pathway engaged in pathogenesis of many cardiovascular diseases, including cardiomyopathy, heart failure and hypertension (73-75). As the central component of RAS, Ang II activated AT1 and AT2 to regulate the physiological functions of cardiomyocytes. Many chemicals targeting RAS, such as angiotensin receptor blockers (valsartan, losartan, etc.), angiotensin converting enzyme inhibitors (captopril, benazepril, enalapril, etc.) and angiotensin receptor-neprilysin inhibitors (Entresto, etc.), were used as medications for

cardiovascular disease therapy. Consequently, the response to Ang II was an essential requirement for a cardiovascular disease model. To the best of our knowledge, only a few studies investigated the physiological effects of Ang II to iPSC-CM. Binah group examined the effects of Ang II on the contraction and intracellular Ca^{2+} transient in hESC-CM (76,77). Harding group and Fukuda group reported that Ang II could induce hypertrophy in iPSC and hESC-derived cardiomyocytes (56,78), while Harding group presented a contrary result that Ang II had no significant effect on cell size in hESC-CM and hiPSC-CM three years later (79). Radisic group used 200 nM Ang II to treat hESC-CM in 3D culture for 7 days, and found no significant differences in cell death and viability in drug-treated and untreated hESC-CM (80). This was the first attempt to study the effects after chronic Ang II exposure. It was unexpected that there were no reports demonstrating the apoptotic effects of Ang II to iPSC-CM. For the first time, we showed evidences that only high concentration Ang II (10 μM , 100 μM and 1 mM) provoked apoptosis after chronic treatment, namely, the apoptosis phenotype was dependent on the concentration and incubation time of Ang II.

The expression of AT1 in iPSC-CM was confirmed by immunofluorescent staining (76). However, according to the RNA-seq data (71,72), the expression level of AT1 was relatively very low compared to GAPDH in iPSC-CM (Figure 4). The low expression level of AT1 might explain short-

term stimuli of Ang II could not induce apoptosis: when the expression of AT2 was missing, even the high concentration Ang II could not trigger the apoptosis pathway through a small amount of AT1. Since Ang II infusion for 28 days could increase the protein levels of AT1 and AT2 in rats (35% and 100% increase, respectively) (81), we assumed that a long-term Ang II incubation could also up-regulate the expression level of AT2 in iPSC-CM, which offered a reasonable explanation for the apoptosis phenotype induced by long-term Ang II treatments: chronic exposure by high concentration Ang II facilitated iPSC-CM to produce enough AT2 so that Ang II could activate AT2 to initiate downstream apoptosis signaling. In conclusion, we proved that Ang II could induce apoptosis in iPSC-CM in a concentration and time-dependent manner. These findings provided necessary supports for iPSC-CM as a powerful tool to investigate cardiovascular diseases *in vitro*.

Method

Cell Culture

Human induced pluripotent stem cells (iPSC) DYR0100 (The American Type Culture Collection, ATCC) were plated on Matrigel matrix (hESC-Qualified, LDEV-Free, Corning, 354277)-coated plates, and then were cultured with StemFlex Medium (Gibco, A3349401). StemFlex Medium was changed every two days. iPSC were passaged every three days or when

the cell culture was 80-90% confluent. During passages, iPSC were rinsed with 1X DPBS (Gibco, 14040133) for one time then were treated with 0.5mM EDTA (Invitrogen, 15575020) in 1X DPBS (Gibco, 14190144) for 10 mins at room temperature. The split ratio was 1:3-1:6. The detailed differentiation protocol was described in the previous published reports (16,82). Briefly, iPSC were treated with small molecule CHIR99021 (Tocris, 4423, final concentration 10 μ M) in the RPMI-BSA medium [RPMI 1640 Medium (HyClone, SH30027.01) supplemented with 213 μ g/ml AA2P (l-ascorbic acid 2-phosphate magnesium) (A8960, Sigma) and 0.1% bovine serum albumin (BSA) (A1470, Sigma)] for 24 hours, then were incubated with RPMI-BSA medium for 48 hours. On differentiation day 4, cells were treated with the small molecule IWP2 (Tocris, 3533, final concentration 5 μ M) in RPMI-BSA medium. After 48 hours, medium was changed to RPMI-BSA medium. Then, RPMI 1640 Medium supplemented with 3% KnockOut Serum Replacement (Gibco, 10828-028) was used to culture the cardiomyocytes in the following experiments. A metabolic selection method was used to purify iPSC-derived cardiomyocytes (16). The metabolic selection medium was prepared as DMEM Medium (No Glucose) (Gibco, 11966-025) supplemented with 0.1% BSA (Sigma, A1470) and 1 $\times$ Linoleic Acid-Oleic Acid-Albumin (Sigma, L9655). Cells were treated with metabolic selection medium for 3-6 days. The medium was changed every 2 days. The purity of cardiomyocytes could reach as

high as 99%. After purification, the cardiomyocyte culture medium supplemented with various amount of Ang II (MedChemExpress, HY-13948) was used for Ang II treatment, and the Ang II medium was changed every 2 days.

Cell Viability Assay

PrestoBlue Cell Viability Reagent (Invitrogen, A13261) was used to detect the viability of cardiomyocytes. PrestoBlue was warmed to room temperature before use. One-tenth volume of PrestoBlue was added to the cardiomyocyte medium to prepare reaction medium. For a 96-well plate, 100 µl reaction media were added to each well. The same amount reaction medium was added to wells without cells as blank controls. Plates were incubated in the 37 °C incubator for 60 mins then fluorescence (excitation wavelength: 560 nm, emission wavelength: 590 nm) was read by Varioskan Flash Multimode Reader (Thermo Scientific). The results were normalized by cell numbers.

Apoptosis Detection Assay

The apoptosis detection kit Annexin V, Alexa Fluor™ 488 conjugate (Invitrogen, V13201) was used to detect the apoptosis status of Ang II-treated and untreated cardiomyocytes. Cardiomyocytes were labeled with

apoptosis marker Annexin V then sorted by the FACS Aria™ II analyzer (BD). FACS results were analyzed by using FlowJo software.

TUNEL Assay

TUNEL assay was performed by using TdT In Situ Apoptosis Detection Kit (R&D Systems, 4812-30-K). Cardiomyocytes were dissociated and replated on the glass coverslips (Fisher Scientific, S175223). We followed the manufacturer's protocol except fixed cells were labeled with DAPI before mounted by fluorescence mounting media (Agilent, S302380-2). Images were taken by the DMI6000 B inverted microscope (Leica) and analyzed by using ImageJ software.

Quantitative PCR

Total RNA was extracted using the UNIQ-10 Column Trizol Total RNA Isolation Kit (Sangon Biotech, B511321-0100) prior to the treatment with DNase I (Sangon Biotech, B618252) for 30 min. mRNA was reverse transcribed using iScript Reverse Transcription Supermix (Bio-Rad, 1708841). Quantitative PCR was performed using a PikoReal Real-Time PCR System (Thermo Fisher) with SsoAdvanced™ Universal SYBR® Green Supermix (Bio-Rad, 1725271). The quantitative PCR primers from the previous report (83) are as followed (from 5' to 3'):

BAX-RT-F: CAAACTGGTGCTCAAGGCC;

352 *BAX*-RT-R: GGGCGTCCCAAAGTAGGAGA;
 353 *BCL2*-RT-F: CTGGTGGACAACATCGCCCT;
 354 *BCL2*-RT-R: TCTTCAGAGACAGCCAGGAGAAAT;
 355 *MKI67*-RT-F: TGTGCCTGCTCGACCCTACA;
 356 *MKI67*-RT-R: TGAAATAGCGATGTGACATGTGCT;
 357 *PCNA*-RT-F: TTTGGTGCAGCTCACCCCTG;
 358 *PCNA*-RT-R: CGCGTTATCTTCGGCCCTTA;
 359 *TP53*-RT-F: GCGTGTTTGTGCCTGTCCTG;
 360 *TP53*-RT-R: TGGTTTCTTCTTTGGCTGGG;
 361 *GADD45A*-RT-F: GATGCCCTGGAGGAAGTGCT;
 362 *GADD45A*-RT-R: GAGCCACATCTCTGTCGTCGT;
 363 *GAPDH*-RT-F: TGGGTGTGAACCATGAGAAG;
 364 *GAPDH*-RT-R: GTGTCGCTGTTGAAGTCAGA.

365

366 RNA-seq data re-analysis

367 Raw fastq files were downloaded from GEO (GSE116574 and GSE137920)
 368 by SRA-Tools (<https://github.com/ncbi/sra-tools>). Low quality bases and
 369 adapters were trimmed by trim_galore
 370 (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/). We
 371 employed Kallisto (v0.46.0) (84) to quantified transcript abundance as
 372 transcripts per million (TPM), using gene annotation in the GENCODE

database (v32) (85). Then we summed TPM of all alternative splicing transcripts of a gene to obtain gene expression levels.

Statistic

Values are expressed as mean $\pm$ SD (standard deviation). Statistical significances were evaluated using one-way ANOVA with Bonferroni correction. $P < 0.05$ was considered statistically significant.

Figure Legend

Figure 1. Cell viability was decreased after long-term Ang II treatment. The effect of a series of Ang II concentrations (1 nM, 10 nM, 100 nM, 1 μ M, 10 μ M, 100 μ M, 1 mM) to iPSC-CM was evaluated by PrestoBlue reagent. The relative fluorescence unit (RFU) fold changes from untreated cardiomyocytes to Ang II-treated cardiomyocytes were calculated in a), short-term (24 hours and 48 hours) treatment set and b), long-term (6 days and 10 days) treatment set (n=7). #, $p < 0.05$; **, $p < 0.001$; unlabeled data, no significant differences.

Figure 2. Long-term Ang II treatment caused apoptosis in iPSC-CM. a) Representative fluorescence-activated cell sorting (FACS) analyses of apoptosis marker Annexin V in iPSC-CM with 0 nM (untreated), 100 μ M and 1 mM Ang II treatment for 10 days. In the unstained group, the DPBS buffer was used instead of the Annexin V staining reagent. b) The

proportion of Annexin V-positive cells (PAP) represented the apoptosis status of iPSC-CM. The fold changes of PAP from untreated cardiomyocytes to Ang II-treated cardiomyocytes at different concentrations were calculated (n=3). **, $p<0.001$; unlabeled data, no significant differences. c) Representative images of TUNEL assay in iPSC-CM with 0 nM (untreated), 100 μ M and 1 mM Ang II treatment for 10 days. Nuclei were stained with DAPI (blue). The white arrows denote TUNEL-positive cardiomyocyte nuclei (green). The scale bar is 100 μ m. d) The proportion of TUNEL-positive cells (TP) also reflected the apoptosis status of iPSC-CM. The fold changes of TP from untreated cardiomyocytes to Ang II-treated cardiomyocytes at 0 nM (untreated), 1 nM, 1 μ M, 100 μ M and 1 mM concentrations were calculated. Approximately 5,000 cells were counted in each group. **, $p<0.001$; unlabeled data, no significant differences.

Figure 3. High concentration Ang II was associated with the expression changes of apoptosis and proliferation related genes. iPSC-CM were treated with 0 nM (untreated), 1 nM, 1 μ M and 1 mM Ang II for 2 days and 10 days. Total mRNA was extracted from each group (n=3). Expression of apoptosis and proliferation related genes in 2-day and 10-day treatments was shown in left and right panels, respectively. #, $p<0.05$; *, $p<0.01$; **, $p<0.001$; NS, no significant differences.

Figure 4. AT1 was the major angiotensin II receptor in iPSC-CM. RNA-seq data about *AGTR1*, *AGTR2*, *ACE*, *REN*, *AGT* and *GAPDH* genes from a) GSE116547 dataset and b) GSE137920 dataset were re-analyzed. Transcripts per million (TPM) were used to evaluate the expression levels of each gene. D+number denoted the days during cardiomyocyte differentiation.

Author contributions

- (I) Conception and design: Jian Zhuang and Jianzheng Cen
- (II) Administrative support: Jian Zhuang, Jianzheng Cen and Jimei Chen
- (III) Provision of study materials or patients: Qiang Gao and Lin Bin
- (IV) Collection and assembly of data: Qiang Gao, Ping Wang, Zhiming Wu and Hailong Qiu
- (V) Data analysis and interpretation: Qiang Gao
- (VI) Manuscript writing: All authors
- (VII) Final approval of manuscript: All authors

Conflicts of Interest

The authors have no conflicts of interest to declare.

Reference

1. Riegel B, Driscoll A, Suwanno J, *et al.* Heart failure self-care in developed and developing

- 438 countries. J Card Fail 2009;15:508-16.
- 439 2. Kannel WB. Incidence and epidemiology of heart failure. Heart Fail Rev 2000;5:167-73.
- 440 3. Ponikowski P, Anker SD, AlHabib KF, *et al.* Heart failure: preventing disease and death worldwide.
- 441 ESC Heart Fail 2014;1:4-25.
- 442 4. Metra M, Teerlink JR. Heart failure. Lancet 2017;390:1981-95.
- 443 5. Bui AL, Horwich TB, Fonarow GC. Epidemiology and risk profile of heart failure. Nat Rev Cardiol
- 444 2011;8:30-41.
- 445 6. Matoba M, Matsui S, Hirakawa T, *et al.* Long-term prognosis of patients with congestive heart
- 446 failure. Jpn Circ J 1990;54:57-61.
- 447 7. Taylor CJ, Roalfe AK, Iles R, *et al.* Ten-year prognosis of heart failure in the community: follow-
- 448 up data from the Echocardiographic Heart of England Screening (ECHOES) study. Eur J
- 449 Heart Fail 2012;14:176-84.
- 450 8. Taylor CJ, Ryan R, Nichols L, *et al.* Survival following a diagnosis of heart failure in primary care.
- 451 Fam Pract 2017;34:161-8.
- 452 9. Cook C, Cole G, Asaria P, *et al.* The annual global economic burden of heart failure. Int J Cardiol
- 453 2014;171:368-76.
- 454 10. Ziaeian B, Fonarow GC. Epidemiology and aetiology of heart failure. Nat Rev Cardiol
- 455 2016;13:368-78.
- 456 11. McMurray JJ, Adamopoulos S, Anker SD, *et al.* [ESC Guidelines for the diagnosis and treatment
- 457 of acute and chronic heart failure 2012]. Turk Kardiyol Dern Ars 2012;40 Suppl 3:77-137.
- 458 12. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult
- 459 fibroblast cultures by defined factors. Cell 2006;126:663-76.
- 460 13. Takahashi K, Tanabe K, Ohnuki M, *et al.* Induction of pluripotent stem cells from adult human
- 461 fibroblasts by defined factors. Cell 2007;131:861-72.
- 462 14. Lian X, Hsiao C, Wilson G, *et al.* Robust cardiomyocyte differentiation from human pluripotent
- 463 stem cells via temporal modulation of canonical Wnt signaling. Proc Natl Acad Sci U S A
- 464 2012;109:E1848-57.
- 465 15. Lian X, Bao X, Zilberter M, *et al.* Chemically defined, albumin-free human cardiomyocyte
- 466 generation. Nat Methods 2015;12:595-6.
- 467 16. Lin B, Lin X, Stachel M, *et al.* Culture in Glucose-Depleted Medium Supplemented with Fatty
- 468 Acid and 3,3',5-Triiodo-L-Thyronine Facilitates Purification and Maturation of Human
- 469 Pluripotent Stem Cell-Derived Cardiomyocytes. Front Endocrinol (Lausanne) 2017;8:253.
- 470 17. Tohyama S, Hattori F, Sano M, *et al.* Distinct metabolic flow enables large-scale purification of
- 471 mouse and human pluripotent stem cell-derived cardiomyocytes. Cell Stem Cell
- 472 2013;12:127-37.
- 473 18. Mehta A, Sequiera GL, Ramachandra CJ, *et al.* Re-trafficking of hERG reverses long QT
- 474 syndrome 2 phenotype in human iPS-derived cardiomyocytes. Cardiovasc Res
- 475 2014;102:497-506.
- 476 19. Malan D, Friedrichs S, Fleischmann BK, *et al.* Cardiomyocytes obtained from induced
- 477 pluripotent stem cells with long-QT syndrome 3 recapitulate typical disease-specific
- 478 features in vitro. Circ Res 2011;109:841-7.
- 479 20. Egashira T, Yuasa S, Suzuki T, *et al.* Disease characterization using LQTS-specific induced
- 480 pluripotent stem cells. Cardiovasc Res 2012;95:419-29.
- 481 21. Priori SG, Napolitano C, Di Pasquale E, *et al.* Induced pluripotent stem cell-derived

cardiomyocytes in studies of inherited arrhythmias. *J Clin Invest* 2013;123:84-91.

22. Vaidyanathan R, Markandeya YS, Kamp TJ, *et al.* IK1-enhanced human-induced pluripotent stem cell-derived cardiomyocytes: an improved cardiomyocyte model to investigate inherited arrhythmia syndromes. *Am J Physiol Heart Circ Physiol* 2016;310:H1611-21.

23. Lan F, Lee AS, Liang P, *et al.* Abnormal calcium handling properties underlie familial hypertrophic cardiomyopathy pathology in patient-specific induced pluripotent stem cells. *Cell Stem Cell* 2013;12:101-13.

24. Tse HF, Ho JC, Choi SW, *et al.* Patient-specific induced-pluripotent stem cells-derived cardiomyocytes recapitulate the pathogenic phenotypes of dilated cardiomyopathy due to a novel DES mutation identified by whole exome sequencing. *Hum Mol Genet* 2013;22:1395-403.

25. Sun N, Yazawa M, Liu J, *et al.* Patient-specific induced pluripotent stem cells as a model for familial dilated cardiomyopathy. *Sci Transl Med* 2012;4:130ra47.

26. Wang G, McCain ML, Yang L, *et al.* Modeling the mitochondrial cardiomyopathy of Barth syndrome with induced pluripotent stem cell and heart-on-chip technologies. *Nat Med* 2014;20:616-23.

27. Guan X, Mack DL, Moreno CM, *et al.* Dystrophin-deficient cardiomyocytes derived from human urine: new biologic reagents for drug discovery. *Stem Cell Res* 2014;12:467-80.

28. Zhang XH, Haviland S, Wei H, *et al.* Ca²⁺ signaling in human induced pluripotent stem cell-derived cardiomyocytes (iPS-CM) from normal and catecholaminergic polymorphic ventricular tachycardia (CPVT)-afflicted subjects. *Cell Calcium* 2013;54:57-70.

29. Tanaka T, Tohyama S, Murata M, *et al.* In vitro pharmacologic testing using human induced pluripotent stem cell-derived cardiomyocytes. *Biochem Biophys Res Commun* 2009;385:497-502.

30. Drawnel FM, Boccardo S, Prummer M, *et al.* Disease modeling and phenotypic drug screening for diabetic cardiomyopathy using human induced pluripotent stem cells. *Cell Rep* 2014;9:810-21.

31. Lu HR, Whittaker R, Price JH, *et al.* High Throughput Measurement of Ca⁺⁺ Dynamics in Human Stem Cell-Derived Cardiomyocytes by Kinetic Image Cytometry: A Cardiac Risk Assessment Characterization Using a Large Panel of Cardioactive and Inactive Compounds. *Toxicol Sci* 2015;148:503-16.

32. Gintant G, Sager PT, Stockbridge N. Evolution of strategies to improve preclinical cardiac safety testing. *Nat Rev Drug Discov* 2016;15:457-71.

33. da Rocha AM, Campbell K, Mironov S, *et al.* hiPSC-CM Monolayer Maturation State Determines Drug Responsiveness in High Throughput Pro-Arrhythmia Screen. *Sci Rep* 2017;7:13834.

34. Kanda Y, Yamazaki D, Osada T, *et al.* Development of torsadogenic risk assessment using human induced pluripotent stem cell-derived cardiomyocytes: Japan iPS Cardiac Safety Assessment (JiCSA) update. *J Pharmacol Sci* 2018;138:233-9.

35. Satsuka A, Kanda Y. Cardiotoxicity assessment of drugs using human iPS cell-derived cardiomyocytes: From proarrhythmia risk to cardiooncology. *Curr Pharm Biotechnol* 2019.

36. Rossignol P, Hernandez AF, Solomon SD, *et al.* Heart failure drug treatment. *Lancet* 2019;393:1034-44.

37. Yancy CW, Jessup M, Bozkurt B, *et al.* 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American

526 College of Cardiology/American Heart Association Task Force on Clinical Practice
527 Guidelines and the Heart Failure Society of America. *Circulation* 2017;136:e137-e61.

528 38. Urata H, Healy B, Stewart RW, *et al.* Angiotensin II-forming pathways in normal and failing
529 human hearts. *Circ Res* 1990;66:883-90.

530 39. Sayer G, Bhat G. The renin-angiotensin-aldosterone system and heart failure. *Cardiol Clin*
531 2014;32:21-32, vii.

532 40. Wang Y, Seto SW, Golledge J. Angiotensin II, sympathetic nerve activity and chronic heart
533 failure. *Heart Fail Rev* 2014;19:187-98.

534 41. Cao J, Wu Y, Liu G, *et al.* [Over-expression of BDNF inhibits angiotensin II-induced apoptosis
535 of cardiomyocytes in SD rats]. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* 2018;34:218-24.

536 42. Ravassa S, Fortuno MA, Gonzalez A, *et al.* Mechanisms of increased susceptibility to
537 angiotensin II-induced apoptosis in ventricular cardiomyocytes of spontaneously
538 hypertensive rats. *Hypertension* 2000;36:1065-71.

539 43. Schroder D, Heger J, Piper HM, *et al.* Angiotensin II stimulates apoptosis via TGF-beta1
540 signaling in ventricular cardiomyocytes of rat. *J Mol Med (Berl)* 2006;84:975-83.

541 44. Ruiz-Ortega M, Ruperez M, Esteban V, *et al.* Molecular mechanisms of angiotensin II-induced
542 vascular injury. *Curr Hypertens Rep* 2003;5:73-9.

543 45. Bascands JL, Girolami JP, Trolly M, *et al.* Angiotensin II induces phenotype-dependent apoptosis
544 in vascular smooth muscle cells. *Hypertension* 2001;38:1294-9.

545 46. Dimmeler S, Rippmann V, Weiland U, *et al.* Angiotensin II induces apoptosis of human
546 endothelial cells. Protective effect of nitric oxide. *Circ Res* 1997;81:970-6.

547 47. Chen J, Chen W, Zhu M, *et al.* Propofol attenuates angiotensin II-induced apoptosis in human
548 coronary artery endothelial cells. *Br J Anaesth* 2011;107:525-32.

549 48. Ding G, Reddy K, Kapasi AA, *et al.* Angiotensin II induces apoptosis in rat glomerular epithelial
550 cells. *Am J Physiol Renal Physiol* 2002;283:F173-80.

551 49. Wang R, Zagariya A, Ibarra-Sunga O, *et al.* Angiotensin II induces apoptosis in human and rat
552 alveolar epithelial cells. *Am J Physiol* 1999;276:L885-9.

553 50. Sun L, Wang W, Xiao W, *et al.* Angiotensin II induces apoptosis in intestinal epithelial cells
554 through the AT2 receptor, GATA-6 and the Bax pathway. *Biochem Biophys Res Commun*
555 2012;424:663-8.

556 51. Bhaskaran M, Reddy K, Radhakrishnan N, *et al.* Angiotensin II induces apoptosis in renal
557 proximal tubular cells. *Am J Physiol Renal Physiol* 2003;284:F955-65.

558 52. Weidekamm C, Hauser P, Hansmann C, *et al.* Effects of AT1 and AT2 receptor blockade on
559 angiotensin II induced apoptosis of human renal proximal tubular epithelial cells. *Wien*
560 *Klin Wochenschr* 2002;114:725-9.

561 53. Lodha S, Dani D, Mehta R, *et al.* Angiotensin II-induced mesangial cell apoptosis: role of
562 oxidative stress. *Mol Med* 2002;8:830-40.

563 54. Lv J, Jia R, Yang D, *et al.* Candesartan attenuates Angiotensin II-induced mesangial cell
564 apoptosis via TLR4/MyD88 pathway. *Biochem Biophys Res Commun* 2009;380:81-6.

565 55. Jia J, Ding G, Zhu J, *et al.* Angiotensin II infusion induces nephrin expression changes and
566 podocyte apoptosis. *Am J Nephrol* 2008;28:500-7.

567 56. Tanaka A, Yuasa S, Mearini G, *et al.* Endothelin-1 induces myofibrillar disarray and contractile
568 vector variability in hypertrophic cardiomyopathy-induced pluripotent stem cell-derived
569 cardiomyocytes. *J Am Heart Assoc* 2014;3:e001263.

570 57. Ponicke K, Heinroth-Hoffmann I, Becker K, *et al.* Trophic effect of angiotensin II in neonatal rat
571 cardiomyocytes: role of endothelin-1 and non-myocyte cells. *Br J Pharmacol*
572 1997;121:118-24.

573 58. Suzuki M, Youle RJ, Tjandra N. Structure of Bax: coregulation of dimer formation and
574 intracellular localization. *Cell* 2000;103:645-54.

575 59. Vaux DL, Cory S, Adams JM. Bcl-2 gene promotes haemopoietic cell survival and cooperates
576 with c-myc to immortalize pre-B cells. *Nature* 1988;335:440-2.

577 60. Korsmeyer SJ. Bcl-2: an antidote to programmed cell death. *Cancer Surv* 1992;15:105-18.

578 61. Salvador JM, Brown-Clay JD, Fornace AJ, Jr. Gadd45 in stress signaling, cell cycle control, and
579 apoptosis. *Adv Exp Med Biol* 2013;793:1-19.

580 62. Aubrey BJ, Kelly GL, Janic A, *et al.* How does p53 induce apoptosis and how does this relate to
581 p53-mediated tumour suppression? *Cell Death Differ* 2018;25:104-13.

582 63. Iatropoulos MJ, Williams GM. Proliferation markers. *Exp Toxicol Pathol* 1996;48:175-81.

583 64. Akazawa H, Yano M, Yabumoto C, *et al.* Angiotensin II type 1 and type 2 receptor-induced cell
584 signaling. *Curr Pharm Des* 2013;19:2988-95.

585 65. Ichiki T. Regulation of angiotensin II receptor expression. *Curr Pharm Des* 2013;19:3013-21.

586 66. Nouet S, Nahmias C. Signal transduction from the angiotensin II AT2 receptor. *Trends*
587 *Endocrinol Metab* 2000;11:1-6.

588 67. Lemarie CA, Schiffrin EL. The angiotensin II type 2 receptor in cardiovascular disease. *J Renin*
589 *Angiotensin Aldosterone Syst* 2010;11:19-31.

590 68. Papp M, Li X, Zhuang J, *et al.* Angiotensin receptor subtype AT(1) mediates alveolar epithelial
591 cell apoptosis in response to ANG II. *Am J Physiol Lung Cell Mol Physiol* 2002;282:L713-
592 8.

593 69. Diep QN, El Mabrouk M, Yue P, *et al.* Effect of AT(1) receptor blockade on cardiac apoptosis in
594 angiotensin II-induced hypertension. *Am J Physiol Heart Circ Physiol* 2002;282:H1635-41.

595 70. Goldenberg I, Grossman E, Jacobson KA, *et al.* Angiotensin II-induced apoptosis in rat
596 cardiomyocyte culture: a possible role of AT1 and AT2 receptors. *J Hypertens*
597 2001;19:1681-9.

598 71. Branco MA, Cotovio JP, Rodrigues CAV, *et al.* Transcriptomic analysis of 3D Cardiac
599 Differentiation of Human Induced Pluripotent Stem Cells Reveals Faster Cardiomyocyte
600 Maturation Compared to 2D Culture. *Sci Rep* 2019;9:9229.

601 72. Lau E, Han Y, Williams DR, *et al.* Splice-Junction-Based Mapping of Alternative Isoforms in the
602 Human Proteome. *Cell Rep* 2019;29:3751-65 e5.

603 73. Johnston CI, Fabris B, Yoshida K. The cardiac renin-angiotensin system in heart failure. *Am*
604 *Heart J* 1993;126:756-60.

605 74. Paul M, Pinto YM, Schunkert H, *et al.* Activation of the renin-angiotensin system in heart failure
606 and hypertrophy--studies in human hearts and transgenic rats. *Eur Heart J* 1994;15 Suppl
607 D:63-7.

608 75. van Thiel BS, van der Pluijm I, te Riet L, *et al.* The renin-angiotensin system and its involvement
609 in vascular disease. *Eur J Pharmacol* 2015;763:3-14.

610 76. Sedan O, Dolnikov K, Zeevi-Levin N, *et al.* 1,4,5-Inositol Trisphosphate-Operated Intracellular
611 Ca²⁺ Stores and Angiotensin-II/Endothelin-1 Signaling Pathway Are Functional in Human
612 Embryonic Stem Cell-Derived Cardiomyocytes. *STEM CELLS* 2008;26:3130-8.

613 77. Sedan O, Dolnikov K, Zeevi-Levin N, *et al.* Human embryonic stem cell-derived cardiomyocytes

614 can mobilize 1,4,5-inositol trisphosphate-operated $[Ca^{2+}]_i$ stores: the functionality of
615 angiotensin-II/endothelin-1 signaling pathways. *Ann N Y Acad Sci* 2010;1188:68-77.

616 78. Foldes G, Mioulane M, Wright JS, *et al.* Modulation of human embryonic stem cell-derived
617 cardiomyocyte growth: a testbed for studying human cardiac hypertrophy? *J Mol Cell*
618 *Cardiol* 2011;50:367-76.

619 79. Foldes G, Matsa E, Kriston-Vizi J, *et al.* Aberrant alpha-adrenergic hypertrophic response in
620 cardiomyocytes from human induced pluripotent cells. *Stem Cell Reports* 2014;3:905-14.

621 80. Nunes SS, Feric N, Pahnke A, *et al.* Human Stem Cell-Derived Cardiac Model of Chronic Drug
622 Exposure. *ACS Biomaterials Science & Engineering* 2017;3:1911-21.

623 81. Mishra JS, More AS, Gopalakrishnan K, *et al.* Testosterone plays a permissive role in angiotensin
624 II-induced hypertension and cardiac hypertrophy in male rats. *Biol Reprod* 2019;100:139-
625 48.

626 82. Shekhar A, Lin X, Lin B, *et al.* ETV1 activates a rapid conduction transcriptional program in
627 rodent and human cardiomyocytes. *Sci Rep* 2018;8:9944.

628 83. Amatori S, Persico G, Fanelli M. Real-time quantitative PCR array to study drug-induced
629 changes of gene expression in tumor cell lines. *Journal of Cancer Metastasis and*
630 *Treatment* 2017;3:90-9.

631 84. Bray NL, Pimentel H, Melsted P, *et al.* Near-optimal probabilistic RNA-seq quantification. *Nat*
632 *Biotechnol* 2016;34:525-7.

633 85. Harrow J, Frankish A, Gonzalez JM, *et al.* GENCODE: the reference human genome annotation
634 for The ENCODE Project. *Genome Res* 2012;22:1760-74.

635

a **b** **c** **d**

Fold Change of expression (Ang II-
treated CM vs. untreated CM)

a**GSE116547** **b****GSE137920**

a **b**