WOX9 functions antagonistic to STF and LAM1 to regulate leaf blade expansion in *Medicago truncatula* and *Nicotiana sylvestris*

Tezera W. Wolabu^{a,b}, Hui Wang^{a,c}, Dimiru Tadesse^a, Fei Zhang^{a,d}, Marjan Behzadrad^a
Varvara E. Tvorogova^{a,c}, Haggagi Abdelmageed^{a,f}, Ye Liu^g, Naichong Chen^{a,h}, Jianghua
Chenⁱ, Randy D. Allen^{a,h}, and Million Tadege^{a,j,l}

^aInstitute for Agricultural Biosciences, Oklahoma State University, Ardmore, OK 73401, USA.

^bNoble Research Institute, LLC, Ardmore, OK 73401, USA.

^cCollege of Grassland Science and Technology, China Agricultural University, Beijing, China.

^dDepartment of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06520-8104, USA.

^eDepartment of Genetics and Biotechnology, St. Petersburg State University, St. Petersburg, Russia.

^fDepartment of Agricultural Botany, Faculty of Agriculture, Cairo University, Giza 12613, Egypt

^gSchool of Life Science, University of Science and Technology of China, Hefei, Anhui 230027, China

^hDepartment of Biochemistry and Molecular Biology, Oklahoma State University, Stillwater, OK, USA

ⁱCAS Key Laboratory of Topical Plant Resources and Sustainable Use, CAS Center for Excellence in Molecular Plant Sciences, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Kunming, Yunnan 650223, China.

^jDepartment of Plant and Soil Sciences, Oklahoma State University, Stillwater, OK, USA

^lTo whom correspondence should be addressed. Email: million.tadege@okstate.edu.

Running Title: WOX9 negatively regulates leaf blade outgrowth

One sentence summary:

WOX9 negatively regulates blade outgrowth antagonizing STF function but directly repressed by STF indicating WOX-mediated homeostasis in cell proliferation and differentiation during leaf morphogenesis.

Corresponding author:

Million Tadege

Institute for Agricultural Biosciences, Oklahoma State University.

3210 Sam Noble Parkway, Ardmore, OK 73401, U.S.A

Tel: 580-224-0629.

Fax: 580-224-0624

Email: million.tadege@okstate.edu

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Million Tadege (million.tadege@okstate.edu).

Abstract

Plant specific WOX family transcription factors are known to regulate embryogenesis, meristem maintenance and lateral organ development. Modern clade WOX genes function through a transcriptional repression mechanism, and the intermediate clade transcriptional activator WOX9 functions with the repressor WOX genes in embryogenesis and meristems maintenance, but the mechanism of this interaction is unclear. *WOX1* homologues *STF* and *LAM1* are required for leaf blade outgrowth in *Medicago truncatula* and *Nicotiana Sylvestris*, respectively. Here we show that *WOX9* negatively regulates leaf blade outgrowth and functions antagonistically to *STF* and *LAM1*. While *NsWOX9* ectopic expression enhances the *lam1* mutant phenotype, and antisense expression partially rescues the *lam1* mutant, both overexpression of *NsWOX9* and knockout by CRISPR/Cas9 genome editing in *N. sylvestris* resulted in a range of severe leaf blade distortions, indicating that controlled negative regulation by *NsWOX9* is required for proper blade development. Our results indicate that direct repression of *WOX9* transcriptional activation activity by the transcriptional repressor *STF/LAM1* is required for correct blade architecture and patterning in *M. truncatula* and *N. sylvestris*. These findings suggest that a balance between transcriptional activation and repression mechanisms by direct interaction of activator and repressor WOX genes may be required for cell proliferation and differentiation homeostasis, and could be an evolutionarily conserved mechanism for the development of complex and diverse morphology in higher plants.

Introduction

WUSCHEL-related homeobox (WOX) factors are plant-specific transcriptional regulator proteins that contain a DNA binding homeodomain similar to WUSCHEL (WUS), the founding member of the family from Arabidopsis. Several elegant studies demonstrated that the WOX family is involved in the regulation of a wide range of key developmental programs ranging from the modulation of zygotic development and embryogenesis by WOX2, WOX8, and WOX9 (Haecker et al., 2004; Breuninger et al., 2008; Ueda et al., 2011) to maintenance of shoot and root apical meristems orchestrated by WUS and WOX5, respectively (Mayer et al., 1998; Sarkar, 2007), along with several other

developmental pathways (Matsumoto and Okada, 2001; Park et al., 2005; Deyhle et al., 2007; Shimizu et al., 2009; Vandenbussche et al., 2009; Hirakawa et al., 2010; Ji et al., 2010; Tadege et al., 2011b; Nakata et al., 2012).

WUS and its homologues in other species, including *TERMINATOR (TER)* in petunia, *ROSULATA (ROA)* in *Antirrhinum*, and *HEADLESS (HDL)* in *Medicago*, are required for shoot apical meristem (SAM) maintenance (Laux et al., 1996; Mayer et al., 1998; Stuurman et al., 2002; Kieffer et al., 2006; Meng et al., 2019; Wang H, 2019). Loss of *WUS* function in the *wus-1* *Arabidopsis* mutant results in premature termination and arrest of the SAM and floral meristem, but the SAM re-establishes itself to resume growth while the process repeats itself, leading to altered plant morphology (Laux et al., 1996; Mayer et al., 1998). In the *hdl* mutant of *M. truncatula*, termination of SAM and axillary meristems is permanent. Since the SAM fails to re-establish itself, these plants only make leaves throughout development (Tadege et al., 2015; Meng et al., 2019; Wang H, 2019). The *hdl* mutant also shows altered leaf shape (Meng et al., 2019; Wang et al., 2019), which was not detected in the *wus* mutant. However, the *wus wox1 prs* triple mutant showed a stronger leaf phenotype than the *wox1 prs* double mutant (Zhang F, 2015), suggesting that *WUS* may also have a redundant function in leaf development. Although *WUS* transcript is specifically expressed in the organizing center (OC) of the SAM (Mayer et al., 1998), it is likely that a non-cell autonomous signal from *WUS* may contribute to blade outgrowth, since the *WUS* protein itself is shown to move from the OC to the stem cell region (Yadav et al., 2011; Daum et al., 2014).

An intimate connection exists between *WUS* and the phytohormone cytokinin, and this appears to be true with some other *WOX* orthologs (Tadege and Mysore, 2011; Tadege, 2016; Wang, 2017). While *WUS* activity is modulated by cytokinin in the SAM and axillary meristem (Wang et al., 2017; Snipes et al., 2018), *WUS* promotes cytokinin activity in the shoot stem cell niche by repressing type-A *ARABIDOPSIS RESPONSE REGULATOR (ARR)* genes *ARR5*, *ARR6*, *ARR7*, and *ARR15* (Leibfried et al., 2005) to activate cell proliferation, and *WUS* physically interacts with the transcriptional co-repressor *TOPELESS (TPL)* to repress target genes that promote cell differentiation (Kieffer et al., 2006; Causier et al., 2012; Yadav et al., 2013). Genes that encode polarity factors which impart adaxial or abaxial identity and maintenance of a differentiated state

to the leaf blade tissues are among the targets directly repressed by WUS (Yadav et al., 2013). This uncovers a mechanism by which WUS maintains undifferentiated stem cells in the SAM, in addition to the well-established CLE peptide signaling (Schoof et al., 2000; Somssich et al., 2016; Hu C, 2018). Although WUS is reported to be a bifunctional transcription factor exhibiting both transcriptional repression and activation activities (Ikeda et al., 2009), its SAM maintenance activity and interaction with cytokinin are shown to be linked to its WUS box (Ikeda et al., 2009; Dolzblasz et al., 2016; Snipes et al., 2018), suggesting that WUS primarily functions as a transcriptional repressor. Interestingly, *WUS* is reported to be activated by *WOX9/STIP*, which is also required for shoot meristem maintenance (Wu et al., 2005) and embryo development (Wu et al., 2007; Ueda et al., 2011). However, *WOX9* is reported to be a strong transcriptional activator (Lin et al., 2013), and it is unclear whether *WUS* and *WOX9* employ the same mechanism in shoot meristem maintenance.

The *WOX9* gain-of-function *stip-D* mutant displays wavy leaf margins indicating problems with cell division in leaf primordium (Wu et al., 2005). But, the *stip* loss-of-function mutant is arrested at the seedling stage (Wu et al., 2005), and it is unclear if *WOX9/STIP* plays a specific role in leaf blade development. However, *WOX* function in leaf development is not uncommon in *Arabidopsis* and several other species. *WOX1* and *PRS/WOX3* in *Arabidopsis* (Vandenbussche et al., 2009; Nakata et al., 2012) and their homologues in maize, rice, petunia, *Medicago* and woodland tobacco regulate leaf blade development (Nardmann et al., 2004; Vandenbussche et al., 2009; Tadege et al., 2011b; Zhuang et al., 2012; Cho et al., 2013; Ishiwata et al., 2013). Unlike leaf polarity factors that are adaxial or abaxial-specific (Waites et al., 1998; Sawa, 1999; Siegfried, 1999; Kerstetter et al., 2001; McConnell et al., 2001; Iwakawa et al., 2002), the *WOX* genes *STF*, *WOX1* and *PRS*, are expressed in the middle at the adaxial-abaxial juxtaposition to control medial-lateral outgrowth of the leaf blade (Tadege et al., 2011a,b; Nakata and Okada, 2012; Nakata et al., 2012), suggesting a novel mechanism for blade expansion.

M. truncatula *STF* or *N. sylvestris* *LAM1* is a transcriptional repressor (Lin, 2013; Lin et al., 2013) and its repression activity is conferred by its WUS box and STF box motifs (Zhang et al., 2014; Zhang and Tadege, 2015). The DNA binding STF homeodomain (HD) and the repression motifs (WUS box and STF box) are critically

required for blade outgrowth function (Lin et al., 2013; Zhang et al., 2014; Zhang et al., 2019). Interestingly, all of the WUS clade Arabidopsis WOX transcription factors (WUS and WOX1-WOX7), which have transcriptional repression activity, can substitute for LAM1 function (Lin et al., 2013), suggesting that modern/WUS clade WOX members have a conserved transcriptional repression mechanism in meristem maintenance and lateral organ development, with specificity conferred by cis elements that drive specific expression patterns.

Intermediate clade WOX members have intact HD but lack repression domains. Here we show that homologues of the intermediate clade transcriptional activator WOX9, namely, MtWOX9-1, MtWOX9-2, and NsWOX9, negatively regulate leaf blade outgrowth in *M. truncatula* and *N. sylvestris*. These factors function antagonistically to STF or LAM1, and exacerbate the *lam1* phenotype. Suppression of *NsWOX9* transcript levels by antisense technology partially rescues the *lam1* mutant leaf blade while the introduction of knockout mutations in the native *NsWOX9* gene by multiplex gRNA genome editing severely affected leaf blade symmetry and expansion. Our results suggest that direct and antagonistic interactions between transcriptional repressor and activator WOX genes may be important to balance cell proliferation with differentiation in acquiring complex morphology in higher plants.

Results

Ectopic expression of *WOX9* enhances *stf* and *lam1* mutant phenotypes.

We have previously shown that, while the WUS clade repressor *WOX* genes of Arabidopsis including *WUS* and *WOX1-WOX7* complement the *lam1* mutant phenotype when driven by the *STF* promoter, the intermediate clade *WOX9* expression exacerbates the *lam1* mutant phenotype (Lin et al., 2013). Therefore, we decided to investigate whether the unique activity of *WOX9* is conserved in *Medicago truncatula* and *Nicotiana sylvestris* and determine its biological significance in leaf development. We isolated orthologous coding sequences for *AtWOX9* from *M. truncatula* (*MtWOX9*) and *N. sylvestris* (*NsWOX9*) and created expression constructs that placed these sequences under control of the *STF* promoter to ectopically express these genes in the *stf* and *lam1* mutant plants. *stf* and *lam1* are severe leaf blade mutants in *M. truncatula* and *N. sylvestris*, respectively, caused by mutation of the *WOX1* orthologs *STF* and *LAM1* (Figures 1B and 1E). The *M. truncatula* genome contains two *WOX9*-like sequences here designated as *MtWOX9-1* and *MtWOX9-2* (Figure S1 and S2). We introduced *MtWOX9-1* driven by the *STF* promoter (*STF::MtWOX9-1*) first into the *stf* *M. truncatula* plants and eight independent transgenic lines were generated. Expression of *MtWOX9-1* in the *stf* mutant background was confirmed by RT-PCR assays. All of these transgenic lines displayed strongly enhanced mutant phenotype with much narrower leaves and thinner stems compared to the *stf* mutant (Figures 1A to 1C). In addition, leaves and stems were significantly shorter in length, leading to a dwarf phenotype that was not characteristic of the *stf* mutant phenotype (Figures 1B and 1C). Similarly, introduction of this construct into the *lam1* mutant background severely affected both leaf length and width, exacerbating the *lam1* mutant phenotype (Figures 1D to 1F). These results indicate that both the *stf* and *lam1* mutants respond similarly to the activity of *MtWOX9-1*, consistent with the effect of *AtWOX9* ectopic expression in the *lam1* mutant (Lin et al., 2013). We also transformed *35S::MtWOX9-1*, *35S::MtWOX9-2* and *35S::NsWOX9* into the *lam1* mutant and obtained severely enhanced mutant phenotypes similar to the *STF::MtWOX9-1* expressing *lam1* lines (Figure 1G to 1I), indicating that these three genes have similar effects on leaf blade outgrowth. In most cases, these transgenic leaves displayed approximately five-fold reduction in leaf length, but this effect appeared to be dependent

Figure 1. Ectopic expression of *WOX9* enhances *stf* and *lam1* mutant phenotypes.

(A) Untransformed *M. truncatula* wild type (WT) (R108) plant. (B) Phenotype of untransformed *stf* mutant. (C) *stf* mutant transformed with *STF::MtWOX9-1*. (D) Untransformed *N. sylvestris* WT plant. (E) Phenotype of untransformed *lam1* mutant. (F) *lam1* mutant transformed with *STF::MtWOX9-1*. (G) *lam1* mutant transformed with *35S::MtWOX9-1*. (H) *lam1* mutant transformed with *35S::MtWOX9-2*. (I) *lam1* mutant transformed with *35S::NsWOX9*. Plants were 10-weeks (E and F) or 5-weeks old (all the rest). Scale bars: 10 cm.

on the level of *WOX9* transgene expression since plants with high level of transgene expression showed more severe phenotypes compared to *lam1* plants with low level of transgene expression (Supplemental Figure 3).

The enhancement of *stf* and *lam1* mutant phenotypes associated with *WOX9* expression suggested to us that *WOX9* acts in opposition to *STF/LAM1* in leaf blade development. Therefore, we introduced an *NsWOX9-antisense* construct into the *lam1* plants to see if reduced levels of *NsWOX9* transcripts could alleviate the mutant phenotype. Indeed, expression of an *NsWOX9-antisense* construct had the opposite effect of *NsWOX9* overexpression, partially rescuing the *lam1* mutant leaf phenotype (Figure 2). However, this partial complementation was limited, and the antisense plants still appeared bushy and failed to make stems. Nonetheless, unlike the untransformed *lam1* mutant control, the *NsWOX9-antisense* leaves showed distinct petioles and blades especially at early stages of development (Figures 2A and 2 B), and the blades were variously branched and curled resulting in unusual leaf structure at maturity (Figures 2C to 2E). These leaf phenotypes suggest that blade outgrowth initiation has significantly progressed in the *NsWOX9-antisense* plants but perhaps aborted before completion.

To more completely evaluate the effects altered *WOX9* expression on the *lam1* mutant phenotype, we carried out structural examination of leaf tissues of WT, *lam1* mutant, *WOX9* overexpressing *lam1* and *WOX9* antisense *lam1* plants (Figures 3A to 3D).

Figure 2. *NsWOX9-antisense* partially rescued *lam1* mutant phenotype.
 (A) Phenotype of *lam1* mutant transformed with *35S::GUS* as control at three weeks of age. Inset on the right is detached leaf close up.
 (B) Partially complemented *lam1* phenotype transformed with *35S::NsWOX9-antisense* construct at three weeks, the inset is close up of a partially complemented leaf blade. Inset on the right is detached leaf close up.
 (C) Partially complemented *lam1* phenotype transformed with *35S::NsWOX9-antisense* construct at seven weeks.
 (D) Representative individual leaves from *35S::NsWOX9-antisense/lam1* plants.
 (E) A magnified view of a leaf in (D). Note the branching and curling of leaves especially in older *35S::NsWOX9-antisense/lam1* plants.
 Scale bars: A-D, 5 cm, E, 1.5 cm.

Transverse sections through the leaf blades showed that the *lam1* leaves had vestigial blade strips at the position of wild type blades (Figure 3F), but these strips were completely absent and blades became fully radialized in *NsWOX9* overexpressing *lam1* lines (Figure 3G). In *NsWOX9-antisense lam1* lines, on the other hand, distinct blade outgrowth was apparent but the nascent blades were not fully expanded compared to the wild type leaves (Figure 3E and 3H), confirming that blade development in the *NsWOX9-antisense* plants initiated more effectively than in the *lam1* plants but was not completed. Taken together, these results indicate that *WOX9* functions oppose those of *STF/LAM1* to negatively regulate leaf blade outgrowth in two unrelated eudicot species *M. truncatula* and *N. sylvestris*.

***WOX9* overexpression severely affects leaf architecture**

To further examine the effect of *WOX9* in leaf blade outgrowth, we introduced *35S::MtWOX9-1*, *35S::MtWOX9-2* and *35S::NsWOX9* into wild type (WT) *N. sylvestris*. Analysis of over 20 independent transgenic lines for each construct revealed that all transgenic lines displayed an array of leaf phenotypes that can generally be grouped into

Figure 3. Transverse section of the leaf blade showing enhancement of the *lam1* blade by 35S::NsWOX9 and partial complementation by 35S::NsWOX9-antisense.

(A) *N. sylvestris* WT leaf blade.

(B) Leaf blade of untransformed *lam1* mutant control.

(C) Leaf blade of *lam1* transformed with 35S::NsWOX9.

(D) Partially complemented leaf blade of *lam1* transformed with 35S::NsWOX9-antisense.

(E) Transverse section of *N. sylvestris* WT leaf blade.

(F) Transverse section of untransformed *lam1* mutant leaf blade.

(G) Transverse sections of *lam1* leaf blade transformed with 35S::NsWOX9 showing radialized blade.

(H) Transverse sections of *lam1* leaf blade transformed with 35S::NsWOX9-antisense showing blade outgrowth. Arrows indicate blade tissue in (E) and (H), vestigial blade stripes in (F) and position of blade in (G). Scale bars: 50 µm.

severe and mild based on the phenotype strength. The wild type *N. sylvestris* leaf blade is a well-expanded flat lamina with smooth margin and distinctive pinnate venation pattern (Figure 4A). In plants with the severe phenotype *WOX9* overexpression completely disrupted this pattern resulting in highly distorted leaf forms. These include narrow and downward curling blades, deep margin serrations, disorganized venation patterns, uneven blade surfaces, as well as retarded plant growth with 2 to 5 tillers and additional leaves leading to a bushy appearance until stem elongation at a later developmental stage (Figures 4B, 4D, 4E, 4G; Supplemental Figure 4). While most overexpressing plants showed this strong phenotype, some exhibited a mild phenotype where the blade margin, shape and venation patterns were largely intact but with puckered and uneven blade surfaces (Figures 4C and 4F; Supplemental Figure 4). *MtWOX9-1* overexpression in *M. truncatula* also produced downward curling and narrower leaves similar to that seen in *N. sylvestris* (Supplemental Figure 5), albeit more mild, with an insignificant effect on tillering. These *WOX9* ectopic expression phenotypes in wild type and in *stf* and *lam1* mutants suggest that *WOX9* could be a negative regulator of leaf blade outgrowth antagonizing the function of *STF/LAM1*.

Deleting *NsWOX9* using multiplex gRNA CRISPR/Cas9 genome editing in *N. sylvestris* alters blade symmetry and expansion

245 To gain insight into the function of the endogenous *WOX9* gene in wild type plants, we
 246 disrupted *NsWOX9* in *N. sylvestris* using CRISPR/Cas9 genome editing technology. We
 247 constructed an *NsWOX9*-multiplex gRNA-CRISPR/Cas9 vector containing three guide
 248 RNAs; gRNA1, gRNA2 and gRNA3 (Figure 5A), and introduced this construct into *N.*
 249 *sylvestris*. A total of 24 transgenic lines were generated and examined for mutations at
 250 the targeted regions. Sixteen putative mutant lines were identified by PCR amplification
 251 of the target regions using specific primers, and Sanger sequencing, indicating a 67%
 252 overall mutagenesis efficiency. Out of these sixteen putative mutants, five representative
 253 lines were selected for further characterization of their mutant phenotypes (Figure 5B).
 254 Target site sequence analysis of the five *NsWOX9*-CRISPR-mutants labeled here as
 255 *NsWOX9-1*, *NsWOX9-2*, *NsWOX9-13*, *NsWOX9-18* and *NsWOX9-22* using SeqMan Pro
 256 15.0.1 (DNASTAR software) revealed five different patterns of deletions ranging from
 257 14 to 183 bp (Figure 5B). *NsWOX9-1* and *NsWOX9-13* had identical 14 bp deletions in
 258 gRNA3. *NsWOX9-2* showed two deletion events at gRNA2 and gRNA3 with 7 and 21
 259 bp deletions, respectively. *NsWOX9-22* showed a 73 bp deletion spanning the upstream

Figure 6. *MtWOX9-1* is weakly expressed in leaf but strongly upregulated in the *stf* mutant leaf of *M. truncatula* and corresponding mutants in *N. sylvestris* and Arabidopsis.

(A) RT-qPCR analysis showing relative expression of *MtWOX9-1* in different tissues *M. truncatula*. leaf, stem, and shoot apex were from 4-week old plants, pods and seeds were 10 days after pollination, flower at anthesis, all the rest were at seedling stage.

(B)-(I) GUS staining in *MtWOX9-1::GUS* transformed lines of *M. truncatula* showing fully expanded leaf (B), shoot apex with folded leaves (C), stem (D), flower with unstained anthers (E), very young pod with seeds (F), older pod with seeds (G), immature seeds (H), and matured seeds (I).

(J) Relative expression of *MtWOX9-1* in 4-week old *stf* mutant leaf in *M. truncatula*.

(K) Relative expression of *NsWOX9* in 4-week old *lam1* mutant leaf of *N. sylvestris*.

(L) Relative expression of *AtWOX9* in the leaf of 4-week old Arabidopsis *wox1 prs* double mutant. Error bars indicate $\pm$ SE (n=3). Asterisks indicate significant difference from the control (*p<0.05, **p<0.01, student t-test).

expression in flowers (Figure 6A). Highest expression of *MtWOX9-2*, on the other hand, was detected in the leaves followed by expression in shoot apices (Supplemental Figure 6). To examine the spatial distribution of expression, we fused a 3kb promoter region of *MtWOX9-1* upstream of the translational start codon to the β -glucuronidase (*GUS*) coding region, and transformed it into *Medicago* R108 leaf explants. GUS staining analysis revealed that expression in the mature leaf was relatively weak with particularly strong expression in the pulvinus at the base of the leaflets (Figure 6B). Strong expression was detected in the shoot apex and immature leaves, followed by flowers and stems (Figures 6C to 6E) but, no staining was detected in the anthers (Figure 6E). Very strong expression of *MtWOX9-1* was detected in immature pods and seeds at early stages of development (Figures 6F to 6I). Interestingly, expression in the seed became progressively restricted as the seed develops, and confined only to the hilum in the mature seed (Figure 6I). These expression patterns suggest that the *MtWOX9-1* function may be more important at the early stages of development, particularly during embryogenesis and leaf morphogenesis.

To investigate the mechanistic relationship between *WOX9* and positive regulators of blade outgrowth, we examined the leaf blade expression levels of *WOX9* in *stf*, *lam1*, and *wox1 prs* mutants of *M. truncatula*, *N. sylvestris*, and *A. thaliana* respectively. RT-qPCR analyses showed that expression of *MtWOX9-1*, *NsWOX9*, and *AtWOX9* was upregulated by 2-4 fold in the leaves of *stf*, *lam1* and *wox1 prs* mutants compared to their respective wild type levels (Figures 6J to 6L), indicating that *WOX9* may be directly or indirectly repressed by the action of STF/LAM1/WOX1 in wild type leaves.

To examine whether *MtWOX9-1* is a direct target of STF, we performed a dexamethazone (DEX) induction experiment using the glucocorticoid (GR) system in the presence of the protein synthesis inhibitor, cycloheximide (CHX). Analysis was performed in 4-week old *stf* mutant plants transformed with the *35S::YFP-GR-STF* construct. The shoot apex and young leaves of the transgenic plants were treated with both DEX and CHX for 3 hours, and *MtWOX9-1* transcript accumulation was monitored by RT-qPCR in the leaves with and without the induction treatment. Our results showed that the expression of *MtWOX9-1* was reduced by approximately 60% in the DEX and CHX treated lines compared to the control CHX alone (Figure 7A). Since new protein synthesis is inhibited by CHX in the treated lines, this result suggests that *MtWOX9-1* may be directly repressed by STF. We repeated this experiment in *N. sylvestris* using *35S::YFP-GR-LAMI* fusion and found approximately 60% repression of *NsWOX9* upon *LAMI* induction by DEX plus CHX treatment (Figure 7B). However, in the reciprocal experiment, induction of *NsWOX9* expression had no significant effect on *LAMI* expression (Figure 7C), indicating that *NsWOX9* does not regulate *LAMI* transcription.

To determine if STF reduces *MtWOX9-1* transcript accumulation by directly targeting its promoter, we performed dual luciferase assay in Arabidopsis protoplasts using the Firefly-Renilla Dual-luciferase assay system (Promega). In the reporter construct, a 1kb promoter region of *MtWOX9-1* upstream of the translation start codon was fused to a mini 35S promoter driving the luciferase reporter gene (Figure 7D), while the effector constructs were made using either STF, or GUS as a negative control, both driven by the 35S promoter (Figure 7D). Consistent with the above results, co-expression of the STF effector in the protoplast almost fully abolished luciferase luminescence

Figure 7. WOX9 expression is directly repressed by GR induction of STF or LAM1 in the presence of cyclohexamide, and by STF in dual luciferase assay.

(A) Relative expression of *MtWOX9-1* in 35S::YFP-GR-STF transformed *M. truncatula* lines with CHX or DEX+CHX treatment for 3 hours in 4-week old leaves.

(B) Relative expression of *NsWOX9* in 2 independently transformed 35S::YFP-GR-LAM1 *N. sylvestris* lines with CHX alone, CHX+DEX or DEX alone treatments for 6 hours in 4-weeks old leaves.

(C) Relative expression of *LAM1* in 35S::YFP-GR-NsWOX9 transformed 2 independent *N. sylvestris* lines with CHX alone, CHX+DEX or DEX alone treatments for 6 hours in 4-weeks old young leaves. Relative gene expression was determined by RT-qPCR analyses.

(D) Schematic representation of reporter and effector constructs used in the dual luciferase assay.

(E) Relative expression of luciferase activity (luminescence) in the presence of 35S::STF effector compared with the 35S::GUS control in Arabidopsis protoplasts. Error bars indicate $\pm$ SE (n=3). Asterisks indicate significant difference from the control (*p<0.05, **p<0.01, ***p<0.001, student t-test).

compared to the GUS effector control (Figure 7E), indicating that this 1 kb region of the *MtWOX9-1* promoter is sufficient for STF-dependent repression of transcription.

To determine if STF indeed binds to the *MtWOX9-1* promoter *in vitro* and *in vivo*, we performed electrophoretic mobility shift assay (EMSA) using biotin labeled probes, and chromatin immunoprecipitation (ChIP) assay, using anti GFP antibody. We previously reported that STF binds preferentially to “AT-rich” DNA elements without a strong consensus sequence (Zhang et al., 2014). We screened six such selected regions in the 3 kb upstream region of the *MtWOX9-1* promoter, and found that the MBP-STF fusion protein was able to bind to three of them (Figure 8A). These STF-binding elements are located at -22, -226 and -491 bp upstream of the *MtWOX9-1* CDS while binding was not detected with the control maltose binding protein (MBP) alone (Figure 8B). Each of these sites were significantly competed by addition of 50-fold excess of the respective unlabeled probes, indicating binding specificity. This shows that, at least *in vitro*, STF can directly bind to multiple sites within the 1 kb fragment of the *MtWOX9-1* promoter, consistent with the dual luciferase assay and GR induction experiments.

Figure 8. STF directly binds to the *MtWOX9-1* promoter in EMSA and ChIP assays.

(A) Schematic representation of the *MtWOX9-1* promoter and CDS regions tested for EMSA and ChIP assays. The 3 promoter regions tested are indicated as P1, P2 and P3 and the *MtWOX9-1* coding region as C1. (B) EMSA showing MBP-STF bound to the biotin-labeled probe at P1, P2 and P3 promoter fragments but not the MBP control alone. Fifty-fold excess of unlabeled P1, P2 or P3 DNA was used to compete with the respective labeled probe (right lanes). (C) *MtWOX9-1* promoter enrichment at P1, P2, P3 regions and CDS C1. Chromatin precipitated with anti-GFP antibody from *35S::STF-GFP* and *35S::GFP* control lines were compared. Purified DNA from the chromatin were used as templates for qPCR. Note that P2 is highly enriched in *35S::STF-GFP* samples. Error bars indicate $\pm$ SE (n=3). Asterisks indicate statistical significance (* p <0.05, *** p <0.001, student t-test).

336 To confirm that STF binds to the *MtWOX9-1* promoter in vivo, we performed
 337 ChIP assays in leaves of Arabidopsis *wox1 prs* double mutant plants transformed with
 338 *pMtWOX9-1::MtWOX9-1* construct. Protoplasts were isolated from these transgenic
 339 plants and transformed with *35S::STF-GFP* fusion from which chromatin was isolated
 340 for analysis. ChIP-qPCR analysis revealed that among the three tested promoter regions
 341 (P1, P2 and P3), STF was highly enriched at P2 with significant enrichment also at P3
 342 (Figure 8C). In contrast, no significant enrichment was detected at P1 or within the
 343 *MtWOX9-1* coding region (C1), indicating that STF binds strongly at the P2 position and
 344 to a limited extent at the P3 region *in planta*. Taken together, these results indicate that
 345 STF directly binds to the proximal region of the *MtWOX9-1* promoter in *M. truncatula*,
 346 and represses its activity, and that STF and WOX9 function antagonistically to regulate
 347 leaf blade outgrowth.

348
 349

Discussion

Plant specific WOX transcription factors regulate a variety of plant developmental programs from embryogenesis to shoot apical meristem maintenance and lateral organ development (Mayer et al., 1998; Schoof et al., 2000; Lohmann et al., 2001; Matsumoto and Okada, 2001; Nardmann et al., 2004; Sarkar, 2007; Breuninger et al., 2008; Shimizu et al., 2009; Vandenbussche et al., 2009; Ji et al., 2010; Tadege et al., 2011b; Nakata et al., 2012). WOX family members are also known for their promiscuous ability to substitute for each other's functions. For example, in Arabidopsis, WUS complements the *prx/wox3* and *wox5* mutants, which are defective in floral organ development and root apical meristem maintenance, respectively (Sarkar, 2007; Shimizu et al., 2009). Conversely, members of the WUS clade WOX genes (WOX1-WOX7), with the exception of WOX4, can substitute for WUS function in stem cell maintenance (Dolzblass et al., 2016). Arabidopsis WUS and WOX1-WOX7 can also complement the *lam1* leaf blade mutant of *N. sylvestris* (Tadege et al., 2011b; Lin et al., 2013). Here we show that the *M. truncatula* and *N. sylvestris* WOX9 homologues, *MtWOX9-1*, *MtWOX9-2* and *NsWOX9* function antagonistically to *STF/LAM1* by negatively regulating blade outgrowth. Ectopic expression of these genes enhanced the *stf* and *lam1* leaf mutant phenotypes, and severely affected blade expansion and morphology in wild type *N. sylvestris* with a range of phenotypes (Figures 1 and 4). Conversely, reducing *NsWOX9* transcript levels in the *lam1* mutant with antisense technology partially complemented the mutant phenotype (Figure 2), indicating that WOX9 antagonizes leaf blade outgrowth in the *STF/LAM1* pathway. However, complete knockout of *NsWOX9* by CRISPR/Cas9 genome editing technology in the wild type background resulted in a range of leaf blade deformations including lack of bilateral symmetry, altered venation patterns, narrow blades and bushy shoots (Figure 5), indicating that WOX9 function is required for proper leaf blade development.

In petunia and tomato, WOX9 homologues are involved in inflorescence development and architecture (Lippman et al., 2008; Rebocho et al., 2008; Costanzo E, 2014). Both the *evergreen* (*evg*) mutant in petunia (Rebocho et al., 2008) and *compound inflorescence* (*s*) in tomato (Lippman et al., 2008), which dramatically alter the wild type inflorescence architecture are caused by lesions in WOX9 homologues. The *s* allele in

tomato results in a highly branched structure with hundreds of flowers, which increases fruit production and may have been selected by breeders ((Lippman et al., 2008). In the *evg* mutant of petunia, on the other hand, the inflorescence stem often fails to bifurcate after the formation of bracts and continues to grow as a single thickened stem without physical separation of the floral meristem (FM) and inflorescence meristem (IM), leading to a fasciated appearance (Rebocho et al., 2008). Unlike *s*, the *evg* FM also fails to produce floral organs suggesting that EVG is required for inflorescence bifurcation and floral organ identity, though *evg* mutants are indistinguishable from wild type during early vegetative growth (Rebocho et al., 2008). Thus, in tomato and petunia, *WOX9* homologues appear to have opposite effects specific to inflorescence development and architecture. However, both tomato and petunia have a cymose inflorescence pattern (determinate growth) and it is unclear whether these inflorescence-associated defects are specific to cymose or are also exhibited by racemose (indeterminate growth) and panicle (mixed inflorescence) inflorescences. At least in *Arabidopsis* (racemose inflorescence), the role of *WOX9* appears not to be restricted to inflorescence development, and in rice (mixed inflorescence), *WOX9* is involved in uniform tiller growth and development (Wang et al., 2014; Fang et al., 2020).

In *Arabidopsis*, *WOX9*, also called *STIMPY* (*STIP*), is required for meristem growth and maintenance and positively regulates *WUS* (Wu et al., 2005). *stip* mutants display arrested growth at an early stage of development but can be fully rescued by sucrose (Wu et al., 2005). *STIP*/*WOX9* is shown to mediate cytokinin signaling during shoot meristem establishment and, together with *WOX2* and *WOX8*, regulates zygote and embryo polarity patterning (Wu et al., 2007; Breuninger et al., 2008; Skylar et al., 2010; Ueda et al., 2011). *WOX9* homologues in other species are also reported to be involved in promoting somatic embryogenesis (Gambino et al., 2011; Tvorogova, 2019). In all of these examples, the function of *WOX9* appears to center on cell proliferation and/or meristematic competence for proper plant growth and development. Our observation of the effect of *WOX9* overexpression and knockout in *Medicago* and woodland tobacco leaf development is consistent with these findings, and may reflect a conserved molecular function in cell proliferation and differentiation during growth and development. For example, the *Arabidopsis* gain-of-function mutant *stip-D* displays

wavy leaf margins and increased number of axillary shoots leading to a bushy phenotype (Wu et al., 2005). A phenotype similar the wavy margins and bushy shoots seen in all *MtWOX9-1*, *MtWOX9-2* and *NsWOX9* overexpressing *N. sylvestris* transgenic lines (Figure 4), which suggests misregulation of cell proliferation in leaf primordia. The observation that both overexpression and knockout of *WOX9* in *N. sylvestris* led to narrow leaf and bushy shoot phenotypes suggests that *WOX9* may be involved in maintaining a balance between cell proliferation and differentiation, necessitating that *WOX9* transcripts be maintained at a required optimum. However, the mechanism of *WOX9* function and control of its steady state transcript levels during leaf morphogenesis and maturation is not well understood. We previously reported *AtWOX9* to be a transcriptional activator, based on its unique effects on the *lam1* mutant of *N. sylvestris* (Lin et al., 2013), which could provide insight into its molecular function.

The Arabidopsis *WOX* family has been divided into three clades based on phylogenetic analysis: the modern/*WUS* clade (*WUS*, *WOX1-7*), intermediate clade (*WOX8, 9, 11, 12*), and ancient clade (*WOX10, 13, 14*) (van der Graaff et al., 2009). This classification is largely consistent in other species as well (Zhang et al., 2010; Hao et al., 2019; Wu, 2020). The *WUS* clade members are characterized by an intact *WUS* box motif (Haecker et al., 2004; Lin et al., 2013), which is a transcriptional repression motif (Ikeda et al., 2009; Lin et al., 2013). Members of this group function primarily as transcriptional repressors, able to complement the *lam1* mutant phenotype (Lin et al., 2013), and are capable of substituting for *WUS* function in maintaining vegetative and floral meristems (Dolzblasz et al., 2016). The *WOX1* homologues *M. truncatula* *STF* and *N. sylvestris* *LAM1* belong to this clade and function as master regulators of leaf blade outgrowth through a transcriptional repression mechanism in association with the co-repressor *TOPLESS* (Tadege et al., 2011b; Lin, 2013; Lin et al., 2013; Zhang et al., 2014; Zhang et al., 2019). The intermediate and ancient clade members have partial or no *WUS* box, and do not have transcriptional repression activity in dual luciferase assays. As a result, they are unable to rescue the *lam1* mutant (Lin et al., 2013) nor substitute for *WUS* function (Dolzblasz et al., 2016). Among the intermediate and ancient clades, *AtWOX9* is unique in that it displays the strongest activation activity in dual luciferase assays, and strongly enhances the *lam1* mutant phenotype, affecting blade outgrowth in both medial-

lateral and proximal-distal axes (Lin et al., 2013), indicating that transcriptional activation activity modulated by AtWOX9 is antagonistic to LAM1 function. This is consistent with the observation that activation activity at the *STF* expression domain antagonizes *STF* function in blade outgrowth (Zhang et al., 2014; Zhang et al., 2019).

The results presented here demonstrated that *WOX9* transcript is upregulated in three leaf blade mutants; *stf* in *M. truncatula*, *lam1* in *N. sylvestris* and *wox1 prs* in Arabidopsis (Figures 6J to 6L), indicating that *WOX9* transcription may be suppressed by STF/LAM1/WOX1 in these species to allow blade outgrowth. Several lines of evidence including a GR inducible system in the presence of DEX and CHX, dual luciferase assay, EMSA, and ChIP confirmed that STF/LAM1 directly binds to the *MtWOX9-1* promoter to repress *WOX9* transcription (Figures 7 and 8), demonstrating that modern clade WOX1/STF/LAM1-mediated repression of intermediate clade *WOX9* is required for proper leaf blade outgrowth in eudicots. In Arabidopsis, *WOX9* functions upstream of *WUS* and is supposed to activate *WUS* to promote vegetative meristem growth (Wu et al., 2005), although it is unclear whether this activation is direct or indirect. *WOX9* is activated by cytokinin signaling (Skylar et al., 2010), and type-B ARRs directly activate *WUS* (Meng et al., 2017; Wang et al., 2017; Zhang et al., 2017; Zubo et al., 2017; Xie, 2018), while *WUS* promotes cytokinin activity by repressing type-A ARRs (Leibfried et al., 2005). Thus, cytokinin signaling provides a potential connection between *WOX9* and *WUS* in Arabidopsis, but whether *WUS* can directly affect *WOX9* activity via negative or positive feedback loop is yet to be determined. Our work clearly demonstrates that in Medicago and woodland tobacco, the *WUS* clade member STF/LAM1 directly represses *MtWOX9-1* or *NsWOX9*, but significant *STF/LAM1* activation by *WOX9* was not detected (Figure 7C). Although *MtWOX9-1* may not activate *STF*, it is likely to activate other targets in leaf development. *WOX9* amino acid sequences from different species show a highly conserved acidic domain at the C-terminus (Supplemental Figure 1), which could mediate transcriptional activation. Thus, our current model is that STF represses the transcription of key targets at the adaxial-abaxial junction to promote cell proliferation, and these targets include *MtWOX9-1* and cell differentiation factors such as *AS2* (Zhang et al., 2014). *WOX9*, on the other hand, may negatively regulate blade outgrowth by directly activating targets independent of STF and/or by activating targets

Figure 9. Schematic representation of hypothetical model for the regulation of leaf blade outgrowth by the interaction of STF/LAM1 and WOX9

STF/LAM1 directly represses *WOX9*, *AS2*, and other unidentified factors to promote leaf blade outgrowth. *WOX9*, on the other hand, negatively regulates leaf blade outgrowth by activating negative regulators of leaf growth and/or directly repressing blade outgrowth processes. The model proposes that STF/LAM1 and *WOX9* may have a common target(s) repressed by STF/LAM1 and activated by *WOX9* to balance cell proliferation with differentiation during leaf morphogenesis.

repressed by STF (Figure 9). The hypothesis that STF and *WOX9* may oppositely regulate common targets providing a critical balance between cell proliferation and differentiation during leaf morphogenesis, explains why *WOX9* ectopic expression enhances the *stf/lam1* mutant phenotype. Since both STF and WUS promote cell proliferation via a transcriptional repression mechanism in the leaf primordium and SAM, respectively, our findings suggest that direct control of *WOX9* activity by WUS may be required for SAM maintenance as well, uncovering a mechanistic framework for *WOX* modulated control of robust plant growth and developmental programs. It would be interesting to investigate if repression of the intermediate and ancient clade *WOX* transcriptional activators by modern clade *WOX* transcriptional repressors is a universal strategy exploited during the evolution of land plants and resulting in the complex morphological architecture of higher plants.

Methods

Plant materials and growth conditions

Plant materials used for this study including *Medicago truncatula* R108, *stf* mutant, and *Nicotiana sylvestris* (woodland tobacco) wild type and *lam1* mutant were grown in one gallon pots in a greenhouse under long-day (LD) conditions with 16/8 hours light/dark cycle at 23-27°C, and in growth room under long-day conditions of 16/8 hours light /dark cycle at 23-25°C, 70-80% relative humidity, and a light intensity of 150 $\mu\text{mol.m}^2$.

Samples collection, RNA extraction

M. truncatula tissue samples (young leaf and shoot) were collected from 4 week old wild type and *stf* mutant plants. For GR induction, four weeks old plants of *p35S::YFP-GR-STF*, *p35S::YFP-GR-LAM1* and *p35S::YFP-GR-NsWOX9* transgenic lines were treated with mock, DEX (10 mM), and CHX (10 mM) for 3hrs in *M. truncatula* and 6 hrs in *N. sylvestris*, before leaf samples were collected. All leaf samples from wild type, mutants and transgenic lines were collected at appropriate times for analyses of gene expression patterns as indicated in figure legends. Collected samples were immediately snap-frozen in liquid nitrogen and stored at -80°C until processing.

Total RNA from shoot apex and young leaf of *M. truncatula* R108 (WT) and *stf* mutant, *N. sylvestris* (WT) and *lam1* mutant, Arabidopsis *Ler* (WT), *wox1/prs* mutant and *p35S::YFP-GR-STF*, *p35S::YFP-GR-LAM1* and *p35S::YFP-GR-NsWOX9* transgenic lines were isolated using TRIzol Reagent (Invitrogen) for cDNA synthesis.

Real time PCR

Expression patterns analyses were performed using quantitative real time PCR (RT-qPCR) and semi-quantitative PCR with specific forward and reverse primers (Supplemental table 1). Reverse transcription (RT) was performed using RNA treated with DNase I (Invitrogen), an oligo (dT) primer, and SuperScript III reverse transcriptase (Invitrogen) according to the manufacturer's instruction. Quantitative RT-PCR assays were performed with three biological repeats and three technical replications of each experiment using SYBR Green real-time PCR Master Mix (Invitrogen). *M. truncatula*, Arabidopsis and *N. sylvestris* actin primers were used as expression standards. All specific forward and reverse primers used for gene cloning and expression and related molecular analyses in this study are listed in Supplemental Table 1.

Gene isolation and transgene construction

MtWOX9-1 and *MtWOX9-2* genes were isolated from the *M. truncatula* genome by BLAST search using the *AtWOX9* sequence. Full-length *MtWOX9-1* & 2 coding sequences were amplified by RT-PCR using total RNA extracted from leaf samples. To generate transgenic plants of *M. truncatula*, the cDNA was sub-cloned into the pDONR207 entry vector (Invitrogen) by BP clonase reaction. The final constructs were produced by an LR clonase reaction between each of the entry vectors and pMDC32 destination vector. The resulting plasmids were transferred into *Agrobacterium tumefaciens* strain *AGL1* and used to transform into *M. truncatula* R108 and *stf* mutant via agro-mediated-transformation using leaf explants as described (Tadege et al., 2011b). *N. sylvestris* WT and *lam1* mutant plants were transformed using *Agrobacterium tumefaciens*, GV2260 strain by leaf disc inoculation method as previously described (Tadege et al., 2011b). Antisense of *N. sylvestris* (*NsWOX9-anti*) construct was made using the Gateway system (Invitrogen). Briefly, *NsWOX9-anti-F* and *NsWOX9-anti-R* primers were designed by reverse attachment attb2 and attb1 to the primers for amplification of the full-length CDS sequence from start and stop codon of the gene. Using such primers, *NsWOX9* was amplified from *NsWOX9-cDNA* and cloned into pDONR207 entry vector (Invitrogen) by BP clonase reaction. Then, the construct was transferred into destination vector pMDC32 by an LR clonase reaction. All generated transgenic lines were confirmed by PCR using specific primers, and reduced expression lines were identified by RT-PCR.

Multiplex gRNA-CRISPR/Cas9 construction

To generate the multiplex gRNA-CRISPR/Cas9 *NsWOX9* construct, we designed three multiplex gRNAs targeting multiple sites of exon 2 (gRNA1-CTTCAAGAATATGGCCA AGT; gRNA2-CTTCAAGAATATGGCCAAGT and gRNA3-TCTCCTGCTGTTATCA CACA) at the upstream of PAM (TGG, TGG, and AGG) sites, respectively, using the web-based tool CRISPR-P (<http://cbi.hzau.edu.cn/cgi-bin/CRISPR>) (Lei, 2014). All designed gRNAs were inserted between tRNA and gRNA scaffolds and clustered in tandem using the Golden Gate assembly method (Engler, 2008). The pGTR plasmid, which contains a tRNA-gRNA fragment, was used as a template to synthesize polycistronic tRNA-gRNA (PTG) (Xie, 2015). The overlapping PCR products

were separated and purified by the Spin Column PCR Product Purification Kit (Wizard SV Gel and PCR Clean-Up System) following manufacturer's instruction (Promega, WI). Then, the chain of multiplex tRNA-gRNA with three *NsWOX9*-spacers was inserted into an optimized vector with AtU6-tRNA-gRNAs-AtUbi10-Cas9-pRGEB31-bar backbone by digestion and ligation using *Fok I* (NEB) and *BsaI* enzymes (Xie, 2015).

CRISPR/Cas plant transformation and analysis

The subsequent multiplex *NsWOX9*spacer-gRNA-CRISPR/Cas9 binary vector construct was transformed into *Agrobacterium* strain GV2260. First, transgenic lines were screened by PCR using genomic DNA and specific primers (PPT-F + PPT-R) of Barsta selection marker. Then, putative *nswox9-CRISPR* mutants were identified through amplification of the target region by PCR using extracted genomic DNA as a template with specific primers designed from the border of the target site. Amplified fragments of the mutated region were sub-cloned into pGEM-T easy plasmid by TA-Cloning and 10 colonies for each locus were subjected to Sanger sequencing. Reads were analyzed by aligning with the reference sequence using the SeqMan Pro 15.0.1 (DNASTAR software for life scientists) (<https://www.dnastar.com/quote-request/>).

Electrophoretic mobility shift assay

The *MtWOX9-I* promoter with 37, 30 and 32 bp oligonucleotides, corresponding to starting sites at -22, -226 and -491, respectively, upstream of the start codon were labeled using the Biotin 3' End DNA Labeling Kit according to the manufacturer's instructions (Thermo Scientific/Pierce). EMSA was performed with the Light Shift Chemiluminescent EMSA Kit (Thermo Scientific/Pierce). Unlabeled probes with a 50-fold higher concentration were used as competitors in each of the competing assays. Purified maltose binding protein (MBP) and MBP-STF were used in the EMSA as described (Zhang et al., 2014). Gel electrophoresis was performed on 10% native polyacrylamide gels. After blotting on a positively charged nylon membrane (Amersham), the DNA was crosslinked using a transilluminator equipped with 312 nm bulbs with the membrane face down for 15 min. The biotin-labeled DNA was detected by Chemiluminescence and exposed to X-ray film (Kodak). The probes and primers used in EMSA assay are listed in Supplemental Table 2.

ChIP Assays

ChIP assays were performed as described previously (Xiong et al., 2013; Chen et al., 2018). Protoplast extracted from 14-d-old *pMtWOX9-1:MtWOX9-1/wox1pr*s transgenic Arabidopsis leaves were transformed with 10 µg of 35S::STF-YFP using the polyethylene glycol-mediated transformation method. Protoplasts were cross-linked by 1% formaldehyde in W5 medium for 20 min and quenched with Glycine (0.2 M) for 5min. The protoplasts were then lysed, and the DNA was sheared on ice with sonication. The sheared chromatin was precleared by salmon sperm-sheared DNA/protein A agarose beads. Precleared chromatins were incubated with 5 µl of anti-GFP antibody (ab290) overnight at 4°C, after which Protein A agarose beads (40 µl) were added, and the samples were incubated at 4°C for 2 hrs. After reversing the crosslinks with Proteinase K at 65°C overnight, DNA was purified and analyzed by q-PCR amplification using specific primers. The input DNA and HA antibody-precipitated DNA were used as PCR templates for the positive and negative controls, respectively. Experiments were repeated three times. The primers used for the ChIP assays are listed in Supplemental Table 1

Dual luciferase assay

For effector plasmids, the coding sequence of STF or GUS was cloned into pDONR207 entry vector and then transferred into p2GW7 using the Gateway system (Invitrogen). Construction of the reporter *pMtWOX9-1-mini-35S-LUC* plasmid containing 1kb of the *MtWOX9-1* promoter was performed as previously described (Zhang et al., 2014). Transient expression assays were performed in Arabidopsis protoplasts as described (Asai et al., 2002). For normalization, 0.5 µg of plasmid pRLC was used as an internal control.

Histological analysis

Leaf samples were fixed in formaldehyde for 48 hrs and dehydrated in an ethanol series (60, 70, 85 and 95%). Then, leaves were embedded in Paraplast (Sigma-Aldrich, St. Louis, MO) and tissue sections (15 µm thick) were cut with a Reichert- Jung 2050 microtome. Specimens were mounted on slides and stained with Safranin O and Light Green as previously described (Tadege et al., 2011b). Images were captured with digital camera mounted on an Olympus BX-51 compound microscope.

Sequence alignment and phylogenetic tree construction

Multiple protein sequence alignment was performed using BioEdit software and the ClustalW program (<http://www.mbio.ncsu.edu/bioedit/bioedit.html>). Species refer to At (*Arabidopsis thaliana*), Ns (*Nicotiana sylvestris*), Pc (*Phaseolus coccineus*), Gm (*Glycine max*), Vv (*Vitis vinifera*), Ph (*Petunia x hybrid*), Cs (*Cucumis sativus*), Sl (*Solanum lycopersicum*) and Ca (*Capsicum annuum*). A neighbor-joining phylogenetic tree was constructed using MEGA-X default settings with 1000 bootstrap replications (<http://www.megasoftware.net/>).

Statistical Analysis

For statistical analysis, Student's *t* test was used as specified in figure legends. Asterisks indicate statistical differences (**P* < 0.05, ***P* < 0.01, ****p* < 0.001).

Accession Numbers

Sequence data from this article can be found in the NCBI (<http://www.ncbi.nlm.nih.gov/>) databases, *M. truncatula* Genome Database (<http://www.medicagogenome.org/>), or Phytozome (<https://phytozome.jgi.doe.gov/pz/portal.html>) under the following accession numbers: MtWOX9-1, Medtr2g015000; MtWOX9-2, Mt7go26130; AtWOX9, AT2G33880; AtWOX8, AT5G45980; NsWOX9, XM_009794999; Ns-LAM1, AEL30893; MtSTF, JF276252; AtWOX1, AT3G18010; AtWOX3, AT2G28610; PcWOX9, ACL11801; GmWOX9, XP_006594207; GmWOX9, XP_003541514; VvWOX9, XP_002273188; CaWOX9, XP_016562050; CsWOX9, XP_004134676; PhWOX9 (evergreen(EG)), ABO93066; PhWOX9 (SOEG); ABO93067; SlWOX9 (COI), NP_001234072; SlWOX9 (COI) isoform, XP_010315848.

Supplemental Data

Supplemental Figure 1. Amino acid sequence alignment of MtWOX9-1/2 and other related eudicot WOX9 sequences.

Supplemental Figure 2. Phylogenetic analysis of MtWOX9-1/2 and other related eudicots WOX9 protein sequences.

Supplemental Figure 3. Ectopic expression of *MtWOX9-1* enhances the *lam1* mutant phenotypes.

Supplemental Figure 4. Leaf phenotypes of *NsWOX9* overexpression in *N. sylvestris*.
Supplemental Figure 5. Phenotype of *MtWOX9-1* overexpression in *M. truncatula*.
Supplemental Figure 6. RT-qPCR analysis of *MtWOX9-2* expression in different tissues of *M. truncatula*.
Supplemental Table 1. List of primers and gRNAs used in this study

Acknowledgements

This work was supported by the National Science Foundation (NSF) grant IOS-1354422, Agriculture and Food Research Initiative Grant No. 2015-67014-22888 from the USDA National Institute of Food and Agriculture, and the Agricultural Experiment Station. Work in VT's laboratory at St Petersburg State University, Russia, was supported by the Russian Science Foundation project no. 16-16-10011 and by grant from the Russian Foundation for Basic Research no. 20-016-00124.

Author Contributions

T.W.W., H.W., and M.T. designed the research. T.W.W., H.W., D.T., F.Z., M.B., H.A., Y.L., and N.C. performed the experiments. T.W.W., H.W., F.Z., J.C., R.A., and M.T. analyzed the data. T.W.W., R.A., and M.T. wrote the manuscript.

Figure legends

Figure. 1. Ectopic expression of *WOX9* enhances *stf* and *lam1* mutant phenotypes.

- (A) Untransformed *M. truncatula* wild type (WT) (R108) plant.
- (B) Phenotype of untransformed *stf* mutant.
- (C) *stf* mutant transformed with *STF::MtWOX9-1*.
- (D) Untransformed *N. sylvestris* WT plant.
- (E) Phenotype of untransformed *lam1* mutant.
- (F) *lam1* mutant transformed with *STF::MtWOX9-1*.
- (G) *lam1* mutant transformed with *35S::MtWOX9-1*.
- (H) *lam1* mutant transformed with *35S::MtWOX9-2*.

(I) *lam1* mutant transformed with 35S::*NsWOX9*. Plants were 10-weeks (E and F) or 5-weeks old (all the rest). Scale bars: 10 cm.

Figure. 2. *NsWOX9-antisense* partially rescued *lam1* mutant phenotype.

(A) Phenotype of *lam1* mutant transformed with 35S::*GUS* as control at three weeks of age. Inset on the right is detached leaf close up.

(B) Partially complemented *lam1* phenotype transformed with 35S::*NsWOX9-antisense* construct at three weeks, the inset is close up of a partially complemented leaf blade. Inset on the right is detached leaf close up.

(C) Partially complemented *lam1* phenotype transformed with 35S::*NsWOX9-antisense* construct at seven weeks.

(D) Representative individual leaves from 35S::*NsWOX9-antisense/lam1* plants.

(E) A magnified view of a leaf in **(D)**. Note the branching and curling of leaves especially in older 35S::*NsWOX9-antisense/lam1* plants. Scale bars: A-D, 5 cm, E, 1.5 cm.

Figure. 3. Transverse section of the leaf blade showing enhancement of the *lam1* blade by 35S::*NsWOX9* and partial complementation by 35S::*NsWOX9-antisense*.

(A) *N. sylvestris* WT leaf blade.

(B) Leaf blade of untransformed *lam1* mutant control.

(C) Leaf blade of *lam1* transformed with 35S::*NsWOX9*.

(D) Partially complemented leaf blade of *lam1* transformed with 35S::*NsWOX9-antisense*.

(E) Transverse section of *N. sylvestris* WT leaf blade.

(F) Transverse section of untransformed *lam1* mutant leaf blade.

(G) Transverse sections of *lam1* leaf blade transformed with 35S::*NsWOX9* showing radialized blade.

(H) Transverse sections of *lam1* leaf blade transformed with 35S::*NsWOX9-antisense* showing blade outgrowth. Arrows indicate blade tissue in **(E)** and **(H)**, vestigial blade stripes in **(F)** and position of blade in **(G)**. Scale bars: 50 μ m.

Figure. 4. *WOX9* ectopic expression in WT *N. sylvestris* alters leaf architecture.

(A) WT *N. sylvestris* control plant at 7 weeks.

(B) Phenotype of *35S::MtWOX9-1/WT* (severe phenotype) at 7 weeks after regeneration.

(C) Phenotype of *35S::MtWOX9-1/WT* (mild-phenotype) at 7 weeks after regeneration.

(D) Phenotypes of different individual representative leaves from different

35S::MtWOX9-1/WT plants showing severe phenotypes at variable stages.

(E) Phenotype of *35S::MtWOX9-2/WT* (severe phenotype) at early growth stage.

(F) Phenotype of *35S::MtWOX9-2/WT* (mild) at early growth stage.

(G) Phenotype of *35S::NsWOX9/WT* (severe) at early growth stage. Scale bars: A-C and

E-G, 10 cm, D, 3 cm.

Figure. 5. Knock out of *NsWOX9* with multiplex *gRNA-CRISPR/Cas9* in *N. sylvestris* alters leaf architecture.

(A) Schematic representation of the three guide RNAs *NsWOX9*-gRNA1, 2 and 3 inserted in pRGEB31-bar-AtUbi10-AtU6-tRNA-gRNA vector.

(B) Mutation events detected at the corresponding target sites of gRNA1, gRNA2 and gRNA3) in five independent CRISPR/Cas lines (*NsWOX9-1*, 2, 13, 18 & 22).

(C-I) Phenotype of CRISPR/Cas9 edited plants and leaves; WT control (C), edited *NsWOX9-13* mutant (D), edited *NsWOX9-22* mutant (E), edited *NsWOX9-18* mutant (F), edited *NsWOX9-2* mutant (G), control WT leaf blade (H), representative individual leaf blades from edited plants; left, half blade deleted (*NsWOX9-22*), and right, narrow and asymmetric blade (*NsWOX9-2*). Red arrows point to blade defects, white arrows show multiple shoots. Scale bars: A-G, 5 cm, H and I, 2.5 cm.

Figure. 6. *MtWOX9-1* is weakly expressed in leaf but strongly upregulated in the *stf* mutant leaf of *M. truncatula* and corresponding mutants in *N. sylvestris* and *Arabidopsis*.

(A) RT-qPCR analysis showing relative expression of *MtWOX9-1* in different tissues *M. truncatula*. leaf, stem, and shoot apex were from 4-week old plants, pods and seeds were 10 days after pollination, flower at anthesis, all the rest were at seedling stage.

(B)-(I) GUS staining in *MtWOX9-1::GUS* transformed lines of *M. truncatula* showing fully expanded leaf **(B)**, shoot apex with folded leaves **(C)**, stem **(D)**, flower with unstained anthers **(E)**, very young pod with seeds **(F)**, older pod with seeds **(G)**, immature seeds **(H)**, and matured seeds **(I)**.

(J) Relative expression of *MtWOX9-1* in 4-week old *stf* mutant leaf in *M. truncatula*.

(K) Relative expression of *NsWOX9* in 4-week old *lam1* mutant leaf of *N. sylvestris*.

(L) Relative expression of *AtWOX9* in the leaf of 4-week old Arabidopsis *wox1 prs* double mutant. Error bars indicate $\pm$ SE (n=3). Asterisks indicate significant difference from the control (*p<0.05, **p<0.01, student t-test).

Figure 7. *WOX9* expression is directly repressed by GR induction of STF or LAM1 in the presence of cyclohexamide, and by STF in dual luciferase assay.

(A) Relative expression of *MtWOX9-1* in *35S::YFP-GR-STF* transformed *M. truncatula* lines with CHX or DEX+CHX treatment for 3 hours in 4-week old leaves.

(B) Relative expression of *NsWOX9* in 2 independently transformed *35S::YFP-GR-LAM1* *N. sylvestris* lines with CHX alone, CHX+DEX or DEX alone treatments for 6 hours in 4-weeks old leaves.

(C) Relative expression of *LAM1* in *35S-YFP-GR-NsWOX9* transformed 2 independent *N. sylvestris* lines with CHX alone, CHX+DEX or DEX alone treatments for 6 hours in 4-weeks old young leaves. Relative gene expression was determined by RT-qPCR analyses.

(D) Schematic representation of reporter and effector constructs used in the dual luciferase assay.

(E) Relative expression of luciferase activity (luminescence) in the presence of *35S::STF* effector compared with the *35S::GUS* control in Arabidopsis protoplasts. Error bars indicate $\pm$ SE (n=3). Asterisks indicate significant difference from the control (*p<0.05, **p<0.01, ***p<0.001, student t-test).

Figure 8. STF directly binds to the *MtWOX9-1* promoter in EMSA and ChIP assays.

(A) Schematic representation of the *MtWOX9-1* promoter and CDS regions tested for EMSA and ChIP assays. The 3 promoter regions tested are indicated as P1, P2 and P3 and the *MtWOX9-1* coding region as C1.

(B) EMSA showing MBP-STF bound to the biotin-labeled probe at P1, P2 and P3 promoter fragments but not the MBP control alone. Fifty-fold excess of unlabeled P1, P2 or P3 DNA was used to compete with the respective labeled probe (right lanes).

(C) *MtWOX9-1* promoter enrichment at P1, P2, P3 regions and CDS C1. Chromatin precipitated with anti-GFP antibody from *35S::STF-GFP* and *35S::GFP* control lines were compared. Purified DNA from the chromatin were used as templates for qPCR. Note that P2 is highly enriched in *35S::STF-GFP* samples. Error bars indicate $\pm$ SE (n=3). Asterisks indicate statistical significance (*p<0.05, ***p<0.001, student t-test).

Figure 9. Schematic representation of hypothetical model for the regulation of leaf blade outgrowth by the interaction of STF/LAM1 and WOX9

STF/LAM1 directly represses *WOX9*, *AS2*, and other unidentified factors to promote leaf blade outgrowth. *WOX9*, on the other hand, negatively regulates leaf blade outgrowth by activating negative regulators of leaf growth and/or directly repressing blade outgrowth processes. The model proposes that STF/LAM1 and *WOX9* may have a common target(s) repressed by STF/LAM1 and activated by *WOX9* to balance cell proliferation with differentiation during leaf morphogenesis.

Parsed Citations

- Asai, T., Boller, T., Gomez-Gomez, L., Sheen, J., Ausubel, F.M., Plotnikova, J., Tena, G., Chiu, W.L., and Willmann, M.R. (2002). MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature (London, England) Nature* 415, 977-983.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Breuninger, H., Rikirsch, E., Hermann, M., Ueda, M., and Laux, T. (2008). Differential expression of WOX genes mediates apical-basal axis formation in the Arabidopsis embryo. *Developmental cell* 14, 867-876.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Causier, B., Ashworth, M., Guo, W., and Davies, B. (2012). The TOPLESS interactome: a framework for gene repression in Arabidopsis. *Plant Physiology* 158, 423-438.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Chen, N., Veerappan, V., Haggag, A., Miyong, K., and Allen, R.D.r.a.o.e. (2018). HSI2/VAL1 Silences AGL15 to Regulate the Developmental Transition from Seed Maturation to Vegetative Growth in Arabidopsis. *Plant Cell* 30, 600-619.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Cho, S.-H., Yoo, S.-C., Zhang, H., Pandeya, D., Koh, H.-J., Hwang, J.-Y., Kim, G.-T., and Paek, N.-C. (2013). The rice narrow leaf2 and narrow leaf3 loci encode WUSCHEL-related homeobox 3A (OsWOX3A) and function in leaf, spikelet, tiller and lateral root development. *The New phytologist* 198, 1071-1084.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Costanzo E, T.C., and Vandenbussche M. (2014). The role of WOX genes in flower development. *Annals of Botany* 114, 1545-1553.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Daum, G., Medzihradsky, A., Suzuki, T., and Lohmann, J.U. (2014). A mechanistic framework for noncell autonomous stem cell induction in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* 111, 14619-14624.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Deyhle, F., Sarkar, A.K., and Tucker, E.J. (2007). WUSCHEL regulates cell differentiation during anther development. *Developmental Biology* 302, 154-159.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Dolzblasz, A, Barry, C., Brendan, D., Elena, C., Eric van der, G., Jinhui, C., Judith, N., Thomas, L., and Wolfgang, W. (2016). Stem Cell Regulation by Arabidopsis WOX Genes. *Molecular plant* 9, 1028-1039.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Engler, C., Kandzia, R., & Marillonnet, S. . (2008). A one pot, one step, precision cloning method with high throughput capability. . *PLoS ONE* 3, e3647.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Fang F, Y.S., Tang J, Bennett MJ, Liang W. (2020). DWT1/DWL2 act together with OsPIP5K1 to regulate plant uniform growth in rice. *New Phytol.* 225, 1234-1246.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Gambino, G., Minuto, M., and Boccacci, P. (2011). Characterization of expression dynamics of WOX homeodomain transcription factors during somatic embryogenesis in Vitis vinifera. *Journal of Experimental Botany* 62, 1089-1101.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Haecker, A, Gross-Hardt, R., Geiges, B., Sarkar, A., Breuninger, H., Herrmann, M., and Laux, T. (2004). Expression dynamics of WOX genes mark cell fate decisions during early embryonic patterning in Arabidopsis thaliana. *Development* 131, 657-668.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Hao, Q., Shan, Z., Yang, Y., Zhang, L., and Zhou, X.-a. (2019). Genome-Wide Analysis of the WOX Gene Family and Function Exploration of GmWOX18 in Soybean. *Plants* 8.**
Pubmed: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)
- Hirakawa, Y., Kondo, Y., and Fukuda, H. (2010). TDIF peptide signaling regulates vascular stem cell proliferation via the WOX4 homeobox gene in Arabidopsis. *The Plant cell* 22, 2618-2629.**

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Hu C, ZY., Cui Y, Cheng K, Liang W, Wei Z, Zhu M, Yin H, Zeng L, Xiao Y, Lv M, Yi J, Hou S, He K, Li J, Gou X. (2018). A group of receptor kinases are essential for CLAVATA signalling to maintain stem cell homeostasis. Nature Plants 4, 205-211.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ikedo, M., Mitsuda, N., and Ohme-Takagi, M. (2009). Arabidopsis WUSCHEL is a bifunctional transcription factor that acts as a repressor in stem cell regulation and as an activator in floral patterning. Plant Cell 21, 3493-3505.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ishiwata, A., Akie, N., Hiroshi, N., Hiro-Yuki, H., Makio, K., Masahiko, M., Misa, O., Misuzu, N., Sae, S.-S., Takahiro, Y., Yusaku, N., and Yutaka, S. (2013). Two WUSCHEL-related homeobox Genes, narrow leaf2 and narrow leaf3, Control Leaf Width in Rice. Plant & cell physiology 54, 779-792.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Iwakawa, H., Ueno, Y., Semiarti, E., Onouchi, H., Kojima, S., Tsukaya, H., Hasebe, M., Soma, T., Ikezaki, M., Machida, C., and Machida, Y. (2002). The ASYMMETRIC LEAVES2 gene of Arabidopsis thaliana, required for formation of a symmetric flat leaf lamina, encodes a member of a novel family of proteins characterized by cysteine repeats and a leucine zipper. Plant & cell physiology 43, 467-478.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ji, J., Strable, J., Shimizu, R., Koenig, D., Sinha, N., and Scanlon, M.J. (2010). WOX4 promotes procambial development. Plant Physiology 152, 1346-1356.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kerstetter, R.A., Bollman, K., Taylor, R.A., Bombles, K., and Poethig, R.S. (2001). KANADI regulates organ polarity in Arabidopsis. Nature 411, 706-709.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kieffer, M., Stern, Y., Cook, H., Clerici, E., Maulbetsch, C., Laux, T., and Davies, B. (2006). Analysis of the transcription factor WUSCHEL and its functional homologue in Antirrhinum reveals a potential mechanism for their roles in meristem maintenance. Plant Cell 18, 560-573.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Laux, T., Mayer, K.F., Berger, J., and Jürgens, G. (1996). The WUSCHEL gene is required for shoot and floral meristem integrity in Arabidopsis. Development 122, 87-96.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lei, Y., Lu, L., Liu, H.Y., Li, S., Xing, F., & Chen, L.L. . (2014). CRISPR-P: a web tool for synthetic single-guide RNA design of CRISPR-system in plants. Molecular Plant 7, 1494-1496.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Leibfried, A., To, J.P., Busch, W., Stehling, S., Kehle, A., Demar, M., Kieber, J.J., and Lohmann, J.U. (2005). WUSCHEL controls meristem function by direct regulation of cytokinin-inducible response regulators. Nature 438, 1172-1175.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lin, H., Niu, L., McHale, N.A., Ohme-Takagi, M., Mysore, K.S., and Tadege, M. (2013). Evolutionarily conserved repressive activity of WOX proteins mediates leaf blade outgrowth and floral organ development in plants. Proceedings of the National Academy of Sciences of the United States of America 110, 366-371.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lin, H., Niu, L., Tadege, M. (2013). STENOFOLIA acts as a repressor in regulating leaf blade outgrowth. Plant Signal Behav 8, pii: e24464.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lippman, Z.B., Cohen, O., and Alvarez, J.P. (2008). The Making of a Compound Inflorescence in Tomato and Related Nightshades. PLoS Biology 6, 2424-2435.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lohmann, J.U., Hong, R.L., Hobe, M., Busch, M.A., Parcy, F., Simon, R., and Weigel, D. (2001). A molecular link between stem cell regulation and floral patterning in Arabidopsis. Cell 105, 793-803.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Matsumoto, N., and Okada, K. (2001). A homeobox gene, PRESSED FLOWER, regulates lateral axis-dependent development of Arabidopsis flowers. Genes Dev. 15, 3355-3364.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Mayer, K.F., Schoof, H., Haecker, A., Lenhard, M., Jürgens, G., and Laux, T. (1998). Role of WUSCHEL in regulating stem cell fate in the Arabidopsis shoot meristem. Cell 95, 805-815.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

McConnell, J.R., Emery, J., Eshed, Y., Bao, N., Bowman, J., and Barton, M.K. (2001). Role of PHABULOSA and PHAVOLUTA in determining radial patterning in shoots. Nature 411, 709-713.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Meng, W., Cheng, Z., Sang, Y., Zhang, M., Rong, X., Wang, Z., Tang, Y., and Zhang, X. (2017). Type-B ARABIDOPSIS RESPONSE REGULATORS Specify the Shoot Stem Cell Niche by Dual Regulation of WUSCHEL. Plant Cell 29, 1357-1372.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Meng, Y., Liu, H., Wang, H., Liu, Y., Zhu, B., Wang, Z., Hou, Y., Zhang, P., Wen, J., Yang, H., Mysore, K.S., Chen, J., Tadege, M., Niu, L., and Lin, H. (2019). HEADLESS, a WUSCHEL homolog, uncovers novel aspects of shoot meristem regulation and leaf blade development in Medicago truncatula. Journal of Experimental Botany 70, 149-163.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Nakata, M., Matsumoto, N., Tsugeki, R., Rikirsch, E., Laux, T., and Okada, K. (2012). Roles of the Middle Domain-Specific WUSCHEL-RELATED HOMEBOX Genes in Early Development of Leaves in Arabidopsis. Plant Cell 24, 519-535.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Nakata, M., Okada, K. (2012). The three-domain model: a new model for the early development of leaves in Arabidopsis thaliana. Plant Signal Behav 7, 1423-1427.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Nardmann, J., Ji, J., Werr, W., and Scanlon, M.J. (2004). The maize duplicate genes narrow sheath1 and narrow sheath2 encode a conserved homeobox gene function in a lateral domain of shoot apical meristems. Development 131, 2827-2839.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Park, S.O., Zheng, Z., Oppenheimer, D.G., and Hauser, B.A. (2005). The PRETTY FEW SEEDS2 gene encodes an Arabidopsis homeodomain protein that regulates ovule development. Development (Cambridge) 132, 841-849.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Rebocho, A.B., Bliet, M., Kusters, E., Castel, R., Procissi, A., Roobeek, I., Souer, E., and Koes, R. (2008). Role of EVERGREEN in the development of the cymose petunia inflorescence. Developmental Cell 15.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Sarkar, A.K., Luijten, M., Miyashima, S., Lenhard, M., Hashimoto, T., Nakajima, K., Scheres, B., Heidstra, R., Laux, T. (2007). Conserved factors regulate signalling in Arabidopsis thaliana shoot and root stem cell organizers. Nature 446, 811-814.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Sawa, S., Watanabe, K., Goto, K., Liu, Y. G., Shibata, D., Kanaya, E., Morita, E. H., Okada, K. (1999). FILAMENTOUS FLOWER, a meristem and organ identity gene of Arabidopsis, encodes a protein with a zinc finger and HMG-related domains. Genes Dev 13, 1079-1088.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Schoof, H., Lenhard, M., Haecker, A., Mayer, K.F., Jürgens, G., and Laux, T. (2000). The stem cell population of Arabidopsis shoot meristems is maintained by a regulatory loop between the CLAVATA and WUSCHEL genes. Cell 100, 635-644.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Shimizu, R., Ji, J., Kelsey, E., Ohtsu, K., Schnable, P.S., and Scanlon, M.J. (2009). Tissue specificity and evolution of meristematic WOX3 function. Plant Physiol. 149, 841-850.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Siegfried, K.R., Eshed, Y., Baum, S. F., Otsuga, D., Drews, G. N., Bowman, J. L. (1999). Members of the YABBY gene family specify

abaxial cell fate in Arabidopsis. Development 126, 4117-4128.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Skylar, A., Hong, F.X., Chory, J., Weigel, D., and Wu, X.L. (2010). STIMPY mediates cytokinin signaling during shoot meristem establishment in Arabidopsis seedlings. Development 137, 541-549.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Snipes, S.A., Rodriguez, K., DeVries, A.E., Miyawaki, K.N., Perales, M., Xie, M., and Reddy, G.V. (2018). Cytokinin stabilizes WUSCHEL by acting on the protein domains required for nuclear enrichment and transcription. PLoS Genetics 14, 1-23.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Somssich, M., Byoung II, J., Simon, R.r.s.u.-d.d., and Jackson, D.j.c.e. (2016). CLAVATA-WUSCHEL signaling in the shoot meristem. Development (09501991) 143, 3238-3248.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Stuurman, J., Jaggi, F., and Kuhlemeier, C. (2002). Shoot meristem maintenance is controlled by a GRAS-gene mediated signal from differentiating cells. Genes Dev 16, 2213-2218.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tadege, M. (2016). WOX3 in the scene: intimacy with hormones. Journal of Experimental Botany 67, 1605-1607.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tadege, M., and Mysore, K.S. (2011). Tnt1 retrotransposon tagging of STF in Medicago truncatula reveals tight coordination of metabolic, hormonal and developmental signals during leaf morphogenesis. Mobile Genetic Elements (2159-2543) 1, 301-303.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tadege, M., Lin, H., Niu, L., and Mysore, K.S. (2011a). Control of dicot leaf blade expansion by a WOX gene, STF. Plant Signal Behav. 6, 1861-1864.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tadege, M., Chen, F., Murray, J., Wen, J., Ratet, P., Udvardi, M., Dixon, R., and Mysore, K. (2015). Control of Vegetative to Reproductive Phase Transition Improves Biomass Yield and Simultaneously Reduces Lignin Content in Medicago truncatula. BioEnergy Research 8, 857-867.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tadege, M., Lin, H., Bedair, M., Berbel, A., Wen, J., Rojas, C.M., Niu, L., Tang, Y., Sumner, L., Ratet, P., McHale, N.A., Madueño, F., and Mysore, K.S. (2011b). STENOFOLIA Regulates Blade Outgrowth and Leaf Vascular Patterning in Medicago truncatula and Nicotiana glauca. Plant Cell 23, 2125-2142.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Tvorogova, V.E., Fedorova, Y.E., Potsenkovskaya, E.A., Kudriashov, A.A., Efremova, E.P., Kvitkovskaya, V.A., Wolabu, T.W., Zhang, F., Tadege, M., & Lutova, L.A. (2019). The WUSCHEL-related homeobox transcription factor MtWOX9-1 stimulates somatic embryogenesis in Medicago truncatula. Plant Cell, Tissue and Organ Culture 3, 517-527.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ueda, M., Zhang, Z.J., and Laux, T. (2011). Transcriptional Activation of Arabidopsis Axis Patterning Genes WOX8/9 Links Zygote Polarity to Embryo Development. Developmental cell 20, 264-270.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

van der Graaff, E., Laux, T., and Rensing, S.A. (2009). The WUS homeobox-containing (WOX) protein family. Genome Biol. 10, 248.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Vandenbussche, M., Horstman, A., Zethof, J., Koes, R., Rijpkema, A.S., and Gerats, T. (2009). Differential recruitment of WOX transcription factors for lateral development and organ fusion in Petunia and Arabidopsis. Plant cell 21, 2269-2283.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Waites, R., Selvadurai, H.R.N., Oliver, I.R., and Hudson, A. (1998). The PHANTASTICA gene encodes a MYB transcription factor involved in growth and dorsoventrality of lateral organs in antirrhinum. Cell 93, 779-789.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wang, H., Niu, L., Fu, C., Meng, Y., Sang, D., Yin, P., Wu, J., Tang, Y., Lu, T., Wang, Z.Y., Tadege, M., Lin, H. (2017). Overexpression of the WOX gene STENOFOLIA improves biomass yield and sugar release in transgenic grasses and display altered cytokinin homeostasis. PLoS Genet. 13, e1006649.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wang H, X.Y., Hong L, Zhang X, Wang X, Zhang J, Ding Z, Meng Z, Wang ZY, Long R, Yang Q, Kong F, Han L, Zhou C. (2019). HEADLESS Regulates Auxin Response and Compound Leaf Morphogenesis in Medicago truncatula. Front Plant Sci. 10, 1024.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wang, J., Caihuan, T., Cui, Z., Bihai, S., Xiuwei, C., Tian-Qi, Z., Zhong, Z., Jia-Wei, W., and Yuling, J. (2017). Cytokinin Signaling Activates WUSCHEL Expression during Axillary Meristem Initiation. Plant Cell 29, 1373-1387.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wang, W., Li, G., Zhao, J., Chu, H., Lin, W., Zhang, D., Wang, Z., and Liang, W.w.s.e.c. (2014). DWARF TILLER1, a WUSCHEL-Related Homeobox Transcription Factor, Is Required for Tiller Growth in Rice. PLoS Genetics 10, 1-15.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wu, C.C.L., F.W. Kramer, E.M. (2020). Large-scale phylogenomic analysis suggests three ancient superclades of the WUSCHEL-RELATED HOMEBOX transcription factor family in plants. PLoS One 14, e0223521.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wu, X., Chory, J., and Weigel, D. (2007). Combinations of WOX activities regulate tissue proliferation during Arabidopsis embryonic development. Developmental biology 309, 306-316.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Wu, X.L., Dabi, T., and Weigel, D. (2005). Requirement of homeobox gene STIMPY/WOX9 for Arabidopsis meristem growth and maintenance. Current Biology 15, 436-440.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Xie, K., Minkenberg, B., & Yang, Y. . (2015). Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. Proceedings of the National Academy of Sciences, USA, 112 (11), 3570–3575. Proceedings of the National Academy of Sciences, USA, 112, 3570-3575.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Xie, M.C., H. Huang, L. O'Neil, RC. Shokhirev, MN. Ecker, JR. (2018). A B-ARR-mediated cytokinin transcriptional network directs hormone cross-regulation and shoot development. Nat Commun. 9, 1604.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Xiong, Y., McCormack, M., Li, L., Hall, Q., Xiang, C., and Sheen, J. (2013). Glucose-TOR signalling reprograms the transcriptome and activates meristems. Nature 496, 181-186.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Yadav, R.K., Perales, M., Gruel, J., Girke, T., Jonsson, H., and Reddy, G.V. (2011). WUSCHEL protein movement mediates stem cell homeostasis in the Arabidopsis shoot apex. Genes & development 25, 2025-2030.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Yadav, R.K., Perales, M., Gruel, J., Ohno, C., Heisler, M., Girke, T., Jonsson, H., and Reddy, G.V. (2013). Plant stem cell maintenance involves direct transcriptional repression of differentiation program. Molecular systems biology 9, 654.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhang, F., May, A., and Irish, V.F. (2017). Type-B ARABIDOPSIS RESPONSE REGULATORs Directly Activate WUSCHEL. Trends in Plant Science 22, 815-817.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhang, F., Wang, Y., Li, G., and Tang, Y.K., EM. Tadege, M. (2014). STENOFOLIA Recruits TOPLESS to Repress ASYMMETRIC LEAVES2 at the Leaf Margin and Promote Leaf Blade Outgrowth in Medicago truncatula. Plant Cell 26, 650-664.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhang, F., Wang, H., Kalve, S., Wolabu, T.W., Nakashima, J., Golz, J.F., and Tadege, M. (2019). Control of leaf blade outgrowth and floral organ development by LEUNIG, ANGUSTIFOLIA3 and WOX transcriptional regulators. New Phytologist 223, 2024-2038.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhang F, T.M. (2015). Repression of AS2 by WOX family transcription factors is required for leaf development in Medicago and Arabidopsis. Plant Signal Behav. 10, e993291.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhang, X., Yin, J., Zhang, D., Zong, J., and Liu, J. (2010). Genome-Wide Analysis of WOX Gene Family in Rice, Sorghum, Maize, Arabidopsis and Poplar [electronic resource]. Journal of integrative plant biology 52, 1016-1026.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zhuang, L.-L., Ambrose, M., Rameau, C., Weng, L., Yang, J., Hu, X.-H., Luo, D., and Li, X. (2012). LATHYROIDES, Encoding a WUSCHEL-Related Homeobox1 Transcription Factor, Controls Organ Lateral Growth, and Regulates Tendril and Dorsal Petal Identities in Garden Pea (Pisum sativum L.). Molecular plant 5, 1333-1345.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Zubo, Y.O., Yamburenko, M.V., Worthen, J.M., Street, I.H., Hill, K., Schaller, G.E., Blakley, I.C., Loraine, A.E., Franco-Zorrilla, J.M., Wenjing, Z., Raines, T., Kieber, J.J., and Solano, R. (2017). Cytokinin induces genome-wide binding of the type-B response regulator ARR10 to regulate growth and development in Arabidopsis. Proceedings of the National Academy of Sciences of the United States of America 114, E5995-E6004.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)