NAD⁺ prevents septic shock by non-canonical inflammasome blockade and IL-10

Jasper Iske^{1,2†}, Rachid El Fatimy^{3†}, Yeqi Nian^{1†}, Siawosh K. Eskandari⁴, Hector Rodriguez Cetina Biefer^{5,6},
Anju Vasudevan⁷ and Abdallah Elkhail^{1*}

¹Division of Transplant Surgery, Department of Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

²Institute of Transplant Immunology, Integrated Research and Treatment Center Transplantation (IFB-Tx), Hannover Medical School, Hannover, Lower Saxony, Germany

³Department of Neurology, Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

⁴Transplantation Research Center, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

⁵Department of Cardiovascular Medicine, Charité Universitätsmedizin Berlin, Berlin, Germany

⁶Deutsches Herzzentrum Berlin (DHZB), Berlin Germany

⁷Angiogenesis and Brain Development Laboratory, Department of Psychiatry, McLean Hospital, Harvard Medical School, Boston, Massachusetts, USA

[†]These authors contributed equally to this work.

Corresponding author:

Abdallah Elkhail, Ph.D.,
Division of Transplant Surgery & Transplant Surgery Research Laboratory,
Brigham and Women's Hospital,
Harvard Medical School,
75 Francis Street,
Boston, MA 02115
E-mail: aelkhail@bwh.harvard.edu

Abstract

Non-canonical inflammasome activation is crucial in the development of septic shock promoting pyroptosis and pro-inflammatory cytokine production via caspase-11 and Gasdermin-D (GSDMD). Here, we show that NAD⁺ treatment protected mice towards bacterial and LPS induced endotoxic shock by blocking the non-canonical inflammasome specifically. NAD⁺ administration impeded systemic IL-1 β and IL-18 production and GSDMD-mediated pyroptosis of macrophages via the IFN- β /STAT-1 signaling machinery. More importantly, NAD⁺ administration not only improved casp-11^{-/-} survival but rendered WT mice completely resistant to septic shock via the IL-10 signaling pathway that was independent from the non-canonical inflammasome. Here, we delineated a two-sided effect of NAD⁺ blocking septic shock through a specific inhibition of the non-canonical inflammasome and promoting immune homeostasis via IL-10, underscoring its unique therapeutic potential.

Summary

NAD⁺ protects against septic shock by blocking the non-canonical inflammasome specifically and via a systemic production of IL-10 cytokine

Introduction

Sepsis is characterized by a systemic inflammatory response syndrome (SIRS)¹ driven by host cells following systemic bacterial infections². The excessive inflammatory response can derail into septic shock resulting in multiple organ failure (MOF), the leading cause of death in intensive care units. Inflammasome activation, which downstream pathways cause the release of proinflammatory cytokines and the induction of an inflammatory cell death termed pyroptosis³, has been pointed out as the major driver of septic shock. Hereby, a two-armed LPS derived induction of the NLRP3-canonical inflammasome, the major source of IL-1 β and IL-18 cytokine production⁴ and the caspase-11 mediated non-canonical inflammasome leading to pyroptosis in monocytes⁵ was determined as the underlying mechanism. Mechanistically, Caspase-11 acts as a pattern recognition receptor for intracellular bacteria⁶ that cleaves gasdermin-D, a membrane-pore forming protein subsequently inducing pyroptotic cell death⁷. The NLRP3-canonical inflammasome in turn, was found to be indispensable⁸ for septic shock induced death. However cross-activation through Caspase-11 promoting cytokine release has been described⁹⁻¹¹, assigning the non-canonical inflammasome a cardinal role¹².

Recent approaches such as anti-proinflammatory cytokine strategies, blocking downstream targets of inflammasomes have been ineffective¹³ while inhibiting inflammatory key regulators such as NF- κ B may promote adverse side-effects¹⁴. Hence, contemporary clinical therapy of septic shock is based on symptomatic treatment rather than curative approaches that clear the cause of the disease itself.

In our previous studies, we have underscored the immunosuppressive properties of NAD⁺ in autoimmune diseases and allo-immunity via the regulation of CD4⁺ T cell fate^{15,16}. More recently, we have shown that NAD⁺ administration protected mice from lethal doses of *Listeria monocytogenes* (*L. m.*) via mast cells (MCs) exclusively and independently of major antigen presenting cells (APCs)¹⁷. However, the underlying mechanism that allows NAD⁺, to concomitantly protect against autoimmune diseases, via its immunosuppressive properties^{15,16}, and against lethal bacterial infection remains unclear.

Therefore, in the current study we investigated whether NAD⁺ protects against bacterial infection by dampening the systemic inflammatory response associated with sepsis or through enhanced bacterial clearance. Although, wild type (WT) mice subjected to NAD⁺ or PBS and lethal doses of pathogenic *Escherichia coli* (*E. coli*) exhibited similar bacterial load in various tissues, mice treated with NAD⁺ displayed a robust survival. Moreover, NAD⁺ protected against LPS-induced death that was associated with a dramatic decrease of systemic IL-1 β and IL-18 levels, two major cytokines involved in the inflammasome signaling machinery. More importantly, we show that NAD⁺ protected from LPS-induced death by targeting specifically the non-canonical inflammasome via a blockade of the STAT-1/IFN- β signaling pathway. Moreover, NAD⁺ treatment rendered not only Caspase-11^{-/-} but WT mice fully resistant to poly(I:C) + LPS induced septic shock, via an inflammasome independent pathway mediated by a systemic IL-10 cytokine production.

Results

NAD⁺ protects mice against septic shock not via bacterial clearance but via inflammasome blockade

Our previous studies have underscored the role of NAD⁺ in regulating CD4⁺ T cell fate and its immunosuppressive properties via IL-10 cytokine production¹⁵⁻¹⁷. More recently, we have shown that NAD⁺ protected mice against lethal doses of *L. m.* independently of major APCs¹⁵. However, it remained unclear whether NAD⁺ protected mice against lethal doses of *L. m.*, a gram-positive bacterium, via a clearance mechanism or by dampening the inflammatory response. Since *L. m.* is known to be an intracellular pathogen, we tested if NAD⁺ protects as well against *E. coli*, a gram-negative bacterium that is well known to induce septic shock¹⁸. Wild type mice were treated with NAD⁺ or PBS for 2 consecutive days followed by a lethal dose (1x10⁹) of *E. coli* or PBS. Notably, mice treated with PBS died within 5 hours after infection, while mice treated with NAD⁺ exhibited an impressive survival (**Fig. 1A**). Moreover, when assessing the bacterial load in liver and kidney (**Fig. 1B**), organs exposed to the infection, by counting CFU in both, NAD⁺ and PBS groups, revealed no significant difference, suggesting that NAD⁺ does not promote bacterial clearance. More importantly, these data suggest that NAD⁺ may reduce the inflammatory response towards bacterial infection. It is well established that the bacterial lipopolysaccharide (LPS) abundant on the outer membrane exhibits a key role in the pathology of *E. coli* derived septic shock¹³. Thus, we further characterize the impact of NAD⁺ on septic shock by subjecting mice to a lethal dose (54mg/kg) of two different LPS serotypes (O111:B4 and O55:B5) described to vary in the antigen lipid A content and to promote distinct hypothermia kinetics¹⁹. Following LPS (O111:B4 and O55:B5) administration, PBS treated control mice displayed severe symptoms of endotoxic shock with a dramatical body temperature decrease (<23°C) within 15 hours. In contrast, mice subjected to NAD⁺ exhibited highly distinct kinetics with a recovery of body temperatures after 15 hours (**Fig. 1C**). When monitoring survival, 100% of PBS treated mice succumbed to LPS after 24 hours while NAD⁺ treated animals exhibited an overall survival >85% (**Fig. 1D**), which was consistent with our bacterial infection model.

LPS-induced death derives from multi-organ failure²⁰. Therefore, lung, kidney, liver, ileum and spleen were harvested 15 hours after LPS administration and tissue damage was assessed by Hematoxylin and Eosin (H&E) staining. Tissue evaluation indicated severe pulmonary hemorrhage, excessive tubular fibrin deposition, hepatocyte cell swelling, disseminated intravascular coagulation (DIC) and ileal villi destruction consistent with a multi-organ-dysfunction syndrome (MODS)²¹ in mice treated with PBS. In contrast, NAD⁺ administration dramatically attenuated signs of organ failure with significantly less pulmonary hemorrhage and DIC, intact liver and kidney tissue architecture and preserved ileal villi (**Fig. 1E, Supp. 1**). To elucidate the protective effects of NAD⁺ systemic levels of IL-1 β and IL-18, two major cytokines implicated in inflammasome activation, were measured 10 and 15 hours after intraperitoneal injection of LPS (**Fig. 1F**). Our findings indicated that LPS injection resulted in a robust systemic increase of IL-1 β and IL-18 in the PBS group, which was almost abolished in NAD⁺ treated mice. Taken together, our results suggest that NAD⁺ protects mice against septic shock not via bacterial clearance but rather via inflammasome blockade.

NAD⁺ specifically inhibits the non-canonical inflammasome

Our data suggest that NAD⁺ protects against septic shock via inflammasome blockade. Monocytes, especially macrophages, have been described as major drivers of inflammasome derived cytokine secretion in the context of septic shock²². Thus, to test the effect of NAD⁺ on inflammasome function, bone marrow derived macrophages (BMDMs) were obtained and both, canonical and non-canonical inflammasomes were stimulated in the presence or absence of 100 μ mol/ml NAD⁺. Activation of the canonical pathway was achieved through LPS priming (1 μ g/ml) followed by ATP stimulation (5 mmol/l). Notably, BMDMs subjected to NAD⁺ or PBS treatment followed by canonical inflammasome activation did not exhibit any significant difference in IL-1 β secretion or pyroptosis that was assessed by LDH release measurement, a marker for cell death²³ (**Fig. 2A**).

To trigger the non-canonical inflammasome pathway, BMDMs were primed with Pam3CSK4, a TLR1/2 agonist, followed by cholera toxin B (CTB) and LPS (2 μ g/ml) administration. The data showed that

NAD⁺ treatment resulted in a robust reduction of IL-1 β release and cell death when compared to the PBS control group (**Fig. 2A**). Furthermore, Western blotting revealed that BMDMs cultured in presence of NAD⁺ exhibited a dramatic decrease of casp-11 expression and its downstream targets including casp-1, IL-1 β and cleaved gasdermin-D (GSDMD) (**Fig 2B**). Moreover, we observed a prominent decrease in casp-1 expression under NAD⁺ treatment that was constant over the time course of 16 hours. In contrast, BMDMs treated with PBS exhibited excessive casp-1 expression at 4 hours that was found to be absent after 16 hours (**Fig. 2C**), which is consistent with the strong cytotoxicity leading to membrane permeabilization and release of Casp-1 into the supernatant. Noteworthy, Pam3CK4 derived BMDM priming was not affected by NAD⁺ since NF- κ B as well as pro-caspase-1 levels had not been altered (**Fig. 2A and Supp. 2**) underlining the specific inhibition of casp-11. To visualize NAD⁺ mediated blockade of pyroptotic macrophage death, BMDMs were treated with PBS or NAD⁺, primed with Pam3CSK4, then stimulated with LPS and CTB and cell viability and apoptosis were monitored using the IncuCyte® live microscopy system. Hereby, we observed distinct longitudinal kinetics over 100 hours with complete disaggregation of cell integrity in the PBS group contrary to overall preserved cell structure in NAD⁺ treated BMDMs (**Fig. 1D, Supp. 3, Mov. 1**). To rule out, that NAD⁺ impairs LPS internalization into cells, BMDMs were stimulated with CTB and LPS that was coupled to a fluorescent reporter (FITC) and transfection effectivity was assessed by fluorescence microscopy and flow cytometry. Our data indicated no significant difference between the PBS and NAD⁺ treated group (**Fig. 2E**), suggesting that NAD⁺ does not alter LPS internalization. Notably, BMDMs only stimulated with LPS showed no internalization of LPS consistent with previous reports¹².

Casp-4 and 5 have been delineated as the human homolog of casp-11 in mice carrying out the same effector functions including pyroptosis induction and IL-1 β secretion²⁴. As clinical relevance, we therefore tested whether NAD⁺ was also able to block the non-canonical pathway in human macrophages. Hence, human macrophages were differentiated from PBMC and treated with NAD⁺ followed by intracellular LPS transfection (Fugene) and IL-1 β secretion and cytotoxicity were quantified. The results

indicated that NAD⁺ treatment significantly dampened both, IL-1 β secretion and pyroptosis (**Fig. 2F**), underscoring its therapeutic potential. Collectively, our results suggest that NAD⁺ acts directly on macrophages by targeting specifically the non-canonical inflammasome signaling machinery.

NAD⁺ inhibits the non-canonical inflammasome through STAT-1/IFN- β pathway

Although our data emphasized that NAD⁺ blocks the non-canonical inflammasome pathway, the underlying mechanisms remained yet to be determined. Therefore, we performed RNA-sequencing of Pam3CSK4 primed BMDMs that were treated with PBS or NAD⁺ and subsequently stimulated with CTB + LPS O111:B4. Interestingly, when blotting gene expression differences in a Venn diagram, we found strikingly more genes commonly expressed in the NAD⁺ and control group when compared to the PBS treated group (**Fig. 3A**). Gene ontology enrichment analysis revealed a significant downregulation of genes involved in the antiviral response in addition to the cellular response to the type-I-IFN, IFN- β , when comparing NAD⁺ and PBS treated groups (**Fig. 3B**). Type-I-IFN are known to promote the expression of over 2,000 IFN-stimulated genes (ISGs), translated into ISGs-induced proteins which have been shown to act by enhancing pathogen detection and restrict their replication²⁵. Recently, it was reported that type-I-IFNs are required for casp-11 expression contributing to non-canonical inflammasome activation^{26,27}. Consistently, LPS-stimulated macrophages from TRIF-deficient mice displayed impaired casp-11 expression, implying a context-dependent role for type-I-IFN in the regulation of caspase-11 activity²⁶. Indeed, when comparing expression of genes involved in IFN- β signaling through cluster analysis we found a tremendous decrease in a myriad of genes in the NAD⁺ treated group (**Fig. 3C**). Most strikingly, GTPases and guanylate binding proteins involved in the downstream signaling of IFN- β were significantly downregulated while IFN- β -receptor expression remained unaffected (**Fig. 3C and Fig. 3D**). Recently, IFN-inducible GTPases and guanylate binding proteins have been assigned a crucial role for the intracellular recognition of LPS and linked caspase-11 activation^{27,28}. Thus, to test if NAD⁺ mediated non-canonical inflammasome blockade via IFN- β , NAD⁺

or PBS treated BMDMs were primed with Pam3CSK4 and subsequently stimulated with LPS O111:B4 + CTB and 1000 U/ml of recombinant IFN- β . Strikingly, administration of recombinant IFN- β resulted in a complete reversal of NAD⁺ mediated blockade of IL-1 β secretion and pyroptosis (**Fig. 3E**). Moreover, IFN- β administration restored casp-11, NLRP3 and GSDMD expression in the NAD⁺ treated group (**Fig. 3F**). It is well established that STAT-1 phosphorylation constitutes the link between intracellular type-I-IFN signaling and the transcription of ISGs through nuclear translocation^{29,30}. Notably, our RNA-seq data indicated a significant downregulation of STAT-1 (**Fig. 3C**). Moreover, we have previously shown that NAD⁺ administration dampens the expression and activation of transcription factors such as STAT-5¹⁶. To test, whether NAD⁺ blocks IFN- β signaling via STAT-1, BMDMs were subjected to NAD⁺ or PBS followed by non-canonical inflammasome stimulation and recombinant IFN- β . After 16 hours STAT-1 expression and phosphorylation were assessed by Western blotting. Consistent with our previous results, NAD⁺ treatment downregulated expression levels of STAT-1 and phospho-STAT-1. In contrast, addition of recombinant IFN- β treatment to NAD⁺ treated BMDMs restored STAT-1 and phospho-STAT-1 expression that was equivalent to the PBS treated group (**Fig. 3G**). Taken together, our data indicate that NAD⁺ impedes non-canonical inflammasome activation via IFN- β /STAT-1 blockade (**Fig. 4**).

NAD⁺ increases Caspase-11^{-/-} mice resistance to endotoxic shock via systemic IL-10 production

Caspase-11^{-/-} mice have been reported to be resistant towards lethal doses of LPS inducing septic shock¹². However, upon priming with TLR3 instead of a TLR4 ligand, Casp-11^{-/-} mice merely exhibit partial resistance towards LPS-induced shock with a 50–60% survival rate^{12,31}. Our data indicate that NAD⁺ prevents LPS-induced cell death via the non-canonical inflammasome pathway and casp-11 blockade. We thus tested whether NAD⁺ could achieve similar protection against septic shock in WT vs casp-11^{-/-} mice. Casp-11^{-/-} mice were intraperitoneally injected with NAD⁺ and PBS and treated with 6 mg/kg poly(I:C) 7 hours prior LPS administration. Consistent with previous studies the results indicated a modest resistance of casp-11^{-/-} mice (40% survival). In high contrast, both WT and casp-11^{-/-} mice subjected to NAD⁺

exhibited 85%-100% survival, respectively when compared to casp-11^{-/-} mice that were treated with PBS, suggesting the existence of an alternative protective pathway against septic shock that is casp-11 independent. WT mice, treated with 6 mg/kg poly(I:C) followed by LPS (54mg/kg) administration not only survived but fully recovered 7 days later (**Mov. 2**), underscoring the unique and robust therapeutic effect of NAD⁺ in septic shock.

Previous studies have reported inferior outcomes of IL-10^{-/-} mice in septic shock^{32,33} pointing out a 20 fold lower lethal dose of LPS compared to WT mice³³. Moreover, IL-10 itself has been shown to prevent mice from septic shock induced death after a single administration³⁴. We have previously delineated immunosuppressive properties of NAD⁺ via a systemic production of IL-10, a robust immunosuppressive cytokine. In addition, we have described the pivotal role of NAD⁺ protecting towards EAE and allograft rejection via an increased frequency of IL-10 producing CD4⁺ T cells^{15,16}. To test if IL-10 plays an additional protective role in the context of NAD⁺ mediated protection towards LPS-induced death, WT mice treated with NAD⁺ or PBS were subjected to intraperitoneal LPS injection (54mg/kg) and IL-10 expression by macrophages, dendritic cells and T cells was assessed 15 hours after LPS administration. Consistent with our previous studies^{15,16}, we found significantly augmented frequencies of IL-10 producing CD4⁺ and CD8⁺ T cells (**Fig. 5A**). Moreover, we detected a dramatic increase of IL-10 production by macrophages, but not the DC population (**Fig. 5B**). Interestingly, IL-10 has been described to inhibit macrophage function and pro-inflammatory cytokine production in both, human³⁵ and mice³⁶. Moreover, autocrine IL-10 secretion of macrophages was found to decrease pro-IL-1 β concentration by promoting signal transducer activator of transcription-3 (STAT-3) expression³⁷. To investigate the potential autocrine impact of an augmented IL-10 production on macrophage self-regulation, we administered combined IL-10 neutralizing antibody and IL-10 receptor antagonist to BMDMs primed with Pam3CSK4 and stimulated with CTB and LPS O111:B4. The results showed that neutralization of the autocrine IL-10 signaling pathway dampened NAD⁺ mediated decrease of IL-1 β secretion and reversed pyroptotic cell death partially (**Fig. 5C**). To further investigate the relevance of our in vitro

findings, IL-10^{-/-} mice were treated with NAD⁺ or PBS, subjected to LPS (54mg/kg) and survival was monitored. Consistent with previous reports^{32,33}, mice lacking IL-10 exhibited an inferior protection against septic shock when compared to WT animals. More importantly, IL-10^{-/-} mice subjected to NAD⁺ exhibited a compromised survival (**Fig. 5D**) suggesting that systemic production of IL-10 following NAD⁺ administration plays a pivotal role in NAD⁺-mediated protection against septic shock.

Discussion

Previously, we have delineated the protective role of NAD⁺ in the context of *L. m.* infection, a gram-positive bacterium¹⁷. However, it remained unclear whether NAD⁺ conveyed resistance towards *L. m.* by an augmented bacterial clearance or rather through its immunosuppressive effects dampening pathological systemic inflammation. Although the cell membrane of *L. m.* has been shown to bear lipoteichoic acids, which resemble the endotoxin LPS from gram-negative bacteria in both, structure and function, it is widely considered as an intracellular bacterium³⁸. In our current study, we administered a lethal dose of pathogenic *E. coli*, that is well known to promote septic shock, and showed that NAD⁺ also protected towards a lethal dose of this gram-negative bacterium. More importantly, we demonstrate that NAD⁺ conveys protection towards septic shock by specifically inhibiting the non-canonical inflammasome but not via bacterial clearance. Mechanistically, NAD⁺ impedes pro-casp-11 and casp-11 expression in macrophages blocking non-canonical derived GSDMD cleavage and NLRP3 inflammasome activation, thus inhibiting pyroptotic cell death and pro-inflammatory cytokine release. The resistance of NAD⁺ treated WT mice against *E. coli* and LPS induced septic shock reflected the robust inhibitory effect observed in vitro of NAD⁺ on the non-canonical inflammasome signaling machinery.

Until now, the exact mechanism how pro-casp-11 expression and its maturation to casp-11 is regulated remains unclear. Given the low basal expression of both pro-casp-11 and casp-11³⁹, a priming signal is required for initiating the non-canonical inflammasome pathway and macrophage sensing of intracellular LPS⁴⁰. Previous work has demonstrated that transcriptional induction of the pro-casp-11 isoforms p42 and p38 in macrophages requires type I IFN stimulation^{39,41} while IFN- β has been shown to promote transcriptional induction and processing of caspase-11²⁶. In line with these findings, CTB treatment of macrophages primed with Pam3csk4 failed to elicit IL-1 β release compared to LPS primed controls while exogenous administration of IFN- β in turn restored CTB-induced IL-1 β production²⁶ underscoring the transcriptional role of type I IFN. Our RNA-seq results indicated a dampened cellular response towards IFN- β while western blotting revealed a significant downregulation of both, pro-casp-11 and casp-11

suggesting a transcriptional downregulation of both enzymes. Consistently, NAD⁺ decreased STAT-1 expression and phosphorylation, which constitutes the mechanistic link between extracellular type I IFN stimulation and transcriptional effects through translocation of phosphorylated STAT-1 to the nucleus inducing ISGs³⁰. Thus, treatment of stimulated macrophages subjected to NAD⁺ with recombinant IFN-β restored STAT-1 signaling, caspase-11 expression and GSDM cleavage which translated into reconstituted IL-1β production and LDH release. Collectively, NAD⁺ mitigates the intracellular response to IFN-β that leads to non-canonical inflammasome induction by suppressing macrophage derived STAT-1 expression and phosphorylation.

Furthermore, we showed that NAD⁺ treatment improved resistance of casp-11^{-/-} mice towards poly I:C primed septic shock. More importantly, WT mice treated with NAD⁺ exhibited 100% survival while casp-11^{-/-} mice treated with PBS exhibited a modest 40% survival, suggesting that NAD⁺ promotes survival beyond non-canonical inflammasome blockade. Our previous studies have delineated the effects of NAD⁺ on various immune cells such as dendritic cells and CD4⁺ T cells including Th1, Th17, regulatory type 1 (Tr1) and Treg cells communicated exclusively through mast cells (MCs)¹⁵⁻¹⁷. Thereby, NAD⁺ treatment promoted MC derived induction of TR1 cells that resulted into increased systemic levels of IL-10. Latter one was found to play a cardinal feature during bacterial infection as MC^{-/-} mice were more susceptible to *L. m.* infection than WT animals when treated with NAD⁺. Here, we found a direct effect of NAD⁺ on macrophages by specifically inhibiting the non-canonical inflammasome and promoting IL-10 production. Polymorphisms in the IL-10 locus or IL-10R deficiencies have been linked to severe intestinal inflammatory diseases in both, human and mice⁴²⁻⁴⁵. More importantly, mice deficient for IL-10 have been shown to display elevated inflammasome activation and IL-1β production resulting in severe colitis⁴⁶ or Ag-induced arthritis⁴⁷. When inhibiting the autocrine pathway for IL-10 through combined receptor antagonization and IL-10 neutralization, we found a pronounced increase of IL-1β production of NAD⁺ treated macrophages stimulated with CTB and LPS (Fig. 4D). This is consistent with previous reports showing that autocrine IL-10 signaling interferes with the transcription of pro-IL-1β³⁷. LDH release in turn, was only restored partly possibly due to missing effects of second party leucocytes

secreting IL-10 in vivo such as Tr1 cells which have been shown to inhibit the transcription of IL-1 β and inflammasome mediated activation of caspase-1⁴⁸.

More recently, casp-8, that plays a central role in apoptosis, has been reported as an important mediator of endotoxemia resistance and LPS-driven systemic inflammation. Since our RNA-sequencing results revealed a dramatically attenuated cellular response towards type-I-IFN with downregulation of various interferon regulatory factors, that have been reported as major regulators of casp-8^{49,50} it is possible that NAD⁺ may exert protection against septic shock by altering caspase-8 expression as well. Although, we have previously reported the protective effect of NAD⁺ against apoptosis of activated CD4⁺ T cells¹⁵, it remains yet to be determined how NAD⁺ impacts executioner proteins of other cell death processes such as apoptosis and necroptosis.

Notably, both casp-8 and casp-11 have been found dispensable in the hematopoietic compartment that produces the pro-inflammatory cytokines necessary to initiate shock⁵¹. Thus, NAD⁺ treatment may improve resistance of casp-11^{-/-} mice to septic shock by also dampening the initiating pro-inflammatory cytokine cascade via its systemic IL-10 cytokine production.

NAD⁺ as a natural component exhibits a powerful and efficient protection towards septic shock.

Moreover, it promotes allograft survival, protects and reverses severe autoimmune diseases and displays beneficial effects in the context of primary immunodeficiency underscoring its potential therapeutic capacity. Importantly, while inhibiting macrophage derived inflammasome function, NAD⁺ does not interfere with NF- κ B signaling which has been shown to promote various inflammatory and autoimmune diseases when dysregulated⁵².

Taken together, we dissected the dichotomous capacity of NAD⁺ to dampen auto- and allo-immunity while concomitantly protecting towards severe bacterial infection, outlining its unique effects in the context of septic shock.

Material and methods

Animals

Young (8-12 weeks) C57BL/6, B6.129P2-IL10tm1Cgn/J and B6.129S4(D2)-Casp4tm1Yuan/J mice were purchased from Charles River Laboratory, Wilmington, MA. All animals were kept in our animal facility in accordance to guidelines established by the Animal Care Committee at Harvard Medical School. Permissions of animal experiments were granted by the Harvard Area Standing Committee on Animals.

Cell culture

8-12-week-old C57BL/6 mice were euthanized by cervical dislocation, sprayed with alcohol and skin was removed to expose femurs. The femur was flushed with ice cold PBS and the obtained bone marrow was filtered through 70um Nylon cell strainer. After washing with PBS, red blood cell lysis was performed using ammonium-chloride-potassium-solution (Fisher scientific) and the reaction was blocked with complete Dulbecco's modified eagle medium (DMEM) (Fisher Scientific) supplemented with 10% endotoxin-free bovine serum and PS. To minimize fibroblast contamination cells were cultured in complete DMEM at 37°C, 5% CO₂ and non-adherent cells were collected after 30 minutes.

Bone marrow cells were then differentiated into macrophages in DMEM supplemented with 10% endotoxin-free bovine serum, PS and 40ng/ml murine GM-CSF (Abcam) for 8 days. Medium was changed every 2 days to remove non-adherent cells. After 8 days of culture the medium was replaced by 40ng/ml GM-CSF containing 100μmol NAD⁺ culture medium. For 2 following days NAD⁺ was added daily until stimulation.

For in vitro experiments BMDMs were cultured overnight in a 24 well plate at 1x10⁶ cells/ml and afterwards primed with 1μg/ml Pam3CSK4 or 1μg/ml LPS O111:B4 (Sigma) for 5-6 hours. Primed BMDMs were then stimulated for 16 hours with 5 mmol ATP or 2μg/ml LPS O111:B4 and 20μg/ml CTB (Sigma) where indicated. IL-1β expression was analyzed in the supernatant by ELISA (Invitrogen) and

cell death was measured by assessing LDH-release using a CytoTox 96 Non-radioactive Cytotoxic Assay (Promega).

For experiments on human cells, PBMC were isolated by performing a density centrifugation in SepMate tubes (Stem cell) using lymphoprep (Stem cell) density gradient medium. PBMC were then plated in DMEM culture medium supplemented with standard antibiotics 10% FCS and human GM-CSF (peprotech) at a density of 1×10^6 cells/ml. The medium was changed every 2-3 days until the cells reached a fully confluence. To induce non-canonical inflammasome activation in human macrophages, cells were then primed with $1 \mu\text{g/ml}$ Pam3CSK4 for 5-6 hours. Subsequently the medium was replaced, and cells were treated with $3 \mu\text{g/ml}$ LPS O111:B4 and 0.25% (vol/vol) Fugene HD Plus (Promega) to cause transfection. Finally, plates were centrifuged at $805 \times g$ for 2 minutes and subsequently cultured for 20 hours at 37°C , 5% CO_2 .

Western blot

For Western blot analysis, proteins were extracted using RIPA buffer and the concentrations determined using Pierce™ BCA Protein Assay Kit. Subsequently, proteins were resolved in SDS-PAGE, transferred to $0.45 \mu\text{m}$ nitrocellulose membranes (BioRad), blocked with 5% non-fat dry milk in PBS with 0.1% Tween 20, and processed for immunodetection. The following primary antibodies were used according to manufacturer's instructions: Pro-Caspase-1 (#ab179515, Abcam), Caspase-1 (#14-9832-82, eBioscience) IL-1 β (AF-401-NA, RD Systems), NLRP3 (#768319, RD Systems), Caspase-11 (#mab8648, RD Systems), Gasdermin D (ab209845, Abcam), P-STAT-1 (#9167S, Cell Signaling), STAT-1 (#9172S, Cell Signaling), NF- κB -p65 (#49445S, Cell Signaling), NF- κB -p52 (#4882S, Cell Signaling), β -Actin (ab3280, Abcam). Antibody detection was performed with HRP-coupled goat secondary anti-mouse or anti-rabbit antibodies (Immunoresearch), followed by ECL reaction (Perkin Elmer) and exposure to Fuji X-ray films. Finally, films were scanned, and signals quantified using the web-based image processing software ImageJ (NIH).

For intracellular detection of LPS, primed BMDMs were stimulated with 20 μ g/ml CTB and FITC-conjugated LPS O111:B4 for 16 hours, washed twice with PBS, fixed in 4% PFA containing PBS for 10 minutes and subsequently analyzed using a confocal microscope and flow cytometry.

Caspase-1 assay

To determine the expression of Caspase-1, primed BMDMs were stimulated with 20 μ g/ml CTB and 2 μ g/ml LPS O111:B4 for 4 and 16 hours respectively, washed twice with PBS, stained using a caspase-1 active staining kit (Abcam) according to the manufacturer's protocol and analyzed using a confocal microscope.

Endotoxic shock model

8-12-week-old C57BL/6 mice were treated with 40mg NAD⁺ for 2 following days before intraperitoneal injection of 54 mg/kg LPS O111:B4 or LPS O55:B5. Where indicated mice were administered 6mg/kg poly(I:C) 6 hours prior to LPS administration. Consequently, survival and body temperature were monitored every 2-4 hours for up to 100 hours. To assess the amount of systemic IL-1 β and IL-18 by Elisa (both Invitrogen), mice were euthanized by decapitation 10 hours and 15 hours after LPS injection serum was isolated from collected blood.

Flow cytometric analysis

To analyze splenic lymphocytes for the intracellular expression IL-10 mice were challenged with 54 mg/kg LPS O111:B4 for 10 hours and euthanized by cervical dislocation subsequently. Spleens were harvested in a sterile environment and single cell suspensions were obtained using a 70 μ m Nylon cell strainer.

Then, 1 $\times$ 10⁶ splenocytes per animal per condition were cultured in RPMI 1640 (#10-040-CV, Corning) supplemented with 10% BenchMark Fetal Bovine Serum (#100-106, Gemini), 1% penicillin/streptomycin (#30-002-CI, Corning), 2 mM L-glutamine (#25-005-CI, Corning), 20 ng/mL phorbol 12-myristate 13-

acetate (PMA, #P8139-1MG, Sigma-Aldrich), 1 µg/mL ionomycin (#I9657-1MG, Sigma-Aldrich), and 0.67 µL/mL BD GolgiStop (#554724, BD Biosciences) for 4 hours at 37°C and 5% CO₂ in 1 mL-volumes in a 12-well plate. After 4 hours, the cells were collected from each 12-well plate well and prepared for flow cytometry by staining the surface epitopes in flow staining buffer consisting of 1× DPBS supplemented with 1.0% (w/v) bovine serum albumin (#A2153, Sigma-Aldrich) and 0.020% sodium azide (#S8032, Sigma-Aldrich) for 25 min at 4°C. Then, the cells were fixed and permeabilized with the eBioscience Foxp3 Fixation/Permeabilization concentrate and diluent cocktail (#00-5523-00, Invitrogen) for 30 min at 4°C. Finally, the intracellular cytokine target was stained in 1× permeabilization buffer diluted from 10× eBioscience Foxp3 Permeabilization Buffer (#00-5523-00, Invitrogen) with deionized water. Finally, the stained samples were analyzed on a FACS Canto II (BD Biosciences, San Jose, CA, United States) flow cytometer, and the resultant flow cytometry standard (FCS) files were analyzed with FlowJo version 10 (Flowjo LLC, Ashland, OR, United States).

Bacterial infection model

Frozen stock suspensions of *Escherichia coli* (Migul) (ATCC, 700928) were obtained from ATCC and cultured in 5ml Luria-Bertani medium at 37°C. Bacterial concentration was determined by plating 100ul, 10-fold serial diluted bacterial samples and counting the colony-forming units (CFU) after overnight incubation at 37°C. One day prior to injection 1ml of culture was reinoculated into 5ml of medium and incubated for 3-5 hours using a 37°C shaker at 250rpm agitation. Bacterial cultures were then centrifuged for 10 minutes at 3000rpm and washed twice with PBS. Mice were previously treated with NAD⁺ for 2 serial days and subsequently infected with *E. coli* by injecting 1ml of 1x10⁹ CFU/ml bacterial suspension intraperitoneally. The survival was monitored. In another set of experiments mice were sacrificed 5h hours after infection and kidneys and liver were harvested. The collected tissues were homogenized in 1ml of sterile PBS and 10-fold serial dilutions plated overnight at 37°C on LB agar plates to determine bacterial load per gram.

Author contributions

JI set up in vitro experiments performing cell culture stimulation, ELISA, cell death assay, live microscopy, cell transfection, RNA sequencing and in vivo experiments. JI analyzed data and wrote the manuscript. RE performed Western blots and NF- κ B microscopy. SE performed flow cytometric analysis. A.V, H.R. and N.Y. analyzed data and edited the manuscript. AE designed experiments, interpreted the data, supervised the work and wrote the manuscript.

Acknowledgements

J.I. was supported by the Biomedical Education Program (BMEP) of the German Academic Exchange Service. Y.N. was supported by the Chinese Scholarship Council (201606370196) and Central South University. H.R.C.B. was supported by the Swiss Society of Cardiac Surgery. A.V. was supported by awards from the National Institute of Mental Health (R01MH110438) and National Institute of Neurological Disorders and Stroke (R01NS100808).

Competing interests

The authors declare no conflict of interest

References

- 1 Kaukonen, K. M., Bailey, M., Pilcher, D., Cooper, D. J. & Bellomo, R. Systemic inflammatory response syndrome criteria in defining severe sepsis. *N. Engl. J. Med.* **372**, 1629-1638, doi:10.1056/NEJMoa1415236 (2015).
- 2 Ramachandran, G. Gram-positive and gram-negative bacterial toxins in sepsis: a brief review. *Virulence* **5**, 213-218, doi:10.4161/viru.27024 (2014).
- 3 Kumar, V. Inflammasomes: Pandora's box for sepsis. *Journal of inflammation research* **11**, 477-502, doi:10.2147/jir.S178084 (2018).
- 4 Lopez-Castejon, G. & Brough, D. Understanding the mechanism of IL-1 β secretion. *Cytokine Growth Factor Rev.* **22**, 189-195, doi:10.1016/j.cytogfr.2011.10.001 (2011).
- 5 Yi, Y. S. Caspase-11 non-canonical inflammasome: a critical sensor of intracellular lipopolysaccharide in macrophage-mediated inflammatory responses. *Immunology* **152**, 207-217, doi:10.1111/imm.12787 (2017).
- 6 Ding, J. & Shao, F. SnapShot: The Noncanonical Inflammasome. *Cell* **168**, 544-544.e541, doi:10.1016/j.cell.2017.01.008 (2017).
- 7 Shi, J. *et al.* Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature* **526**, 660-665, doi:10.1038/nature15514 (2015).
- 8 Man, S. M. *et al.* Differential roles of caspase-1 and caspase-11 in infection and inflammation. *Sci. Rep.* **7**, 45126, doi:10.1038/srep45126 (2017).
- 9 Kayagaki, N. *et al.* Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature* **526**, 666-671, doi:10.1038/nature15541 (2015).
- 10 Kayagaki, N. *et al.* Non-canonical inflammasome activation targets caspase-11. *Nature* **479**, 117-121, doi:10.1038/nature10558 (2011).
- 11 Yang, D., He, Y., Munoz-Planillo, R., Liu, Q. & Nunez, G. Caspase-11 Requires the Pannexin-1 Channel and the Purinergic P2X7 Pore to Mediate Pyroptosis and Endotoxic Shock. *Immunity* **43**, 923-932, doi:10.1016/j.immuni.2015.10.009 (2015).
- 12 Kayagaki, N. *et al.* Noncanonical inflammasome activation by intracellular LPS independent of TLR4. *Science* **341**, 1246-1249, doi:10.1126/science.1240248 (2013).
- 13 Angus, D. C. & van der Poll, T. Severe sepsis and septic shock. *N. Engl. J. Med.* **369**, 840-851, doi:10.1056/NEJMra1208623 (2013).
- 14 Fraser, C. C. Exploring the positive and negative consequences of NF-kappaB inhibition for the treatment of human disease. *Cell cycle (Georgetown, Tex.)* **5**, 1160-1163, doi:10.4161/cc.5.11.2773 (2006).
- 15 Tullius, S. G. *et al.* NAD⁺ protects against EAE by regulating CD4⁺ T-cell differentiation. *Nat Commun* **5**, 5101, doi:10.1038/ncomms6101 (2014).
- 16 Elkhail, A. *et al.* NAD(+) regulates Treg cell fate and promotes allograft survival via a systemic IL-10 production that is CD4(+) CD25(+) Foxp3(+) T cells independent. *Sci. Rep.* **6**, 22325, doi:10.1038/srep22325 (2016).
- 17 Rodriguez Cetina Biefer, H. *et al.* Mast cells regulate CD4(+) T-cell differentiation in the absence of antigen presentation. *J Allergy Clin Immunol* **142**, 1894-1908.e1897, doi:10.1016/j.jaci.2018.01.038 (2018).
- 18 Mellata, M., Mitchell, N. M., Schodel, F., Curtiss, R. R. & Pier, G. B. Novel vaccine antigen combinations elicit protective immune responses against Escherichia coli sepsis. *Vaccine* **34**, 656-662, doi:10.1016/j.vaccine.2015.12.014 (2016).
- 19 Dogan, M. D., Ataoglu, H. & Akarsu, E. S. Effects of different serotypes of Escherichia coli lipopolysaccharides on body temperature in rats. *Life Sci* **67**, 2319-2329, doi:10.1016/s0024-3205(00)00821-3 (2000).
- 20 Bullock, B. & Benham, M. D. in *StatPearls* (StatPearls PublishingStatPearls Publishing LLC., 2019).

- 21 Rossaint, J. & Zarbock, A. Pathogenesis of Multiple Organ Failure in Sepsis. *Crit. Rev. Immunol.* **35**, 277-291 (2015).
- 22 Evans, T. J. The role of macrophages in septic shock. *Immunobiology* **195**, 655-659, doi:10.1016/s0171-2985(96)80029-5 (1996).
- 23 Chan, F. K., Moriwaki, K. & De Rosa, M. J. Detection of necrosis by release of lactate dehydrogenase activity. *Methods Mol. Biol.* **979**, 65-70, doi:10.1007/978-1-62703-290-2_7 (2013).
- 24 Shi, J. *et al.* Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature* **514**, 187-192, doi:10.1038/nature13683 (2014).
- 25 Schneider, W. M., Chevillotte, M. D. & Rice, C. M. Interferon-stimulated genes: a complex web of host defenses. *Annu. Rev. Immunol.* **32**, 513-545, doi:10.1146/annurev-immunol-032713-120231 (2014).
- 26 Rathinam, V. A. *et al.* TRIF licenses caspase-11-dependent NLRP3 inflammasome activation by gram-negative bacteria. *Cell* **150**, 606-619, doi:10.1016/j.cell.2012.07.007 (2012).
- 27 Tang, Y. *et al.* TRIF signaling is required for caspase-11-dependent immune responses and lethality in sepsis. *Mol. Med.* **24**, 66, doi:10.1186/s10020-018-0065-y (2018).
- 28 Pilla, D. M. *et al.* Guanylate binding proteins promote caspase-11-dependent pyroptosis in response to cytoplasmic LPS. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 6046-6051, doi:10.1073/pnas.1321700111 (2014).
- 29 Stark, G. R. & Darnell, J. E., Jr. The JAK-STAT pathway at twenty. *Immunity* **36**, 503-514, doi:10.1016/j.immuni.2012.03.013 (2012).
- 30 Ivashkiv, L. B. & Donlin, L. T. Regulation of type I interferon responses. *Nat. Rev. Immunol.* **14**, 36-49, doi:10.1038/nri3581 (2014).
- 31 Hagar, J. A. *et al.* Lipopolysaccharide Potentiates Insulin-Driven Hypoglycemic Shock. *J Immunol* **199**, 3634-3643, doi:10.4049/jimmunol.1700820 (2017).
- 32 Latifi, S. Q., O'Riordan, M. A. & Levine, A. D. Interleukin-10 controls the onset of irreversible septic shock. *Infect. Immun.* **70**, 4441-4446, doi:10.1128/iai.70.8.4441-4446.2002 (2002).
- 33 Berg, D. J. *et al.* Interleukin-10 is a central regulator of the response to LPS in murine models of endotoxic shock and the Shwartzman reaction but not endotoxin tolerance. *J. Clin. Invest.* **96**, 2339-2347, doi:10.1172/jci118290 (1995).
- 34 Howard, M., Muchamuel, T., Andrade, S. & Menon, S. Interleukin 10 protects mice from lethal endotoxemia. *J. Exp. Med.* **177**, 1205-1208, doi:10.1084/jem.177.4.1205 (1993).
- 35 de Waal Malefyt, R., Abrams, J., Bennett, B., Figdor, C. G. & de Vries, J. E. Interleukin 10(IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J. Exp. Med.* **174**, 1209-1220, doi:10.1084/jem.174.5.1209 (1991).
- 36 Fiorentino, D. F., Zlotnik, A., Mosmann, T. R., Howard, M. & O'Garra, A. IL-10 inhibits cytokine production by activated macrophages. *J. Immunol.* **147**, 3815-3822 (1991).
- 37 Sun, Y. *et al.* Interleukin-10 inhibits interleukin-1 β production and inflammasome activation of microglia in epileptic seizures. *J Neuroinflammation* **16**, 66, doi:10.1186/s12974-019-1452-1 (2019).
- 38 Farber, J. M. & Peterkin, P. I. *Listeria monocytogenes*, a food-borne pathogen. *Microbiol. Rev.* **55**, 476-511 (1991).
- 39 Schauvliege, R., Vanrobaeys, J., Schotte, P. & Beyaert, R. Caspase-11 gene expression in response to lipopolysaccharide and interferon-gamma requires nuclear factor-kappa B and signal transducer and activator of transcription (STAT) 1. *J. Biol. Chem.* **277**, 41624-41630, doi:10.1074/jbc.M207852200 (2002).
- 40 Yang, J., Zhao, Y. & Shao, F. Non-canonical activation of inflammatory caspases by cytosolic LPS in innate immunity. *Curr. Opin. Immunol.* **32**, 78-83, doi:10.1016/j.coi.2015.01.007 (2015).
- 41 Yen, J. H. & Ganea, D. Interferon beta induces mature dendritic cell apoptosis through caspase-11/caspase-3 activation. *Blood* **114**, 1344-1354, doi:10.1182/blood-2008-12-196592 (2009).

- 42 Franke, A. *et al.* Sequence variants in IL10, ARPC2 and multiple other loci contribute to ulcerative colitis susceptibility. *Nat Genet* **40**, 1319-1323, doi:10.1038/ng.221 (2008).
- 43 Franke, A. *et al.* Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nat Genet* **42**, 1118-1125, doi:10.1038/ng.717 (2010).
- 44 Kuhn, R., Lohler, J., Rennick, D., Rajewsky, K. & Muller, W. Interleukin-10-deficient mice develop chronic enterocolitis. *Cell* **75**, 263-274, doi:10.1016/0092-8674(93)80068-p (1993).
- 45 Begue, B. *et al.* Defective IL10 signaling defining a subgroup of patients with inflammatory bowel disease. *Am J Gastroenterol* **106**, 1544-1555, doi:10.1038/ajg.2011.112 (2011).
- 46 Zhang, J., Fu, S., Sun, S., Li, Z. & Guo, B. Inflammasome activation has an important role in the development of spontaneous colitis. *Mucosal Immunol* **7**, 1139-1150, doi:10.1038/mi.2014.1 (2014).
- 47 Greenhill, C. J. *et al.* Interleukin-10 regulates the inflammasome-driven augmentation of inflammatory arthritis and joint destruction. *Arthritis Res Ther* **16**, 419, doi:10.1186/s13075-014-0419-y (2014).
- 48 Yao, Y. *et al.* Tr1 Cells, but Not Foxp3+ Regulatory T Cells, Suppress NLRP3 Inflammasome Activation via an IL-10-Dependent Mechanism. *J Immunol* **195**, 488-497, doi:10.4049/jimmunol.1403225 (2015).
- 49 Apelbaum, A., Yarden, G., Warszawski, S., Harari, D. & Schreiber, G. Type I interferons induce apoptosis by balancing cFLIP and caspase-8 independent of death ligands. *Mol Cell Biol* **33**, 800-814, doi:10.1128/mcb.01430-12 (2013).
- 50 Newton, K. *et al.* Cleavage of RIPK1 by caspase-8 is crucial for limiting apoptosis and necroptosis. *Nature*, doi:10.1038/s41586-019-1548-x (2019).
- 51 Mandal, P. *et al.* Caspase-8 Collaborates with Caspase-11 to Drive Tissue Damage and Execution of Endotoxic Shock. *Immunity* **49**, 42-55.e46, doi:10.1016/j.immuni.2018.06.011 (2018).
- 52 Liu, T., Zhang, L., Joo, D. & Sun, S. C. NF-kappaB signaling in inflammation. *Signal transduction and targeted therapy* **2**, doi:10.1038/sigtrans.2017.23 (2017).

Figure Legends

Figure 1. NAD⁺ protects mice from lethal bacterial infection and endotoxic shock by dampening systemic inflammation

(A) C57BL/6 mice were treated with PBS or NAD⁺ for 2 days prior to administration of a lethal dose of either pathogenic *E. coli* or LPS (O55:B5 /O111:B4) by intraperitoneal injection. (B) Kidneys and Livers were removed after 5 hours of infection, homogenized, plated on LB agar plates and bacterial load was determined by counting CFU (C) Survival was monitored over 48 hours after bacterial infection and (D) LPS injection of both serotypes. In addition, body temperature was monitored in the kinetics of up to 100 hours (E) Lungs, Kidneys and Livers were removed after 15 hours and subsequently IHC was performed staining for H&E (F) Systemic levels (serum) of IL-1β and IL-18 were assessed by ELISA.

Column plots display mean with standard deviation. Statistical significance was determined by using Student's T-test or One-Way-ANOVA while survival data were compared using log-rank Mantel-Cox test. Asterisks indicate p-values * = p<0.05, ** = p<0.01 and *** = p<0.001, only significant values are shown.

Figure 2. *NAD⁺ specifically inhibits the non-canonical inflammasome by targeting caspase 11*

Bone marrow was isolated from C57BL/6 mice and BMDMs were differentiated in vitro. Subsequently, BMDMs were cultured in the presence of 100 μ mol NAD⁺ or PBS for 2 following days. BMDMs were then primed with either 1 μ g/ml Pam3CSK4 or 1 μ g/ml LPS O111:B4 for 5-6 hours. Next primed BMDMs were stimulated for 16 hours with 5 mmol ATP or 2 μ g/ml LPS O111:B4 and 20 μ g/ml CTB. (A) Pro-casp-1, Pro-casp-11, Casp-11, NLRP3, Casp-1, IL-1 β and GSDMD expression were determined using Western blot and (B) IL-1 β secretion and LDH release were assessed in the supernatant. (C) Time dependent Caspase-1 expression was determined via active staining and assessed using a confocal microscope. (D) Cell viability and apoptosis were monitored using the IncuCyte® live microscopy system (E) LPS transfection was visualized by using FITC-coupled LPS O111:B4 and quantified by confocal microscopy and flow cytometry. (F) For human experiments macrophages were differentiated from PBMC, primed with 1 μ g/ml Pam3CSK4 for 5-6 hours and subsequently transfected with LPS O111:B4 and 0.25% (vol/vol) Eugene HD Plus.

Column plots display mean with standard deviation. Statistical significance was determined by using

Student's T-test or One-Way-ANOVA. Asterisks indicate p-values * = $p < 0.05$, ** = $p < 0.01$ and *** = $p < 0.001$, only significant values are shown.

Figure 3. NAD⁺ mediated inhibition of the non-canonical inflammasome is based on an impaired response to IFN- β

Differentiated BMDMs were cultured in the presence of 100 μ mol NAD⁺ or PBS for 2 following days. BMDMs were then primed with 1 μ g/ml Pam3CSK4, subsequently stimulated with 2 μ g/ml LPS O111:B4 and 20 μ g/ml CTB and RNA sequencing was performed. Unstimulated BMDMs served as controls. (A) Venn diagram plotting common gene expression between all 3 groups (B) Gene ontology enrichment analysis displaying the highest significant pathways differing when comparing NAD⁺ and PBS treated BMDMs (C) Expression cluster analysis of genes involved in IFN- β signaling through cluster analysis

depicted in a heat map (D) Vulcano plot displaying the most significant genes up or downregulated comparing NAD⁺ and PBS treated BMDMs (E) Stimulated BMDMs were additionally treated with recombinant INF- β , and IL-1 β and LDH release were measured. (F) Moreover, pro-Casp-1, Casp-11, NLRP3, GSDMD, (G) STAT-1 and Phospho-STAT-1 expression were assessed by Western blot.

Column plots display mean with standard deviation. Statistical significance was determined by using Student's T-test or One-Way-ANOVA. Asterisks indicate p-values * = $p < 0.05$, ** = $p < 0.01$ and *** = $p < 0.001$, only significant values are shown.

Figure 4. Inhibitory effects of NAD⁺ on IFN- β downstream signaling and inflammasome activation

NAD⁺ inhibits STAT-1 expression and phosphorylation, thus compromising the intracellular response to IFN- β . Subsequently, stimulation of the IFNAR receptor by IFN- β leads to a decreased transcription of pro-caspase-11 as well as ISGs (IFN-inducible GTPases and GBPs). Due to diminished caspase-11 levels, non-canonical inflammasome activation through intracellular, gram negative bacteria opsonization by GBPs is significantly inhibited.

Figure 5. *IL-10 constitutes an additional pathway mediating the protective capacities of NAD⁺ in the context of septic shock*

(A) Caspase-11^{-/-} mice were treated with NAD⁺ or PBS for 2 following days. Subsequently mice were subjected to 4mg/kg Poly(I:C) 6 hours prior to LPS injection and survival was monitored. C57BL/6 mice treated with either NAD⁺ or PBS were injected with LPS and after 10 hours, Splenic frequencies of IL-10 producing (B) Macrophages and Dendritic cells (C) and CD4⁺ and CD8⁺ T cells were assessed by flow cytometry. (D) BMDMs treated with NAD⁺ or PBS were stimulated with LPS and CTB in the presence of 1μg/ml IL-10 neutralizing antibodies and 1μg/ml IL-10 receptor antagonists. Subsequently IL-1β and LDH release were assessed. (E) IL-10^{-/-} mice treated with NAD⁺ or PBS were challenged with 54mg/kg LPS O111:B4 and survival was monitored

Column plots display mean with standard deviation. Statistical significance was determined by using Student's T-test or One-Way-ANOVA while survival data were compared using log-rank Mantel-Cox test. Asterisks indicate p-values * = p<0.05, ** = p<0.01 and *** = p<0.001, only significant values are shown.

Supplementary Figure 1. NAD⁺ preserves ileal villi structure and reduces splenic hemorrhage during LPS induced septic shock

C57BL/6 mice were treated with PBS or NAD⁺ for 2 days prior to administration of a lethal dose of LPS (O55:B5 /O111:B4) by intraperitoneal injection. Ileum and Spleen were removed after 15 hours and subsequently IHC was performed staining for H&E.

Supplementary Figure 2. NAD⁺ does not alter BMDM derived NF-κB expression or phosphorylation

Differentiated BMDMs were cultured in the presence of 100μmol NAD⁺ or PBS for 2 following days. BMDMs were then primed with 1μg/ml Pam3CSK4 and subsequently stimulated with 2μg/ml LPS O111:B4 and 20μg/ml CTB. Unstimulated BMDMs served as controls. (A) P52 and p65 expression was determined using Western blot. (B) stimulated BMDMs were stained with p52, p65 and phospho-p65 and expression levels assessed using confocal microscopy.

Supplementary Figure 3. Unstimulated BMDM cell viability and apoptosis

Differentiated BMDMs were cultured in the presence of 100 μ mol NAD⁺ or PBS for 2 following days. BMDMs were then primed with 1 μ g/ml Pam3CSK4, subsequently stimulated with 2 μ g/ml LPS O111:B4 and 20 μ g/ml CTB and cell viability and apoptosis were monitored for 100 hours using the IncuCyte® live microscopy system.